

MATH 322 – Graph Theory

Fall Term 2021

Notes for Lecture 10

Thursday, October 7

Important results from last time

Proposition 1

Let G be a connected graph of order ≥ 3 .

An edge e in G is NOT a bridge of G if and only if e belongs to some cycle contained in G .

Remark. Equivalently we could state the proposition as follows: "Let H be a connected graph of order ≥ 2 . Then e is a bridge of H if and only if e does NOT belong to any cycle contained in H ."

Important results from last time

Proposition 1

Let G be a connected graph of order ≥ 3 .

An edge e in G is NOT a bridge of G if and only if e belongs to some cycle contained in G .

Remark. Equivalently we could state the proposition as follows: "Let H be a connected graph of order ≥ 2 . Then e is a bridge of H if and only if e does NOT belong to any cycle contained in H ."

Proposition 2

Let G be a connected graph of order ≥ 2 , and let v be a vertex of G such that $\deg(v) = 1$.

Then v cannot be a cutvertex.

Important results from last time (cont.)

Proposition 3

Let T be a tree on at least two vertices. Then, for every two different vertices u, v of T , there is a **unique** path in T starting at u and ending at v .

Important results from last time (cont.)

Proposition 3

Let T be a tree on at least two vertices. Then, **for every two different vertices u, v of T** , there is a **unique** path in T starting at u and ending at v .

Proposition 4

Let T be a tree **of order ≥ 3** . Then T has vertices of degree ≥ 2 , and every such vertex is a cutvertex of T .

Proof of Proposition 4 (1st half from last time)

Consider a tree T of order ≥ 3 . First we need to check that T has at least one vertex of degree ≥ 2 .

Write n for the order of T , that is, for the number of vertices in T . By Theorem 1a from Lecture 7, we know that $e(T) = n - 1$.

Proof of Proposition 4 (1st half from last time)

Consider a tree T of order ≥ 3 . First we need to check that T has at least one vertex of degree ≥ 2 .

Write n for the order of T , that is, for the number of vertices in T . By Theorem 1a from Lecture 7, we know that $e(T) = n - 1$.

Assume towards a contradiction that T contained only leaves, that is, vertices of degree 1.

Proof of Proposition 4 (1st half from last time)

Consider a tree T of order ≥ 3 . First we need to check that T has at least one vertex of degree ≥ 2 .

Write n for the order of T , that is, for the number of vertices in T . By Theorem 1a from Lecture 7, we know that $e(T) = n - 1$.

Assume towards a contradiction that T contained only leaves, that is, vertices of degree 1.

Then we would have

$$\sum_{v \in V(T)} \deg(v) = \sum_{v \in V(T)} 1 = |T| = n < 2(n - 1) = 2e(T),$$

where the (strict) inequality holds because $n \geq 3$. This contradicts the Handshaking Lemma.

Proof of Proposition 4 (1st half from last time)

Consider a tree T of order ≥ 3 . First we need to check that T has at least one vertex of degree ≥ 2 .

Write n for the order of T , that is, for the number of vertices in T . By Theorem 1a from Lecture 7, we know that $e(T) = n - 1$.

Assume towards a contradiction that T contained only leaves, that is, vertices of degree 1.

Then we would have

$$\sum_{v \in V(T)} \deg(v) = \sum_{v \in V(T)} 1 = |T| = n < 2(n - 1) = 2e(T),$$

where the (strict) inequality holds because $n \geq 3$. This contradicts the Handshaking Lemma.

Thus T must contain at least one vertex of degree ≥ 2 .

Proof of Proposition 4 (cont.)

Consider now a vertex v_0 of T which has degree ≥ 2 . We will show that $T - v_0$ is a disconnected subgraph.

Proof of Proposition 4 (cont.)

Consider now a vertex v_0 of T which has degree ≥ 2 . We will show that $T - v_0$ is a disconnected subgraph.

In fact, we will show that **the number of connected components of $T - v_0$ is equal to the number of neighbours of v_0 in T** , and thus that $T - v_0$ has at least 2 connected components.

Proof of Proposition 4 (cont.)

Consider now a vertex v_0 of T which has degree ≥ 2 . We will show that $T - v_0$ is a disconnected subgraph.

In fact, we will show that **the number of connected components of $T - v_0$ is equal to the number of neighbours of v_0 in T** , and thus that $T - v_0$ has at least 2 connected components.

Indeed, let u_1, w_1 be two different neighbours of v_0 in T . Then the path $P_0 : u_1 v_0 w_1$ is a path of length 2 contained in T which starts at u_1 and ends at w_1 .

Proof of Proposition 4 (cont.)

Consider now a vertex v_0 of T which has degree ≥ 2 . We will show that $T - v_0$ is a disconnected subgraph.

In fact, we will show that **the number of connected components of $T - v_0$ is equal to the number of neighbours of v_0 in T** , and thus that $T - v_0$ has at least 2 connected components.

Indeed, let u_1, w_1 be two different neighbours of v_0 in T . Then the path $P_0 : u_1 v_0 w_1$ is a path of length 2 contained in T which starts at u_1 and ends at w_1 .

Recall now Proposition 3 that we proved last time (and that we just restated): it implies that there is NO OTHER path in T which starts at u_1 and ends at w_1 .

In other words, we cannot find a path P_1 in $T - v_0$ which starts at u_1 and ends at w_1 (because such a path would also be a path in T , and it would be different from the path P_0).

Proof of Proposition 4 (cont.)

Consider now a vertex v_0 of T which has degree ≥ 2 . We will show that $T - v_0$ is a disconnected subgraph.

In fact, we will show that **the number of connected components of $T - v_0$ is equal to the number of neighbours of v_0 in T** , and thus that $T - v_0$ has at least 2 connected components.




Indeed, let u_1, w_1 be two different neighbours of v_0 in T . Then the path $P_0 : u_1 v_0 w_1$ is a path of length 2 contained in T which starts at u_1 and ends at w_1 .

Recall now Proposition 3 that we proved last time (and that we just restated): it implies that there is NO OTHER path in T which starts at u_1 and ends at w_1 .

In other words, we cannot find a path P_1 in $T - v_0$ which starts at u_1 and ends at w_1 (because such a path would also be a path in T , and it would be different from the path P_0).

We conclude that each of the neighbours of v_0 in T will end up in a different connected component of $T - v_0$.

Checking these on examples



Question. In each of the graphs above, which of the vertices are cutvertices, and which are not?

Reminder: Whitney's Theorem tested on 'extreme' examples

- In the case of the complete graph K_n ($n \geq 2$) we have seen that:
 - $\kappa(K_n) = n - 1$ by definition,
 - $\lambda(K_n) = n - 1$ by the analysis we did last time,
 - and clearly $\delta(K_n) = n - 1$.
- In the case of a tree T_n on n vertices (where $n \geq 3$, because a tree on 2 vertices would just be K_2 and we have already handled this example), we have so far seen:

Reminder: Whitney's Theorem tested on 'extreme' examples

- In the case of the complete graph K_n ($n \geq 2$) we have seen that:
 - $\kappa(K_n) = n - 1$ by definition,
 - $\lambda(K_n) = n - 1$ by the analysis we did last time,
 - and clearly $\delta(K_n) = n - 1$.
- In the case of a tree T_n on n vertices (where $n \geq 3$, because a tree on 2 vertices would just be K_2 and we have already handled this example), we have so far seen:
 - that T_n must have leaves, and therefore $\delta(T_n) = 1$,

Reminder: Whitney's Theorem tested on 'extreme' examples

- In the case of the complete graph K_n ($n \geq 2$) we have seen that:
 - $\kappa(K_n) = n - 1$ by definition,
 - $\lambda(K_n) = n - 1$ by the analysis we did last time,
 - and clearly $\delta(K_n) = n - 1$.
- In the case of a tree T_n on n vertices (where $n \geq 3$, because a tree on 2 vertices would just be K_2 and we have already handled this example), we have so far seen:
 - that T_n must have leaves, and therefore $\delta(T_n) = 1$,
 - and that every edge is a bridge, and thus $\lambda(T_n) = 1$.

Reminder: Whitney's Theorem tested on 'extreme' examples

- In the case of the complete graph K_n ($n \geq 2$) we have seen that:
 - $\kappa(K_n) = n - 1$ by definition,
 - $\lambda(K_n) = n - 1$ by the analysis we did last time,
 - and clearly $\delta(K_n) = n - 1$.
- In the case of a tree T_n on n vertices (where $n \geq 3$, because a tree on 2 vertices would just be K_2 and we have already handled this example), we have so far seen:
 - that T_n must have leaves, and therefore $\delta(T_n) = 1$,
 - and that every edge is a bridge, and thus $\lambda(T_n) = 1$.
 - Finally, such a tree will have vertices of degree ≥ 2 , and each such vertex will be a cutvertex. Thus we will have $\kappa(T_n) = 1$.

Whitney's Theorem

Theorem 1 (Whitney, 1932)

For every connected graph G (of order ≥ 2), we have

$$\kappa(G) \leq \lambda(G) \leq \delta(G).$$

Whitney's Theorem

Theorem 1 (Whitney, 1932)

For every connected graph G (of order ≥ 2), we have

$$\kappa(G) \leq \lambda(G) \leq \delta(G).$$

We have already proved (see Lecture 8) part of the theorem, that is, that $\lambda(G) \leq \delta(G)$.

Whitney's Theorem

Theorem 1 (Whitney, 1932)

For every connected graph G (of order ≥ 2), we have

$$\kappa(G) \leq \lambda(G) \leq \delta(G).$$

We have already proved (see Lecture 8) part of the theorem, that is, that $\lambda(G) \leq \delta(G)$.

Before we justify the rest of the theorem's conclusion, we are going to take a detour, and gather a few more characterisations for trees which can be justified using the most recent results.

Characterisations of trees

Reminder: Theorem 1a from Lecture 7

Let T be a graph of order n . The following two statements are equivalent.

- (i) T is a tree (that is, T is acyclic and connected).
- (ii) T is connected and has precisely $n - 1$ edges.

Characterisations of trees

Reminder: Theorem 1a from Lecture 7

Let T be a graph of order n . The following two statements are equivalent.

- (i) T is a tree (that is, T is acyclic and connected).
- (ii) T is connected and has precisely $n - 1$ edges.

Theorem 2a

Let T be a graph of order n . The following statements are equivalent.

- (i) T is a tree.

Characterisations of trees

Reminder: Theorem 1a from Lecture 7

Let T be a graph of order n . The following two statements are equivalent.

- (i) T is a tree (that is, T is acyclic and connected).
- (ii) T is connected and has precisely $n - 1$ edges.

Theorem 2a

Let T be a graph of order n . The following statements are equivalent.

- (i) T is a tree.
- (ii') T is connected and every edge is a bridge.

Characterisations of trees

Reminder: Theorem 1a from Lecture 7

Let T be a graph of order n . The following two statements are equivalent.

- (i) T is a tree (that is, T is acyclic and connected).
- (ii) T is connected and has precisely $n - 1$ edges.

Theorem 2a

Let T be a graph of order n . The following statements are equivalent.

- (i) T is a tree.
- (ii') T is connected and every edge is a bridge.
- (iii') T is acyclic and has precisely $n - 1$ edges.

Characterisations of trees

Reminder: Theorem 1a from Lecture 7

Let T be a graph of order n . The following two statements are equivalent.

- (i) T is a tree (that is, T is acyclic and connected).
- (ii) T is connected and has precisely $n - 1$ edges.

Theorem 2a

Let T be a graph of order n . The following statements are equivalent.

- (i) T is a tree.
- (ii') T is connected and every edge is a bridge.
- (iii') T is acyclic and has precisely $n - 1$ edges.
- (iv') T has $n - 1$ edges and every edge is a bridge.

Characterisations of trees

Reminder: Theorem 1a from Lecture 7

Let T be a graph of order n . The following two statements are equivalent.

- (i) T is a tree (that is, T is acyclic and connected).
- (ii) T is connected and has precisely $n - 1$ edges.

Theorem 2a

Let T be a graph of order n . The following statements are equivalent.

- (i) T is a tree.
- (ii') T is connected and every edge is a bridge.
- (iii') T is acyclic and has precisely $n - 1$ edges.
- (iv') T has $n - 1$ edges and every edge is a bridge.
- (v') For every two different vertices u, v of T , there is a **unique** path in T starting at u and ending at v .

Characterisations of trees

Reminder: Theorem 1a from Lecture 7

Let T be a graph of order n . The following two statements are equivalent.

- (i) T is a tree (that is, T is acyclic and connected).
- (ii) T is connected and has precisely $n - 1$ edges.

Theorem 2a

Let T be a graph of order n . The following statements are equivalent.

- (i) T is a tree.
- (ii') T is connected and every edge is a bridge.
- (iii') T is acyclic and has precisely $n - 1$ edges.
- (iv') T has $n - 1$ edges and every edge is a bridge.
- (v') For every two different vertices u, v of T , there is a **unique** path in T starting at u and ending at v .
- (vi') T is acyclic, and adding to it a single new edge creates precisely one cycle (regardless of where this edge is added).

Proof of Theorem 2a

We will verify the desired equivalences by proving the following:

$$(i) \Rightarrow (iv') \Rightarrow (iii') \Rightarrow (ii') \Rightarrow (i)$$

Proof of Theorem 2a

We will verify the desired equivalences by proving the following:

$$(i) \Rightarrow (iv') \Rightarrow (iii') \Rightarrow (ii') \Rightarrow (i) \quad \text{and} \quad (i) \Rightarrow (v') \Rightarrow (vi') \Rightarrow (i).$$

Proof of Theorem 2a

We will verify the desired equivalences by proving the following:

$$(i) \Rightarrow (iv') \Rightarrow (iii') \Rightarrow (ii') \Rightarrow (i) \quad \text{and} \quad (i) \Rightarrow (v') \Rightarrow (vi') \Rightarrow (i).$$

First let's recall what we have already shown:

- 1 If T is a tree on n vertices, then T has precisely $n - 1$ edges (*this was shown within the proof of Theorem 1a*).

Proof of Theorem 2a

We will verify the desired equivalences by proving the following:

$$(i) \Rightarrow (iv') \Rightarrow (iii') \Rightarrow (ii') \Rightarrow (i) \quad \text{and} \quad (i) \Rightarrow (v') \Rightarrow (vi') \Rightarrow (i).$$

First let's recall what we have already shown:

- 1 If T is a tree on n vertices, then T has precisely $n - 1$ edges (*this was shown within the proof of Theorem 1a*).
- 2 An edge e is a bridge of T if and only if e does NOT belong to any cycle contained in T (*this is Prop 1 from last time*).

Proof of Theorem 2a

We will verify the desired equivalences by proving the following:

$$(i) \Rightarrow (iv') \Rightarrow (iii') \Rightarrow (ii') \Rightarrow (i) \quad \text{and} \quad (i) \Rightarrow (v') \Rightarrow (vi') \Rightarrow (i).$$

First let's recall what we have already shown:

- 1 If T is a tree on n vertices, then T has precisely $n - 1$ edges (*this was shown within the proof of Theorem 1a*).
- 2 An edge e is a bridge of T if and only if e does NOT belong to any cycle contained in T (*this is Prop 1 from last time*).
- 3 If T is a tree, then, for every two different vertices u, v of T , there is a **unique** path in T starting at u and ending at v (*this is Prop 3 from last time*).

Proof of Theorem 2a

We will verify the desired equivalences by proving the following:

$$(i) \Rightarrow (iv') \Rightarrow (iii') \Rightarrow (ii') \Rightarrow (i) \quad \text{and} \quad (i) \Rightarrow (v') \Rightarrow (vi') \Rightarrow (i).$$

First let's recall what we have already shown:

- 1 If T is a tree on n vertices, then T has precisely $n - 1$ edges (*this was shown within the proof of Theorem 1a*).
- 2 An edge e is a bridge of T if and only if e does NOT belong to any cycle contained in T (*this is Prop 1 from last time*).
- 3 If T is a tree, then, for every two different vertices u, v of T , there is a **unique** path in T starting at u and ending at v (*this is Prop 3 from last time; notice also that it is $(i) \Rightarrow (v')$*).

Proof of Theorem 2a

We will verify the desired equivalences by proving the following:

$$(i) \Rightarrow (iv') \Rightarrow (iii') \Rightarrow (ii') \Rightarrow (i) \quad \text{and} \quad (i) \Rightarrow (v') \Rightarrow (vi') \Rightarrow (i).$$

First let's recall what we have already shown:

- 1 If T is a tree on n vertices, then T has precisely $n - 1$ edges (*this was shown within the proof of Theorem 1a*).
- 2 An edge e is a bridge of T if and only if e does NOT belong to any cycle contained in T (*this is Prop 1 from last time*).
- 3 If T is a tree, then, for every two different vertices u, v of T , there is a **unique** path in T starting at u and ending at v (*this is Prop 3 from last time; notice also that it is $(i) \Rightarrow (v')$*).

$(i) \Rightarrow (iv')$ Since T is a tree on n vertices, we know that it has $n - 1$ edges. Moreover, T contains no cycles, and thus none of its edges belongs to a cycle. By Prop 1 from last time, this implies that each of the edges of T is a bridge.

Proof of Theorem 2a

We will verify the desired equivalences by proving the following:

$$(i) \Rightarrow (iv') \Rightarrow (iii') \Rightarrow (ii') \Rightarrow (i) \quad \text{and} \quad (i) \Rightarrow (v') \Rightarrow (vi') \Rightarrow (i).$$

First let's recall what we have already shown:

- 1 If T is a tree on n vertices, then T has precisely $n - 1$ edges (*this was shown within the proof of Theorem 1a*).
- 2 An edge e is a bridge of T if and only if e does NOT belong to any cycle contained in T (*this is Prop 1 from last time*).
- 3 If T is a tree, then, for every two different vertices u, v of T , there is a **unique** path in T starting at u and ending at v (*this is Prop 3 from last time; notice also that it is $(i) \Rightarrow (v')$*).

$(i) \Rightarrow (iv')$ Since T is a tree on n vertices, we know that it has $n - 1$ edges. Moreover, T contains no cycles, and thus none of its edges belongs to a cycle. By Prop 1 from last time, this implies that each of the edges of T is a bridge.

$(iv') \Rightarrow (iii')$ Already by our assumptions we know that T has $n - 1$ edges. Moreover, we know that every edge of T is a bridge. Note now that, if T contained cycles, then all the edges belonging to any of these cycles would NOT be bridges of T . Thus the latter assumption about T implies that T is acyclic.

Proof of Theorem 2a (cont.)

(iii') \Rightarrow (ii') Again, because T is assumed acyclic, we get that every edge of T is a bridge. Moreover, T is either a tree or a forest. In the former case, we immediately get that T is connected.

We will now check that the latter case is not possible given our assumptions.

Proof of Theorem 2a (cont.)

(iii') \Rightarrow (ii') Again, because T is assumed acyclic, we get that every edge of T is a bridge. Moreover, T is either a tree or a forest. In the former case, we immediately get that T is connected.

We will now check that the latter case is not possible given our assumptions.

Assume that we could have a forest T satisfying the given assumptions. In other words, assume that T is an acyclic graph with at least two connected components, total number of vertices n and total number of edges $n - 1$. Write T_1, T_2, \dots, T_k for the connected components of T (where $k \geq 2$).

Proof of Theorem 2a (cont.)

(iii') \Rightarrow (ii') Again, because T is assumed acyclic, we get that every edge of T is a bridge. Moreover, T is either a tree or a forest. In the former case, we immediately get that T is connected.

We will now check that the latter case is not possible given our assumptions.

Assume that we could have a forest T satisfying the given assumptions. In other words, assume that T is an acyclic graph with at least two connected components, total number of vertices n and total number of edges $n - 1$. Write T_1, T_2, \dots, T_k for the connected components of T (where $k \geq 2$).

Since T is a forest, each of its connected components T_i is a tree, and thus we know that $e(T_i) = |T_i| - 1$ (that is, the size of T_i is equal to the order of T_i minus 1).

Proof of Theorem 2a (cont.)

(iii') \Rightarrow (ii') Again, because T is assumed acyclic, we get that every edge of T is a bridge. Moreover, T is either a tree or a forest. In the former case, we immediately get that T is connected.

We will now check that the latter case is not possible given our assumptions.

Assume that we could have a forest T satisfying the given assumptions. In other words, assume that T is an acyclic graph with at least two connected components, total number of vertices n and total number of edges $n - 1$. Write T_1, T_2, \dots, T_k for the connected components of T (where $k \geq 2$).

Since T is a forest, each of its connected components T_i is a tree, and thus we know that $e(T_i) = |T_i| - 1$ (that is, the size of T_i is equal to the order of T_i minus 1).

But then we have

$$\begin{aligned} n - 1 &= e(T) = e(T_1) + e(T_2) + \cdots + e(T_k) \\ &= (|T_1| - 1) + (|T_2| - 1) + \cdots + (|T_k| - 1) \\ &= (|T_1| + |T_2| + \cdots + |T_k|) - k = |T| - k = n - k, \end{aligned}$$

which contradicts the assumption that $k > 1$.

Proof of Theorem 2a (cont.)

(iii') \Rightarrow (ii') Again, because T is assumed acyclic, we get that every edge of T is a bridge. Moreover, T is either a tree or a forest. In the former case, we immediately get that T is connected.

We will now check that the latter case is not possible given our assumptions.

Assume that we could have a forest T satisfying the given assumptions. In other words, assume that T is an acyclic graph with at least two connected components, total number of vertices n and total number of edges $n - 1$. Write T_1, T_2, \dots, T_k for the connected components of T (where $k \geq 2$).

Since T is a forest, each of its connected components T_i is a tree, and thus we know that $e(T_i) = |T_i| - 1$ (that is, the size of T_i is equal to the order of T_i minus 1).

But then we have

$$\begin{aligned} n - 1 &= e(T) = e(T_1) + e(T_2) + \cdots + e(T_k) \\ &= (|T_1| - 1) + (|T_2| - 1) + \cdots + (|T_k| - 1) \\ &= (|T_1| + |T_2| + \cdots + |T_k|) - k = |T| - k = n - k, \end{aligned}$$

which contradicts the assumption that $k > 1$. Thus this assumption was incorrect.

Proof of Theorem 2a (cont.)

(iii') \Rightarrow (ii') Again, because T is assumed acyclic, we get that every edge of T is a bridge. Moreover, T is either a tree or a forest. In the former case, we immediately get that T is connected.

We will now check that the latter case is not possible given our assumptions.

Assume that we could have a forest T satisfying the given assumptions. In other words, assume that T is an acyclic graph with at least two connected components, total number of vertices n and total number of edges $n - 1$. Write T_1, T_2, \dots, T_k for the connected components of T (where $k \geq 2$).

Since T is a forest, each of its connected components T_i is a tree, and thus we know that $e(T_i) = |T_i| - 1$ (that is, the size of T_i is equal to the order of T_i minus 1).

But then we have

$$\begin{aligned} n - 1 &= e(T) = e(T_1) + e(T_2) + \dots + e(T_k) \\ &= (|T_1| - 1) + (|T_2| - 1) + \dots + (|T_k| - 1) \\ &= (|T_1| + |T_2| + \dots + |T_k|) - k = |T| - k = n - k, \end{aligned}$$

which contradicts the assumption that $k > 1$. Thus this assumption was incorrect.

(ii') \Rightarrow (i) Since we assumed that every edge of T is a bridge, T must be acyclic. Moreover, T is connected by assumption. Thus T is a tree.

Proof of Theorem 2a (cont.)

$(i) \Rightarrow (v')$ This is precisely Proposition 3 from last time.

Proof of Theorem 2a (cont.)

$(i) \Rightarrow (v')$ This is precisely Proposition 3 from last time.

$(v') \Rightarrow (vi')$ Assume that we know that, for every two different vertices u, v of T , there is a **unique** path in T starting at u and ending at v .

Proof of Theorem 2a (cont.)

$(i) \Rightarrow (v')$ This is precisely Proposition 3 from last time.

$(v') \Rightarrow (vi')$ Assume that we know that, for every two different vertices u, v of T , there is a **unique** path in T starting at u and ending at v .

First of all, this implies that T has no cycles (because, if we had a cycle, then for every pair of different vertices u_1, v_1 on the cycle, there would be two $u_1 - v_1$ paths contained in the cycle itself).

Proof of Theorem 2a (cont.)

$(i) \Rightarrow (v')$ This is precisely Proposition 3 from last time.

$(v') \Rightarrow (vi')$ Assume that we know that, for every two different vertices u, v of T , there is a **unique** path in T starting at u and ending at v .

First of all, this implies that T has no cycles (because, if we had a cycle, then for every pair of different vertices u_1, v_1 on the cycle, there would be two $u_1 - v_1$ paths contained in the cycle itself).

Consider now two different vertices u_2, v_2 in T which are not already joined by an edge in T (*clearly this excludes the cases where $T = K_1$ or $T = K_2$, for which statement (vi') is true in a vacuous manner, since we simply cannot add any (more) edges*),

Proof of Theorem 2a (cont.)

$(i) \Rightarrow (v')$ This is precisely Proposition 3 from last time.

$(v') \Rightarrow (vi')$ Assume that we know that, for every two different vertices u, v of T , there is a **unique** path in T starting at u and ending at v .

First of all, this implies that T has no cycles (because, if we had a cycle, then for every pair of different vertices u_1, v_1 on the cycle, there would be two $u_1 - v_1$ paths contained in the cycle itself).

Consider now two different vertices u_2, v_2 in T which are not already joined by an edge in T (clearly this excludes the cases where $T = K_1$ or $T = K_2$, for which statement (vi') is true in a vacuous manner, since we simply cannot add any (more) edges), and add the edge $e_2 = \{u_2, v_2\}$ to get a supergraph \tilde{T} of T .

Proof of Theorem 2a (cont.)

$(i) \Rightarrow (v')$ This is precisely Proposition 3 from last time.

$(v') \Rightarrow (vi')$ Assume that we know that, for every two different vertices u, v of T , there is a **unique** path in T starting at u and ending at v .

First of all, this implies that T has no cycles (because, if we had a cycle, then for every pair of different vertices u_1, v_1 on the cycle, there would be two $u_1 - v_1$ paths contained in the cycle itself).

Consider now two different vertices u_2, v_2 in T which are not already joined by an edge in T (clearly this excludes the cases where $T = K_1$ or $T = K_2$, for which statement (vi') is true in a vacuous manner, since we simply cannot add any (more) edges), and add the edge $e_2 = \{u_2, v_2\}$ to get a supergraph \tilde{T} of T .

By our assumption, there was already a $u_2 - v_2$ path P_0 in T . Note that this path has length ≥ 2 (since u_2 and v_2 are not joined in T), and thus, with the addition of the new edge, we now have a cycle C_0 in \tilde{T} which contains the vertices u_2, v_2 .

Proof of Theorem 2a (cont.)

$(i) \Rightarrow (v')$ This is precisely Proposition 3 from last time.

$(v') \Rightarrow (vi')$ Assume that we know that, for every two different vertices u, v of T , there is a **unique** path in T starting at u and ending at v .

First of all, this implies that T has no cycles (because, if we had a cycle, then for every pair of different vertices u_1, v_1 on the cycle, there would be two $u_1 - v_1$ paths contained in the cycle itself).

Consider now two different vertices u_2, v_2 in T which are not already joined by an edge in T (clearly this excludes the cases where $T = K_1$ or $T = K_2$, for which statement (vi') is true in a vacuous manner, since we simply cannot add any (more) edges), and add the edge $e_2 = \{u_2, v_2\}$ to get a supergraph \tilde{T} of T .

By our assumption, there was already a $u_2 - v_2$ path P_0 in T . Note that this path has length ≥ 2 (since u_2 and v_2 are not joined in T), and thus, with the addition of the new edge, we now have a cycle C_0 in \tilde{T} which contains the vertices u_2, v_2 .

It remains to show that C_0 is the only cycle of \tilde{T} .

- Clearly we can't have a cycle in \tilde{T} which does not contain the edge e_2 , because then this would also be a cycle in the original graph T .

Proof of Theorem 2a (cont.)

$(i) \Rightarrow (v')$ This is precisely Proposition 3 from last time.

$(v') \Rightarrow (vi')$ Assume that we know that, for every two different vertices u, v of T , there is a **unique** path in T starting at u and ending at v .

First of all, this implies that T has no cycles (because, if we had a cycle, then for every pair of different vertices u_1, v_1 on the cycle, there would be two $u_1 - v_1$ paths contained in the cycle itself).

Consider now two different vertices u_2, v_2 in T which are not already joined by an edge in T (clearly this excludes the cases where $T = K_1$ or $T = K_2$, for which statement (vi') is true in a vacuous manner, since we simply cannot add any (more) edges), and add the edge $e_2 = \{u_2, v_2\}$ to get a supergraph \tilde{T} of T .

By our assumption, there was already a $u_2 - v_2$ path P_0 in T . Note that this path has length ≥ 2 (since u_2 and v_2 are not joined in T), and thus, with the addition of the new edge, we now have a cycle C_0 in \tilde{T} which contains the vertices u_2, v_2 .

It remains to show that C_0 is the only cycle of \tilde{T} .

- Clearly we can't have a cycle in \tilde{T} which does not contain the edge e_2 , because then this would also be a cycle in the original graph T .
- If C_1 were another cycle in \tilde{T} which contained the edge e_2 , and, say, we could write $C_1 : u_2 z_1 z_2 \cdots z_{k-1} v_2 u_2$, then the path $P_1 : u_2 z_1 z_2 \cdots z_{k-1} v_2$ should be a $u_2 - v_2$ path in T different from P_0 . This contradicts our assumption about the uniqueness of P_0 .

Proof of Theorem 2a (cont.)

Finally, we verify that

$$(vi') \Rightarrow (i)$$

Proof of Theorem 2a (cont.)

Finally, we verify that

$(vi') \Rightarrow (i)$ That is, we assume that T is acyclic, and moreover, that, if we add a single new edge, this creates precisely one cycle (regardless of where this edge is added), and we show that T is a tree.

Proof of Theorem 2a (cont.)

Finally, we verify that

$(vi') \Rightarrow (i)$ That is, we assume that T is acyclic, and moreover, that, if we add a single new edge, this creates precisely one cycle (regardless of where this edge is added), and we show that T is a tree.

We already have that T is acyclic. Let's assume that T is NOT connected. Then T has at least two connected components, say components T_1 and T_2 .

Consider a vertex u in T_1 and a vertex v in T_2 , and add the edge $e_0 = \{u, v\}$.

Proof of Theorem 2a (cont.)

Finally, we verify that

$(vi') \Rightarrow (i)$ That is, we assume that T is acyclic, and moreover, that, if we add a single new edge, this creates precisely one cycle (regardless of where this edge is added), and we show that T is a tree.

We already have that T is acyclic. Let's assume that T is NOT connected. Then T has at least two connected components, say components T_1 and T_2 .

Consider a vertex u in T_1 and a vertex v in T_2 , and add the edge $e_0 = \{u, v\}$. Then, by our assumption, this creates precisely one cycle (which must contain the edge e_0 , because otherwise T would already contain a cycle).

Proof of Theorem 2a (cont.)

Finally, we verify that

$(vi') \Rightarrow (i)$ That is, we assume that T is acyclic, and moreover, that, if we add a single new edge, this creates precisely one cycle (regardless of where this edge is added), and we show that T is a tree.

We already have that T is acyclic. Let's assume that T is NOT connected. Then T has at least two connected components, say components T_1 and T_2 .

Consider a vertex u in T_1 and a vertex v in T_2 , and add the edge $e_0 = \{u, v\}$. Then, by our assumption, this creates precisely one cycle (which must contain the edge e_0 , because otherwise T would already contain a cycle).

But then this shows that there is a $u - v$ path in the original graph T (since we can 'travel' from u to v on the cycle without traversing the edge e_0), which contradicts the assumption that u and v belong to different connected components of the original graph T .

Proof of Theorem 2a (cont.)

Finally, we verify that

$(vi') \Rightarrow (i)$ That is, we assume that T is acyclic, and moreover, that, if we add a single new edge, this creates precisely one cycle (regardless of where this edge is added), and we show that T is a tree.

We already have that T is acyclic. Let's assume that T is NOT connected. Then T has at least two connected components, say components T_1 and T_2 .

Consider a vertex u in T_1 and a vertex v in T_2 , and add the edge $e_0 = \{u, v\}$. Then, by our assumption, this creates precisely one cycle (which must contain the edge e_0 , because otherwise T would already contain a cycle).

But then this shows that there is a $u - v$ path in the original graph T (since we can 'travel' from u to v on the cycle without traversing the edge e_0), which contradicts the assumption that u and v belong to different connected components of the original graph T .

We conclude that T must be connected, and thus that T is a tree.

Recap of the characterisations for trees

Theorem 1a from Lecture 7

Let T be a graph of order n . The following two statements are equivalent.

- (i) T is a tree (that is, T is acyclic and connected).
- (ii) T is connected and has precisely $n - 1$ edges.

Theorem 2a

Let T be a graph of order n . The following statements are equivalent.

- (i) T is a tree.
- (ii') T is connected and every edge is a bridge.
- (iii') T is acyclic and has precisely $n - 1$ edges.
- (iv') T has $n - 1$ edges and every edge is a bridge.
- (v') For every two different vertices u, v of T , there is a **unique** path in T starting at u and ending at v .
- (vi') T is acyclic, and adding to it a single new edge creates precisely one cycle (regardless of where this edge is added).

Back to Whitney's theorem

Theorem 1 (Whitney, 1932)

For every connected graph G (of order ≥ 2), we have

$$\kappa(G) \leq \lambda(G) \leq \delta(G).$$

We have proved so far (see Lecture 8) that $\lambda(G) \leq \delta(G)$.

Proving that $\kappa(G) \leq \lambda(G)$

Consider a connected graph G on at least 2 vertices, and let n be the order of G (as we just said, $n \geq 2$).

Consider also an edge cut $\{e_1, e_2, \dots, e_{t_0}\}$ of G such that $t_0 = \lambda(G)$. In other words, $G - \{e_1, e_2, \dots, e_{t_0}\}$ is a disconnected subgraph, while t_0 , the cardinality of this edge cut, is as small as possible. By what we have already showed, $t_0 \leq \delta(G) \leq n - 1$.

Proving that $\kappa(G) \leq \lambda(G)$

Consider a connected graph G on at least 2 vertices, and let n be the order of G (as we just said, $n \geq 2$).

Consider also an edge cut $\{e_1, e_2, \dots, e_{t_0}\}$ of G such that $t_0 = \lambda(G)$. In other words, $G - \{e_1, e_2, \dots, e_{t_0}\}$ is a disconnected subgraph, while t_0 , the cardinality of this edge cut, is as small as possible. By what we have already showed, $t_0 \leq \delta(G) \leq n - 1$.

'Easier' case: $t_0 = \lambda(G) = n - 1$. Then, we must have $\delta(G) = n - 1$ too, and thus $G = K_n$. But then, according to our definitions, $\kappa(G) = n - 1$ too, and the inequality $\kappa(G) \leq \lambda(G)$ holds (in fact, we have equality).

Proving that $\kappa(G) \leq \lambda(G)$

Consider a connected graph G on at least 2 vertices, and let n be the order of G (as we just said, $n \geq 2$).

Consider also an edge cut $\{e_1, e_2, \dots, e_{t_0}\}$ of G such that $t_0 = \lambda(G)$. In other words, $G - \{e_1, e_2, \dots, e_{t_0}\}$ is a disconnected subgraph, while t_0 , the cardinality of this edge cut, is as small as possible. By what we have already showed, $t_0 \leq \delta(G) \leq n - 1$.

'Easier' case: $t_0 = \lambda(G) = n - 1$. Then, we must have $\delta(G) = n - 1$ too, and thus $G = K_n$. But then, according to our definitions, $\kappa(G) = n - 1$ too, and the inequality $\kappa(G) \leq \lambda(G)$ holds (in fact, we have equality).

Remaining cases: Assume now that $t_0 \leq n - 2$. Since $G - \{e_1, e_2, \dots, e_{t_0}\}$ is a disconnected subgraph, we can find vertices u_1, v_1 of G which fall into different connected components of $G - \{e_1, e_2, \dots, e_{t_0}\}$.

Proving that $\kappa(G) \leq \lambda(G)$ (cont.)

Remaining cases: Assume now that $t_0 \leq n - 2$. Since $G - \{e_1, e_2, \dots, e_{t_0}\}$ is a disconnected subgraph, we can find vertices u_1, v_1 of G which fall into different connected components of $G - \{e_1, e_2, \dots, e_{t_0}\}$.

We consider two subcases:

Subcase 1: u_1, v_1 are NOT joined by an edge in G . Then any path in G connecting u_1 and v_1 must have length ≥ 2 (or in other words, it must contain intermediate vertices too).

Proving that $\kappa(G) \leq \lambda(G)$ (cont.)

Remaining cases: Assume now that $t_0 \leq n - 2$. Since $G - \{e_1, e_2, \dots, e_{t_0}\}$ is a disconnected subgraph, we can find vertices u_1, v_1 of G which fall into different connected components of $G - \{e_1, e_2, \dots, e_{t_0}\}$.

We consider two subcases:

Subcase 1: u_1, v_1 are NOT joined by an edge in G . Then any path in G connecting u_1 and v_1 must have length ≥ 2 (or in other words, it must contain intermediate vertices too).

Moreover, every path in G connecting u_1 and v_1 must traverse one or more of the edges e_1, e_2, \dots, e_{t_0} .

Proving that $\kappa(G) \leq \lambda(G)$ (cont.)

Remaining cases: Assume now that $t_0 \leq n - 2$. Since $G - \{e_1, e_2, \dots, e_{t_0}\}$ is a disconnected subgraph, we can find vertices u_1, v_1 of G which fall into different connected components of $G - \{e_1, e_2, \dots, e_{t_0}\}$.

We consider two subcases:

Subcase 1: u_1, v_1 are NOT joined by an edge in G . Then any path in G connecting u_1 and v_1 must have length ≥ 2 (or in other words, it must contain intermediate vertices too).

Moreover, every path in G connecting u_1 and v_1 must traverse one or more of the edges e_1, e_2, \dots, e_{t_0} . (Indeed, if we could also find a $u_1 - v_1$ path in G which did not contain any of the edges e_1, e_2, \dots, e_{t_0} , then this would be a path in the subgraph $G - \{e_1, e_2, \dots, e_{t_0}\}$ too, and thus u_1, v_1 would not belong to different connected components of the subgraph.)

Proving that $\kappa(G) \leq \lambda(G)$ (cont.)

Remaining cases: Assume now that $t_0 \leq n - 2$. Since $G - \{e_1, e_2, \dots, e_{t_0}\}$ is a disconnected subgraph, we can find vertices u_1, v_1 of G which fall into different connected components of $G - \{e_1, e_2, \dots, e_{t_0}\}$.

We consider two subcases:

Subcase 1: u_1, v_1 are NOT joined by an edge in G . Then any path in G connecting u_1 and v_1 must have length ≥ 2 (or in other words, it must contain intermediate vertices too).

Moreover, every path in G connecting u_1 and v_1 must traverse one or more of the edges e_1, e_2, \dots, e_{t_0} . (Indeed, if we could also find a $u_1 - v_1$ path in G which did not contain any of the edges e_1, e_2, \dots, e_{t_0} , then this would be a path in the subgraph $G - \{e_1, e_2, \dots, e_{t_0}\}$ too, and thus u_1, v_1 would not belong to different connected components of the subgraph.)

Now, for each of the edges e_i , pick one of its endvertices **which is different from both u_1 and v_1** (this is always possible here because the edge $\{u_1, v_1\}$ does not exist in G), and denote it by w_i .

Proving that $\kappa(G) \leq \lambda(G)$ (cont.)

Remaining cases: Assume now that $t_0 \leq n - 2$. Since $G - \{e_1, e_2, \dots, e_{t_0}\}$ is a disconnected subgraph, we can find vertices u_1, v_1 of G which fall into different connected components of $G - \{e_1, e_2, \dots, e_{t_0}\}$.

We consider two subcases:

Subcase 1: u_1, v_1 are NOT joined by an edge in G . Then any path in G connecting u_1 and v_1 must have length ≥ 2 (or in other words, it must contain intermediate vertices too).

Moreover, every path in G connecting u_1 and v_1 must traverse one or more of the edges e_1, e_2, \dots, e_{t_0} . (Indeed, if we could also find a $u_1 - v_1$ path in G which did not contain any of the edges e_1, e_2, \dots, e_{t_0} , then this would be a path in the subgraph $G - \{e_1, e_2, \dots, e_{t_0}\}$ too, and thus u_1, v_1 would not belong to different connected components of the subgraph.)

Now, for each of the edges e_i , pick one of its endvertices **which is different from both u_1 and v_1** (this is always possible here because the edge $\{u_1, v_1\}$ does not exist in G), and denote it by w_i . Then the vertex set $\{w_1, w_2, \dots, w_{t_0}\}$ is a vertex cut of G , since

$$G - \{w_1, w_2, \dots, w_{t_0}\} \subseteq G - \{e_1, e_2, \dots, e_{t_0}\},$$

and thus the vertices u_1, v_1 , which are still left in $G - \{w_1, w_2, \dots, w_{t_0}\}$, will again be separated.

One more side note here: $\{w_1, w_2, \dots, w_{t_0}\}$ contains **at most** t_0 vertices (because some vertex here might be a common endvertex of two or more of the edges e_1, e_2, \dots, e_{t_0} , and might have been picked more than once).

This shows that $\kappa(G) \leq t_0 = \lambda(G)$ in this subcase.

Proving that $\kappa(G) \leq \lambda(G)$ (cont.)

Subcase 2: u_1, v_1 are joined by an edge in G . Then, because u_1, v_1 are separated in $G - \{e_1, e_2, \dots, e_{t_0}\}$, one of the edges that we remove must be the edge $\{u_1, v_1\}$.

Proving that $\kappa(G) \leq \lambda(G)$ (cont.)

Subcase 2: u_1, v_1 are joined by an edge in G . Then, because u_1, v_1 are separated in $G - \{e_1, e_2, \dots, e_{t_0}\}$, one of the edges that we remove must be the edge $\{u_1, v_1\}$. Say without loss of generality that $\{u_1, v_1\} = e_{t_0}$.

Proving that $\kappa(G) \leq \lambda(G)$ (cont.)

Subcase 2: u_1, v_1 are joined by an edge in G . Then, because u_1, v_1 are separated in $G - \{e_1, e_2, \dots, e_{t_0}\}$, one of the edges that we remove must be the edge $\{u_1, v_1\}$. Say without loss of generality that $\{u_1, v_1\} = e_{t_0}$.

Then, as before, for each of the edges e_i with $i \neq t_0$, pick one of its endvertices **which is different from both u_1 and v_1** , and denote it by w_i .

Proving that $\kappa(G) \leq \lambda(G)$ (cont.)

Subcase 2: u_1, v_1 are joined by an edge in G . Then, because u_1, v_1 are separated in $G - \{e_1, e_2, \dots, e_{t_0}\}$, one of the edges that we remove must be the edge $\{u_1, v_1\}$. Say without loss of generality that $\{u_1, v_1\} = e_{t_0}$.

Then, as before, for each of the edges e_i with $i \neq t_0$, pick one of its endvertices **which is different from both u_1 and v_1** , and denote it by w_i .

Note now that

$$(G - \{w_1, w_2, \dots, w_{t_0-1}\}) - e_{t_0} \subseteq G - \{e_1, e_2, \dots, e_{t_0}\},$$

and thus u_1, v_1 are separated in the smaller subgraph too.

Note also that the subgraph $(G - \{w_1, w_2, \dots, w_{t_0-1}\}) - e_{t_0}$ contains at least $n - (t_0 - 1) = n - t_0 + 1 \geq n + 1 - (n - 2) = 3$ vertices (where we're using the initial assumption that $t_0 \leq n - 2$). Thus it contains at least one more vertex z_0 different from u_1 and v_1 .

Proving that $\kappa(G) \leq \lambda(G)$ (cont.)

Subcase 2: u_1, v_1 are joined by an edge in G . Then, because u_1, v_1 are separated in $G - \{e_1, e_2, \dots, e_{t_0}\}$, one of the edges that we remove must be the edge $\{u_1, v_1\}$. Say without loss of generality that $\{u_1, v_1\} = e_{t_0}$.

Then, as before, for each of the edges e_i with $i \neq t_0$, pick one of its endvertices **which is different from both u_1 and v_1** , and denote it by w_i .

Note now that

$$(G - \{w_1, w_2, \dots, w_{t_0-1}\}) - e_{t_0} \subseteq G - \{e_1, e_2, \dots, e_{t_0}\},$$

and thus u_1, v_1 are separated in the smaller subgraph too.

Note also that the subgraph $(G - \{w_1, w_2, \dots, w_{t_0-1}\}) - e_{t_0}$ contains **at least** $n - (t_0 - 1) = n - t_0 + 1 \geq n + 1 - (n - 2) = 3$ vertices (where we're using the initial assumption that $t_0 \leq n - 2$). Thus it contains at least one more vertex z_0 different from u_1 and v_1 .

- If z_0 and u_1 are in the same connected component of $(G - \{w_1, w_2, \dots, w_{t_0-1}\}) - e_{t_0}$, **which implies that z_0 and v_1 are in different connected components**, then, instead of removing the edge e_{t_0} , remove the vertex u_1 . In the graph $G - \{w_1, w_2, \dots, w_{t_0-1}, u_1\}$, the vertices z_0 and v_1 are in different connected components, and thus the subset $\{w_1, w_2, \dots, w_{t_0-1}, u_1\}$ is a vertex cut of G with cardinality $\leq t_0 = \lambda(G)$.

Proving that $\kappa(G) \leq \lambda(G)$ (cont.)

Subcase 2: u_1, v_1 are joined by an edge in G . Then, because u_1, v_1 are separated in $G - \{e_1, e_2, \dots, e_{t_0}\}$, one of the edges that we remove must be the edge $\{u_1, v_1\}$. Say without loss of generality that $\{u_1, v_1\} = e_{t_0}$.

Then, as before, for each of the edges e_i with $i \neq t_0$, pick one of its endvertices **which is different from both u_1 and v_1** , and denote it by w_i .

Note now that

$$(G - \{w_1, w_2, \dots, w_{t_0-1}\}) - e_{t_0} \subseteq G - \{e_1, e_2, \dots, e_{t_0}\},$$





and thus u_1, v_1 are separated in the smaller subgraph too.

Note also that the subgraph $(G - \{w_1, w_2, \dots, w_{t_0-1}\}) - e_{t_0}$ contains **at least** $n - (t_0 - 1) = n - t_0 + 1 \geq n + 1 - (n - 2) = 3$ vertices (where we're using the initial assumption that $t_0 \leq n - 2$). Thus it contains at least one more vertex z_0 different from u_1 and v_1 .


- If z_0 and u_1 are in the same connected component of $(G - \{w_1, w_2, \dots, w_{t_0-1}\}) - e_{t_0}$, **which implies that z_0 and v_1 are in different connected components**, then, instead of removing the edge e_{t_0} , remove the vertex u_1 . In the graph $G - \{w_1, w_2, \dots, w_{t_0-1}, u_1\}$, the vertices z_0 and v_1 are in different connected components, and thus the subset $\{w_1, w_2, \dots, w_{t_0-1}, u_1\}$ is a vertex cut of G with cardinality $\leq t_0 = \lambda(G)$.
- If instead z_0 and u_1 are in **different connected components** of $(G - \{w_1, w_2, \dots, w_{t_0-1}\}) - e_{t_0}$, then analogously, instead of the edge e_{t_0} , remove the vertex v_1 . Now, in the graph $G - \{w_1, w_2, \dots, w_{t_0-1}, v_1\}$, it's the vertices z_0 and u_1 which are in different connected components.

Checking Whitney's theorem (and the proof constructions) on examples

For each of the graphs G below, find $\delta(G)$, $\lambda(G)$ and $\kappa(G)$, as well as edge cuts and vertex cuts which 'capture' the parameters.




One more practice example



Same question as before: What is $\delta(G)$, $\lambda(G)$ and $\kappa(G)$ here?

Past exam problem

Consider the following connected graph G_0 .



- (a) Show that $\kappa(G_0) = 2$. Give a full justification.
- (b) What is $\lambda(G_0)$? Determine it precisely, and justify your answer fully.