

MATH 322 – Graph Theory

Fall Term 2021

Notes for Lecture 8

Tuesday, September 28

A detour: 'Forbidden' subgraphs

Many important results in Graph Theory are stated as follows: a graph G has a certain property if and only if we cannot find some 'not so nice' graphs $H_1, H_2, \dots, H_n, \dots$ among the subgraphs of G , or, even more 'strictly' sometimes, if we cannot find some given graphs $H_1, H_2, \dots, H_n, \dots$ among the induced subgraphs of G .
(we'll see some initial examples in the next two slides)

A detour: 'Forbidden' subgraphs

Many important results in Graph Theory are stated as follows: a graph G has a certain property if and only if we cannot find some 'not so nice' graphs $H_1, H_2, \dots, H_n, \dots$ among the subgraphs of G , or, even more 'strictly' sometimes, if we cannot find some given graphs $H_1, H_2, \dots, H_n, \dots$ among the induced subgraphs of G .
(we'll see some initial examples in the next two slides)

Special Terminology. We call a graph H a forbidden subgraph for a property P of graphs if the following holds true: given any graph G ,
property P holds true for G **if and only if**
 G does not contain H as an **induced subgraph**
(in other words, if H is not isomorphic to an induced subgraph of G).

Some examples

- A graph G is a tree or a forest if and only if G does not contain any cycles.

Some examples



- A graph G is a tree or a forest if and only if G does not contain any cycles.
- A criterion for bipartite graphs:

A graph G is (a subgraph of) a bipartite graph if and only if G does not contain any odd cycles (that is, if and only if none of the subgraphs of G is an odd cycle).

Some examples

- A graph G is a tree or a forest if and only if G does not contain any cycles.
- A criterion for bipartite graphs:

A graph G is (a subgraph of) a bipartite graph if and only if G does not contain any odd cycles (that is, if and only if none of the subgraphs of G is an odd cycle).



No odd cycles;
can view this as a subgraph of $K_{6,6}$ (how?)

Update on Terminology


Given a bipartite graph $K_{m,n}$, we will call any of its subgraphs a bipartite graph too.

Update on Terminology


Given a bipartite graph $K_{m,n}$, we will call any of its subgraphs a bipartite graph too. Moreover, most of the time we will call $K_{m,n}$ a complete bipartite graph to distinguish from the rest of its subgraphs.

Update on Terminology

Given a bipartite graph $K_{m,n}$, we will call **any of its subgraphs a bipartite graph** too. Moreover, most of the time we will call $K_{m,n}$ a **complete bipartite graph** to distinguish from the rest of its subgraphs.




Graph $K_{6,4}$




Bipartite Graph; Subgraph of $K_{10,7}$

Update on Terminology


Given a bipartite graph $K_{m,n}$, we will call **any of its subgraphs a bipartite graph** too. Moreover, most of the time we will call $K_{m,n}$ a **complete bipartite graph** to distinguish from the rest of its subgraphs.



Graph $K_{6,4}$



Bipartite Graph; Subgraph of $K_{10,7}$



No odd cycles;
viewed as a subgraph of $K_{6,6}$

Criterion for bipartite graphs

A graph G is (a subgraph of) a bipartite graph
if and only if G does not contain any **odd** cycles
(that is, if and only if none of the subgraphs of G is an odd cycle).

Criterion for bipartite graphs

A graph G is (a subgraph of) a bipartite graph
if and only if G does not contain any **odd** cycles
(that is, if and only if none of the subgraphs of G is an odd cycle).

Justification. We can directly check (say, by induction on the number k of vertices of a cycle that we consider each time) that a complete bipartite graph does NOT contain any odd cycles, so each of its subgraphs will also not contain any odd cycles.

*(In other words, first check that any $K_{m,n}$ does not contain 3-cycles, and then that it cannot contain 5-cycles and so on; this is because the parts of $K_{m,n}$ that the vertices of a cycle belong to have to be alternating, that is, if we start from a vertex in, say, the 'upper' part of $K_{m,n}$, we then **have to** move to a vertex in the 'lower' part, and so on.)*

Criterion for bipartite graphs

A graph G is (a subgraph of) a bipartite graph
if and only if G does not contain any **odd** cycles
(that is, if and only if none of the subgraphs of G is an odd cycle).

Justification. To show the converse, consider a graph G that does NOT contain any odd cycles.

– Note that it suffices to show that each connected component of G is a bipartite graph (attention: not necessarily complete), **therefore we can assume that G is connected** (because otherwise we focus our attention on one connected component of G at a time).

Criterion for bipartite graphs

A graph G is (a subgraph of) a bipartite graph
if and only if G does not contain any **odd** cycles
(that is, if and only if none of the subgraphs of G is an odd cycle).

Justification. To show the converse, consider a graph G that does NOT contain any odd cycles.

– Note that it suffices to show that each connected component of G is a bipartite graph (attention: not necessarily complete), **therefore we can assume that G is connected** (because otherwise we focus our attention on one connected component of G at a time).

Fix a vertex v_0 of G , and ‘colour’ it blue. Since G is connected, if we consider another vertex w of G , we can find at least one $v_0 - w$ path in G .

Criterion for bipartite graphs

A graph G is (a subgraph of) a bipartite graph
if and only if G does not contain any **odd** cycles
(that is, if and only if none of the subgraphs of G is an odd cycle).

Justification. To show the converse, consider a graph G that does NOT contain any odd cycles.

– Note that it suffices to show that each connected component of G is a bipartite graph (attention: not necessarily complete), **therefore we can assume that G is connected** (because otherwise we focus our attention on one connected component of G at a time).

Fix a vertex v_0 of G , and ‘colour’ it blue. Since G is connected, if we consider another vertex w of G , we can find at least one $v_0 - w$ path in G .

Idea: For each vertex w of G , choose a $v_0 - w$ path in G which has **the shortest possible length**. Denote this length by ℓ_w .

Criterion for bipartite graphs

A graph G is (a subgraph of) a bipartite graph
if and only if G does not contain any **odd** cycles
(that is, if and only if none of the subgraphs of G is an odd cycle).

Justification. To show the converse, consider a graph G that does NOT contain any odd cycles.

– Note that it suffices to show that each connected component of G is a bipartite graph (attention: not necessarily complete), **therefore we can assume that G is connected** (because otherwise we focus our attention on one connected component of G at a time).

Fix a vertex v_0 of G , and ‘colour’ it blue. Since G is connected, if we consider another vertex w of G , we can find at least one $v_0 - w$ path in G .

Idea: For each vertex w of G , choose a $v_0 - w$ path in G which has **the shortest possible length**. Denote this length by ℓ_w .

- If ℓ_w is an even integer, then ‘colour’ the vertex w blue.

Criterion for bipartite graphs

A graph G is (a subgraph of) a bipartite graph
if and only if G does not contain any **odd** cycles
(that is, if and only if none of the subgraphs of G is an odd cycle).

Justification. To show the converse, consider a graph G that does NOT contain any odd cycles.

– Note that it suffices to show that each connected component of G is a bipartite graph (attention: not necessarily complete), **therefore we can assume that G is connected** (because otherwise we focus our attention on one connected component of G at a time).

Fix a vertex v_0 of G , and ‘colour’ it blue. Since G is connected, if we consider another vertex w of G , we can find at least one $v_0 - w$ path in G .

Idea: For each vertex w of G , choose a $v_0 - w$ path in G which has **the shortest possible length**. Denote this length by ℓ_w .

- If ℓ_w is an even integer, then ‘colour’ the vertex w blue.
- If ℓ_w is an odd integer, then ‘colour’ the vertex w red.

Criterion for bipartite graphs

A graph G is (a subgraph of) a bipartite graph
if and only if G does not contain any **odd** cycles
(that is, if and only if none of the subgraphs of G is an odd cycle).

Justification. To show the converse, consider a graph G that does NOT contain any odd cycles.

– Note that it suffices to show that each connected component of G is a bipartite graph (attention: not necessarily complete), **therefore we can assume that G is connected** (because otherwise we focus our attention on one connected component of G at a time).

Fix a vertex v_0 of G , and ‘colour’ it blue. Since G is connected, if we consider another vertex w of G , we can find at least one $v_0 - w$ path in G .

Idea: For each vertex w of G , choose a $v_0 - w$ path in G which has **the shortest possible length**. Denote this length by ℓ_w .

- If ℓ_w is an even integer, then ‘colour’ the vertex w blue.
- If ℓ_w is an odd integer, then ‘colour’ the vertex w red.

Claim. The blue vertices of G form one part, and the red vertices another part. There is NO edge in G joining two blue vertices, and also NO edge joining two red vertices, so G can be viewed as a bipartite graph.

Proof of the claim. Suppose that two (different) blue vertices w_1, w_2 in G were joined.

Proof of the claim. Suppose that two (different) blue vertices w_1, w_2 in G were joined. Note first of all that v_0 cannot be any of these vertices (say, it cannot be vertex w_1), because then we would have that $\ell_{w_2} = 1$, and thus w_2 would have been 'coloured' red.

Proof of the claim. Suppose that two (different) blue vertices w_1, w_2 in G were joined. Note first of all that v_0 cannot be any of these vertices (say, it cannot be vertex w_1), because then we would have that $\ell_{w_2} = 1$, and thus w_2 would have been 'coloured' red. Find a $v_0 - w_1$ path P_1 in G which has length ℓ_{w_1} (that is, the smallest possible), and similarly find a $v_0 - w_2$ path P_2 in G which has length ℓ_{w_2} . **Recall that both ℓ_{w_1} and ℓ_{w_2} are even.**

Proof of the claim. Suppose that two (different) blue vertices w_1, w_2 in G were joined. Note first of all that v_0 cannot be any of these vertices (say, it cannot be vertex w_1), because then we would have that $\ell_{w_2} = 1$, and thus w_2 would have been 'coloured' red.

Find a $v_0 - w_1$ path P_1 in G which has length ℓ_{w_1} (that is, the smallest possible), and similarly find a $v_0 - w_2$ path P_2 in G which has length ℓ_{w_2} . **Recall that both ℓ_{w_1} and ℓ_{w_2} are even.**

Clearly, P_1 and P_2 have common vertices (definitely the vertex v_0). Let z_0 be the 'last' vertex of P_1 which also appears in P_2 , that is, the one that is closest to w_1 .

Proof of the claim. Suppose that two (different) blue vertices w_1, w_2 in G were joined. Note first of all that v_0 cannot be any of these vertices (say, it cannot be vertex w_1), because then we would have that $\ell_{w_2} = 1$, and thus w_2 would have been 'coloured' red.

Find a $v_0 - w_1$ path P_1 in G which has length ℓ_{w_1} (that is, the smallest possible), and similarly find a $v_0 - w_2$ path P_2 in G which has length ℓ_{w_2} . **Recall that both ℓ_{w_1} and ℓ_{w_2} are even.**

Clearly, P_1 and P_2 have common vertices (definitely the vertex v_0). Let z_0 be the 'last' vertex of P_1 which also appears in P_2 , that is, the one that is closest to w_1 .

Important Observation. Let Q_1 be the $v_0 - z_0$ subpath of P_1 , and let Q_2 be the $v_0 - z_0$ subpath of P_2 . Then both Q_1 and Q_2 have the same length, which is equal to ℓ_{z_0} (indeed, these are $v_0 - z_0$ paths in G , so they cannot have length $< \ell_{z_0}$; also, if e.g. Q_1 had length $> \ell_{z_0}$, then we could replace Q_1 by a $v_0 - z_0$ path which has shortest possible length, and this would give us a shorter $v_0 - w_1$ path than P_1 , contradicting our choice of P_1).

Proof of the claim. Suppose that two (different) blue vertices w_1, w_2 in G were joined. Note first of all that v_0 cannot be any of these vertices (say, it cannot be vertex w_1), because then we would have that $\ell_{w_2} = 1$, and thus w_2 would have been 'coloured' red.

Find a $v_0 - w_1$ path P_1 in G which has length ℓ_{w_1} (that is, the smallest possible), and similarly find a $v_0 - w_2$ path P_2 in G which has length ℓ_{w_2} . Recall that both ℓ_{w_1} and ℓ_{w_2} are even.

Clearly, P_1 and P_2 have common vertices (definitely the vertex v_0). Let z_0 be the 'last' vertex of P_1 which also appears in P_2 , that is, the one that is closest to w_1 .

Important Observation. Let Q_1 be the $v_0 - z_0$ subpath of P_1 , and let Q_2 be the $v_0 - z_0$ subpath of P_2 . Then both Q_1 and Q_2 have the same length, which is equal to ℓ_{z_0} (indeed, these are $v_0 - z_0$ paths in G , so they cannot have length $< \ell_{z_0}$; also, if e.g. Q_1 had length $> \ell_{z_0}$, then we could replace Q_1 by a $v_0 - z_0$ path which has shortest possible length, and this would give us a shorter $v_0 - w_1$ path than P_1 , contradicting our choice of P_1).

Thus, if we set Q'_1 to be the $z_0 - w_1$ subpath of P_1 (that is, the remaining part of P_1 after 'deleting' Q_1 except for the vertex z_0), then Q'_1 has length $\ell_{w_1} - \ell_{z_0}$. Similarly, if we set Q'_2 to be the $z_0 - w_2$ subpath of P_2 , then Q'_2 has length $\ell_{w_2} - \ell_{z_0}$.

- Note that either both $\ell_{w_1} - \ell_{z_0}$ and $\ell_{w_2} - \ell_{z_0}$ are even, or both of them are odd.

Proof of the claim. Suppose that two (different) blue vertices w_1, w_2 in G were joined. Note first of all that v_0 cannot be any of these vertices (say, it cannot be vertex w_1), because then we would have that $\ell_{w_2} = 1$, and thus w_2 would have been 'coloured' red.

Find a $v_0 - w_1$ path P_1 in G which has length ℓ_{w_1} (that is, the smallest possible), and similarly find a $v_0 - w_2$ path P_2 in G which has length ℓ_{w_2} . Recall that both ℓ_{w_1} and ℓ_{w_2} are even.

Clearly, P_1 and P_2 have common vertices (definitely the vertex v_0). Let z_0 be the 'last' vertex of P_1 which also appears in P_2 , that is, the one that is closest to w_1 .

Important Observation. Let Q_1 be the $v_0 - z_0$ subpath of P_1 , and let Q_2 be the $v_0 - z_0$ subpath of P_2 . Then both Q_1 and Q_2 have the same length, which is equal to ℓ_{z_0} (indeed, these are $v_0 - z_0$ paths in G , so they cannot have length $< \ell_{z_0}$; also, if e.g. Q_1 had length $> \ell_{z_0}$, then we could replace Q_1 by a $v_0 - z_0$ path which has shortest possible length, and this would give us a shorter $v_0 - w_1$ path than P_1 , contradicting our choice of P_1).

Thus, if we set Q'_1 to be the $z_0 - w_1$ subpath of P_1 (that is, the remaining part of P_1 after 'deleting' Q_1 except for the vertex z_0), then Q'_1 has length $\ell_{w_1} - \ell_{z_0}$. Similarly, if we set Q'_2 to be the $z_0 - w_2$ subpath of P_2 , then Q'_2 has length $\ell_{w_2} - \ell_{z_0}$.

- Note that either both $\ell_{w_1} - \ell_{z_0}$ and $\ell_{w_2} - \ell_{z_0}$ are even, or both of them are odd.
- Moreover, Q'_1 and Q'_2 don't have any common vertices anymore, except for their initial vertex z_0 .

Proof of the claim. Suppose that two (different) blue vertices w_1, w_2 in G were joined. Note first of all that v_0 cannot be any of these vertices (say, it cannot be vertex w_1), because then we would have that $\ell_{w_2} = 1$, and thus w_2 would have been 'coloured' red.

Find a $v_0 - w_1$ path P_1 in G which has length ℓ_{w_1} (that is, the smallest possible), and similarly find a $v_0 - w_2$ path P_2 in G which has length ℓ_{w_2} . Recall that both ℓ_{w_1} and ℓ_{w_2} are even.

Clearly, P_1 and P_2 have common vertices (definitely the vertex v_0). Let z_0 be the 'last' vertex of P_1 which also appears in P_2 , that is, the one that is closest to w_1 .

Important Observation. Let Q_1 be the $v_0 - z_0$ subpath of P_1 , and let Q_2 be the $v_0 - z_0$ subpath of P_2 . Then both Q_1 and Q_2 have the same length, which is equal to ℓ_{z_0} (indeed, these are $v_0 - z_0$ paths in G , so they cannot have length $< \ell_{z_0}$; also, if e.g. Q_1 had length $> \ell_{z_0}$, then we could replace Q_1 by a $v_0 - z_0$ path which has shortest possible length, and this would give us a shorter $v_0 - w_1$ path than P_1 , contradicting our choice of P_1).

Thus, if we set Q'_1 to be the $z_0 - w_1$ subpath of P_1 (that is, the remaining part of P_1 after 'deleting' Q_1 except for the vertex z_0), then Q'_1 has length $\ell_{w_1} - \ell_{z_0}$. Similarly, if we set Q'_2 to be the $z_0 - w_2$ subpath of P_2 , then Q'_2 has length $\ell_{w_2} - \ell_{z_0}$.

- Note that either both $\ell_{w_1} - \ell_{z_0}$ and $\ell_{w_2} - \ell_{z_0}$ are even, or both of them are odd.
- Moreover, Q'_1 and Q'_2 don't have any common vertices anymore, except for their initial vertex z_0 .

We can thus form a cycle C_0 in G by traversing Q'_1 , then the edge $w_1 w_2$, and then Q'_2 in the reverse direction (that is, going from w_2 to z_0).

Proof of the claim. Suppose that two (different) blue vertices w_1, w_2 in G were joined. Note first of all that v_0 cannot be any of these vertices (say, it cannot be vertex w_1), because then we would have that $\ell_{w_2} = 1$, and thus w_2 would have been 'coloured' red.

Find a $v_0 - w_1$ path P_1 in G which has length ℓ_{w_1} (that is, the smallest possible), and similarly find a $v_0 - w_2$ path P_2 in G which has length ℓ_{w_2} . Recall that both ℓ_{w_1} and ℓ_{w_2} are even.

Clearly, P_1 and P_2 have common vertices (definitely the vertex v_0). Let z_0 be the 'last' vertex of P_1 which also appears in P_2 , that is, the one that is closest to w_1 .

Important Observation. Let Q_1 be the $v_0 - z_0$ subpath of P_1 , and let Q_2 be the $v_0 - z_0$ subpath of P_2 . Then both Q_1 and Q_2 have the same length, which is equal to ℓ_{z_0} (indeed, these are $v_0 - z_0$ paths in G , so they cannot have length $< \ell_{z_0}$; also, if e.g. Q_1 had length $> \ell_{z_0}$, then we could replace Q_1 by a $v_0 - z_0$ path which has shortest possible length, and this would give us a shorter $v_0 - w_1$ path than P_1 , contradicting our choice of P_1).

Thus, if we set Q'_1 to be the $z_0 - w_1$ subpath of P_1 (that is, the remaining part of P_1 after 'deleting' Q_1 except for the vertex z_0), then Q'_1 has length $\ell_{w_1} - \ell_{z_0}$. Similarly, if we set Q'_2 to be the $z_0 - w_2$ subpath of P_2 , then Q'_2 has length $\ell_{w_2} - \ell_{z_0}$.

- Note that either both $\ell_{w_1} - \ell_{z_0}$ and $\ell_{w_2} - \ell_{z_0}$ are even, or both of them are odd.
- Moreover, Q'_1 and Q'_2 don't have any common vertices anymore, except for their initial vertex z_0 .

We can thus form a cycle C_0 in G by traversing Q'_1 , then the edge $w_1 w_2$, and then Q'_2 in the reverse direction (that is, going from w_2 to z_0). C_0 has size (that is, number of edges) equal to

$$\ell_{w_1} - \ell_{z_0} + 1 + \ell_{w_2} - \ell_{z_0}.$$

This is an odd number, contradicting the assumption that G contains NO odd cycles. Thus the 'blue' vertices w_1, w_2 cannot be joined by an edge.

Proof of the claim. Suppose that two (different) blue vertices w_1, w_2 in G were joined. Note first of all that v_0 cannot be any of these vertices (say, it cannot be vertex w_1), because then we would have that $\ell_{w_2} = 1$, and thus w_2 would have been 'coloured' red.

Find a $v_0 - w_1$ path P_1 in G which has length ℓ_{w_1} (that is, the smallest possible), and similarly find a $v_0 - w_2$ path P_2 in G which has length ℓ_{w_2} . Recall that both ℓ_{w_1} and ℓ_{w_2} are even.

Clearly, P_1 and P_2 have common vertices (definitely the vertex v_0). Let z_0 be the 'last' vertex of P_1 which also appears in P_2 , that is, the one that is closest to w_1 .

Important Observation. Let Q_1 be the $v_0 - z_0$ subpath of P_1 , and let Q_2 be the $v_0 - z_0$ subpath of P_2 . Then both Q_1 and Q_2 have the same length, which is equal to ℓ_{z_0} (indeed, these are $v_0 - z_0$ paths in G , so they cannot have length $< \ell_{z_0}$; also, if e.g. Q_1 had length $> \ell_{z_0}$, then we could replace Q_1 by a $v_0 - z_0$ path which has shortest possible length, and this would give us a shorter $v_0 - w_1$ path than P_1 , contradicting our choice of P_1).

Thus, if we set Q'_1 to be the $z_0 - w_1$ subpath of P_1 (that is, the remaining part of P_1 after 'deleting' Q_1 except for the vertex z_0), then Q'_1 has length $\ell_{w_1} - \ell_{z_0}$. Similarly, if we set Q'_2 to be the $z_0 - w_2$ subpath of P_2 , then Q'_2 has length $\ell_{w_2} - \ell_{z_0}$.

- Note that either both $\ell_{w_1} - \ell_{z_0}$ and $\ell_{w_2} - \ell_{z_0}$ are even, or both of them are odd.
- Moreover, Q'_1 and Q'_2 don't have any common vertices anymore, except for their initial vertex z_0 .

We can thus form a cycle C_0 in G by traversing Q'_1 , then the edge $w_1 w_2$, and then Q'_2 in the reverse direction (that is, going from w_2 to z_0). C_0 has size (that is, number of edges) equal to

$$\ell_{w_1} - \ell_{z_0} + 1 + \ell_{w_2} - \ell_{z_0}.$$

This is an odd number, contradicting the assumption that G contains NO odd cycles.

Thus the 'blue' vertices w_1, w_2 cannot be joined by an edge.


Similarly, we can show that two 'red' vertices cannot be joined.

A criterion for line graphs

A criterion for line graphs

Theorem (Beineke, 1968)

A graph G is the *line graph* of some other graph H **if and only if** the following 9 graphs are forbidden subgraphs for G (in other words, if and only if **none** of the following graphs is an induced subgraph of G).



from the Balakrishnan-Ranganathan book

Back to Connectivity

Definitions

Let G be a connected graph.

- 1 A vertex v of G is called a cutvertex of G if we have that

$$G - v$$

is no longer connected.

- 2 An edge e in G is called a bridge (or a cutedge) of G if we have that

$$G - e$$

is no longer connected.

Definitions

Let G be a connected graph.

- 1 A vertex v of G is called a cutvertex of G if we have that

$$G - v$$

is no longer connected.

- 2 An edge e in G is called a bridge (or a cutedge) of G if we have that

$$G - e$$

is no longer connected.

Definition again, more generally formulated

If we start with a graph H which is not necessarily connected, then

- 1 a vertex \tilde{v} of H is called a cutvertex of H if, by deleting \tilde{v} (and of course all the edges \tilde{v} is incident with), we increase the number of connected components of H .
- 2 an edge \tilde{e} in H is called a bridge of H if, by deleting \tilde{e} , we increase the number of connected components of H .

Connectivity (cont.)

Definitions

Let $G = (V, E)$ be a connected graph.

- 1 A subset V' of the vertex set V of G is called a vertex cut if we have that

$$G - V'$$

is disconnected. We call it a k -vertex cut if the cardinality $|V'|$ of V' is equal to k (that is, if V' contains k vertices of G).

V' is also called a separating set of vertices of G .

Connectivity (cont.)

Definitions

Let $G = (V, E)$ be a connected graph.

- 1 A subset V' of the vertex set V of G is called a vertex cut if we have that

$$G - V'$$

is disconnected. We call it a k -vertex cut if the cardinality $|V'|$ of V' is equal to k (that is, if V' contains k vertices of G).

V' is also called a *separating set of vertices* of G .

- 2 A subset E' of the edge set E of G is called an edge cut if we have that

$$G - E'$$

is disconnected. We call it a k -edge cut if $|E'| = k$.

E' is also called a *separating set of edges* of G .

The parameters of Vertex and Edge Connectivity

**The parameters of
Vertex and Edge Connectivity**




Edge Connectivity

The parameter $\lambda(G)$

Let G be a connected graph of order ≥ 2 . We define the edge connectivity $\lambda(G)$ of G to be the minimum cardinality of an edge cut of G .

Testing our understanding on examples

Question. What is $\lambda(G)$ for each of the following graphs G ?



A “flower” graph

A very important and useful observation

Part of an important theorem we will state shortly

Let G be a connected graph of order ≥ 2 (which implies that G has at least one edge).

We have that: $\lambda(G) \leq \delta(G)$.

A very important and useful observation

Part of an important theorem we will state shortly

Let G be a connected graph of order ≥ 2 (which implies that G has at least one edge).

We have that: $\lambda(G) \leq \delta(G)$.

Justification. Let v_0 be a vertex in G which has minimum degree, that is, $\deg_G(v_0) = \delta(G)$. Denote this degree by d_0 .

A very important and useful observation

Part of an important theorem we will state shortly

Let G be a connected graph of order ≥ 2 (which implies that G has at least one edge).

We have that: $\lambda(G) \leq \delta(G)$.

Justification. Let v_0 be a vertex in G which has minimum degree, that is, $\deg_G(v_0) = \delta(G)$. Denote this degree by d_0 .

Then v_0 has d_0 neighbours in G , say the vertices w_1, w_2, \dots, w_{d_0} , and is incident with exactly d_0 edges of G , the edges

$$e_1 = v_0 w_1, e_2 = v_0 w_2, \dots, e_{d_0} = v_0 w_{d_0}.$$

A very important and useful observation

Part of an important theorem we will state shortly

Let G be a connected graph of order ≥ 2 (which implies that G has at least one edge).

We have that: $\lambda(G) \leq \delta(G)$.

Justification. Let v_0 be a vertex in G which has minimum degree, that is, $\deg_G(v_0) = \delta(G)$. Denote this degree by d_0 .

Then v_0 has d_0 neighbours in G , say the vertices w_1, w_2, \dots, w_{d_0} , and is incident with exactly d_0 edges of G , the edges

$$e_1 = v_0 w_1, e_2 = v_0 w_2, \dots, e_{d_0} = v_0 w_{d_0}.$$


If we delete the edges e_1, e_2, \dots, e_{d_0} , then v_0 is an isolated vertex in the resulting subgraph $G - \{e_1, e_2, \dots, e_{d_0}\}$ (which still has all the other vertices too), and thus $G - \{e_1, e_2, \dots, e_{d_0}\}$ is disconnected.

This shows that $\{e_1, e_2, \dots, e_{d_0}\}$ is an edge cut of G , and since it has cardinality $d_0 = \delta(G)$, we must have

$$\lambda(G) := \text{minimum cardinality of an edge cut of } G \leq \delta(G).$$

Examples (cont.)

Question. What is the edge connectivity of a complete graph?
That is, what is $\lambda(K_n)$?



Complete graph on vertices $\{1, 2, 3, 4, 5, 6, 7\}$

An equivalent way of thinking about the last question

Given that every graph on n vertices is a subgraph of the complete graph K_n , and given that we are asking for the minimum number of edges that we would have to remove from K_n in order to get a disconnected graph, we could ask this question in a 'complementary' way too:

What is the maximum size (that is, largest number of edges) of a disconnected subgraph of K_n which contains all n vertices?

Or more simply, what is the maximum size of a disconnected graph on n vertices?

An equivalent way of thinking about the last question

Given that every graph on n vertices is a subgraph of the complete graph K_n , and given that we are asking for the minimum number of edges that we would have to remove from K_n in order to get a disconnected graph, we could ask this question in a 'complementary' way too:

What is the maximum size (that is, largest number of edges) of a disconnected subgraph of K_n which contains all n vertices?

Or more simply, what is the maximum size of a disconnected graph on n vertices?

How does the answer to the above question relate to $\lambda(K_n)$?

An equivalent way of thinking about the last question

Given that every graph on n vertices is a subgraph of the complete graph K_n , and given that we are asking for the minimum number of edges that we would have to remove from K_n in order to get a disconnected graph, we could ask this question in a 'complementary' way too:

What is the maximum size (that is, largest number of edges) of a disconnected subgraph of K_n which contains all n vertices?

Or more simply, what is the maximum size of a disconnected graph on n vertices?

How does the answer to the above question relate to $\lambda(K_n)$?

It's worth comparing this question with the 'opposite' question we discussed in the last lectures, about **what the minimum size of a connected graph H on n vertices is**. Recall that we have found this minimum size to be $n - 1$ (because any graph with size $< n - 1$ will be disconnected, whereas paths (or any other trees) on n vertices are connected (by definition) and have precisely $n - 1$ edges as we showed).

Answer to our two 'complementary' questions

What is the maximum size (that is, largest number of edges) of a disconnected subgraph of K_n which contains all n vertices?

Or more simply, what is the maximum size of a disconnected graph on n vertices?

Also, how does the answer to the above question relate to $\lambda(K_n)$?

Answer to our two ‘complementary’ questions

What is the maximum size (that is, largest number of edges) of a disconnected subgraph of K_n which contains all n vertices?

Or more simply, what is the maximum size of a disconnected graph on n vertices?

Also, how does the answer to the above question relate to $\lambda(K_n)$?

Answer. We know that the size $e(K_n)$ of K_n is

$$\binom{n}{2} = \frac{n(n-1)}{2}$$

(since the edge set of K_n contains all 2-element subsets of the vertex set of K_n); if e_{\max} is the maximum size of a disconnected subgraph of K_n which contains all n vertices, then

$$\lambda(K_n) = \binom{n}{2} - e_{\max}.$$

Maximum size of a disconnected graph on n vertices

Consider a graph G of order n which is disconnected and has the maximum possible number of edges. We can make the following observations:

Maximum size of a disconnected graph on n vertices

Consider a graph G of order n which is disconnected and has the maximum possible number of edges. We can make the following observations:

- G must have exactly two connected components.

Maximum size of a disconnected graph on n vertices

Consider a graph G of order n which is disconnected and has the maximum possible number of edges. We can make the following observations:

- G must have exactly two connected components. Indeed, if G had at least three connected components, and say G_1, G_2, G_3 were three of these components, then no vertices from G_2 would be joined with vertices from G_3 ; however if we started adding edges which join such vertices, we would increase the number of edges in G , while G would still be disconnected (since none of the vertices in G_1 would be joined with a vertex in G_2 or G_3); but we have assumed that G has the maximum possible number of edges among disconnected graphs, so we shouldn't be able to add any more edges to it without it becoming connected.

Maximum size of a disconnected graph on n vertices

Consider a graph G of order n which is disconnected and has the maximum possible number of edges. We can make the following observations:

- G must have exactly two connected components. Indeed, if G had at least three connected components, and say G_1, G_2, G_3 were three of these components, then no vertices from G_2 would be joined with vertices from G_3 ; however if we started adding edges which join such vertices, we would increase the number of edges in G , while G would still be disconnected (since none of the vertices in G_1 would be joined with a vertex in G_2 or G_3); but we have assumed that G has the maximum possible number of edges among disconnected graphs, so we shouldn't be able to add any more edges to it without it becoming connected.
- If G_1 and G_2 are the two connected components of G , then every vertex in G_1 has to be joined with every other vertex in G_1 , and similarly for the vertices in G_2 .

Maximum size of a disconnected graph on n vertices

Consider a graph G of order n which is disconnected and has the maximum possible number of edges. We can make the following observations:

- G must have **exactly two** connected components. Indeed, if G had at least three connected components, and say G_1, G_2, G_3 were three of these components, then no vertices from G_2 would be joined with vertices from G_3 ; however if we started adding edges which join such vertices, we would increase the number of edges in G , while G would still be disconnected (since none of the vertices in G_1 would be joined with a vertex in G_2 or G_3); but we have assumed that G has the maximum possible number of edges among disconnected graphs, so we shouldn't be able to add any more edges to it without it becoming connected.
- If G_1 and G_2 are the two connected components of G , then every vertex in G_1 has to be joined with every other vertex in G_1 , and similarly for the vertices in G_2 . Indeed, if we could find two vertices in G_1 (or two vertices in G_2) which are not joined already by an edge, we could add that edge and hence increase the number of edges in G , without ending up with a connected graph; again this would contradict the assumption that G has the maximum possible number of edges among disconnected graphs.

Maximum size of a disconnected graph on n vertices

Consider a graph G of order n which is disconnected and has the maximum possible number of edges. We can make the following observations:

- G must have exactly two connected components. Indeed, if G had at least three connected components, and say G_1, G_2, G_3 were three of these components, then no vertices from G_2 would be joined with vertices from G_3 ; however if we started adding edges which join such vertices, we would increase the number of edges in G , while G would still be disconnected (since none of the vertices in G_1 would be joined with a vertex in G_2 or G_3); but we have assumed that G has the maximum possible number of edges among disconnected graphs, so we shouldn't be able to add any more edges to it without it becoming connected.
- If G_1 and G_2 are the two connected components of G , then every vertex in G_1 has to be joined with every other vertex in G_1 , and similarly for the vertices in G_2 . Indeed, if we could find two vertices in G_1 (or two vertices in G_2) which are not joined already by an edge, we could add that edge and hence increase the number of edges in G , without ending up with a connected graph; again this would contradict the assumption that G has the maximum possible number of edges among disconnected graphs.

Based on these, we see that G is the disjoint union of two complete graphs:

$$G = K_k \oplus K_{n-k}$$

for some $1 \leq k \leq n-1$.

Maximum size of a disconnected graph on n vertices (cont.)

We have already confirmed that

$$G = K_k \oplus K_{n-k}$$

for some $1 \leq k \leq n - 1$.

Maximum size of a disconnected graph on n vertices (cont.)

We have already confirmed that

$$G = K_k \oplus K_{n-k}$$

for some $1 \leq k \leq n-1$.

But then

$$e(G) = \binom{k}{2} + \binom{n-k}{2} = \frac{k(k-1)}{2} + \frac{(n-k)(n-k-1)}{2}.$$

Maximum size of a disconnected graph on n vertices (cont.)

We have already confirmed that

$$G = K_k \oplus K_{n-k}$$

for some $1 \leq k \leq n-1$.

But then

$$e(G) = \binom{k}{2} + \binom{n-k}{2} = \frac{k(k-1)}{2} + \frac{(n-k)(n-k-1)}{2}.$$

It remains to check that, when we only consider $k \in \{1, 2, \dots, n-2, n-1\}$, this expression for $e(G)$ takes its maximum value when $k = 1$ or $k = n-1$ (in fact, the function $k \mapsto \binom{k}{2} + \binom{n-k}{2}$ is decreasing for $1 \leq k \leq \lfloor \frac{n}{2} \rfloor$, and then it becomes increasing).

Maximum size of a disconnected graph on n vertices (cont.)

We have already confirmed that

$$G = K_k \oplus K_{n-k}$$

for some $1 \leq k \leq n-1$.

But then

$$e(G) = \binom{k}{2} + \binom{n-k}{2} = \frac{k(k-1)}{2} + \frac{(n-k)(n-k-1)}{2}.$$

It remains to check that, when we only consider $k \in \{1, 2, \dots, n-2, n-1\}$, this expression for $e(G)$ takes its **maximum value when $k=1$ or $k=n-1$** (in fact, the function $k \mapsto \binom{k}{2} + \binom{n-k}{2}$ is decreasing for $1 \leq k \leq \lfloor \frac{n}{2} \rfloor$, and then it becomes increasing).

Plugging $k = n-1$ above, we obtain that

$$e(G) = \binom{n-1}{2}$$

is the maximum possible size of a disconnected graph on n vertices. In fact, we have also found that such a graph should be **the disjoint union of a complete graph on $n-1$ vertices and an isolated vertex**.

Maximum size of a disconnected graph on n vertices (cont.)

We have already confirmed that

$$G = K_k \oplus K_{n-k}$$

for some $1 \leq k \leq n-1$.

But then

$$e(G) = \binom{k}{2} + \binom{n-k}{2} = \frac{k(k-1)}{2} + \frac{(n-k)(n-k-1)}{2}.$$

It remains to check that, when we only consider $k \in \{1, 2, \dots, n-2, n-1\}$, this expression for $e(G)$ takes its **maximum value when $k=1$ or $k=n-1$** (in fact, the function $k \mapsto \binom{k}{2} + \binom{n-k}{2}$ is decreasing for $1 \leq k \leq \lfloor \frac{n}{2} \rfloor$, and then it becomes increasing).

Plugging $k = n-1$ above, we obtain that

$$e(G) = \binom{n-1}{2}$$

is the maximum possible size of a disconnected graph on n vertices. In fact, we have also found that such a graph should be **the disjoint union of a complete graph on $n-1$ vertices and an isolated vertex**.

We also conclude that $\lambda(K_n) = e(K_n) - \binom{n-1}{2} = \binom{n}{2} - \binom{n-1}{2} = n-1$.

Conclusions from all related discussions so far

- The maximum size of a connected graph on n vertices is $\binom{n}{2}$.

Question 1. Which connected graphs of order n have this size?

Conclusions from all related discussions so far

- The maximum size of a connected graph on n vertices is $\binom{n}{2}$.

Question 1. Which connected graphs of order n have this size? Only the complete graph K_n has this size.

Conclusions from all related discussions so far

- The maximum size of a connected graph on n vertices is $\binom{n}{2}$.

Question 1. Which connected graphs of order n have this size? Only the complete graph K_n has this size.

- The minimum size of a connected graph on n vertices is $n - 1$.

Question 2. Which connected graphs of order n have this size?

Conclusions from all related discussions so far

- The maximum size of a connected graph on n vertices is $\binom{n}{2}$.

Question 1. Which connected graphs of order n have this size? Only the complete graph K_n has this size.

- The minimum size of a connected graph on n vertices is $n - 1$.

Question 2. Which connected graphs of order n have this size? We have seen that only trees on n vertices have this size.

Conclusions from all related discussions so far

- The maximum size of a connected graph on n vertices is $\binom{n}{2}$.

Question 1. Which connected graphs of order n have this size? Only the complete graph K_n has this size.

- The minimum size of a connected graph on n vertices is $n - 1$.

Question 2. Which connected graphs of order n have this size? We have seen that only trees on n vertices have this size.

- The maximum size of a disconnected graph on n vertices is $\binom{n}{2} - (n-1) = \binom{n-1}{2}$.

Conclusions from all related discussions so far

- The maximum size of a connected graph on n vertices is $\binom{n}{2}$.

Question 1. Which connected graphs of order n have this size? Only the complete graph K_n has this size.

- The minimum size of a connected graph on n vertices is $n - 1$.

Question 2. Which connected graphs of order n have this size? We have seen that only trees on n vertices have this size.

- The maximum size of a disconnected graph on n vertices is $\binom{n}{2} - (n-1) = \binom{n-1}{2}$.

- The minimum size of a disconnected graph on n vertices is 0 (and the only example here is the Null Graph N_n).

Conclusions from all related discussions so far

- The maximum size of a connected graph on n vertices is $\binom{n}{2}$.

Question 1. Which connected graphs of order n have this size? Only the complete graph K_n has this size.

- The minimum size of a connected graph on n vertices is $n - 1$.

Question 2. Which connected graphs of order n have this size? We have seen that only trees on n vertices have this size.

- The maximum size of a disconnected graph on n vertices is $\binom{n}{2} - (n-1) = \binom{n-1}{2}$.

- The minimum size of a disconnected graph on n vertices is 0 (and the only example here is the Null Graph N_n).

More generally, we can show...

Theorem

Let G be a graph of order n which has exactly k connected components (where $1 \leq k \leq n$).

Then the maximum possible size of G is $\binom{n-k+1}{2}$, and the minimum possible size is $n - k$.

See video lecture 5.2 (Edge estimates) from Dr. Seidon Alsaody's list.

One more conclusion

Let G be a connected graph of order n that is different from K_n .

One more conclusion

Let G be a connected graph of order n that is different from K_n .
Then $\lambda(G) < n - 1$ (why?).

One more conclusion

Let G be a connected graph of order n that is different from K_n .
Then $\lambda(G) < n - 1$ (why?).

We can reach this conclusion by relying on the useful observation we stated earlier: we have that

- if G is not the complete graph on n vertices, then G has at least one vertex which is not connected to every other vertex, and hence $\delta(G) < n - 1$.

One more conclusion


Let G be a connected graph of order n that is different from K_n .
Then $\lambda(G) < n - 1$ (why?).

We can reach this conclusion by relying on the useful observation we stated earlier: we have that

- if G is not the complete graph on n vertices, then G has at least one vertex which is not connected to every other vertex, and hence $\delta(G) < n - 1$.
- By the observation, we get that $\lambda(G) \leq \delta(G) < n - 1$.

The other extreme


Question. Consider a tree T on n vertices. What is $\lambda(T)$? Does it depend on which tree on n vertices we start with?



Trees of order 6 or less;
from the Harris-Hirst-Mossinghoff book

The other extreme

Question. Consider a tree T on n vertices. What is $\lambda(T)$? Does it depend on which tree on n vertices we start with?




Trees of order 6 or less;
from the Harris-Hirst-Mossinghoff book

We always have $\lambda(T) = 1$ for any tree T with at least two vertices. Moreover, we have that every edge of T is a bridge (or equivalently, a cutedge).

The other extreme

Question. Consider a tree T on n vertices. What is $\lambda(T)$? Does it depend on which tree on n vertices we start with?



Trees of order 6 or less;
from the Harris-Hirst-Mossinghoff book


We always have $\lambda(T) = 1$ for any tree T with at least two vertices. Moreover, we have that every edge of T is a bridge (or equivalently, a cutedge). (this is because, by deleting an arbitrary edge e from T , we are left with a subgraph $T - e$ on n vertices which has $n - 2$ edges)

What about Vertex Connectivity?


What about Vertex Connectivity?

*Completely analogously to edge connectivity, **we would like** to define the parameter of 'vertex connectivity' of a connected graph G to be the minimum cardinality of a vertex cut of G .*

But...
complete graphs don't have vertex cuts



But...
complete graphs don't have vertex cuts




Important Remark. Given $n \geq 2$, we have that, for every (proper) subset V' of the vertex set V of K_n , the graph $K_n - V'$ is again a complete graph (on the vertices $V \setminus V'$ now), and hence it cannot be disconnected.


\leadsto a complete graph does not have any vertex cuts.

In all other cases however...

Any connected graph of order ≥ 2 which is not a complete graph will have vertex cuts. *(For each of the examples below, find a vertex cut; if possible, try to find one with smallest possible cardinality.)*



Graph G_1



Graph G_2

**How do we reconcile these two facts,
in order to give a useful definition?**

**How do we reconcile these two facts,
in order to give a useful definition?**

to be discussed next time