

MATH 322 – Graph Theory

Fall Term 2021

Notes for Lecture 6

Tuesday, September 21

Reminder: Operations on Graphs

- disjoint union of graphs
- complement of a graph
- induced subgraph

Reminder: Operations on Graphs

From last time

- disjoint union of graphs
- complement of a graph
- induced subgraph
- join of graphs

Reminder: Operations on Graphs

From last time

- disjoint union of graphs
- complement of a graph
- induced subgraph
- join of graphs
- line graph of a graph

Reminder: Operations on Graphs

From last time

- disjoint union of graphs
- complement of a graph
- induced subgraph
- join of graphs
- line graph of a graph
- *vertex deletion*

Reminder: Operations on Graphs

From last time

- disjoint union of graphs
- complement of a graph
- induced subgraph
- join of graphs
- line graph of a graph
- vertex deletion
- edge deletion

Reminder: Disjoint Union of Graphs

If $G_1 = (V_1, E_1)$ and $G_2 = (V_2, E_2)$ are two graphs with **disjoint vertex sets**, namely such that

$$V_1 \cap V_2 = \emptyset,$$

then the ordered pair

$$(V_1 \cup V_2, E_1 \cup E_2)$$

is a new graph whose vertices consist of all the vertices of G_1 and all the vertices of G_2 .

Reminder: Disjoint Union of Graphs

If $G_1 = (V_1, E_1)$ and $G_2 = (V_2, E_2)$ are two graphs with disjoint vertex sets, namely such that

$$V_1 \cap V_2 = \emptyset,$$

then the ordered pair

$$(V_1 \cup V_2, E_1 \cup E_2)$$


is a new graph whose vertices consist of all the vertices of G_1 and all the vertices of G_2 .

We denote this new graph by $G_1 \oplus G_2$,
and we call it the disjoint union of G_1 and G_2 .


(Note also that, in the graph $G_1 \oplus G_2$, none of the vertices in V_1 is joined with a vertex in V_2 , and vice versa; this is because $E_1 \subseteq [V_1]^2$ and $E_2 \subseteq [V_2]^2$, so each edge of $G_1 \oplus G_2$ is either an unordered pair of elements of V_1 , or an unordered pair of elements of V_2 .)

Examples

The graphs below can be viewed as disjoint unions of their connected components.



Disjoint union of 3 graphs



Disjoint union of two 3-cycles

Reminder: Complement of a Graph

Let $G = (V, E)$ be a graph. Recall that E is a subset of the set of 2-element subsets of V (sometimes we denote this set by $[V]^2$).

We can construct a new graph H on the same set of vertices by setting

$$H = (V, [V]^2 \setminus E),$$

that is, by setting the edge set of H to be the **complement** of E in $[V]^2$.

In essence, what we are doing is removing any edges/‘connections’ we have in G , and then we are joining any two vertices that were not joined in G .

Reminder: Complement of a Graph

Let $G = (V, E)$ be a graph. Recall that E is a subset of the set of 2-element subsets of V (sometimes we denote this set by $[V]^2$).

We can construct a new graph H on the same set of vertices by setting

$$H = (V, [V]^2 \setminus E),$$

that is, by setting the edge set of H to be the **complement** of E in $[V]^2$.

In essence, what we are doing is removing any edges/‘connections’ we have in G , and then we are joining any two vertices that were not joined in G .

Definition. The new graph is called the complement of G , and is denoted by \overline{G} .

Reminder: Subgraphs and Induced Subgraphs

Let $G = (V, E)$ be a graph.

Definition. A subgraph H of G is an ordered pair (V', E')

- where $\emptyset \neq V' \subseteq V$ (that is, V' is a non-empty subset of V),
- and where $E' \subseteq E$ with every edge $e \in E'$ having both endvertices in V' .

In this case, we write $H \subseteq G$.

Reminder: Subgraphs and Induced Subgraphs

Let $G = (V, E)$ be a graph.

Definition. A subgraph H of G is an ordered pair (V', E')

- where $\emptyset \neq V' \subseteq V$ (that is, V' is a non-empty subset of V),
- and where $E' \subseteq E$ with every edge $e \in E'$ having both endvertices in V' .

In this case, we write $H \subseteq G$.

Definition. If $H = (V', E')$ is a subgraph of G , and E' contains **all the edges of G which have both endvertices in V'** , then we say that H is the subgraph of G that is induced or spanned by V' .

We denote this induced subgraph by $G[V']$.

Reminder: Join of Two Graphs

Consider two graphs $G_1 = (V_1, E_1)$ and $G_2 = (V_2, E_2)$ with disjoint vertex sets, namely such that

$$V_1 \cap V_2 = \emptyset.$$

Reminder: Join of Two Graphs

Consider two graphs $G_1 = (V_1, E_1)$ and $G_2 = (V_2, E_2)$ with disjoint vertex sets, namely such that

$$V_1 \cap V_2 = \emptyset.$$

The join of G_1 and G_2 is a new graph with vertex set

$$V_1 \cup V_2$$

and edge set

$$E_1 \cup E_2 \cup \{\{v, w\} : v \in V_1, w \in V_2\}.$$

We denote the join of G_1 and G_2 by $G_1 \vee G_2$.

Reminder: Join of Two Graphs

Consider two graphs $G_1 = (V_1, E_1)$ and $G_2 = (V_2, E_2)$ with disjoint vertex sets, namely such that

$$V_1 \cap V_2 = \emptyset.$$

The join of G_1 and G_2 is a new graph with vertex set

$$V_1 \cup V_2$$

and edge set

$$E_1 \cup E_2 \cup \{\{v, w\} : v \in V_1, w \in V_2\}.$$

We denote the join of G_1 and G_2 by $G_1 \vee G_2$.

Observe that the vertices of $G_1 \vee G_2$ consist of all the vertices of G_1 and of G_2 , and two such vertices v, w are connected in $G_1 \vee G_2$ exactly when:

Reminder: Join of Two Graphs

Consider two graphs $G_1 = (V_1, E_1)$ and $G_2 = (V_2, E_2)$ with disjoint vertex sets, namely such that

$$V_1 \cap V_2 = \emptyset.$$

The join of G_1 and G_2 is a new graph with vertex set

$$V_1 \cup V_2$$

and edge set

$$E_1 \cup E_2 \cup \{\{v, w\} : v \in V_1, w \in V_2\}.$$

We denote the join of G_1 and G_2 by $G_1 \vee G_2$.

Observe that the vertices of $G_1 \vee G_2$ consist of all the vertices of G_1 and of G_2 , and two such vertices v, w are connected in $G_1 \vee G_2$ exactly when:

- both v, w are vertices of G_1 , and there is an edge in G_1 joining v and w ,

Reminder: Join of Two Graphs

Consider two graphs $G_1 = (V_1, E_1)$ and $G_2 = (V_2, E_2)$ with disjoint vertex sets, namely such that

$$V_1 \cap V_2 = \emptyset.$$

The join of G_1 and G_2 is a new graph with vertex set

$$V_1 \cup V_2$$

and edge set

$$E_1 \cup E_2 \cup \{\{v, w\} : v \in V_1, w \in V_2\}.$$

We denote the join of G_1 and G_2 by $G_1 \vee G_2$.

Observe that the vertices of $G_1 \vee G_2$ consist of all the vertices of G_1 and of G_2 , and two such vertices v, w are connected in $G_1 \vee G_2$ exactly when:

- both v, w are vertices of G_1 , and there is an edge in G_1 joining v and w ,
- or both v, w are vertices of G_2 , and there is an edge in G_2 joining v and w ,

Reminder: Join of Two Graphs

Consider two graphs $G_1 = (V_1, E_1)$ and $G_2 = (V_2, E_2)$ with disjoint vertex sets, namely such that

$$V_1 \cap V_2 = \emptyset.$$

The join of G_1 and G_2 is a new graph with vertex set

$$V_1 \cup V_2$$

and edge set


$$E_1 \cup E_2 \cup \{\{v, w\} : v \in V_1, w \in V_2\}.$$

We denote the join of G_1 and G_2 by $G_1 \vee G_2$.

Observe that the vertices of $G_1 \vee G_2$ consist of all the vertices of G_1 and of G_2 , and two such vertices v, w are connected in $G_1 \vee G_2$ exactly when:

- both v, w are vertices of G_1 , and there is an edge in G_1 joining v and w ,
- or both v, w are vertices of G_2 , and there is an edge in G_2 joining v and w ,
- or finally v is a vertex of G_1 and w is a vertex of G_2 (or conversely, v is a vertex of G_2 and w a vertex of G_1).


Examples



The bipartite graph $K_{6,4}$ can be viewed as the join of the null graphs N_6 and N_4 :


$$K_{6,4} = N_6 \vee N_4.$$

Examples




The bipartite graph $K_{6,4}$ can be viewed as the join of the null graphs N_6 and N_4 :

$$K_{6,4} = N_6 \vee N_4.$$




Examples



The bipartite graph $K_{6,4}$ can be viewed as the join of the null graphs N_6 and N_4 :

$$K_{6,4} = N_6 \vee N_4.$$



The wheel graph W_{10} can be viewed as the join of the cycle C_9 and the null graph N_1 :

$$W_{10} = C_9 \vee N_1.$$

Reminder: Line Graph of a Graph

Let $G = (V, E)$ be a graph, which has a **non-empty** edge set E (in other words, there is at least one edge in G).

Reminder: Line Graph of a Graph

Let $G = (V, E)$ be a graph, which has a **non-empty** edge set E (in other words, there is at least one edge in G).

Then we can define a new graph, called the line graph of G , which will be denoted by $L(G)$, as follows:

- the vertex set of $L(G)$ is the edge set of G ; in other words, $V(L(G)) = E(G)$;

Reminder: Line Graph of a Graph


Let $G = (V, E)$ be a graph, which has a **non-empty** edge set E (in other words, there is at least one edge in G).

Then we can define a new graph, called the line graph of G , which will be denoted by $L(G)$, as follows:

- the vertex set of $L(G)$ is the edge set of G ; in other words, $V(L(G)) = E(G)$;
- two 'vertices' in $L(G)$ are joined if, when we view them as edges of G , they are adjacent, or in other words, they have a common endvertex.


An example

Recall the graph G_1 :




An example

Recall the graph G_1 :



Then $L(G_1)$ is the following graph:



'Deleting' Vertices

Let $G = (V, E)$ be a graph, and consider a (nonempty) subset V' of the vertex set V of G .

'Deleting' Vertices

Let $G = (V, E)$ be a graph, and consider a (nonempty) subset V' of the vertex set V of G .

As we just recalled, the induced subgraph of G with vertex set V' is the maximal subgraph of G with vertex set V' , or in other words, it is the subgraph of G on the vertices in V' which contains all the (possible) edges of G joining two vertices in V' .

We denote the induced subgraph of G on the vertices in V' by $G[V']$.

'Deleting' Vertices

Let $G = (V, E)$ be a graph, and consider a (nonempty) subset V' of the vertex set V of G .

As we just recalled, the induced subgraph of G with vertex set V' is the maximal subgraph of G with vertex set V' , or in other words, it is the subgraph of G on the vertices in V' which contains all the (possible) edges of G joining two vertices in V' .

We denote the induced subgraph of G on the vertices in V' by $G[V']$.

Consider now a **proper** subset V'' of the vertex set V of G .

We can construct a new graph, which will also be a subgraph of G , **if we simply delete the vertices of G which are contained in V'' , and of course also remove all the edges of G which have at least one endvertex in V'' .**

'Deleting' Vertices

Let $G = (V, E)$ be a graph, and consider a (nonempty) subset V' of the vertex set V of G .

As we just recalled, the induced subgraph of G with vertex set V' is the maximal subgraph of G with vertex set V' , or in other words, it is the subgraph of G on the vertices in V' which contains all the (possible) edges of G joining two vertices in V' .

We denote the induced subgraph of G on the vertices in V' by $G[V']$.

Consider now a **proper** subset V'' of the vertex set V of G .

We can construct a new graph, which will also be a subgraph of G , **if we simply delete the vertices of G which are contained in V'' , and of course also remove all the edges of G which have at least one endvertex in V'' .**

Observe that the new graph is precisely the induced subgraph with vertex set $V \setminus V''$, or in other words the subgraph $G[V \setminus V'']$.

'Deleting' Vertices

Let $G = (V, E)$ be a graph, and consider a (nonempty) subset V' of the vertex set V of G .

As we just recalled, the induced subgraph of G with vertex set V' is the maximal subgraph of G with vertex set V' , or in other words, it is the subgraph of G on the vertices in V' which contains all the (possible) edges of G joining two vertices in V' .

We denote the induced subgraph of G on the vertices in V' by $G[V']$.

Consider now a **proper** subset V'' of the vertex set V of G .

We can construct a new graph, which will also be a subgraph of G , **if we simply delete the vertices of G which are contained in V'' , and of course also remove all the edges of G which have at least one endvertex in V'' .**

Observe that the new graph is precisely the induced subgraph with vertex set $V \setminus V''$, or in other words the subgraph $G[V \setminus V'']$.

We will also use the notation $G - V''$ for this graph.

Moreover, when V'' contains **only one vertex of G** , say vertex v_0 , we will also sometimes write $G - v_0$ instead of $G - \{v_0\}$.

'Deleting' Edges

Let $G = (V, E)$ be a graph, and consider a subset E' of the edge set E of G .

'Deleting' Edges

Let $G = (V, E)$ be a graph, and consider a subset E' of the edge set E of G .

We can construct a new graph, which will also be a subgraph of G , **if we simply delete the edges of G which are contained in the edge subset E'** . In other words, we can consider the subgraph

$$(V, E \setminus E')$$

of G , which we will denote by $G - E'$.

Again, if E' contains **only one element, in this case one edge of G** , say edge e_0 , we will more simply write $G - e_0$ instead of $G - \{e_0\}$.

Next Main Topic:
Results about Connected Graphs
and about Connectivity

One such result:
Complement of a disconnected graph

Proposition 1

Let G be a disconnected graph, that is, a graph that has at least two connected components.

Then the complement \overline{G} of G must be connected.

Proof of Proposition 1

According to our assumption, G is a disconnected graph, so it has at least two connected components, say, components G_1 and G_2 .

Proof of Proposition 1

According to our assumption, G is a disconnected graph, so it has at least two connected components, say, components G_1 and G_2 .

Consider two different vertices $u, v \in V(\overline{G}) = V(G)$. We need to show that there is a path in \overline{G} starting at u and ending at v .

We distinguish two cases:

Proof of Proposition 1

According to our assumption, G is a disconnected graph, so it has at least two connected components, say, components G_1 and G_2 .

Consider two different vertices $u, v \in V(\overline{G}) = V(G)$. We need to show that there is a path in \overline{G} starting at u and ending at v .

We distinguish two cases:

Case 1: u, v are in different connected components of G . Then we certainly have that u, v are not joined by an edge in G , and hence, by the way we define \overline{G} , u and v must be joined by an edge in \overline{G} .

Proof of Proposition 1

According to our assumption, G is a disconnected graph, so it has at least two connected components, say, components G_1 and G_2 .

Consider two different vertices $u, v \in V(\overline{G}) = V(G)$. We need to show that there is a path in \overline{G} starting at u and ending at v .

We distinguish two cases:

Case 1: u, v are in different connected components of G . Then we certainly have that u, v are not joined by an edge in G , and hence, by the way we define \overline{G} , u and v must be joined by an edge in \overline{G} .

Case 2: u, v belong to the same connected component of G . Then at least one of the components G_1, G_2 that we considered above must be different from the connected component of G which contains both u and v ; say, G_1 is different from the component of u and v .

Consider a vertex w in G_1 . Since u and v are not contained in G_1 , while G_1 contains all the neighbours of w (given the way we define the connected components of G), we see that $uw \notin E(G)$ and similarly $vw \notin E(G)$.

But then $uw \in E(\overline{G})$ and similarly $vw \in E(\overline{G})$. It follows that the path

$$(\{u, w, v\}, \{uw, wv\})$$

is a path in \overline{G} connecting u and v .

Proof of Proposition 1

According to our assumption, G is a disconnected graph, so it has at least two connected components, say, components G_1 and G_2 .

Consider two different vertices $u, v \in V(\overline{G}) = V(G)$. We need to show that there is a path in \overline{G} starting at u and ending at v .

We distinguish two cases:

Case 1: u, v are in different connected components of G . Then we certainly have that u, v are not joined by an edge in G , and hence, by the way we define \overline{G} , u and v must be joined by an edge in \overline{G} .

Case 2: u, v belong to the same connected component of G . Then at least one of the components G_1, G_2 that we considered above must be different from the connected component of G which contains both u and v ; say, G_1 is different from the component of u and v .

Consider a vertex w in G_1 . Since u and v are not contained in G_1 , while G_1 contains all the neighbours of w (given the way we define the connected components of G), we see that $uw \notin E(G)$ and similarly $vw \notin E(G)$.

But then $uw \in E(\overline{G})$ and similarly $vw \in E(\overline{G})$. It follows that the path

$$(\{u, w, v\}, \{uw, wv\})$$

is a path in \overline{G} connecting u and v .

We conclude that \overline{G} is connected.

Two important parameters of graphs

Let $G = (V, E)$ be a finite graph.

- We will denote by $\delta(G)$ the minimum degree of a vertex in G ,

Two important parameters of graphs

Let $G = (V, E)$ be a finite graph.


- We will denote by $\delta(G)$ the minimum degree of a vertex in G ,
- and by $\Delta(G)$ the maximum degree of a vertex in G .

Two important parameters of graphs


Let $G = (V, E)$ be a finite graph.

- We will denote by $\delta(G)$ the minimum degree of a vertex in G ,
- and by $\Delta(G)$ the maximum degree of a vertex in G .

Two examples.



Here $\delta(G) = 2$ and $\Delta(G) = 4$



Here $\delta(G) = 1$ and $\Delta(G) = 4$

One example of the significance of $\delta(G)$

Proposition 2

Let G be a finite graph of order n , and assume that $\delta(G) \geq \frac{n-1}{2}$.
Then G **must** be connected.

One example of the significance of $\delta(G)$

Proposition 2

Let G be a finite graph of order n , and assume that $\delta(G) \geq \frac{n-1}{2}$. Then G **must** be connected.

Practice Exercise for you! Justify with an example that the lower bound $\frac{n-1}{2}$ for $\delta(G)$ is **best possible** and we cannot make it smaller; that is, come up with a graph G' which

- satisfies $\delta(G') \geq \frac{n-2}{2}$,
- but is still **disconnected**.

Proof of Proposition 2

Assume towards a contradiction that G is disconnected. Then G has at least two connected components, and if we consider one of those connected components of G which has smallest possible order, say, component G_1 , then we can be certain that G_1 contains $\leq \frac{n}{2}$ vertices of G (*why?*).

Proof of Proposition 2

Assume towards a contradiction that G is disconnected. Then G has at least two connected components, and if we consider one of those connected components of G which has smallest possible order, say, component G_1 , then we can be certain that G_1 contains $\leq \frac{n}{2}$ vertices of G (*why?*).

At the same time, $\delta(G_1) \geq \delta(G) \geq \frac{n-1}{2}$. This is because, for every vertex u contained in G_1 , we also have all its neighbours belonging to G_1 (given that G_1 is a maximal connected subgraph of G), and hence

$$\deg_{G_1}(u) = \deg_G(u).$$

Proof of Proposition 2

Assume towards a contradiction that G is disconnected. Then G has at least two connected components, and if we consider one of those connected components of G which has smallest possible order, say, component G_1 , then we can be certain that G_1 contains $\leq \frac{n}{2}$ vertices of G (*why?*).

At the same time, $\delta(G_1) \geq \delta(G) \geq \frac{n-1}{2}$. This is because, for every vertex u contained in G_1 , we also have all its neighbours belonging to G_1 (given that G_1 is a maximal connected subgraph of G), and hence

$$\deg_{G_1}(u) = \deg_G(u).$$

We now reach a contradiction:

- since G_1 has order $\leq \frac{n}{2}$, its maximum degree $\Delta(G_1)$ must be $\leq \frac{n}{2} - 1 = \frac{n-2}{2}$;

Proof of Proposition 2

Assume towards a contradiction that G is disconnected. Then G has at least two connected components, and if we consider one of those connected components of G which has smallest possible order, say, component G_1 , then we can be certain that G_1 contains $\leq \frac{n}{2}$ vertices of G (*why?*).

At the same time, $\delta(G_1) \geq \delta(G) \geq \frac{n-1}{2}$. This is because, for every vertex u contained in G_1 , we also have all its neighbours belonging to G_1 (given that G_1 is a maximal connected subgraph of G), and hence

$$\deg_{G_1}(u) = \deg_G(u).$$

We now reach a contradiction:

- since G_1 has order $\leq \frac{n}{2}$, its maximum degree $\Delta(G_1)$ must be $\leq \frac{n}{2} - 1 = \frac{n-2}{2}$;
- at the same time, $\Delta(G_1) \geq \delta(G_1) \geq \frac{n-1}{2}$, which contradicts the previous conclusion.

Proof of Proposition 2

Assume towards a contradiction that G is disconnected. Then G has at least two connected components, and if we consider one of those connected components of G which has smallest possible order, say, component G_1 , then we can be certain that G_1 contains $\leq \frac{n}{2}$ vertices of G (*why?*).

At the same time, $\delta(G_1) \geq \delta(G) \geq \frac{n-1}{2}$. This is because, for every vertex u contained in G_1 , we also have all its neighbours belonging to G_1 (given that G_1 is a maximal connected subgraph of G), and hence

$$\deg_{G_1}(u) = \deg_G(u).$$

We now reach a contradiction:

- since G_1 has order $\leq \frac{n}{2}$, its maximum degree $\Delta(G_1)$ must be $\leq \frac{n}{2} - 1 = \frac{n-2}{2}$;
- at the same time, $\Delta(G_1) \geq \delta(G_1) \geq \frac{n-1}{2}$, which contradicts the previous conclusion.

The above show that the assumption that G has at least two connected components was incorrect.

Another such result

One of the problems to be given in HW2

Let k be a positive integer, and let G be a graph satisfying $\delta(G) \geq k$.

- Show that G contains a path of length at least k .
- If $k \geq 2$, show that G contains a cycle of order at least $k + 1$.

One more result about connected graphs

Proposition 3

Let G be a graph of order n , and suppose that G has size strictly less than $n - 1$.

Then G **must** be disconnected.

One more result about connected graphs

Proposition 3

Let G be a graph of order n , and suppose that G has size strictly less than $n - 1$.

Then G **must** be disconnected.

We'll justify this very soon.

Important Subgraphs and related notions

Recall the families of **paths** and of **cycle graphs**, and recall that we have seen examples where we look for graphs from these families among the subgraphs of a given graph G (with, say, vertex set $V = \{v_1, v_2, \dots, v_n\}$).

- **paths**
- **cycles**

Important Subgraphs and related notions

Recall the families of **paths** and of **cycle graphs**, and recall that we have seen examples where we look for graphs from these families among the subgraphs of a given graph G (with, say, vertex set $V = \{v_1, v_2, \dots, v_n\}$). Related notions are the following:

- **paths**
- **cycles**

Important Subgraphs and related notions

Recall the families of **paths** and of **cycle graphs**, and recall that we have seen examples where we look for graphs from these families among the subgraphs of a given graph G (with, say, vertex set $V = \{v_1, v_2, \dots, v_n\}$). Related notions are the following:

- **walks**
- **paths**
- **cycles**

Important Subgraphs and related notions

Recall the families of **paths** and of **cycle graphs**, and recall that we have seen examples where we look for graphs from these families among the subgraphs of a given graph G (with, say, vertex set $V = \{v_1, v_2, \dots, v_n\}$). Related notions are the following:

- **walks** A *walk* of length k in G is a sequence of (not necessarily distinct) vertices $v_{i_0}, v_{i_1}, v_{i_2}, \dots, v_{i_k}$ from V , such that $v_i v_{i+1} \in E(G)$ for every $i = 0, 1, 2, \dots, k-1$. The vertices v_{i_0} and v_{i_k} are called the *endvertices* of the walk, and we sometimes say that this is a $v_{i_0} - v_{i_k}$ walk.
- **paths**
- **cycles**

Important Subgraphs and related notions

Recall the families of **paths** and of **cycle graphs**, and recall that we have seen examples where we look for graphs from these families among the subgraphs of a given graph G (with, say, vertex set $V = \{v_1, v_2, \dots, v_n\}$). Related notions are the following:

- **walks** A *walk* of length k in G is a sequence of (not necessarily distinct) vertices $v_{i_0}, v_{i_1}, v_{i_2}, \dots, v_{i_k}$ from V , such that $v_i v_{i+1} \in E(G)$ for every $i = 0, 1, 2, \dots, k-1$. The vertices v_{i_0} and v_{i_k} are called the *endvertices* of the walk, and we sometimes say that this is a $v_{i_0} - v_{i_k}$ walk.

Since G is a graph (and thus, according to our convention, it does not contain multiple edges), we can completely describe the walk by writing the vertices one next to the other in the correct order: $v_{i_0} v_{i_1} v_{i_2} \dots v_{i_k}$.

- **paths**
- **cycles**

Important Subgraphs and related notions

Recall the families of **paths** and of **cycle graphs**, and recall that we have seen examples where we look for graphs from these families among the subgraphs of a given graph G (with, say, vertex set $V = \{v_1, v_2, \dots, v_n\}$). Related notions are the following:

- **walks** A *walk* of length k in G is a sequence of (not necessarily distinct) vertices $v_{i_0}, v_{i_1}, v_{i_2}, \dots, v_{i_k}$ from V , such that $v_i v_{i+1} \in E(G)$ for every $i = 0, 1, 2, \dots, k-1$. The vertices v_{i_0} and v_{i_k} are called the *endvertices* of the walk, and we sometimes say that this is a $v_{i_0} - v_{i_k}$ walk.

Since G is a graph (and thus, according to our convention, it does not contain multiple edges), we can completely describe the walk by writing the vertices one next to the other in the correct order: $v_{i_0} v_{i_1} v_{i_2} \dots v_{i_k}$.

- **paths** Recall that a path contained in the graph G is simply a walk in which **all the vertices are distinct**.
- **cycles**

Important Subgraphs and related notions

Recall the families of **paths** and of **cycle graphs**, and recall that we have seen examples where we look for graphs from these families among the subgraphs of a given graph G (with, say, vertex set $V = \{v_1, v_2, \dots, v_n\}$). Related notions are the following:

- **walks** A *walk* of length k in G is a sequence of (not necessarily distinct) vertices $v_{i_0}, v_{i_1}, v_{i_2}, \dots, v_{i_k}$ from V , such that $v_i v_{i+1} \in E(G)$ for every $i = 0, 1, 2, \dots, k-1$. The vertices v_{i_0} and v_{i_k} are called the *endvertices* of the walk, and we sometimes say that this is a $v_{i_0} - v_{i_k}$ walk.

Since G is a graph (and thus, according to our convention, it does not contain multiple edges), we can completely describe the walk by writing the vertices one next to the other in the correct order: $v_{i_0} v_{i_1} v_{i_2} \dots v_{i_k}$.

- **paths** Recall that a path contained in the graph G is simply a walk in which **all the vertices are distinct**.
- **cycles** If the endvertices v_{i_0} and v_{i_k} of the walk coincide, but all the other vertices are distinct and also different from v_{i_0} , then we get a 'closed path', which we have called a *cycle*.

Important Subgraphs and related notions

Recall the families of **paths** and of **cycle graphs**, and recall that we have seen examples where we look for graphs from these families among the subgraphs of a given graph G (with, say, vertex set $V = \{v_1, v_2, \dots, v_n\}$). Related notions are the following:

- **walks** A *walk* of length k in G is a sequence of (not necessarily distinct) vertices $v_{i_0}, v_{i_1}, v_{i_2}, \dots, v_{i_k}$ from V , such that $v_i v_{i+1} \in E(G)$ for every $i = 0, 1, 2, \dots, k-1$. The vertices v_{i_0} and v_{i_k} are called the *endvertices* of the walk, and we sometimes say that this is a $v_{i_0} - v_{i_k}$ walk.

Since G is a graph (and thus, according to our convention, it does not contain multiple edges), we can completely describe the walk by writing the vertices one next to the other in the correct order: $v_{i_0} v_{i_1} v_{i_2} \dots v_{i_k}$.

- **paths** Recall that a path contained in the graph G is simply a walk in which **all the vertices are distinct**.
- **cycles** If the endvertices v_{i_0} and v_{i_k} of the walk coincide, but all the other vertices are distinct and also different from v_{i_0} , then we get a 'closed path', which we have called a *cycle*.
- **trails**

Important Subgraphs and related notions

Recall the families of **paths** and of **cycle graphs**, and recall that we have seen examples where we look for graphs from these families among the subgraphs of a given graph G (with, say, vertex set $V = \{v_1, v_2, \dots, v_n\}$). Related notions are the following:

- **walks** A *walk* of length k in G is a sequence of (not necessarily distinct) vertices $v_{i_0}, v_{i_1}, v_{i_2}, \dots, v_{i_k}$ from V , such that $v_i v_{i+1} \in E(G)$ for every $i = 0, 1, 2, \dots, k-1$. The vertices v_{i_0} and v_{i_k} are called the *endvertices* of the walk, and we sometimes say that this is a $v_{i_0} - v_{i_k}$ walk.
Since G is a graph (and thus, according to our convention, it does not contain multiple edges), we can completely describe the walk by writing the vertices one next to the other in the correct order: $v_{i_0} v_{i_1} v_{i_2} \dots v_{i_k}$.
- **paths** Recall that a path contained in the graph G is simply a walk in which **all the vertices are distinct**.
- **cycles** If the endvertices v_{i_0} and v_{i_k} of the walk coincide, but all the other vertices are distinct and also different from v_{i_0} , then we get a 'closed path', which we have called a *cycle*.
- **trails** If **all the edges in a walk are distinct** (but not necessarily all the vertices), we call this walk a *trail*.

Important Subgraphs and related notions

Recall the families of **paths** and of **cycle graphs**, and recall that we have seen examples where we look for graphs from these families among the subgraphs of a given graph G (with, say, vertex set $V = \{v_1, v_2, \dots, v_n\}$). Related notions are the following:


- **walks** A *walk* of length k in G is a sequence of (not necessarily distinct) vertices $v_{i_0}, v_{i_1}, v_{i_2}, \dots, v_{i_k}$ from V , such that $v_i v_{i+1} \in E(G)$ for every $i = 0, 1, 2, \dots, k-1$. The vertices v_{i_0} and v_{i_k} are called the *endvertices* of the walk, and we sometimes say that this is a $v_{i_0} - v_{i_k}$ walk.
Since G is a graph (and thus, according to our convention, it does not contain multiple edges), we can completely describe the walk by writing the vertices one next to the other in the correct order: $v_{i_0} v_{i_1} v_{i_2} \dots v_{i_k}$.
- **paths** Recall that a path contained in the graph G is simply a walk in which **all the vertices are distinct**.
- **cycles** If the endvertices v_{i_0} and v_{i_k} of the walk coincide, but all the other vertices are distinct and also different from v_{i_0} , then we get a 'closed path', which we have called a *cycle*.
- **trails** If **all the edges in a walk are distinct** (but not necessarily all the vertices), we call this walk a *trail*.
- **circuits**

Important Subgraphs and related notions


Recall the families of **paths** and of **cycle graphs**, and recall that we have seen examples where we look for graphs from these families among the subgraphs of a given graph G (with, say, vertex set $V = \{v_1, v_2, \dots, v_n\}$). Related notions are the following:

- **walks** A *walk* of length k in G is a sequence of (not necessarily distinct) vertices $v_{i_0}, v_{i_1}, v_{i_2}, \dots, v_{i_k}$ from V , such that $v_i v_{i+1} \in E(G)$ for every $i = 0, 1, 2, \dots, k-1$. The vertices v_{i_0} and v_{i_k} are called the *endvertices* of the walk, and we sometimes say that this is a $v_{i_0} - v_{i_k}$ walk.
Since G is a graph (and thus, according to our convention, it does not contain multiple edges), we can completely describe the walk by writing the vertices one next to the other in the correct order: $v_{i_0} v_{i_1} v_{i_2} \dots v_{i_k}$.
- **paths** Recall that a path contained in the graph G is simply a walk in which **all the vertices are distinct**.
- **cycles** If the endvertices v_{i_0} and v_{i_k} of the walk coincide, but all the other vertices are distinct and also different from v_{i_0} , then we get a 'closed path', which we have called a *cycle*.
- **trails** If **all the edges in a walk are distinct** (but not necessarily all the vertices), we call this walk a *trail*.
- **circuits** A *circuit* is a 'closed trail', that is, a walk in which **all edges are distinct**, and also the endvertices coincide.

Practice on examples




Graph G_1




Graph G_2

Practice on examples




Graph G_1




Graph G_2

The $a - e$ walk in graph G_1 given by $a b g d f c b d e$

Practice on examples




Graph G_1




Graph G_2

The $a - e$ walk in graph G_1 given by $a b g d f c b d e$ is a trail of length 8, but it is not a path (because the edges are all distinct, but the vertices b and d appear in the sequence more than once).

Practice on examples



Graph G_1




Graph G_2


The $a - e$ walk in graph G_1 given by $a b g d f c b d e$ is a trail of length 8, but it is not a path (because the edges are all distinct, but the vertices b and d appear in the sequence more than once).

The walk in graph G_2 given by $a c d e g f d a$

Practice on examples



Graph G_1



Graph G_2

The $a - e$ walk in graph G_1 given by $abgdfcbde$ is a trail of length 8, but it is not a path (because the edges are all distinct, but the vertices b and d appear in the sequence more than once).

The walk in graph G_2 given by $acdegfd a$ is a circuit of length 7, but it is not a cycle (because the edges are all distinct, and the initial and terminal vertex coincide (so it is a 'closed trail'), but at the same time the vertex d , which is not the initial or terminal vertex, also appears twice in the sequence).

Relating walks and paths

Theorem

Let G be a graph, and let u, v be two different vertices of G .
Any $u - v$ walk in G contains a $u - v$ path.

Relating walks and paths

Theorem

Let G be a graph, and let u, v be two different vertices of G .
Any $u - v$ walk in G contains a $u - v$ path.

Proof of the theorem. We will use induction on the length k of the path.

Relating walks and paths

Theorem

Let G be a graph, and let u, v be two different vertices of G .
Any $u - v$ walk in G contains a $u - v$ path.

Proof of the theorem. We will use induction on the length k of the path.

Base case. If $k = 1$, then the walk $u v$ is clearly a path.

Relating walks and paths

Theorem

Let G be a graph, and let u, v be two different vertices of G . Any $u - v$ walk in G contains a $u - v$ path.

Proof of the theorem. We will use induction on the length k of the path.

Base case. If $k = 1$, then the walk uv is clearly a path. In fact, in this setting, even when $k = 2$, we automatically have that the given $u - v$ walk is a $u - v$ path: this is because we have assumed that u and v are different vertices, so if the walk is uwv with uw and $wv \in E(G)$, then clearly the vertex w has to be different both from u and from v , and thus all the vertices appearing in the walk appear only once.

Relating walks and paths

Theorem

Let G be a graph, and let u, v be two different vertices of G . Any $u - v$ walk in G contains a $u - v$ path.

Proof of the theorem. We will use induction on the length k of the path.

Base case. If $k = 1$, then the walk $u v$ is clearly a path. In fact, in this setting, even when $k = 2$, we automatically have that the given $u - v$ walk is a $u - v$ path: this is because we have assumed that u and v are different vertices, so if the walk is $u w v$ with uw and $wv \in E(G)$, then clearly the vertex w has to be different both from u and from v , and thus all the vertices appearing in the walk appear only once.

Induction Step. Consider now some $k \geq 1$ (or $k \geq 2$), and assume that we already know that every $u - v$ walk of length $\leq k$ contains a $u - v$ path.

Consider a $u - v$ walk of length $k + 1$, which we can write as

$$u = w_0 w_1 w_2 \cdots w_k w_{k+1} = v.$$

Relating walks and paths

Theorem

Let G be a graph, and let u, v be two different vertices of G .
Any $u - v$ walk in G contains a $u - v$ path.

Proof of the theorem. We will use induction on the length k of the path.

Base case. If $k = 1$, then the walk $u v$ is clearly a path. In fact, in this setting, even when $k = 2$, we automatically have that the given $u - v$ walk is a $u - v$ path: this is because we have assumed that u and v are different vertices, so if the walk is $u w v$ with uw and $wv \in E(G)$, then clearly the vertex w has to be different both from u and from v , and thus all the vertices appearing in the walk appear only once.

Induction Step. Consider now some $k \geq 1$ (or $k \geq 2$), and assume that we already know that every $u - v$ walk of length $\leq k$ contains a $u - v$ path.

Consider a $u - v$ walk of length $k + 1$, which we can write as

$$u = w_0 w_1 w_2 \cdots w_k w_{k+1} = v.$$

- If all the vertices in this walk are distinct, then this is a $u - v$ path already.

Relating walks and paths

Theorem

Let G be a graph, and let u, v be two different vertices of G . Any $u - v$ walk in G contains a $u - v$ path.

Proof of the theorem. We will use induction on the length k of the path.

Base case. If $k = 1$, then the walk uv is clearly a path. In fact, in this setting, even when $k = 2$, we automatically have that the given $u - v$ walk is a $u - v$ path: this is because we have assumed that u and v are different vertices, so if the walk is uwv with uw and $wv \in E(G)$, then clearly the vertex w has to be different both from u and from v , and thus all the vertices appearing in the walk appear only once.

Induction Step. Consider now some $k \geq 1$ (or $k \geq 2$), and assume that we already know that every $u - v$ walk of length $\leq k$ contains a $u - v$ path.

Consider a $u - v$ walk of length $k + 1$, which we can write as

$$u = w_0 w_1 w_2 \cdots w_k w_{k+1} = v.$$

- If all the vertices in this walk are distinct, then this is a $u - v$ path already.
- If there are vertices in the sequence which are equal, then we can find $j, r \in \{0, 1, 2, \dots, k, k + 1\}$ with $j < r$ such that $w_j = w_r$. But then the sequence

$$u = w_0 w_1 w_2 \cdots w_{j-1} w_j \rightsquigarrow w_{r+1} w_{r+2} \cdots w_k w_{k+1} = v,$$

in which we go directly from the vertex w_j to the vertex w_{r+1} (which we can do because w_{r+1} is adjacent to $w_r = w_j$), is also a $u - v$ walk which has length $< k + 1$, and thus by the Inductive Hypothesis it will contain a $u - v$ path.

Proof of Proposition 3

Recall that we have a graph G of order n and of size $< n - 1$, and we are asked to show that G is disconnected.

Proof of Proposition 3

Recall that we have a graph G of order n and of size $< n - 1$, and we are asked to show that G is disconnected.

Assume towards a contradiction that we have considered such a graph G which is connected, and suppose that $V(G) = \{v_1, v_2, \dots, v_n\}$.

Proof of Proposition 3

Recall that we have a graph G of order n and of size $< n - 1$, and we are asked to show that G is disconnected.

Assume towards a contradiction that we have considered such a graph G which is connected, and suppose that $V(G) = \{v_1, v_2, \dots, v_n\}$.

Since G has been assumed connected,

- we can find a path starting at vertex v_1 and ending at vertex v_2 ,
- and similarly we can find a path starting at vertex v_2 and ending at vertex v_3 ,
- and so on, until finally we find a path which starts at vertex v_{n-1} and ends at vertex v_n .

By simply traversing all these paths one after the other, we can construct a walk in G which starts at vertex v_1 , ends at vertex v_n , and passes by every vertex of G .

Proof of Proposition 3 (cont.)

Claim. The $v_1 - v_n$ walk described above must contain at least $n - 1$ different edges.

Proof of Proposition 3 (cont.)

Claim. The $v_1 - v_n$ walk described above must contain at least $n - 1$ different edges.

Indeed, reorder the vertices v_2, v_3, \dots, v_n as $v_{i_2}, v_{i_3}, \dots, v_{i_n}$ according to which of them appears first in the walk, which one appears second, and so on.

- Note that the vertex v_{i_2} must come right after the initial vertex v_1 of the walk, and that the edge $e_1 = v_1 v_{i_2}$ is the first edge of the walk.

Proof of Proposition 3 (cont.)

Claim. The $v_1 - v_n$ walk described above must contain at least $n - 1$ different edges.

Indeed, reorder the vertices v_2, v_3, \dots, v_n as $v_{i_2}, v_{i_3}, \dots, v_{i_n}$ according to which of them appears first in the walk, which one appears second, and so on.

- Note that the vertex v_{i_2} must come right after the initial vertex v_1 of the walk, and that the edge $e_1 = v_1 v_{i_2}$ is the first edge of the walk.
- Note also that when we end up at vertex v_{i_3} for the first time, then we have travelled along a ‘new’ edge e_2 , that is, an edge which is different from $e_1 = v_1 v_{i_2}$ (given that v_{i_3} is incident with e_2 , whereas it is not incident with e_1).

Proof of Proposition 3 (cont.)

Claim. The $v_1 - v_n$ walk described above must contain at least $n - 1$ different edges.

Indeed, reorder the vertices v_2, v_3, \dots, v_n as $v_{i_2}, v_{i_3}, \dots, v_{i_n}$ according to which of them appears first in the walk, which one appears second, and so on.

- Note that the vertex v_{i_2} must come right after the initial vertex v_1 of the walk, and that the edge $e_1 = v_1 v_{i_2}$ is the first edge of the walk.
- Note also that when we end up at vertex v_{i_3} for the first time, then we have travelled along a ‘new’ edge e_2 , that is, an edge which is different from $e_1 = v_1 v_{i_2}$ (given that v_{i_3} is incident with e_2 , whereas it is not incident with e_1).
- Continuing like this, we see that the first time we end up at vertex v_{i_j} (for any $j \in \{2, \dots, n\}$), we have travelled along a ‘new’ edge e_{j-1} which is different from the previous edges we have found, given that v_{i_j} is incident with e_{j-1} , but it is not incident with any of these previous edges (as this is the first time we come to vertex v_{i_j} while traversing our walk).

Proof of Proposition 3 (cont.)

Claim. The $v_1 - v_n$ walk described above must contain at least $n - 1$ different edges.

Indeed, reorder the vertices v_2, v_3, \dots, v_n as $v_{i_2}, v_{i_3}, \dots, v_{i_n}$ according to which of them appears first in the walk, which one appears second, and so on.

- Note that the vertex v_{i_2} must come right after the initial vertex v_1 of the walk, and that the edge $e_1 = v_1 v_{i_2}$ is the first edge of the walk.
- Note also that when we end up at vertex v_{i_3} for the first time, then we have travelled along a ‘new’ edge e_2 , that is, an edge which is different from $e_1 = v_1 v_{i_2}$ (given that v_{i_3} is incident with e_2 , whereas it is not incident with e_1).
- Continuing like this, we see that the first time we end up at vertex v_{i_j} (for any $j \in \{2, \dots, n\}$), we have travelled along a ‘new’ edge e_{j-1} which is different from the previous edges we have found, given that v_{i_j} is incident with e_{j-1} , but it is not incident with any of these previous edges (as this is the first time we come to vertex v_{i_j} while traversing our walk).

By the claim, we get that the graph G itself must contain at least $n - 1$ edges, which contradicts the assumption that the size of G is $< n - 1$. Thus our assumption that G is connected was incorrect.

Cyclic/Acyclic Graphs – Trees and Forests

Definition

- 1 A graph G is called cyclic if it contains at least one cycle. It is called acyclic otherwise.

Cyclic/Acyclic Graphs – Trees and Forests

Definition

- 1 A graph G is called cyclic if it contains at least one cycle. It is called acyclic otherwise.
- 2 A connected acyclic graph is called a tree.

Cyclic/Acyclic Graphs – Trees and Forests

Definition


- 1 A graph G is called cyclic if it contains at least one cycle. It is called acyclic otherwise.
- 2 A connected acyclic graph is called a tree.
- 3 On the other hand, if an acyclic graph is disconnected, then it is called a forest (in other words, *forests* are disjoint unions of two or more *trees*).

Cyclic/Acyclic Graphs – Trees and Forests

Definition

- 1 A graph G is called cyclic if it contains at least one cycle. It is called acyclic otherwise.
- 2 A connected acyclic graph is called a tree.
- 3 On the other hand, if an acyclic graph is disconnected, then it is called a forest (in other words, *forests* are disjoint unions of two or more *trees*).
- 4 A vertex of degree 1 in a tree or a forest is called a leaf.


Examples



from the Harris-Hirst-Mossinghoff book


Question. Which of the above graphs are trees? Which are not, and why? Which vertices are leaves in the graphs that are trees?

Example of a cyclic graph



Map of the Paris Subway-Regional Train system

Example of an acyclic graph



One more example of a tree capturing a real-life event/situation




FIGURE 1.35. The 2008 Men's Sweet 16.

from the Harris-Hirst-Mossinghoff book