

# **MATH 322 – Graph Theory**

## **Fall Term 2021**

### **Notes for Lecture 25**

Wednesday, December 8

# **Crossing Number of a Graph, and Plane/Planar Graphs**

From last time

## Definition 1

Given a (finite) graph  $G$ , we define its crossing number to be the minimum number of edge crossings that a (permissible) pictorial representation of  $G$  can have.

We usually denote this number by  $cr(G)$ .

From last time

### Definition 1

Given a (finite) graph  $G$ , we define its crossing number to be the minimum number of edge crossings that a (permissible) pictorial representation of  $G$  can have.

We usually denote this number by  $cr(G)$ .


### Definition 2

A graph  $G = (V, E)$  is called planar if we can draw it in the plane without any edge crossings. In other words, if  $cr(G) = 0$ .

Any such pictorial representation of  $G$  will be called a planar embedding, or equivalently a plane representation, of  $G$ .


## Faces of a planar graph

Consider now a plane representation  $G_0$  of a planar finite graph (note that  $G_0$  for now denotes the specific plane representation we are considering); for instance,  $G_0$  could be



## Faces of a planar graph


Consider now a plane representation  $G_0$  of a planar finite graph (note that  $G_0$  for now denotes the specific plane representation we are considering); for instance,  $G_0$  could be



Consider also a point  $p$  which is NOT one of the vertices of  $G_0$ , NOR is it found in the relative interior of any of the edges of  $G_0$ . Then  $p \in \mathbb{R}^2 \setminus G_0$ , which is an open set. Thus we can also find a maximal connected open subset of  $\mathbb{R}^2 \setminus G_0$  which contains  $p$ .

## Faces of a planar graph

Consider now a plane representation  $G_0$  of a planar finite graph (note that  $G_0$  for now denotes the specific plane representation we are considering); for instance,  $G_0$  could be




Consider also a point  $p$  which is NOT one of the vertices of  $G_0$ , NOR is it found in the relative interior of any of the edges of  $G_0$ . Then  $p \in \mathbb{R}^2 \setminus G_0$ , which is an open set. Thus we can also find a maximal connected open subset of  $\mathbb{R}^2 \setminus G_0$  which contains  $p$ .

### Definition 3

If  $G_0$  is a plane representation of a planar finite graph, then the maximal connected open subsets of  $\mathbb{R}^2 \setminus G_0$  are called the faces of  $G_0$ .

## Faces of a planar graph

Consider now a plane representation  $G_0$  of a planar finite graph (note that  $G_0$  for now denotes the specific plane representation we are considering); for instance,  $G_0$  could be



Consider also a point  $p$  which is NOT one of the vertices of  $G_0$ , NOR is it found in the relative interior of any of the edges of  $G_0$ . Then  $p \in \mathbb{R}^2 \setminus G_0$ , which is an open set. Thus we can also find a maximal connected open subset of  $\mathbb{R}^2 \setminus G_0$  which contains  $p$ .





### Definition 3

If  $G_0$  is a plane representation of a planar finite graph, then the maximal connected open subsets of  $\mathbb{R}^2 \setminus G_0$  are called the faces of  $G_0$ .


**Note** that since  $G_0$  is finite, it will have exactly one unbounded face, the area 'outside' of/surrounding the graph/diagram, while all the other faces will be bounded sets.




## Testing the definition on examples



# Testing the definition on examples (cont.)



Examples and non-examples  
of trees and forests



The forbidden subgraphs  
from Beineke's thm (see Lec 8)

## Results about the faces

### Definition 4

Let  $G_0$  be a plane representation of a planar finite graph, and let  $f_0$  be a face of  $G_0$ . We define the length  $\ell(f_0)$  of the face  $f_0$  to be

the length of a closed walk which bounds  $f_0$  (in other words, which traces the boundary of  $f_0$ ) and has minimum possible length.


# Results about the faces

## Definition 4

Let  $G_0$  be a plane representation of a planar finite graph, and let  $f_0$  be a face of  $G_0$ . We define the length  $\ell(f_0)$  of the face  $f_0$  to be

the length of a closed walk which bounds  $f_0$  (in other words, which traces the boundary of  $f_0$ ) and has minimum possible length.

**Example.** The plane graph below has 3 faces: the unbounded face  $f_1$ , the face  $f_2$  which is bounded by the 3-cycle  $b c f b$ , and the face  $f_3$  which is contained within the 4-cycle  $c d e f c$ .




# Results about the faces

## Definition 4

Let  $G_0$  be a plane representation of a planar finite graph, and let  $f_0$  be a face of  $G_0$ . We define the length  $\ell(f_0)$  of the face  $f_0$  to be

the length of a closed walk which bounds  $f_0$  (in other words, which traces the boundary of  $f_0$ ) and has minimum possible length.

**Example.** The plane graph below has 3 faces: the unbounded face  $f_1$ , the face  $f_2$  which is bounded by the 3-cycle  $b c f b$ , and the face  $f_3$  which is contained within the 4-cycle  $c d e f c$ .



– Note that a closed walk tracing the boundary of  $f_2$  is the 3-cycle  $b c f b$  (and this is also best possible), thus  $f_2$  has length 3.


# Results about the faces

## Definition 4

Let  $G_0$  be a plane representation of a planar finite graph, and let  $f_0$  be a face of  $G_0$ . We define the length  $\ell(f_0)$  of the face  $f_0$  to be

the length of a closed walk which bounds  $f_0$  (in other words, which traces the boundary of  $f_0$ ) and has minimum possible length.

**Example.** The plane graph below has 3 faces: the unbounded face  $f_1$ , the face  $f_2$  which is bounded by the 3-cycle  $b c f b$ , and the face  $f_3$  which is contained within the 4-cycle  $c d e f c$ .



- Note that a closed walk tracing the boundary of  $f_2$  is the 3-cycle  $b c f b$  (and this is also best possible), thus  $f_2$  has length 3.
- On the other hand, a closed walk tracing the entire boundary of  $f_3$  is the walk

$$g d c f e d g.$$

Note also that we cannot find any other closed walks (tracing the boundary of  $f_3$ ) which will have even shorter length, and thus  $\ell(f_3) = \text{length of the above walk} = 6$ .


# Results about the faces

## Definition 4

Let  $G_0$  be a plane representation of a planar finite graph, and let  $f_0$  be a face of  $G_0$ . We define the length  $\ell(f_0)$  of the face  $f_0$  to be

the length of a closed walk which bounds  $f_0$  (in other words, which traces the boundary of  $f_0$ ) and has minimum possible length.

**Example.** The plane graph below has 3 faces: the unbounded face  $f_1$ , the face  $f_2$  which is bounded by the 3-cycle  $b c f b$ , and the face  $f_3$  which is contained within the 4-cycle  $c d e f c$ .



- Note that a closed walk tracing the boundary of  $f_2$  is the 3-cycle  $b c f b$  (and this is also best possible), thus  $f_2$  has length 3.
- On the other hand, a closed walk tracing the entire boundary of  $f_3$  is the walk

$$g d c f e d g.$$

Note also that we cannot find any other closed walks (tracing the boundary of  $f_3$ ) which will have even shorter length, and thus  $\ell(f_3) = \text{length of the above walk} = 6$ .

- Finally, similarly we can find that  $\ell(f_1) = 7$ . (which closed walk would you use here?)

## Results about the faces

### Theorem 1

Let  $G_0$  be a plane representation of a planar finite graph, and let  $f_1, f_2, \dots, f_s$  be the faces of  $G_0$ . Then we have that

$$\sum_{i=1}^s \ell(f_i) = 2e(G_0).$$

*Observe that this result is reminiscent of the Handshaking Lemma (and its proof is also done in the same spirit).*



## Results about the faces

### Theorem 1

Let  $G_0$  be a plane representation of a planar finite graph, and let  $f_1, f_2, \dots, f_s$  be the faces of  $G_0$ . Then we have that

$$\sum_{i=1}^s \ell(f_i) = 2e(G_0).$$

*Observe that this result is reminiscent of the Handshaking Lemma (and its proof is also done in the same spirit).*

### Theorem 2 (Euler's formula, 1758)

Let  $G_0$  be a plane representation of a connected planar finite graph, and write  $n$  for the number of vertices of  $G_0$ ,  $e$  for the number of edges of  $G_0$ , and  $\tilde{f}$  for the number of faces of  $G_0$ .

## Results about the faces

### Theorem 1

Let  $G_0$  be a plane representation of a planar finite graph, and let  $f_1, f_2, \dots, f_s$  be the faces of  $G_0$ . Then we have that

$$\sum_{i=1}^s \ell(f_i) = 2e(G_0).$$

*Observe that this result is reminiscent of the Handshaking Lemma (and its proof is also done in the same spirit).*

### Theorem 2 (Euler's formula, 1758)

Let  $G_0$  be a plane representation of a connected planar finite graph, and write  $n$  for the number of vertices of  $G_0$ ,  $e$  for the number of edges of  $G_0$ , and  $\tilde{f}$  for the number of faces of  $G_0$ . Then (regardless of what exactly  $G_0$  looks like), we will always have that

$$n - e + \tilde{f} = 2.$$

## Proof of Euler's formula

We will be relying on a very famous theorem from set-theoretic topology:

### The Jordan curve theorem

A curve  $\mathcal{C}$  in  $\mathbb{R}^2$  which is **simple** and **closed** divides its complement  $\mathbb{R}^2 \setminus \mathcal{C}$  into two maximal connected open subsets: a bounded (or 'interior') region (which is enclosed by  $\mathcal{C}$ ), and an unbounded (or 'exterior') region.

Furthermore, we will proceed by induction in the number  $e$  of edges. Note that, since  $G_0$  is connected, we must have  $e = e(G_0) \geq n - 1$ .

Furthermore, we will proceed by induction in the number  $e$  of edges. Note that, since  $G_0$  is connected, we must have  $e = e(G_0) \geq n - 1$ .

**Base Case:**  $e = n - 1$ .

Furthermore, we will proceed by induction in the number  $e$  of edges. **Note that, since  $G_0$  is connected, we must have  $e = e(G_0) \geq n - 1$ .**

**Base Case:**  $e = n - 1$ . Then we have that  $G_0$  is a tree, and thus (as we also verified on examples) it will only have one face, the unbounded one. But then  $\tilde{f} = 1$ , and  $n - e + \tilde{f} = n - (n - 1) + 1 = 2$ , as we wanted.

Furthermore, we will proceed by induction in the number  $e$  of edges. **Note that, since  $G_0$  is connected, we must have  $e = e(G_0) \geq n - 1$ .**

**Base Case:**  $e = n - 1$ . Then we have that  $G_0$  is a tree, and thus (as we also verified on examples) it will only have one face, the unbounded one. But then  $\tilde{f} = 1$ , and  $n - e + \tilde{f} = n - (n - 1) + 1 = 2$ , as we wanted.

**Induction Step:** Assume now that we have already confirmed Euler's formula for connected plane graphs with  $n$  vertices and  $\tilde{e}$  edges, where  $\tilde{e}$  is some integer  $\geq n - 1$ . Consider then a connected plane graph  $\tilde{G}_0$  with  $n$  vertices and  $\tilde{e} + 1$  edges.

Furthermore, we will proceed by induction in the number  $e$  of edges. **Note that, since  $G_0$  is connected, we must have  $e = e(G_0) \geq n - 1$ .**

**Base Case:**  $e = n - 1$ . Then we have that  $G_0$  is a tree, and thus (as we also verified on examples) it will only have one face, the unbounded one. But then  $\tilde{f} = 1$ , and  $n - e + \tilde{f} = n - (n - 1) + 1 = 2$ , as we wanted.

**Induction Step:** Assume now that we have already confirmed Euler's formula for connected plane graphs with  $n$  vertices and  $\tilde{e}$  edges, where  $\tilde{e}$  is some integer  $\geq n - 1$ . Consider then a connected plane graph  $\tilde{G}_0$  with  $n$  vertices and  $\tilde{e} + 1$  edges. Then we definitely have  $e(\tilde{G}_0) > n - 1$ , and  $\tilde{G}_0$  is not a tree. Thus we can find an edge  $z_0$  of  $\tilde{G}_0$  which is NOT a bridge of  $\tilde{G}_0$  (in other words,  $z_0$  is contained in some cycle of  $\tilde{G}_0$ , which also implies that  $z_0$  is part of the boundary of at least one bounded face  $f_0$  of  $\tilde{G}_0$ ).



Furthermore, we will proceed by induction in the number  $e$  of edges. **Note that, since  $G_0$  is connected, we must have  $e = e(G_0) \geq n - 1$ .**

**Base Case:**  $e = n - 1$ . Then we have that  $G_0$  is a tree, and thus (as we also verified on examples) it will only have one face, the unbounded one. But then  $\tilde{f} = 1$ , and  $n - e + \tilde{f} = n - (n - 1) + 1 = 2$ , as we wanted.

**Induction Step:** Assume now that we have already confirmed Euler's formula for connected plane graphs with  $n$  vertices and  $\tilde{e}$  edges, where  $\tilde{e}$  is some integer  $\geq n - 1$ . Consider then a connected plane graph  $\tilde{G}_0$  with  $n$  vertices and  $\tilde{e} + 1$  edges. Then we definitely have  $e(\tilde{G}_0) > n - 1$ , and  $\tilde{G}_0$  is not a tree. Thus we can find an edge  $z_0$  of  $\tilde{G}_0$  which is NOT a bridge of  $\tilde{G}_0$  (in other words,  $z_0$  is contained in some cycle of  $\tilde{G}_0$ , which also implies that  **$z_0$  is part of the boundary of at least one bounded face  $f_0$  of  $\tilde{G}_0$** ). But then  $\tilde{G}_0 - z_0$  is a connected subgraph of  $\tilde{G}_0$  (thus it is also a plane graph), and moreover we have that

- $\tilde{G}_0 - z_0$  has the same number of vertices as  $\tilde{G}_0$  (in fact, it has the same vertices),

Furthermore, we will proceed by induction in the number  $e$  of edges. **Note that, since  $G_0$  is connected, we must have  $e = e(G_0) \geq n - 1$ .**

**Base Case:**  $e = n - 1$ . Then we have that  $G_0$  is a tree, and thus (as we also verified on examples) it will only have one face, the unbounded one. But then  $\tilde{f} = 1$ , and  $n - e + \tilde{f} = n - (n - 1) + 1 = 2$ , as we wanted.

**Induction Step:** Assume now that we have already confirmed Euler's formula for connected plane graphs with  $n$  vertices and  $\tilde{e}$  edges, where  $\tilde{e}$  is some integer  $\geq n - 1$ . Consider then a connected plane graph  $\tilde{G}_0$  with  $n$  vertices and  $\tilde{e} + 1$  edges. Then we definitely have  $e(\tilde{G}_0) > n - 1$ , and  $\tilde{G}_0$  is not a tree. Thus we can find an edge  $z_0$  of  $\tilde{G}_0$  which is NOT a bridge of  $\tilde{G}_0$  (in other words,  $z_0$  is contained in some cycle of  $\tilde{G}_0$ , which also implies that  **$z_0$  is part of the boundary of at least one bounded face  $f_0$  of  $\tilde{G}_0$** ). But then  $\tilde{G}_0 - z_0$  is a connected subgraph of  $\tilde{G}_0$  (thus it is also a plane graph), and moreover we have that

- $\tilde{G}_0 - z_0$  has the same number of vertices as  $\tilde{G}_0$  (in fact, it has the same vertices),
- and  $e(\tilde{G}_0 - z_0) = e(\tilde{G}_0) - 1 = \tilde{e}$ .

Furthermore, we will proceed by induction in the number  $e$  of edges. **Note that, since  $G_0$  is connected, we must have  $e = e(G_0) \geq n - 1$ .**

**Base Case:**  $e = n - 1$ . Then we have that  $G_0$  is a tree, and thus (as we also verified on examples) it will only have one face, the unbounded one. But then  $\tilde{f} = 1$ , and  $n - e + \tilde{f} = n - (n - 1) + 1 = 2$ , as we wanted.

**Induction Step:** Assume now that we have already confirmed Euler's formula for connected plane graphs with  $n$  vertices and  $\tilde{e}$  edges, where  $\tilde{e}$  is some integer  $\geq n - 1$ . Consider then a connected plane graph  $\tilde{G}_0$  with  $n$  vertices and  $\tilde{e} + 1$  edges. Then we definitely have  $e(\tilde{G}_0) > n - 1$ , and  $\tilde{G}_0$  is not a tree. Thus we can find an edge  $z_0$  of  $\tilde{G}_0$  which is NOT a bridge of  $\tilde{G}_0$  (in other words,  $z_0$  is contained in some cycle of  $\tilde{G}_0$ , which also implies that  **$z_0$  is part of the boundary of at least one bounded face  $f_0$  of  $\tilde{G}_0$** ). But then  $\tilde{G}_0 - z_0$  is a connected subgraph of  $\tilde{G}_0$  (thus it is also a plane graph), and moreover we have that

- $\tilde{G}_0 - z_0$  has the same number of vertices as  $\tilde{G}_0$  (in fact, it has the same vertices),
- and  $e(\tilde{G}_0 - z_0) = e(\tilde{G}_0) - 1 = \tilde{e}$ .
- Finally, observe that, by removing edge  $z_0$ , we definitely broke the cycle that contained it (in fact, any cycle that contained it), and we have also united two faces of  $\tilde{G}_0$  into one maximal connected open subset now. In other words, we have that

$$\tilde{f}(\tilde{G}_0 - z_0) = \tilde{f}(\tilde{G}_0) - 1$$

*(note that this exact relation about the faces of the original graph and its connected subgraph can be rigorously justified by the Jordan curve theorem).*

Furthermore, we will proceed by induction in the number  $e$  of edges. **Note that, since  $G_0$  is connected, we must have  $e = e(G_0) \geq n - 1$ .**

**Base Case:**  $e = n - 1$ . Then we have that  $G_0$  is a tree, and thus (as we also verified on examples) it will only have one face, the unbounded one. But then  $\tilde{f} = 1$ , and  $n - e + \tilde{f} = n - (n - 1) + 1 = 2$ , as we wanted.

**Induction Step:** Assume now that we have already confirmed Euler's formula for connected plane graphs with  $n$  vertices and  $\tilde{e}$  edges, where  $\tilde{e}$  is some integer  $\geq n - 1$ . Consider then a connected plane graph  $\tilde{G}_0$  with  $n$  vertices and  $\tilde{e} + 1$  edges. Then we definitely have  $e(\tilde{G}_0) > n - 1$ , and  $\tilde{G}_0$  is not a tree. Thus we can find an edge  $z_0$  of  $\tilde{G}_0$  which is NOT a bridge of  $\tilde{G}_0$  (in other words,  $z_0$  is contained in some cycle of  $\tilde{G}_0$ , which also implies that  **$z_0$  is part of the boundary of at least one bounded face  $f_0$  of  $\tilde{G}_0$** ). But then  $\tilde{G}_0 - z_0$  is a connected subgraph of  $\tilde{G}_0$  (thus it is also a plane graph), and moreover we have that

- $\tilde{G}_0 - z_0$  has the same number of vertices as  $\tilde{G}_0$  (in fact, it has the same vertices),
- and  $e(\tilde{G}_0 - z_0) = e(\tilde{G}_0) - 1 = \tilde{e}$ .
- Finally, observe that, by removing edge  $z_0$ , we definitely broke the cycle that contained it (in fact, any cycle that contained it), and we have also united two faces of  $\tilde{G}_0$  into one maximal connected open subset now. In other words, we have that

$$\tilde{f}(\tilde{G}_0 - z_0) = \tilde{f}(\tilde{G}_0) - 1$$

*(note that this exact relation about the faces of the original graph and its connected subgraph can be rigorously justified by the Jordan curve theorem).*

But then, by the Inductive Hypothesis, we can write

$$2 = |\tilde{G}_0 - z_0| - e(\tilde{G}_0 - z_0) + \tilde{f}(\tilde{G}_0 - z_0)$$

Furthermore, we will proceed by induction in the number  $e$  of edges. **Note that, since  $G_0$  is connected, we must have  $e = e(G_0) \geq n - 1$ .**

**Base Case:**  $e = n - 1$ . Then we have that  $G_0$  is a tree, and thus (as we also verified on examples) it will only have one face, the unbounded one. But then  $\tilde{f} = 1$ , and  $n - e + \tilde{f} = n - (n - 1) + 1 = 2$ , as we wanted.

**Induction Step:** Assume now that we have already confirmed Euler's formula for connected plane graphs with  $n$  vertices and  $\tilde{e}$  edges, where  $\tilde{e}$  is some integer  $\geq n - 1$ . Consider then a connected plane graph  $\tilde{G}_0$  with  $n$  vertices and  $\tilde{e} + 1$  edges. Then we definitely have  $e(\tilde{G}_0) > n - 1$ , and  $\tilde{G}_0$  is not a tree. Thus we can find an edge  $z_0$  of  $\tilde{G}_0$  which is NOT a bridge of  $\tilde{G}_0$  (in other words,  $z_0$  is contained in some cycle of  $\tilde{G}_0$ , which also implies that  **$z_0$  is part of the boundary of at least one bounded face  $f_0$  of  $\tilde{G}_0$** ). But then  $\tilde{G}_0 - z_0$  is a connected subgraph of  $\tilde{G}_0$  (thus it is also a plane graph), and moreover we have that

- $\tilde{G}_0 - z_0$  has the same number of vertices as  $\tilde{G}_0$  (in fact, it has the same vertices),
- and  $e(\tilde{G}_0 - z_0) = e(\tilde{G}_0) - 1 = \tilde{e}$ .
- Finally, observe that, by removing edge  $z_0$ , we definitely broke the cycle that contained it (in fact, any cycle that contained it), and we have also united two faces of  $\tilde{G}_0$  into one maximal connected open subset now. In other words, we have that

$$\tilde{f}(\tilde{G}_0 - z_0) = \tilde{f}(\tilde{G}_0) - 1$$

*(note that this exact relation about the faces of the original graph and its connected subgraph can be rigorously justified by the Jordan curve theorem).*

But then, by the Inductive Hypothesis, we can write

$$2 = |\tilde{G}_0 - z_0| - e(\tilde{G}_0 - z_0) + \tilde{f}(\tilde{G}_0 - z_0) = n - (e(\tilde{G}_0) - 1) + (\tilde{f}(\tilde{G}_0) - 1) = n - e(\tilde{G}_0) + \tilde{f}(\tilde{G}_0),$$

as we wanted.





## One immediate and very useful consequence of Euler's formula

### Important corollary of Euler's formula










Let  $G = (V, E)$  be a (connected) finite planar graph.

Then the number of faces of a planar embedding of  $G$  does NOT depend on the particular embedding we will consider. In other words, we can talk about the faces of  $G$  even before drawing  $G$  in the plane in a way that results in no edge crossings.

## Testing Euler's formula on our examples



# Testing Euler's formula on our examples (cont.)





## Other important consequences of Euler's formula

### Theorem 3

It  $G$  is a planar graph with at least 3 vertices, then

$$e(G) \leq 3|G| - 6.$$

## Other important consequences of Euler's formula

### Theorem 3

It  $G$  is a planar graph with at least 3 vertices, then

$$e(G) \leq 3|G| - 6.$$

Moreover, if  $G$  is also **triangle-free**, then

$$e(G) \leq 2|G| - 4.$$

## Other important consequences of Euler's formula

### Theorem 3

It  $G$  is a planar graph with at least 3 vertices, then

$$e(G) \leq 3|G| - 6.$$

Moreover, if  $G$  is also **triangle-free**, then

$$e(G) \leq 2|G| - 4.$$

*Justification.* It suffices to consider connected graphs (because otherwise we could just add some edges (without introducing any edge crossings, or any new cycles) in order to get a connected supergraph  $\tilde{G}$ ; but then we would have  $|\tilde{G}| = |G|$ , while  $e(G) \leq e(\tilde{G})$ , and thus if we could establish the inequalities for  $\tilde{G}$ , then the corresponding inequalities for  $G$  would also hold).

## Other important consequences of Euler's formula

### Theorem 3

It  $G$  is a planar graph with at least 3 vertices, then

$$e(G) \leq 3|G| - 6.$$

Moreover, if  $G$  is also **triangle-free**, then

$$e(G) \leq 2|G| - 4.$$

*Justification.* It suffices to consider connected graphs (because otherwise we could just add some edges (without introducing any edge crossings, or any new cycles) in order to get a connected supergraph  $\tilde{G}$ ; but then we would have  $|\tilde{G}| = |G|$ , while  $e(G) \leq e(\tilde{G})$ , and thus if we could establish the inequalities for  $\tilde{G}$ , then the corresponding inequalities for  $G$  would also hold).

Moreover, if we have any vertex in  $G$  which is a *leaf* (that is, it has degree 1), then by removing it we only decrease the quantities  $3|G| - e(G) - 6$  or  $2|G| - e(G) - 4$  (now calculated for the resulting subgraph we get).

## Other important consequences of Euler's formula

### Theorem 3

It  $G$  is a planar graph with at least 3 vertices, then

$$e(G) \leq 3|G| - 6.$$

Moreover, if  $G$  is also **triangle-free**, then

$$e(G) \leq 2|G| - 4.$$

*Justification.* It suffices to consider connected graphs (because otherwise we could just add some edges (without introducing any edge crossings, or any new cycles) in order to get a connected supergraph  $\tilde{G}$ ; but then we would have  $|\tilde{G}| = |G|$ , while  $e(G) \leq e(\tilde{G})$ , and thus if we could establish the inequalities for  $\tilde{G}$ , then the corresponding inequalities for  $G$  would also hold).

Moreover, if we have any vertex in  $G$  which is a *leaf* (that is, it has degree 1), then by removing it we only decrease the quantities  $3|G| - e(G) - 6$  or  $2|G| - e(G) - 4$  (now calculated for the resulting subgraph we get). Thus, if we could still show that these quantities are  $\geq 0$ , then the original inequalities we want would also hold.

## Other important consequences of Euler's formula

### Theorem 3

It  $G$  is a planar graph with at least 3 vertices, then

$$e(G) \leq 3|G| - 6.$$

Moreover, if  $G$  is also **triangle-free**, then

$$e(G) \leq 2|G| - 4.$$

*Justification.* It suffices to consider connected graphs (because otherwise we could just add some edges (without introducing any edge crossings, or any new cycles) in order to get a connected supergraph  $\tilde{G}$ ; but then we would have  $|\tilde{G}| = |G|$ , while  $e(G) \leq e(\tilde{G})$ , and thus if we could establish the inequalities for  $\tilde{G}$ , then the corresponding inequalities for  $G$  would also hold).

Moreover, if we have any vertex in  $G$  which is a *leaf* (that is, it has degree 1), then by removing it we only decrease the quantities  $3|G| - e(G) - 6$  or  $2|G| - e(G) - 4$  (now calculated for the resulting subgraph we get). Thus, if we could still show that these quantities are  $\geq 0$ , then the original inequalities we want would also hold.

**Important Note.** If we started with a graph  $G$  which has exactly 3 vertices, then we shouldn't be able to remove any of those vertices (*attention to our assumptions!*). But for graphs with exactly 3 vertices, we can verify the desired inequalities simply by inspection.

Based on the above, it suffices to verify the desired inequalities

for an arbitrary connected planar graph  $G$   
which has at least 3 vertices, and which has NO leaves.

Based on the above, it suffices to verify the desired inequalities

for an arbitrary connected planar graph  $G$   
which has at least 3 vertices, and which has NO leaves.

- Since we are now assuming  $G$  is connected, it will have more than one face (thus it will have at least one bounded face).



Based on the above, it suffices to verify the desired inequalities

for an arbitrary connected planar graph  $G$   
which has at least 3 vertices, and which has NO leaves.

- Since we are now assuming  $G$  is connected, it will have more than one face (thus it will have at least one bounded face).
- Moreover, the boundary of each bounded face will contain at least 3 edges, and thus the same will be true for the unbounded face.

Based on the above, it suffices to verify the desired inequalities

for an arbitrary connected planar graph  $G$   
which has at least 3 vertices, and which has NO leaves.

- Since we are now assuming  $G$  is connected, it will have more than one face (thus it will have at least one bounded face).
- Moreover, the boundary of each bounded face will contain at least 3 edges, and thus the same will be true for the unbounded face. But then the length of each face is also at least 3.

Based on the above, it suffices to verify the desired inequalities

for an arbitrary connected planar graph  $G$   
which has at least 3 vertices, and which has NO leaves.

- Since we are now assuming  $G$  is connected, it will have more than one face (thus it will have at least one bounded face).
- Moreover, the boundary of each bounded face will contain at least 3 edges, and thus the same will be true for the unbounded face. But then the length of each face is also at least 3.

Let  $f_1, f_2, \dots, f_s$  be the faces of  $G$ . Then, by Theorem 1 we have that

$$\begin{aligned} 2e(G) &= \sum_{i=1}^s \ell(f_i) \geq \sum_{i=1}^s 3 = 3\tilde{f}(G) \\ \Rightarrow \quad \tilde{f}(G) &\leq \frac{2}{3}e(G). \end{aligned} \tag{1}$$

Based on the above, it suffices to verify the desired inequalities

for an arbitrary connected planar graph  $G$   
which has at least 3 vertices, and which has NO leaves.

- Since we are now assuming  $G$  is connected, it will have more than one face (thus it will have at least one bounded face).
- Moreover, the boundary of each bounded face will contain at least 3 edges, and thus the same will be true for the unbounded face. But then the length of each face is also at least 3.

Let  $f_1, f_2, \dots, f_s$  be the faces of  $G$ . Then, by Theorem 1 we have that

$$\begin{aligned} 2e(G) &= \sum_{i=1}^s \ell(f_i) \geq \sum_{i=1}^s 3 = 3\tilde{f}(G) \\ \Rightarrow \quad \tilde{f}(G) &\leq \frac{2}{3}e(G). \end{aligned} \tag{1}$$

But then, combining the last inequality with Euler's formula, we get

$$|G| - e(G) + \frac{2}{3}e(G) \geq |G| - e(G) + \tilde{f}(G) = 2,$$

which implies the 1st inequality in Theorem 3.

Based on the above, it suffices to verify the desired inequalities

for an arbitrary connected planar graph  $G$   
which has at least 3 vertices, and which has NO leaves.

- Since we are now assuming  $G$  is connected, it will have more than one face (thus it will have at least one bounded face).
- Moreover, the boundary of each bounded face will contain at least 3 edges, and thus the same will be true for the unbounded face. But then the length of each face is also at least 3.

Let  $f_1, f_2, \dots, f_s$  be the faces of  $G$ . Then, by Theorem 1 we have that

$$\begin{aligned} 2e(G) &= \sum_{i=1}^s \ell(f_i) \geq \sum_{i=1}^s 3 = 3\tilde{f}(G) \\ \Rightarrow \quad \tilde{f}(G) &\leq \frac{2}{3}e(G). \end{aligned} \tag{1}$$

But then, combining the last inequality with Euler's formula, we get

$$|G| - e(G) + \frac{2}{3}e(G) \geq |G| - e(G) + \tilde{f}(G) = 2,$$

which implies the 1st inequality in Theorem 3.

Finally, if  $G$  is also triangle-free, then each face will instead have length at least 4, and thus we will be able to replace (1) by the stronger

$$2e(G) = \sum_{i=1}^s \ell(f_i) \geq \sum_{i=1}^s 4 = 4\tilde{f}(G).$$

Based on the above, it suffices to verify the desired inequalities

for an arbitrary connected planar graph  $G$   
which has at least 3 vertices, and which has NO leaves.

- Since we are now assuming  $G$  is connected, it will have more than one face (thus it will have at least one bounded face).
- Moreover, the boundary of each bounded face will contain at least 3 edges, and thus the same will be true for the unbounded face. But then the length of each face is also at least 3.

Let  $f_1, f_2, \dots, f_s$  be the faces of  $G$ . Then, by Theorem 1 we have that

$$\begin{aligned} 2e(G) &= \sum_{i=1}^s \ell(f_i) \geq \sum_{i=1}^s 3 = 3\tilde{f}(G) \\ \Rightarrow \quad \tilde{f}(G) &\leq \frac{2}{3}e(G). \end{aligned} \tag{1}$$

But then, combining the last inequality with Euler's formula, we get

$$|G| - e(G) + \frac{2}{3}e(G) \geq |G| - e(G) + \tilde{f}(G) = 2,$$

which implies the 1st inequality in Theorem 3.

Finally, if  $G$  is also triangle-free, then each face will instead have length at least 4, and thus we will be able to replace (1) by the stronger

$$2e(G) = \sum_{i=1}^s \ell(f_i) \geq \sum_{i=1}^s 4 = 4\tilde{f}(G).$$

Then, continuing as before, we will obtain the 2nd inequality of the theorem.

## Immediate Corollaries of Thm 3

### Corollary 1

If  $G$  is a planar (not-necessarily complete) bipartite graph with order  $|G| = n \geq 3$ , then  $G$  has at most  $2n - 4$  edges.

## Immediate Corollaries of Thm 3

### Corollary 1

If  $G$  is a planar (not-necessarily complete) bipartite graph with order  $|G| = n \geq 3$ , then  $G$  has at most  $2n - 4$  edges.

### Corollary 2

Neither of the graphs  $K_{3,3}$  and  $K_5$  is planar.



## Immediate Corollaries of Thm 3



### Corollary 1

If  $G$  is a planar (not-necessarily complete) bipartite graph with order  $|G| = n \geq 3$ , then  $G$  has at most  $2n - 4$  edges.

### Corollary 2

Neither of the graphs  $K_{3,3}$  and  $K_5$  is planar.

In other words,  $\text{cr}(K_{3,3}) \geq 1$ , and similarly  $\text{cr}(K_5) \geq 1$ .



## These examples of non-planar graphs are the most characteristic ones

Reminder from last time: Definition of the graph operation of  
'*subdivision*'

Let  $G = (V, E)$  be a graph. A subdivision of  $G$  is a graph  $H$  which satisfies the following:

- $H$  contains all the vertices of  $G$ , and perhaps a few more vertices;
- every edge  $e = \{v_1, v_2\}$  in  $G$ 
  - either appears as an edge of  $H$  as well,

## These examples of non-planar graphs are the most characteristic ones

Reminder from last time: Definition of the graph operation of  
'subdivision'

Let  $G = (V, E)$  be a graph. A subdivision of  $G$  is a graph  $H$  which satisfies the following:

- $H$  contains all the vertices of  $G$ , and perhaps a few more vertices;
- every edge  $e = \{v_1, v_2\}$  in  $G$ 
  - either appears as an edge of  $H$  as well,
  - or is replaced in  $H$  by the path  $v_1 w_1 w_2 \dots w_{s-1} w_s v_2$ , where  $w_1 (= w_1(v_1, v_2)), w_2 (= w_2(v_1, v_2)), \dots, w_s (= w_s(v_1, v_2))$  are new vertices that we add to the vertex set of  $H$  (and are only meant as internal vertices of this new path in  $H$ , that is, they will only be used in relation to vertices  $v_1$  and  $v_2$ , to 'elongate' the previous edge  $\{v_1, v_2\}$ ).

## These examples of non-planar graphs are the most characteristic ones






Reminder from last time: Definition of the graph operation of  
'subdivision'

Let  $G = (V, E)$  be a graph. A subdivision of  $G$  is a graph  $H$  which satisfies the following:

- $H$  contains all the vertices of  $G$ , and perhaps a few more vertices;
- every edge  $e = \{v_1, v_2\}$  in  $G$ 
  - either appears as an edge of  $H$  as well,
  - or is replaced in  $H$  by the path  $v_1 w_1 w_2 \dots w_{s-1} w_s v_2$ , where  $w_1 (= w_1(v_1, v_2)), w_2 (= w_2(v_1, v_2)), \dots, w_s (= w_s(v_1, v_2))$  are new vertices that we add to the vertex set of  $H$  (and are only meant as internal vertices of this new path in  $H$ , that is, they will only be used in relation to vertices  $v_1$  and  $v_2$ , to 'elongate' the previous edge  $\{v_1, v_2\}$ ).

In other words, in the second case,  $e = \{v_1, v_2\}$  is replaced by the edges  $\{v_1, w_1\}, \{w_1, w_2\}, \dots, \{w_{s-1}, w_s\}$  and  $\{w_s, v_2\}$ .

## Some subdivisions of $K_{3,3}$ and of $K_5$





## Kuratowski's theorem on planar graphs

### Theorem (Kuratowski, 1930)

A finite graph  $G$  is planar **if and only if** none of its subgraphs is a subdivision of  $K_{3,3}$  or of  $K_5$ .

# Kuratowski's theorem on planar graphs

## Theorem (Kuratowski, 1930)

A finite graph  $G$  is planar **if and only if** none of its subgraphs is a subdivision of  $K_{3,3}$  or of  $K_5$ .

## Terminology

If  $G$  contains a subgraph  $H$  which is a subdivision of  $K_{3,3}$  or of  $K_5$ , then  $H$  is called a Kuratowski subgraph of  $G$ .



# Kuratowski's theorem on planar graphs

## Theorem (Kuratowski, 1930)

A finite graph  $G$  is planar **if and only if** none of its subgraphs is a subdivision of  $K_{3,3}$  or of  $K_5$ .

## Terminology

If  $G$  contains a subgraph  $H$  which is a subdivision of  $K_{3,3}$  or of  $K_5$ , then  $H$  is called a Kuratowski subgraph of  $G$ .

With this terminology at hand, Kuratowski's theorem can be more simply stated as:

**a finite graph  $G$  is planar if and only if  
 $G$  does not have any Kuratowski subgraphs.**

