

MATH 322 – Graph Theory

Fall Term 2021

Notes for Lecture 1

Thursday, September 2

Outline for Lecture 1

- ① Introduction to the main object/concept of the course
- ② Motivating examples/problems

Outline for Lecture 1

- ① Introduction to the main object/concept of the course
- ② Motivating examples/problems

Question: What is Graph Theory? What is it about?

Outline for Lecture 1

- ① Introduction to the main object/concept of the course
- ② Motivating examples/problems

Question: What is Graph Theory? What is it about?

Most Obvious/Easy/Direct Answer: It is the area of Mathematics that studies *Graphs*!

Outline for Lecture 1

- ① Introduction to the main object/concept of the course
- ② Motivating examples/problems

Question: What is Graph Theory? What is it about?

Most Obvious/Easy/Direct Answer: It is the area of Mathematics that studies *Graphs*!

- But what is a *graph*?
- From the syllabus posted on eClass: “*Graphs are constructions that encode “links” (or lack of a link) between objects of a certain set*”.

Definition of the notion of “graph”

Let us now make this clearer by giving a mathematical definition.

We will first give the most basic definition of the main concept/object of study in this course, which is also the one we will spend most of our time on this term.

Definition of the notion of “graph”

Let us now make this clearer by giving a mathematical definition.

We will first give the most basic definition of the main concept/object of study in this course, which is also the one we will spend most of our time on this term.

To define a graph G , we need two things:

- a nonempty set of objects $V(G)$ that will be the set of vertices or nodes of G ;

Definition of the notion of “graph”

Let us now make this clearer by giving a mathematical definition.

We will first give the most basic definition of the main concept/object of study in this course, which is also the one we will spend most of our time on this term.

To define a graph G , we need two things:

- a nonempty set of objects $V(G)$ that will be the set of vertices or nodes of G ;
- a set $E(G)$ of the edges of G : namely a set containing all the pairs of vertices of G that are connected/joined in G .

Definition of the notion of “graph”

To define a graph G , we need two things:

- a nonempty set of objects $V(G)$ that will be the set of vertices or nodes of G ;
- a set $E(G)$ of the edges of G : namely a set containing all the pairs of vertices of G that are connected/joined in G .

Definition of the notion of “graph”

To define a graph G , we need two things:

- a nonempty set of objects $V(G)$ that will be the set of vertices or nodes of G ;
- a set $E(G)$ of the edges of G : namely a set containing all the pairs of vertices of G that are connected/joined in G .

Important point to pay attention to here:

- the set $V(G)$ can be any (nonempty) set we want (see examples below);

Definition of the notion of “graph”

To define a graph G , we need two things:

- a nonempty set of objects $V(G)$ that will be the set of vertices or nodes of G ;
- a set $E(G)$ of the edges of G : namely a set containing all the pairs of vertices of G that are connected/joined in G .

Important point to pay attention to here:

- the set $V(G)$ can be any (nonempty) set we want (see examples below);
- on the other hand, the set $E(G)$ of edges of G has to contain (unordered) pairs of elements of $V(G)$, namely it has to be a subset of the set of 2-element subsets of $V(G)$.

(Abstract) Example 1

If $V(G) = \{v_1, v_2, v_3, v_4\}$, then $E(G)$ has to be a subset of the set

$$\left\{ \{v_1, v_2\}, \{v_1, v_3\}, \{v_1, v_4\}, \{v_2, v_3\}, \{v_2, v_4\}, \{v_3, v_4\} \right\}.$$

(Abstract) Example 1

If $V(G) = \{v_1, v_2, v_3, v_4\}$, then $E(G)$ has to be a subset of the set

$$\left\{ \{v_1, v_2\}, \{v_1, v_3\}, \{v_1, v_4\}, \{v_2, v_3\}, \{v_2, v_4\}, \{v_3, v_4\} \right\}.$$

E.g. we could have

$$E(G) = \left\{ \{v_1, v_2\}, \{v_1, v_4\}, \{v_2, v_3\}, \{v_2, v_4\}, \{v_3, v_4\} \right\}$$

(here we just removed one 2-element subset of $V(G)$ from above).

(Abstract) Example 1

If $V(G) = \{v_1, v_2, v_3, v_4\}$, then $E(G)$ has to be a subset of the set

$$\left\{ \{v_1, v_2\}, \{v_1, v_3\}, \{v_1, v_4\}, \{v_2, v_3\}, \{v_2, v_4\}, \{v_3, v_4\} \right\}.$$

E.g. we could have

$$E(G) = \left\{ \{v_1, v_2\}, \{v_1, v_4\}, \{v_2, v_3\}, \{v_2, v_4\}, \{v_3, v_4\} \right\}$$

(here we just removed one 2-element subset of $V(G)$ from above).

Then our graph is

$$\begin{aligned} G &= (V, E) \\ &= \left(\{v_1, v_2, v_3, v_4\}, \left\{ \{v_1, v_2\}, \{v_1, v_4\}, \{v_2, v_3\}, \{v_2, v_4\}, \{v_3, v_4\} \right\} \right). \end{aligned}$$

(Abstract) Example 1 (cont.)

Our graph is

$$\begin{aligned} G &= (V, E) \\ &= \left(\{v_1, v_2, v_3, v_4\}, \left\{ \{v_1, v_2\}, \{v_1, v_4\}, \{v_2, v_3\}, \{v_2, v_4\}, \{v_3, v_4\} \right\} \right). \end{aligned}$$

(Abstract) Example 1 (cont.)

Our graph is

$$G = (V, E) \\ = \left(\{v_1, v_2, v_3, v_4\}, \left\{ \{v_1, v_2\}, \{v_1, v_4\}, \{v_2, v_3\}, \{v_2, v_4\}, \{v_3, v_4\} \right\} \right).$$

Much easier to convey all this information by drawing G !


How do we do this?

(Abstract) Example 1 (cont.)

Our graph is

$$G = (V, E) \\ = \left(\{v_1, v_2, v_3, v_4\}, \left\{ \{v_1, v_2\}, \{v_1, v_4\}, \{v_2, v_3\}, \{v_2, v_4\}, \{v_3, v_4\} \right\} \right).$$

Much easier to convey all this information by drawing G !




(Abstract) Example 1 (cont.)

Remark. We will soon see that the “names” of the vertices are not important pieces of information in **such a graph**, but rather what is most important to keep track of here is how the vertices are connected to each other. Thus, as we will explain, in many cases it will make sense to omit these “names” or labels.

(Abstract) Example 1 (cont.)

Remark. We will soon see that the “names” of the vertices are not important pieces of information in **such a graph**, but rather what is most important to keep track of here is how the vertices are connected to each other. Thus, as we will explain, in many cases it will make sense to omit these “names” or labels.

Let's draw again the graph G as an unlabelled graph:



(Abstract) Example 2

Suppose $V(G) = \{a, b, c, d, e, f, g, h\}$ and

$$E(G) = \left\{ \{a, d\}, \{a, e\}, \{b, c\}, \{b, e\}, \{b, g\}, \{c, f\}, \{d, f\}, \{d, g\}, \{g, h\} \right\}.$$

Then we can draw G as follows:




Figure: from the Harris-Hirst-Mossinghoff book

(Abstract) Example 3

Let $G = (\{1, 2, 3, 4, 5, 6\}, \{12, 14, 16, 25, 34, 36, 45, 56\})$.

We can draw G as follows:




Figure: from Bollobás' book

(Abstract) Example 3

Let $G = (\{1, 2, 3, 4, 5, 6\}, \{12, 14, 16, 25, 34, 36, 45, 56\})$.

We can draw G as follows:




Figure: from Bollobás' book

Notes. (i) Omit the curly brackets (that is, write 12 instead of $\{1, 2\}$ only when it's clear there won't be confusion.

(Abstract) Example 3

Let $G = (\{1, 2, 3, 4, 5, 6\}, \{12, 14, 16, 25, 34, 36, 45, 56\})$.

We can draw G as follows:




Figure: from Bollobás' book

Notes. (i) Omit the curly brackets (that is, write 12 instead of $\{1, 2\}$ only when it's clear there won't be confusion.

(ii) Here 12 is the same as 21, since the edges are unordered pairs.

A more complicated example




Figure: from Marcus' book

A more complicated example




Figure: from Marcus' book

Besides the fact that it's more practical ('less of an effort') to draw an unlabelled graph instead of a labelled one, we will soon also give some mathematical intuition regarding why in certain cases it makes more sense to work with unlabelled graphs.

One more example: is this a graph???




Figure: from the Balakrishnan-Ranganathan book

Answer reveal

Answer reveal

Not according to the definition we have given. Here

- there are *loops* (see at the top of the stick figure),
- and also there are sets of *multiple edges* or *parallel edges* (see the edges that form the “ears” of the stick figure).

Answer reveal

Not according to the definition we have given. Here

- there are *loops* (see at the top of the stick figure),
- and also there are sets of *multiple edges* or *parallel edges* (see the edges that form the “ears” of the stick figure).

Now let's go back to the definition of a graph that we gave: we see that the set $E(G)$ of edges of a graph G has to be

or in other words **a set of unordered pairs,**
 a set of 2-element subsets of $V(G)$.

Answer reveal

Not according to the definition we have given. Here

- there are *loops* (see at the top of the stick figure),
- and also there are sets of *multiple edges* or *parallel edges* (see the edges that form the “ears” of the stick figure).

Now let's go back to the definition of a graph that we gave: we see that the set $E(G)$ of edges of a graph G has to be

a set of unordered pairs,
or in other words **a set of 2-element subsets of $V(G)$.**

- Since $E(G)$ is a set, its elements cannot be repeated \rightsquigarrow we cannot have multiple edges in a graph G (that is, two or more edges with the same endpoints);

Answer reveal


Not according to the definition we have given. Here

- there are *loops* (see at the top of the stick figure),
- and also there are sets of *multiple edges* or *parallel edges* (see the edges that form the “ears” of the stick figure).

Now let's go back to the definition of a graph that we gave: we see that the set $E(G)$ of edges of a graph G has to be

a set of unordered pairs,
or in other words **a set of 2-element subsets of $V(G)$.**

- Since $E(G)$ is a set, its elements cannot be repeated \rightsquigarrow we cannot have multiple edges in a graph G (that is, two or more edges with the same endpoints);
- since $E(G)$ only contains 2-element subsets of $V(G)$, we cannot have loops!



Such a construction is still very closely related to graphs; in fact, it is one of the main “variations” of the concept of ‘graph’ that is also studied in Graph Theory.

In this course, we will call such a construction a *multigraph*.

Real-life examples: Transportation Networks

Real-life examples: Transportation Networks




Figure: Map of the Paris Subway-Regional Train system

The Paris Subway-Regional Train system

There is a natural way to think of this as a graph:

- the vertices are the different stations;
- an edge exists between two stations if you can get from one station to the other by taking one of the trains (one of the different lines) without stopping at any other station.

The Paris Subway-Regional Train system

There is a natural way to think of this as a graph:

- the vertices are the different stations;
- an edge exists between two stations if you can get from one station to the other by taking one of the trains (one of the different lines) without stopping at any other station.

By studying the map, we can see that we can get from any given station to any other given station; we will see that a graph that has this property is called **connected**.

The Paris Subway-Regional Train system

By studying the map, we can see that we can get from any given station to any other given station; such a graph is called connected.

Given this property, useful questions to ask here are (and these indeed come up in Graph Theory):

The Paris Subway-Regional Train system

By studying the map, we can see that we can get from any given station to any other given station; such a graph is called connected.

Given this property, useful questions to ask here are (and these indeed come up in Graph Theory):

- given two stations, what is the shortest route or *path* (a term which we will properly define soon) that takes us from one station to the other? How long is this path? (That is, how many intermediate stations would we have to pass through?)

The Paris Subway-Regional Train system

By studying the map, we can see that we can get from any given station to any other given station; such a graph is called connected.

Given this property, useful questions to ask here are (and these indeed come up in Graph Theory):

- given two stations, what is the shortest route or *path* (a term which we will properly define soon) that takes us from one station to the other? How long is this path? (That is, how many intermediate stations would we have to pass through?)
- Obviously for some pairs of stations this shortest path will be quite long (say, if you start from a station in Northwest Paris, and want to go to a station in Southeast Paris). But how long (at the most) is this shortest path for the average pair of stations/vertices (or for a large enough portion of pairs of vertices)?

Real-life examples: Social Media Platforms

Example 1: Facebook! We can think of Facebook as a graph:

Real-life examples: Social Media Platforms

Example 1: Facebook! We can think of Facebook as a graph:

- our set of vertices are the individual Facebook users (or individual accounts);

Real-life examples: Social Media Platforms

Example 1: Facebook! We can think of Facebook as a graph:

- our set of vertices are the individual Facebook users (or individual accounts);
- two users/vertices are connected by an edge if they are Facebook friends.

Real-life examples: Social Media Platforms

Example 1: Facebook! We can think of Facebook as a graph:

- our set of vertices are the individual Facebook users (or individual accounts);
- two users/vertices are connected by an edge if they are Facebook friends.

Example 2: Twitter! Similarly, we could view the Twitter platform as a graph:

Real-life examples: Social Media Platforms

Example 1: Facebook! We can think of Facebook as a graph:

- our set of vertices are the individual Facebook users (or individual accounts);
- two users/vertices are connected by an edge if they are Facebook friends.

Example 2: Twitter! Similarly, we could view the Twitter platform as a graph:

- the set of vertices here are the different accounts;

Real-life examples: Social Media Platforms

Example 1: Facebook! We can think of Facebook as a graph:

- our set of vertices are the individual Facebook users (or individual accounts);
- two users/vertices are connected by an edge if they are Facebook friends.

Example 2: Twitter! Similarly, we could view the Twitter platform as a graph:

- the set of vertices here are the different accounts;
- two accounts/vertices are connected by an edge if one of them follows the other.

Facebook vs Twitter?

Question: Are the two constructions of the exact same type?

That is, do they **both** satisfy the definition that we gave above, or is it more natural to say that one of them does not fit the definition so well?