

MATH 322 – Graph Theory

Fall Term 2021

Notes for Lecture 9

Tuesday, October 5

Connectivity; Reminder of Basic Definitions

Let $G = (V, E)$ be a connected graph.

- 1 A vertex v of G is called a cutvertex of G if we have that

$$G - v$$

is no longer connected.

- 2 More generally, a subset V' of the vertex set V of G is called a vertex cut if we have that

$$G - V'$$

is disconnected. We call it a k -vertex cut if the cardinality $|V'|$ of V' is equal to k (that is, if V' contains k vertices of G). V' is also called a separating set of vertices of G .

Connectivity; Reminder of Basic Definitions

Let $G = (V, E)$ be a connected graph.

- 1 A vertex v of G is called a cutvertex of G if we have that

$$G - v$$

is no longer connected.

- 2 More generally, a subset V' of the vertex set V of G is called a vertex cut if we have that

$$G - V'$$

is disconnected. We call it a k -vertex cut if the cardinality $|V'|$ of V' is equal to k (that is, if V' contains k vertices of G). V' is also called a separating set of vertices of G .

- 1 An edge e in G is called a bridge (or a cutedge) of G if we have that

$$G - e$$

is no longer connected.

- 2 More generally, a subset E' of the edge set E of G is called an edge cut if we have that

$$G - E'$$

is disconnected. We call it a k -edge cut if $|E'| = k$. E' is also called a separating set of edges of G .

The parameter of Edge Connectivity

The parameter $\lambda(G)$


Let G be a connected graph of order ≥ 2 . We define the edge connectivity $\lambda(G)$ of G to be the minimum cardinality of an edge cut of G .

What about Vertex Connectivity?


What about Vertex Connectivity?

*Completely analogously to edge connectivity, **we would like** to define the parameter of 'vertex connectivity' of a connected graph G to be the minimum cardinality of a vertex cut of G .*

But...
complete graphs don't have vertex cuts



But...
complete graphs don't have vertex cuts




Important Remark. Given $n \geq 2$, we have that, for every (proper) subset V' of the vertex set V of K_n , the graph $K_n - V'$ is again a complete graph (on the vertices $V \setminus V'$ now), and hence it cannot be disconnected.


\leadsto a complete graph does not have any vertex cuts.

In all other cases however...

Any connected graph of order ≥ 2 which is not a complete graph will have vertex cuts. (*Practice Exercise from last time: For each of the examples below, find a vertex cut; if possible, try to find one with smallest possible cardinality.*)



Graph G_1



Graph G_2

How to define Vertex Connectivity

Definition 1

A connected graph G is said to be k -vertex connected if G has **at least $k + 1$ vertices**, and **there does not exist** a vertex cut of G of cardinality $k - 1$.

How to define Vertex Connectivity

Definition 1

A connected graph G is said to be k -vertex connected if G has **at least $k + 1$ vertices**, and **there does not exist** a vertex cut of G of cardinality $k - 1$.

Useful Remark. Note that,

- if a connected graph H of order $n \geq 3$ has an s -vertex cut with $s \leq n - 2$,

How to define Vertex Connectivity

Definition 1

A connected graph G is said to be k -vertex connected if G has **at least $k + 1$ vertices**, and **there does not exist** a vertex cut of G of cardinality $k - 1$.

Useful Remark. Note that,

- if a connected graph H of order $n \geq 3$ has an s -vertex cut with $s \leq n - 2$,
- then it also has t -vertex cuts for each cardinality t between s and $n - 2$
(this is because if, by removing certain s vertices of H , we end up with a disconnected graph, then clearly by removing $t = s + (t - s)$ vertices of H in a suitable way (that is, by including the s vertices from the s -vertex cut we found before) we will end up again with a disconnected graph.

How to define Vertex Connectivity

Definition 1

A connected graph G is said to be k -vertex connected if G has **at least $k + 1$ vertices**, and **there does not exist** a vertex cut of G of cardinality $k - 1$.

Useful Remark. Note that,

- if a connected graph H of order $n \geq 3$ has an s -vertex cut with $s \leq n - 2$,
- then it also has t -vertex cuts for each cardinality t between s and $n - 2$
(this is because if, by removing certain s vertices of H , we end up with a disconnected graph, then clearly by removing $t = s + (t - s)$ vertices of H in a suitable way (that is, by including the s vertices from the s -vertex cut we found before) we will end up again with a disconnected graph).

Thus a k -vertex connected graph will have no s -vertex cuts for any $s < k - 1$ either. In other words, a k -vertex connected graph G is also a t -vertex connected graph for every $0 < t \leq k$.

How to define Vertex Connectivity

Definition 1

A connected graph G is said to be k -vertex connected if G has **at least $k + 1$ vertices**, and **there does not exist** a vertex cut of G of cardinality $k - 1$.

Useful Remark. Note that,

- if a connected graph H of order $n \geq 3$ has an s -vertex cut with $s \leq n - 2$,
- then it also has t -vertex cuts for each cardinality t between s and $n - 2$
(this is because if, by removing certain s vertices of H , we end up with a disconnected graph, then clearly by removing $t = s + (t - s)$ vertices of H in a suitable way (that is, by including the s vertices from the s -vertex cut we found before) we will end up again with a disconnected graph.

Thus a k -vertex connected graph will have no s -vertex cuts for any $s < k - 1$ either. In other words, a k -vertex connected graph G is also a t -vertex connected graph for every $0 < t \leq k$.

According to how we gave Definition 1, we get that

the complete graph K_n is $(n - 1)$ -vertex connected.

Vertex Connectivity (cont.)

Definition 2: the parameter $\kappa(G)$

Let G be a connected graph of order ≥ 2 . We define the vertex connectivity $\kappa(G)$ of G to be the maximum integer k such that G is k -vertex connected.

Note that, since we start with a connected graph G with at least two vertices, we will have that G is 1-vertex connected, and hence $\kappa(G) \geq 1$.

Vertex Connectivity (cont.)

Definition 2: the parameter $\kappa(G)$

Let G be a connected graph of order ≥ 2 . We define the vertex connectivity $\kappa(G)$ of G to be the maximum integer k such that G is k -vertex connected.

Note that, since we start with a connected graph G with at least two vertices, we will have that G is 1-vertex connected, and hence $\kappa(G) \geq 1$.

By convention $\kappa(K_1) = 0$.

Vertex Connectivity (cont.)

Definition 2: the parameter $\kappa(G)$

Let G be a connected graph of order ≥ 2 . We define the vertex connectivity $\kappa(G)$ of G to be the maximum integer k such that G is k -vertex connected.

Note that, since we start with a connected graph G with at least two vertices, we will have that G is 1-vertex connected, and hence $\kappa(G) \geq 1$.

By convention $\kappa(K_1) = 0$.

Very Useful Remark

Let G be a connected graph on n vertices which is different from K_n . Then the vertex connectivity $\kappa(G)$ of G coincides with the minimum cardinality of a vertex cut of G .

We can now state...

a very useful result in Connectivity which relates the parameters.

We can now state...

a very useful result in Connectivity which relates the parameters.

Theorem 1 (H. Whitney, 1932)

For every connected graph G (of order ≥ 2), we have

$$\kappa(G) \leq \lambda(G) \leq \delta(G).$$

We can now state...

a very useful result in Connectivity which relates the parameters.

Theorem 1 (H. Whitney, 1932)

For every connected graph G (of order ≥ 2), we have

$$\kappa(G) \leq \lambda(G) \leq \delta(G).$$

Remarks. 1. We have already proved (see Lecture 8) part of the theorem, that is, that $\lambda(G) \leq \delta(G)$.

We can now state...

a very useful result in Connectivity which relates the parameters.

Theorem 1 (H. Whitney, 1932)

For every connected graph G (of order ≥ 2), we have

$$\kappa(G) \leq \lambda(G) \leq \delta(G).$$

- Remarks.** 1. We have already proved (see Lecture 8) part of the theorem, that is, that $\lambda(G) \leq \delta(G)$.
2. We have seen by now that the conclusion of the theorem is true when $G = K_n$ with $n \geq 2$.

The conclusion of the theorem tested on 'extreme' examples

- In the case of the complete graph K_n ($n \geq 2$) we have seen that:
 - $\kappa(K_n) = n - 1$ by definition,

The conclusion of the theorem tested on 'extreme' examples

- In the case of the complete graph K_n ($n \geq 2$) we have seen that:
 - $\kappa(K_n) = n - 1$ by definition,
 - $\lambda(K_n) = n - 1$ by the analysis we did last time,

The conclusion of the theorem tested on 'extreme' examples

- In the case of the complete graph K_n ($n \geq 2$) we have seen that:
 - $\kappa(K_n) = n - 1$ by definition,
 - $\lambda(K_n) = n - 1$ by the analysis we did last time,
 - and clearly $\delta(K_n) = n - 1$.

The conclusion of the theorem tested on 'extreme' examples

- In the case of the complete graph K_n ($n \geq 2$) we have seen that:
 - $\kappa(K_n) = n - 1$ by definition,
 - $\lambda(K_n) = n - 1$ by the analysis we did last time,
 - and clearly $\delta(K_n) = n - 1$.
- In the case of a tree T_n on n vertices (where $n \geq 3$, because a tree on 2 vertices would just be K_2 and we have already handled this example), we have so far seen:

The conclusion of the theorem tested on 'extreme' examples

- In the case of the complete graph K_n ($n \geq 2$) we have seen that:
 - $\kappa(K_n) = n - 1$ by definition,
 - $\lambda(K_n) = n - 1$ by the analysis we did last time,
 - and clearly $\delta(K_n) = n - 1$.
- In the case of a tree T_n on n vertices (where $n \geq 3$, because a tree on 2 vertices would just be K_2 and we have already handled this example), we have so far seen:
 - that T_n must have leaves, and therefore $\delta(T_n) = 1$,

The conclusion of the theorem tested on 'extreme' examples

- In the case of the complete graph K_n ($n \geq 2$) we have seen that:
 - $\kappa(K_n) = n - 1$ by definition,
 - $\lambda(K_n) = n - 1$ by the analysis we did last time,
 - and clearly $\delta(K_n) = n - 1$.
- In the case of a tree T_n on n vertices (where $n \geq 3$, because a tree on 2 vertices would just be K_2 and we have already handled this example), we have so far seen:
 - that T_n must have leaves, and therefore $\delta(T_n) = 1$,
 - and that every edge is a bridge, and thus $\lambda(T_n) = 1$.

The conclusion of the theorem tested on 'extreme' examples

- In the case of the complete graph K_n ($n \geq 2$) we have seen that:
 - $\kappa(K_n) = n - 1$ by definition,
 - $\lambda(K_n) = n - 1$ by the analysis we did last time,
 - and clearly $\delta(K_n) = n - 1$.
- In the case of a tree T_n on n vertices (where $n \geq 3$, **because a tree on 2 vertices would just be K_2 and we have already handled this example**), we have so far seen:
 - that T_n must have leaves, and therefore $\delta(T_n) = 1$,
 - and that every edge is a bridge, and thus $\lambda(T_n) = 1$.

Regarding the latter, we will now give one more justification, along with a more general result which characterises edges that are bridges in any connected graph G .

The conclusion of the theorem tested on 'extreme' examples

- In the case of the complete graph K_n ($n \geq 2$) we have seen that:
 - $\kappa(K_n) = n - 1$ by definition,
 - $\lambda(K_n) = n - 1$ by the analysis we did last time,
 - and clearly $\delta(K_n) = n - 1$.
- In the case of a tree T_n on n vertices (where $n \geq 3$, **because a tree on 2 vertices would just be K_2 and we have already handled this example**), we have so far seen:
 - that T_n must have leaves, and therefore $\delta(T_n) = 1$,
 - and that every edge is a bridge, and thus $\lambda(T_n) = 1$.

Regarding the latter, we will now give one more justification, along with a more general result which characterises edges that are bridges in any connected graph G .

Also, what can we say about $\kappa(T_n)$?

Useful facts about connectivity

Useful facts about connectivity

Proposition 1

Let G be a connected graph of order ≥ 3 .

Useful facts about connectivity

Proposition 1

Let G be a connected graph of order ≥ 3 .

An edge e in G is NOT a bridge of G if and only if e belongs to some cycle contained in G .

Useful facts about connectivity

Proposition 1

Let G be a connected graph of order ≥ 3 .

An edge e in G is NOT a bridge of G if and only if e belongs to some cycle contained in G .

Remark. Equivalently we could say that: “ e is a bridge of G if and only if e does NOT belong to any cycle contained in G ”.

Useful facts about connectivity

Proposition 1

Let G be a connected graph of order ≥ 3 .

An edge e in G is NOT a bridge of G if and only if e belongs to some cycle contained in G .

Remark. Equivalently we could say that: “ e is a bridge of G if and only if e does NOT belong to any cycle contained in G ”.

As a consequence, we get that every edge of an acyclic graph (that is, a graph which is a tree or a forest) is a bridge.

Proof of Proposition 1

We have to show two things:

- 1 If an edge e of G belongs to some cycle contained in G , then e is NOT a bridge.

Proof of Proposition 1

We have to show two things:

- ① If an edge e of G belongs to some cycle contained in G , then e is NOT a bridge.
- ② If an edge e of G is NOT a bridge, then it belongs to some cycle contained in G .

Proof of Proposition 1

We have to show two things:

- 1 If an edge e of G belongs to some cycle contained in G , then e is NOT a bridge.
- 2 If an edge e of G is NOT a bridge, then it belongs to some cycle contained in G .

Recall Theorem 1a from Lecture 7:

Let T be a graph of order n . The following two statements are equivalent.

- (i) T is a tree (that is, T is acyclic and connected).
- (ii) T is connected and has precisely $n - 1$ edges.

Proof of Proposition 1

We have to show two things:

- ① If an edge e of G belongs to some cycle contained in G , then e is NOT a bridge.
- ② If an edge e of G is NOT a bridge, then it belongs to some cycle contained in G .

Recall Theorem 1a from Lecture 7:

Let T be a graph of order n . The following two statements are equivalent.

- (i) T is a tree (that is, T is acyclic and connected).
- (ii) T is connected and has precisely $n - 1$ edges.

We had essentially given the proof of direction ① within the proof of Theorem 1a. Let's recall the argument here:

Proof of Proposition 1

We have to show two things:

- ① If an edge e of G belongs to some cycle contained in G , then e is NOT a bridge.
- ② If an edge e of G is NOT a bridge, then it belongs to some cycle contained in G .

Recall Theorem 1a from Lecture 7:

Let T be a graph of order n . The following two statements are equivalent.

- (i) T is a tree (that is, T is acyclic and connected).
- (ii) T is connected and has precisely $n - 1$ edges.

We had essentially given the proof of direction ① within the proof of Theorem 1a.

Let's recall the argument here: *Suppose $w_1 w_2 \cdots w_k w_1$ is a k -cycle in G (for some $k \geq 3$), and e_0 is, say, the edge $\{w_1, w_2\}$. Then $G - e_0$ is connected too.*

Proof of Proposition 1

We have to show two things:

- ① If an edge e of G belongs to some cycle contained in G , then e is NOT a bridge.
- ② If an edge e of G is NOT a bridge, then it belongs to some cycle contained in G .

Recall Theorem 1a from Lecture 7:

Let T be a graph of order n . The following two statements are equivalent.

- (i) T is a tree (that is, T is acyclic and connected).
- (ii) T is connected and has precisely $n - 1$ edges.

We had essentially given the proof of direction ① within the proof of Theorem 1a. Let's recall the argument here: *Suppose $w_1 w_2 \cdots w_k w_1$ is a k -cycle in G (for some $k \geq 3$), and e_0 is, say, the edge $\{w_1, w_2\}$. Then $G - e_0$ is connected too.*

Indeed, for any two different vertices u, v in $V(G - e_0) = V(G)$, we can find a $u - v$ path in G . But then, if this path contains the edge $e_0 = \{w_1, w_2\}$, we can replace it by the path $w_2 \cdots w_k w_1$ (or, if needed, by the reverse path $w_1 w_k w_{k-1} \cdots w_2$); this sub-path would contain only different edges from e_0 , but it would take us again from w_2 to w_1 (or from w_1 to w_2).

Proof of Proposition 1

We have to show two things:

- ① If an edge e of G belongs to some cycle contained in G , then e is NOT a bridge.
- ② If an edge e of G is NOT a bridge, then it belongs to some cycle contained in G .

Recall Theorem 1a from Lecture 7:

Let T be a graph of order n . The following two statements are equivalent.

- (i) T is a tree (that is, T is acyclic and connected).
- (ii) T is connected and has precisely $n - 1$ edges.

We had essentially given the proof of direction ① within the proof of Theorem 1a. Let's recall the argument here: *Suppose $w_1 w_2 \cdots w_k w_1$ is a k -cycle in G (for some $k \geq 3$), and e_0 is, say, the edge $\{w_1, w_2\}$. Then $G - e_0$ is connected too.*

Indeed, for any two different vertices u, v in $V(G - e_0) = V(G)$, we can find a $u - v$ path in G . But then, if this path contains the edge $e_0 = \{w_1, w_2\}$, we can replace it by the path $w_2 \cdots w_k w_1$ (or, if needed, by the reverse path $w_1 w_k w_{k-1} \cdots w_2$); this sub-path would contain only different edges from e_0 , but it would take us again from w_2 to w_1 (or from w_1 to w_2). Thus, by inserting this sub-path into the original $u - v$ path **wherever the edge e_0 appears**, we would get a $u - v$ **walk** in $G - e_0$, as needed.

Proof of Proposition 1 (cont.)

Proof of direction ②. Let us now assume that e_0 is an edge of G which is NOT a bridge.

Proof of Proposition 1 (cont.)

Proof of direction ②. Let us now assume that e_0 is an edge of G which is NOT a bridge.

Then by definition we know that $G - e_0$ is a connected subgraph of G . Suppose that the endvertices of e_0 are the vertices $u_0, v_0 \in V(G) = V(G - e_0)$.

Proof of Proposition 1 (cont.)

Proof of direction ②. Let us now assume that e_0 is an edge of G which is NOT a bridge.

Then by definition we know that $G - e_0$ is a connected subgraph of G . Suppose that the endvertices of e_0 are the vertices $u_0, v_0 \in V(G) = V(G - e_0)$.

Given that $G - e_0$ is connected, we can find a $u_0 - v_0$ path P in $G - e_0$. Suppose that

P is the path $u_0 w_1 w_2 \cdots w_{k-1} v_0$

where w_1, w_2, \dots, w_{k-1} are $k - 1$ vertices of G different from both u_0 and v_0 .

Proof of Proposition 1 (cont.)

Proof of direction ②. Let us now assume that e_0 is an edge of G which is NOT a bridge.

Then by definition we know that $G - e_0$ is a connected subgraph of G . Suppose that the endvertices of e_0 are the vertices $u_0, v_0 \in V(G) = V(G - e_0)$.

Given that $G - e_0$ is connected, we can find a $u_0 - v_0$ path P in $G - e_0$. Suppose that

P is the path $u_0 w_1 w_2 \cdots w_{k-1} v_0$

where w_1, w_2, \dots, w_{k-1} are $k - 1$ vertices of G different from both u_0 and v_0 .

Note that in $G - e_0$ the endvertices u_0, v_0 of the path P are not joined by an edge, but in the original graph G they are joined by the edge e_0 . Thus in the original graph G we can 'close' the path P and get the cycle

$u_0 w_1 w_2 \cdots w_{k-1} v_0 u_0$.

Proof of Proposition 1 (cont.)

Proof of direction ②. Let us now assume that e_0 is an edge of G which is NOT a bridge.

Then by definition we know that $G - e_0$ is a connected subgraph of G . Suppose that the endvertices of e_0 are the vertices $u_0, v_0 \in V(G) = V(G - e_0)$.

Given that $G - e_0$ is connected, we can find a $u_0 - v_0$ path P in $G - e_0$. Suppose that

P is the path $u_0 w_1 w_2 \cdots w_{k-1} v_0$

where w_1, w_2, \dots, w_{k-1} are $k - 1$ vertices of G different from both u_0 and v_0 .





Note that in $G - e_0$ the endvertices u_0, v_0 of the path P are not joined by an edge, but in the original graph G they are joined by the edge e_0 . Thus in the original graph G we can 'close' the path P and get the cycle

$u_0 w_1 w_2 \cdots w_{k-1} v_0 u_0$.

In other words, the edge $e_0 = \{u_0, v_0\}$ of G (which, as was assumed, is NOT a bridge of G) is contained in some cycle of G .

Checking this on examples

Find the edges that are bridges and the edges that are not.



Useful facts about connectivity (cont.)

Proposition 2

Let G be a connected graph of order ≥ 2 , and let v be a vertex of G such that $\deg(v) = 1$.

Then v cannot be a cutvertex.

Useful facts about connectivity (cont.)

Proposition 2

Let G be a connected graph of order ≥ 2 , and let v be a vertex of G such that $\deg(v) = 1$.

Then v cannot be a cutvertex.

We had essentially given the proof of this fact too within the proof of Theorem 1a. Let us recall it here.

Useful facts about connectivity (cont.)

Proposition 2

Let G be a connected graph of order ≥ 2 , and let v be a vertex of G such that $\deg(v) = 1$.

Then v cannot be a cutvertex.

We had essentially given the proof of this fact too within the proof of Theorem 1a. Let us recall it here.

Proof. We need to show that $G - v$ is a connected graph.

Easy Case: If G is a connected graph of order 2 exactly, then $G = K_2$ and neither of its two vertices is a cutvertex (moreover, $K_2 - v$ is essentially K_1 , no matter which vertex v of K_2 we consider here).

Useful facts about connectivity (cont.)

Proposition 2

Let G be a connected graph of order ≥ 2 , and let v be a vertex of G such that $\deg(v) = 1$.

Then v cannot be a cutvertex.

We had essentially given the proof of this fact too within the proof of Theorem 1a. Let us recall it here.

Proof. We need to show that $G - v$ is a connected graph.

Easy Case: If G is a connected graph of order 2 exactly, then $G = K_2$ and neither of its two vertices is a cutvertex (moreover, $K_2 - v$ is essentially K_1 , no matter which vertex v of K_2 we consider here).

For the rest of the proof, assume that $|G| \geq 3$, and consider two different vertices u_1, w_1 of G which are also different from v . Then, since G is connected, we can find a $u_1 - w_1$ path in G .

Useful facts about connectivity (cont.)

Proposition 2

Let G be a connected graph of order ≥ 2 , and let v be a vertex of G such that $\deg(v) = 1$.

Then v cannot be a cutvertex.

We had essentially given the proof of this fact too within the proof of Theorem 1a. Let us recall it here.

Proof. We need to show that $G - v$ is a connected graph.



Easy Case: If G is a connected graph of order 2 exactly, then $G = K_2$ and neither of its two vertices is a cutvertex (moreover, $K_2 - v$ is essentially K_1 , no matter which vertex v of K_2 we consider here).

For the rest of the proof, assume that $|G| \geq 3$, and consider two different vertices u_1, w_1 of G which are also different from v . Then, since G is connected, we can find a $u_1 - w_1$ path in G .

Note that v cannot appear in this path, because otherwise it would be one of the intermediate vertices, so it would have at least two neighbours. This contradicts the assumption that $\deg_G(v) = 1$.

Thus the $u_1 - w_1$ path we have considered is also a path of the subgraph $G - v$. Since u_1, w_1 were arbitrary vertices of $G - v$, the proof is complete.

Checking this on examples



What about examples that are trees?

We now know that none of the leaves of a tree T on 2 or more vertices is a cutvertex of T .

What about examples that are trees?

We now know that none of the leaves of a tree T on 2 or more vertices is a cutvertex of T .

By the way, how many leaves **at the very least** does such a tree have? It's easy to see directly that, if $|T| = 2$, then $T = K_2$ and both its two vertices are leaves.

What about examples that are trees?

We now know that none of the leaves of a tree T on 2 or more vertices is a cutvertex of T .

By the way, how many leaves **at the very least** does such a tree have? It's easy to see directly that, if $|T| = 2$, then $T = K_2$ and both its two vertices are leaves.

But what about a tree T on 3 or more vertices? All we have said is that T contains at least one leaf;

What about examples that are trees?

We now know that none of the leaves of a tree T on 2 or more vertices is a cutvertex of T .

By the way, how many leaves **at the very least** does such a tree have? It's easy to see directly that, if $|T| = 2$, then $T = K_2$ and both its two vertices are leaves.

But what about a tree T on 3 or more vertices? All we have said is that T contains at least one leaf; **we can now improve on this.**

Corollary of Theorem 1a from Lecture 7

Let T be a tree with at least 2 vertices.

Then T contains at least two leaves (namely it has at least two vertices of degree 1).

What about examples that are trees?

We now know that none of the leaves of a tree T on 2 or more vertices is a cutvertex of T .

By the way, how many leaves **at the very least** does such a tree have? It's easy to see directly that, if $|T| = 2$, then $T = K_2$ and both its two vertices are leaves.

But what about a tree T on 3 or more vertices? All we have said is that T contains at least one leaf; **we can now improve on this.**

Corollary of Theorem 1a from Lecture 7

Let T be a tree with at least 2 vertices.

Then T contains at least two leaves (namely it has at least two vertices of degree 1).

Proof. Again, note that, if $|T| = 2$, then by simple inspection we get that both vertices of T are leaves.

What about examples that are trees?

We now know that none of the leaves of a tree T on 2 or more vertices is a cutvertex of T .

By the way, how many leaves at the very least does such a tree have? It's easy to see directly that, if $|T| = 2$, then $T = K_2$ and both its two vertices are leaves.

But what about a tree T on 3 or more vertices? All we have said is that T contains at least one leaf; we can now improve on this.

Corollary of Theorem 1a from Lecture 7

Let T be a tree with at least 2 vertices.

Then T contains at least two leaves (namely it has at least two vertices of degree 1).

Proof. Again, note that, if $|T| = 2$, then by simple inspection we get that both vertices of T are leaves.

Assume now that $|T| \geq 3$. Let's write n for the number of vertices of T , and let's write V' for the subset of the vertices of T which have degree ≥ 2 .

Let us also set $m = |V'|$; then the number of leaves in T (the number that we are interested in) is $n - m$.

Proof of the corollary (cont.)

We now recall some of our key observations from the proof of Theorem 1a:

- T is connected and has 3 or more vertices, so we can't have $\delta(T) = 0$.
- Also, we can't have $\delta(T) \geq 2$, because in such a case T would contain at least one cycle (see HW2, Problem 5(ii)).

Thus $\delta(T) = 1$, and T contains at least one leaf. In other words, $n - m \geq 1$.

Proof of the corollary (cont.)

We now recall some of our key observations from the proof of Theorem 1a:

- T is connected and has 3 or more vertices, so we can't have $\delta(T) = 0$.
- Also, we can't have $\delta(T) \geq 2$, because in such a case T would contain at least one cycle (see HW2, Problem 5(ii)).

Thus $\delta(T) = 1$, and T contains at least one leaf. In other words, $n - m \geq 1$.

By Theorem 1a again, we know that T contains exactly $n - 1$ edges, that is, $e(T) = n - 1$. Then, the Handshaking Lemma gives us

$$2n - 2 = 2e(T) = \sum_{v \in V(T)} \deg(v).$$

Proof of the corollary (cont.)

We now recall some of our key observations from the proof of Theorem 1a:

- T is connected and has 3 or more vertices, so we can't have $\delta(T) = 0$.
- Also, we can't have $\delta(T) \geq 2$, because in such a case T would contain at least one cycle (see HW2, Problem 5(ii)).

Thus $\delta(T) = 1$, and T contains at least one leaf. In other words, $n - m \geq 1$.

By Theorem 1a again, we know that T contains exactly $n - 1$ edges, that is, $e(T) = n - 1$. Then, the Handshaking Lemma gives us

$$2n - 2 = 2e(T) = \sum_{v \in V(T)} \deg(v).$$

At the same time, $\sum_{v \in V(T)} \deg(v) > \sum_{u \in V'} \deg(u) \geq 2|V'| = 2m$

(here the first inequality is strict because there is at least one leaf in T , and hence the first sum has more (positive) terms than the second one).

Proof of the corollary (cont.)

We now recall some of our key observations from the proof of Theorem 1a:

- T is connected and has 3 or more vertices, so we can't have $\delta(T) = 0$.
- Also, we can't have $\delta(T) \geq 2$, because in such a case T would contain at least one cycle (see HW2, Problem 5(ii)).

Thus $\delta(T) = 1$, and T contains at least one leaf. In other words, $n - m \geq 1$.

By Theorem 1a again, we know that T contains exactly $n - 1$ edges, that is, $e(T) = n - 1$. Then, the Handshaking Lemma gives us

$$2n - 2 = 2e(T) = \sum_{v \in V(T)} \deg(v).$$

$$\text{At the same time, } \sum_{v \in V(T)} \deg(v) > \sum_{u \in V'} \deg(u) \geq 2|V'| = 2m$$

(here the first inequality is strict because there is at least one leaf in T , and hence the first sum has more (positive) terms than the second one).

Combining the two, we see that

$$2n - 2 > 2m \Rightarrow 2(n - m) > 2 \Rightarrow n - m > 1 \Rightarrow \mathbf{n - m \geq 2}.$$

Another useful characterisation of trees

Proposition 3

Let T be a tree on at least two vertices. Then, for every two different vertices u, v of T , there is a **unique** path in T starting at u and ending at v .

In other words, in a tree there is only one way to 'travel' from one fixed vertex to another fixed vertex.

Another useful characterisation of trees

Proposition 3

Let T be a tree on at least two vertices. Then, for every two different vertices u, v of T , there is a **unique** path in T starting at u and ending at v .

In other words, in a tree there is only one way to 'travel' from one fixed vertex to another fixed vertex.

Proof. Fix two different vertices u_0, v_0 of T , and assume towards a contradiction that we can find two different paths P_1, P_2 in T which both start at u_0 and end at v_0 . Suppose that

$$P_1 : u_0 w_1 w_2 \cdots w_{k-1} v_0 \quad \text{while} \quad P_2 : u_0 z_1 z_2 \cdots z_{s-1} v_0.$$

Another useful characterisation of trees

Proposition 3

Let T be a tree on at least two vertices. Then, for every two different vertices u, v of T , there is a **unique** path in T starting at u and ending at v .

In other words, in a tree there is only one way to 'travel' from one fixed vertex to another fixed vertex.

Proof. Fix two different vertices u_0, v_0 of T , and assume towards a contradiction that we can find two different paths P_1, P_2 in T which both start at u_0 and end at v_0 . Suppose that

$$P_1 : u_0 w_1 w_2 \cdots w_{k-1} v_0 \quad \text{while} \quad P_2 : u_0 z_1 z_2 \cdots z_{s-1} v_0.$$

It's better to consider two possibilities here.

Case 1: u_0, v_0 are joined by an edge e_0 in T .

Another useful characterisation of trees

Proposition 3

Let T be a tree on at least two vertices. Then, **for every two different vertices u, v of T** , there is a **unique** path in T starting at u and ending at v .

In other words, in a tree there is only one way to 'travel' from one fixed vertex to another fixed vertex.

Proof. Fix two different vertices u_0, v_0 of T , and assume towards a contradiction that we can find two different paths P_1, P_2 in T which both start at u_0 and end at v_0 . Suppose that

$$P_1 : u_0 w_1 w_2 \cdots w_{k-1} v_0 \quad \text{while} \quad P_2 : u_0 z_1 z_2 \cdots z_{s-1} v_0.$$

It's better to consider two possibilities here.

Case 1: u_0, v_0 are joined by an edge e_0 in T . Then, one more path starting at u_0 and ending at v_0 is the (very short) path $P_0 : u_0 v_0$ (which has length 1).

Another useful characterisation of trees

Proposition 3

Let T be a tree on at least two vertices. Then, **for every two different vertices u, v of T** , there is a **unique** path in T starting at u and ending at v .

In other words, in a tree there is only one way to 'travel' from one fixed vertex to another fixed vertex.

Proof. Fix two different vertices u_0, v_0 of T , and assume towards a contradiction that we can find two different paths P_1, P_2 in T which both start at u_0 and end at v_0 . Suppose that

$$P_1 : u_0 w_1 w_2 \cdots w_{k-1} v_0 \quad \text{while} \quad P_2 : u_0 z_1 z_2 \cdots z_{s-1} v_0.$$

It's better to consider two possibilities here.

Case 1: u_0, v_0 are joined by an edge e_0 in T . Then, one more path starting at u_0 and ending at v_0 is the (very short) path $P_0 : u_0 v_0$ (which has length 1).

Since P_1 and P_2 are different paths, at least one of them will be different from P_0 , thus it will contain at least one more vertex in it; let's say that P_1 here has length ≥ 2 .

Another useful characterisation of trees

Proposition 3

Let T be a tree on at least two vertices. Then, **for every two different vertices u, v of T** , there is a **unique** path in T starting at u and ending at v .

In other words, in a tree there is only one way to 'travel' from one fixed vertex to another fixed vertex.

Proof. Fix two different vertices u_0, v_0 of T , and assume towards a contradiction that we can find two different paths P_1, P_2 in T which both start at u_0 and end at v_0 . Suppose that

$$P_1 : u_0 w_1 w_2 \cdots w_{k-1} v_0 \quad \text{while} \quad P_2 : u_0 z_1 z_2 \cdots z_{s-1} v_0.$$

It's better to consider two possibilities here.

Case 1: u_0, v_0 are joined by an edge e_0 in T . Then, one more path starting at u_0 and ending at v_0 is the (very short) path $P_0 : u_0 v_0$ (which has length 1).

Since P_1 and P_2 are different paths, at least one of them will be different from P_0 , thus it will contain at least one more vertex in it; let's say that P_1 here has length ≥ 2 .

But then the walk $u_0 w_1 w_2 \cdots w_{k-1} v_0 u_0$ contains at least 3 different vertices, and thus it is a cycle on $k + 1 \geq 3$ vertices **contained in T** .

This contradicts the assumption that T is a tree (and thus an acyclic graph).

Proof of Proposition 3 (cont.)

Case 2: u_0, v_0 are NOT joined by an edge in T .

Proof of Proposition 3 (cont.)

Case 2: u_0, v_0 are NOT joined by an edge in T . Then both paths P_1, P_2 (recall that

$$P_1 : u_0 w_1 w_2 \cdots w_{k-1} v_0 \quad \text{while} \quad P_2 : u_0 z_1 z_2 \cdots z_{s-1} v_0)$$

have length ≥ 2 . We can also assume without loss of generality that $k = \text{length}(P_1) \geq s = \text{length}(P_2)$.

Proof of Proposition 3 (cont.)

Case 2: u_0, v_0 are NOT joined by an edge in T . Then both paths P_1, P_2 (recall that

$$P_1 : u_0 w_1 w_2 \cdots w_{k-1} v_0 \quad \text{while} \quad P_2 : u_0 z_1 z_2 \cdots z_{s-1} v_0)$$

have length ≥ 2 . We can also assume without loss of generality that $k = \text{length}(P_1) \geq s = \text{length}(P_2)$.

Clearly the initial vertex in each path is the same, the vertex u_0 .

Proof of Proposition 3 (cont.)

Case 2: u_0, v_0 are NOT joined by an edge in T . Then both paths P_1, P_2 (recall that

$$P_1 : u_0 w_1 w_2 \cdots w_{k-1} v_0 \quad \text{while} \quad P_2 : u_0 z_1 z_2 \cdots z_{s-1} v_0)$$

have length ≥ 2 . We can also assume without loss of generality that $k = \text{length}(P_1) \geq s = \text{length}(P_2)$.

Clearly the initial vertex in each path is the same, the vertex u_0 .

We could also have $w_1 = z_2$, and even $w_2 = z_2$. **However**, given that the paths have been assumed different, **we can find some $i \geq 1$ such that $w_j = z_j$ for all $j \leq i - 1$ but $w_i \neq z_i$.**

Proof of Proposition 3 (cont.)

Case 2: u_0, v_0 are NOT joined by an edge in T . Then both paths P_1, P_2 (recall that

$$P_1 : u_0 w_1 w_2 \cdots w_{k-1} v_0 \quad \text{while} \quad P_2 : u_0 z_1 z_2 \cdots z_{s-1} v_0)$$

have length ≥ 2 . We can also assume without loss of generality that $k = \text{length}(P_1) \geq s = \text{length}(P_2)$.

Clearly the initial vertex in each path is the same, the vertex u_0 .

We could also have $w_1 = z_2$, and even $w_2 = z_2$. **However**, given that the paths have been assumed different, **we can find some $i \geq 1$ such that $w_j = z_j$ for all $j \leq i - 1$ but $w_i \neq z_i$.**

Note then that the walk

$$z_{i-1} = w_{i-1} \quad w_i \quad w_{i+1} \cdots w_{k-1} v_0 \quad z_{s-1} z_{s-2} \cdots z_i \quad z_{i-1}$$

is closed, and even if there are any 'repeated' vertices here (except for the 'outer' vertices which we know are equal, as we wanted), once we remove them and 'glue the remaining sequence back together' to get a shorter walk, we will end up with a cycle on 3 or more vertices.

Proof of Proposition 3 (cont.)

Case 2: u_0, v_0 are NOT joined by an edge in T . Then both paths P_1, P_2 (recall that

$$P_1 : u_0 w_1 w_2 \cdots w_{k-1} v_0 \quad \text{while} \quad P_2 : u_0 z_1 z_2 \cdots z_{s-1} v_0)$$

have length ≥ 2 . We can also assume without loss of generality that $k = \text{length}(P_1) \geq s = \text{length}(P_2)$.

Clearly the initial vertex in each path is the same, the vertex u_0 .

We could also have $w_1 = z_2$, and even $w_2 = z_2$. **However**, given that the paths have been assumed different, **we can find some $i \geq 1$ such that $w_j = z_j$ for all $j \leq i - 1$ but $w_i \neq z_i$.**

Note then that the walk

$$z_{i-1} = w_{i-1} \quad w_i \quad w_{i+1} \cdots w_{k-1} v_0 \quad z_{s-1} z_{s-2} \cdots z_i \quad z_{i-1}$$

is closed, and even if there are any 'repeated' vertices here (except for the 'outer' vertices which we know are equal, as we wanted), once we remove them and 'glue the remaining sequence back together' to get a shorter walk, we will end up with a cycle on 3 or more vertices.

Again, this will contradict the assumption that T is a tree.

Proof of Proposition 3 (cont.)

Case 2: u_0, v_0 are NOT joined by an edge in T . Then both paths P_1, P_2 (recall that

$$P_1 : u_0 w_1 w_2 \cdots w_{k-1} v_0 \quad \text{while} \quad P_2 : u_0 z_1 z_2 \cdots z_{s-1} v_0)$$

have length ≥ 2 . We can also assume without loss of generality that $k = \text{length}(P_1) \geq s = \text{length}(P_2)$.

Clearly the initial vertex in each path is the same, the vertex u_0 .

We could also have $w_1 = z_2$, and even $w_2 = z_2$. **However**, given that the paths have been assumed different, **we can find some $i \geq 1$ such that $w_j = z_j$ for all $j \leq i - 1$ but $w_i \neq z_i$.**

Note then that the walk

$$z_{i-1} = w_{i-1} \quad w_i \quad w_{i+1} \cdots w_{k-1} v_0 \quad z_{s-1} z_{s-2} \cdots z_i \quad z_{i-1}$$

is closed, and even if there are any 'repeated' vertices here (except for the 'outer' vertices which we know are equal, as we wanted), once we remove them and 'glue the remaining sequence back together' to get a shorter walk, we will end up with a cycle on 3 or more vertices.

Again, this will contradict the assumption that T is a tree.

Case 1 and **Case 2** combined complete the proof.

Cutvertices in trees

Proposition 4

Let T be a tree **of order ≥ 3** . Then T has vertices of degree ≥ 2 , and every such vertex is a cutvertex of T .

Cutvertices in trees

Proposition 4

Let T be a tree **of order ≥ 3** . Then T has vertices of degree ≥ 2 , and every such vertex is a cutvertex of T .

Proof. First of all we check that T has at least one vertex of degree 2.

Let n be the number of vertices of T . By Theorem 1a from Lecture 7, we know that $e(T) = n - 1$.

Cutvertices in trees

Proposition 4

Let T be a tree **of order ≥ 3** . Then T has vertices of degree ≥ 2 , and every such vertex is a cutvertex of T .

Proof. First of all we check that T has at least one vertex of degree 2.

Let n be the number of vertices of T . By Theorem 1a from Lecture 7, we know that $e(T) = n - 1$. Assume towards a contradiction that T contained only leaves, that is, vertices of degree 1.

Cutvertices in trees

Proposition 4

Let T be a tree **of order ≥ 3** . Then T has vertices of degree ≥ 2 , and every such vertex is a cutvertex of T .

Proof. First of all we check that T has at least one vertex of degree 2.

Let n be the number of vertices of T . By Theorem 1a from Lecture 7, we know that $e(T) = n - 1$. Assume towards a contradiction that T contained only leaves, that is, vertices of degree 1.

Then we would have

$$\sum_{v \in V(T)} \deg(v) = \sum_{v \in V(T)} 1 = |T| = n < 2(n - 1) = 2e(T),$$

where the (strict) inequality holds because $n \geq 3$. This contradicts the Handshaking Lemma.

Cutvertices in trees

Proposition 4

Let T be a tree **of order ≥ 3** . Then T has vertices of degree ≥ 2 , and every such vertex is a cutvertex of T .

Proof. First of all we check that T has at least one vertex of degree 2.

Let n be the number of vertices of T . By Theorem 1a from Lecture 7, we know that $e(T) = n - 1$. Assume towards a contradiction that T contained only leaves, that is, vertices of degree 1.

Then we would have

$$\sum_{v \in V(T)} \deg(v) = \sum_{v \in V(T)} 1 = |T| = n < 2(n - 1) = 2e(T),$$

where the (strict) inequality holds because $n \geq 3$. This contradicts the Handshaking Lemma.

Thus T must contain at least one vertex of degree 2.

Proof of Proposition 4 (cont.)

Proposition 4

Let T be a tree **of order ≥ 3** . Then T has vertices of degree ≥ 2 , and every such vertex is a cutvertex of T .

Proof of Proposition 4 (cont.)

Proposition 4

Let T be a tree **of order ≥ 3** . Then T has vertices of degree ≥ 2 , and every such vertex is a cutvertex of T .

Consider now a vertex v_0 of T which has degree ≥ 2 . We will show that $T - v_0$ is a disconnected subgraph.

Proof of Proposition 4 (cont.)

Proposition 4

Let T be a tree **of order ≥ 3** . Then T has vertices of degree ≥ 2 , and every such vertex is a cutvertex of T .

Consider now a vertex v_0 of T which has degree ≥ 2 . We will show that $T - v_0$ is a disconnected subgraph.

In fact, we will show that **the number of connected components of $T - v_0$ is equal to the number of neighbours of v_0 in T** , and thus that $T - v_0$ has at least 2 connected components.

Proof of Proposition 4 (cont.)

Proposition 4

Let T be a tree **of order ≥ 3** . Then T has vertices of degree ≥ 2 , and every such vertex is a cutvertex of T .

Consider now a vertex v_0 of T which has degree ≥ 2 . We will show that $T - v_0$ is a disconnected subgraph.

In fact, we will show that the number of connected components of $T - v_0$ is equal to the number of neighbours of v_0 in T , and thus that $T - v_0$ has at least 2 connected components.

Indeed, let u_1, w_1 be two different neighbours of v_0 in T . Then the path $P_0 : u_1 v_0 w_1$ is a path of length 2 contained in T which starts at u_1 and ends at w_1 .

Proof of Proposition 4 (cont.)

Proposition 4

Let T be a tree **of order ≥ 3** . Then T has vertices of degree ≥ 2 , and every such vertex is a cutvertex of T .

Consider now a vertex v_0 of T which has degree ≥ 2 . We will show that $T - v_0$ is a disconnected subgraph.

In fact, we will show that **the number of connected components of $T - v_0$ is equal to the number of neighbours of v_0 in T** , and thus that $T - v_0$ has at least 2 connected components.

Indeed, let u_1, w_1 be two different neighbours of v_0 in T . Then the path $P_0 : u_1 v_0 w_1$ is a path of length 2 contained in T which starts at u_1 and ends at w_1 .

Recall now Proposition 3 that we just showed: it implies that there is NO OTHER path in T which starts at u_1 and ends at w_1 . In other words, we cannot find a path P_1 in $T - v_0$ which starts at u_1 and ends at w_1 (*because such a path would also be a path in T , and it would be different from the path P_0*).

Proof of Proposition 4 (cont.)

Proposition 4

Let T be a tree **of order ≥ 3** . Then T has vertices of degree ≥ 2 , and every such vertex is a cutvertex of T .

Consider now a vertex v_0 of T which has degree ≥ 2 . We will show that $T - v_0$ is a disconnected subgraph.




In fact, we will show that the number of connected components of $T - v_0$ is equal to the number of neighbours of v_0 in T , and thus that $T - v_0$ has at least 2 connected components.

Indeed, let u_1, w_1 be two different neighbours of v_0 in T . Then the path $P_0 : u_1 v_0 w_1$ is a path of length 2 contained in T which starts at u_1 and ends at w_1 .

Recall now Proposition 3 that we just showed: it implies that there is NO OTHER path in T which starts at u_1 and ends at w_1 . In other words, we cannot find a path P_1 in $T - v_0$ which starts at u_1 and ends at w_1 (*because such a path would also be a path in T , and it would be different from the path P_0*).

We conclude that each of the neighbours of v_0 in T will end up in a different connected component of $T - v_0$.

Checking these on examples



Question. In each of the graphs above, which of the vertices are cutvertices, and which are not?

Back to Whitney's Theorem tested on 'extreme' examples

- In the case of the complete graph K_n ($n \geq 2$) we have seen that:
 - $\kappa(K_n) = n - 1$ by definition,
 - $\lambda(K_n) = n - 1$ by the analysis we did last time,
 - and clearly $\delta(K_n) = n - 1$.
- In the case of a tree T_n on n vertices (where $n \geq 3$, because a tree on 2 vertices would just be K_2 and we have already handled this example), we have so far seen:

Back to Whitney's Theorem tested on 'extreme' examples

- In the case of the complete graph K_n ($n \geq 2$) we have seen that:
 - $\kappa(K_n) = n - 1$ by definition,
 - $\lambda(K_n) = n - 1$ by the analysis we did last time,
 - and clearly $\delta(K_n) = n - 1$.
- In the case of a tree T_n on n vertices (where $n \geq 3$, because a tree on 2 vertices would just be K_2 and we have already handled this example), we have so far seen:
 - that T_n must have leaves, and therefore $\delta(T_n) = 1$,

Back to Whitney's Theorem tested on 'extreme' examples

- In the case of the complete graph K_n ($n \geq 2$) we have seen that:
 - $\kappa(K_n) = n - 1$ by definition,
 - $\lambda(K_n) = n - 1$ by the analysis we did last time,
 - and clearly $\delta(K_n) = n - 1$.
- In the case of a tree T_n on n vertices (where $n \geq 3$, because a tree on 2 vertices would just be K_2 and we have already handled this example), we have so far seen:
 - that T_n must have leaves, and therefore $\delta(T_n) = 1$,
 - and that every edge is a bridge, and thus $\lambda(T_n) = 1$.

Back to Whitney's Theorem tested on 'extreme' examples

- In the case of the complete graph K_n ($n \geq 2$) we have seen that:
 - $\kappa(K_n) = n - 1$ by definition,
 - $\lambda(K_n) = n - 1$ by the analysis we did last time,
 - and clearly $\delta(K_n) = n - 1$.
- In the case of a tree T_n on n vertices (where $n \geq 3$, because a tree on 2 vertices would just be K_2 and we have already handled this example), we have so far seen:
 - that T_n must have leaves, and therefore $\delta(T_n) = 1$,
 - and that every edge is a bridge, and thus $\lambda(T_n) = 1$.
 - Finally, such a tree will have vertices of degree ≥ 2 , and each such vertex will be a cutvertex. Thus we will have $\kappa(T_n) = 1$.