

Simulation of laser-induced rectification in a nano-scale diode



Presenter: XJ Xu

Daniel Kidd,¹ Xiaojia Xu,¹ Cody Covington,¹ Kazuyuki Watanabe,² Kalman Varga¹

¹*Department of Physics and Astronomy, Vanderbilt University*

²*Department of Physics, Tokyo University of Science*

Mar 6, 2018

Modern age of electronics

ENIAC (1956)

- 20,000 vacuum-tubes
- 5,000 Hz operations on 10 digit numbers
- Power: 150 kW

Sunway TaihuLight (2017)


- 40,960 SW26010 processors
- 3.06 TFLOPS per processor
- Power: 15 MW

Lower power consumption by scale, higher reliability, and more intuitive circuit design.




Pushing for the petahertz range

- Limited electron transport velocity in semiconductor transistors.
- Recent interest in electron photoemission from metal nanotips.
- Ultrafast laser-guidance of electrons.
- This inspired research pointing back to vacuum transport to take advantage of higher vacuum transport velocities.




Diamant, E. Halahmi, L. Kronik, J. Levy, R. Naaman, and J. Roulston, Appl. Phys. Lett. **92**, 262903 (2008).
J.-W. Han, J. S. Oh, and M. Meyyappan, Appl. Phys. Lett. **100**, 213505 (2012).
J.-W. Han, D.-I. Moon, and M. Meyyappan, Nano Lett. **17**, 2146 (2017).

Jellium model



Optical microscopic image of opposing tungsten nanotips.
Higuchi et al. (2015)



$$D = 30 \text{ \AA}$$

Homogeneous laser field

The form of the laser field is a variation of the smooth turn-on pulse.

- Wavelength of 780 nm
- Polarized parallel to the axis of symmetry

$$\mathbf{E}_{\text{laser}}(t) = \begin{cases} -\mathbf{E}_0 \sin\left(\frac{\pi t}{2T_r}\right) \sin(\omega t + \varphi), & \text{if } 0 \leq t \leq T_r, \\ -\mathbf{E}_0 \sin(\omega t + \varphi), & \text{otherwise,} \end{cases}$$

Where φ is the phase of the laser and T_r is the ramping time.
The vector potential is then:

$$\mathbf{A}(t) = - \int_0^t \mathbf{E}_{\text{laser}}(t') dt'.$$

RT-TDDFT

We solve the time-dependent Kohn-Sham equation

$$i\hbar \frac{\partial \phi_j(\mathbf{r}, t)}{\partial t} = \left[-\frac{\hbar^2}{2m} \nabla^2 + V_{\text{KS}}[\rho](\mathbf{r}, t) \right] \phi_j(\mathbf{r}, t)$$


The Kohn-Sham potential is a functional of the density and consists of 3 terms:

- 1) the Hartree potential
- 2) the exchange-correlation potential (using LDA)
- 3) the external potential (includes V_{laser})

$$V_{\text{laser}}(\mathbf{r}, t) = \frac{1}{2m} |\mathbf{A}(t)|^2 - \frac{i\hbar}{m} \mathbf{A}(t) \cdot \nabla$$

Time propagation using split-operator.

$$\Psi(t + \Delta t) = e^{i\hat{H}\Delta t/\hbar} \Psi(t)$$



Induced
current \mathbf{j} and
flux Φ

The induced oscillating current is:

$$\mathbf{j}(\mathbf{r}, t) = 2 \sum_{j=1}^N \frac{\hbar}{2mi} (\phi_j^* \nabla \phi_j - \phi_j \nabla \phi_j^*) + \frac{1}{m} \mathbf{A} \rho$$

The resulting flux at the plane bisecting the two edges of the jellium model:


$$\Phi(z_0, t) = \int \mathbf{j}(\mathbf{r}, t) \delta(z - z_0) d\mathbf{r}$$

- The flux at the boundary of is integrated with respect to time


$$N_{\text{tr}}(t) = \int_0^t \Phi(t') dt'$$


- We find the **probability density transferred N_{tr}**
- The **probability density transfer rate k_{tr}** is determined with linear regression of $t \leq 15$ fs
- k_{tr} levels off $15 \leq t \leq 50$
- Parameters:
 - Separation distance: 30 Å
 - Field intensity: 1.33×10^{13} W/cm²
 - Ramping time $T_r = 2.48$ fs

Calculating the results



20 Å diode with 30 Å separation






Phase shift dependence

- Symmetric cylinder jellium model with ramping time (control run)
 $T_r = 2.48$ fs
- Diode jellium model with ramping time
 $T_r = 2.48$ fs
- Diode jellium model with ramping time
 $T_r = 9.94$ fs


- In fig (a), the short ramping time makes the system sensitive to phase shifts.
- Positive offset in fig (b) as opposed to negligible offset in fig (a)
- Phase shift becomes insignificant if the ramping time is long enough.

Dependence of separation distance, D



- We simply change the size of the computational box to vary D
- Parameters kept constant:
 $I = 1.33 \times 10^{13} \text{ W/cm}^2$
 $T_r = 9.94 \text{ fs}$
- Results from simulations with phases of 0 and π were averaged to **eliminate phase dependence**
- Approximately exponential decay

Dependence of intensity, I



- Phase shift dependence is eliminated in a similar fashion
- Parameters kept constant:
 $D = 30 \text{ \AA}$
 $T_r = 9.94 \text{ fs}$
- For sufficiently high intensities, the local near-field enhancement becomes negligible.

Conclusions

- We have computationally demonstrated laser-induced rectification by simulating the effects of increased electron emission due to near-field enhancement within a periodic jellium system with geometrical asymmetry.
- Opening new doors for new nanoscale “vacuum-tube-based” devices
- Approximately exponential increase in transport rate for small separations
- Convergence of transport rate for higher laser intensities
- The paper is available at:

Daniel Kidd, Xiaojia Xu, Cody Covington, Kazuyuki Watanabe, Kalman Varga.

Simulation of laser-induced rectification in a nanoscale vacuum-tube diode

Journal of Applied Physics **123**, 054501 (2018)

Acknowledgements

This work was supported by the National Science Foundation (NSF) under Grant Nos. OISE 1261117, PHY 1314463, and OISE 1613549.

My trip to LA was made possible by the support from the College of Arts and Science and the Department of Physics.

Many thank to Daniel Kidd, Cody Covington, Kalman Varga, Kazuyuki Watanabe, Yasumitsu Suzuki and Elena Silaeva.



VANDERBILT
College of Arts and Science

