




c

Sweden	15.4% [5.7, 83.1]	10.3% [1.6, 64.4]	1.6% [1.1, 5.8]	0.3% [0.0, 6.3]	9.3% [0.2, 62.9]	4.4% [0.1, 24.8]
United Kingdom	27.0% [11.2, 78.8]	12.3% [1.7, 87.7]	3.3% [0.9, 6.3]	-0.5% [-0.2, 92.5]	11.1% [0.0, 87.1]	5.8% [-0.2, 30.1]
Australia	14.6% [10.4, 86.7]	2.3% [1.2, 74.7]	2.0% [1.0, 4.8]	0.0% [0.0, 16.3]	0.4% [0.1, 73.6]	1.0% [0.0, 27.3]
China	2.6% [1.9, 21.7]	5.7% [2.7, 74.7]	2.0% [1.4, 10.8]	-0.1% [-0.2, 68.8]	3.8% [0.1, 73.5]	2.3% [0.2, 23.8]
Japan	7.1% [2.3, 29.4]	1.0% [1.5, 84.3]	3.0% [1.4, 6.3]	-0.2% [-0.1, 56.8]	-1.4% [-0.8, 83.8]	2.4% [-0.1, 22.0]
New Zealand	20.2% [13.0, 91.0]	2.3% [1.1, 70.0]	2.0% [1.1, 7.9]	0.0% [-0.1, 53.7]	0.3% [0.0, 68.0]	0.6% [-0.3, 29.8]
Thailand	26.5% [11.7, 47.1]	17.8% [3.7, 47.5]	1.1% [1.4, 9.0]	16.1% [0.0, 36.4]	16.7% [0.3, 46.2]	1.3% [-0.3, 20.1]
Brazil	22.6% [12.1, 76.5]	12.2% [2.0, 75.2]	1.1% [0.9, 5.3]	5.8% [0.0, 67.2]	11.3% [0.3, 74.4]	-6.3% [-0.5, 26.0]
United States	14.7% [10.9, 89.6]	4.4% [1.7, 73.9]	1.3% [1.1, 5.8]	0.0% [0.0, 15.3]	3.3% [0.3, 72.3]	2.1% [0.2, 28.9]
	Routine schedule timeliness	Infant-exposure package	Pregnancy Tdap scale-up	Adolescent booster scale-up	Close-contact adult adjunct	Targeted high-risk PEP

Programme strategy

Relative reduction vs current practice (%)