

NATIONAL SENIOR CERTIFICATE

GRADE 12

JUNE 2023

TECHNICAL MATHEMATICS P2

MARKS: 150

TIME: 3 hours

This question paper consists of 16 pages, including 2-page information sheet and a special answer book.

INSTRUCTIONS AND INFORMATION


Read the following instructions carefully before answering the questions.

- 1. This question paper consists of ELEVEN questions.
- 2. Answer ALL the questions in the SPECIAL ANSWER BOOK provided.
- 3. Clearly show ALL calculations, diagrams, graphs, et cetera which you have used in determining the answers.
- 4. Answers only will NOT necessarily be awarded full marks.
- 5. You may use an approved scientific calculator (non-programmable and non-graphical) unless stated otherwise.
- 6. If necessary, round off your answers to TWO decimal places, unless stated otherwise.
- 7. Diagrams are NOT necessarily drawn to scale.
- 8. An information sheet with formulae is included at the end of the question paper.
- 9. Write neatly and legibly.

In the diagram below ABC is a triangle with vertices A(-2; 5); B(8; p) and C.

The inclination angle of AB is 135°.

AC is parallel to the *y*-axis and BC is parallel to the *x*-axis.



1.1 Determine the gradient of AB. (2)

1.2 Show that p = -5. (3)

1.3 Determine the coordinates of M, the midpoint of AB. (2)

1.4 Write down the equation of BC. (1)


1.5 Write down the coordinates of C. (2)

1.6 Show that $CM \perp AB$. (3)

1.7 Determine the equation of the straight-line parallel to CM and which passes through point A. (3)


[16]

2.1 The diagram below shows the circle with equation $x^2 + y^2 = 20$. A is a contact point of a tangent to the circle.



- 2.1.1 Write down the radius of the circle in simplified surd form. (1)
- 2.1.2 Determine the equation of the tangent to the circle at point A in the form y = ... (4)
- 2.1.3 Write down the coordinates of another point where the line AO intersects with the circle. (2)
- 2.2 Sketch the graph of $\frac{x^2}{16} + \frac{y^2}{25} = 1$. Clearly indicate the intercepts. (3)

3.1 In the diagram below, P(8;k) is a point on the Cartesian plane. OP forms a reflex angle q with the positive x-axis with OP equal to 10 units.



Determine the value of the following, WITHOUT using a calculator:

$$3.1.1 \quad \cos\theta$$
 (1)

$$3.1.2 k$$
 (3)

$$\frac{\tan \theta}{\csc \theta} \tag{3}$$

3.2 Determine the values of
$$x$$
, if $3\cos x - 1 = -1.5$ for $x \in [0^\circ; 360^\circ]$ [11]

4.1 Simplify:

$$(1+\cos x)(1-\cos x) \tag{2}$$

4.2 Simplify:

$$\frac{\cos^2(2\pi - x)\tan^2 x}{\sin(180^\circ + x)\csc(180^\circ - x)}$$
(6)

4.3 Prove that:

$$\cot x + \tan x = \csc x \cdot \sec x \tag{4}$$

[12]

Copyright reserved


Given the functions defined by $f(x) = \cos 2x$ and $g(x) = \sin(x-30^\circ)$ for $x \in [0^\circ; 180^\circ]$.

- 5.1 Write down the period of f. (1)
- 5.2 Write down the amplitude of g. (1)
- On the same axes given in your SPECIAL ANSWER BOOK draw the graphs of f and g. Clearly show the turning points, endpoints, and the intercepts with the axes. (8)
- 5.4 Use your graphs to determine for which values of x is:

$$5.4.1 f(x) \le 0 (2)$$

5.4.2 $f(x).g(x) \ge 0$ in the second quadrant (2) [14]

- 6.1 Write down the cosine rule for $\triangle PQR$. (1)
- 6.2 In the diagram below, AC is the diameter of the circle ABC. AC is produced to D such that DC = CD = 6 cm. $A = 54^{\circ}$



Determine:


6.2.1 The size of
$$\stackrel{\wedge}{ABC}$$
, stating a reason (2)

6.2.2 The size of
$$\stackrel{\wedge}{BCD}$$
, stating a reason (2)

6.2.5 The area of
$$\triangle ABC$$
 (3) [14]

Copyright reserved

In the diagram below, ABC is a circle with centre O. OD = 3 cm, BC = 11 cm and $OD \perp BC$. BO, AO and OC are joined.




- Determine the length of BD, stating a reason. 7.1 (2)
- 7.2 Calculate the length of OB. (2)
- 7.3 Show that $\triangle ABD \equiv \triangle ACD$. (4)
- Calculate the size of \hat{B}_1 . 7.4 (2)
- Hence, calculate the size A, stating reasons. 7.5 (4)

[14]

In the diagram below, a circle with centre O is given.

HE and HD are tangents to the circle such that $\hat{E}_1=49^\circ$ and $\hat{F}_1=65^\circ$.



8.1 Give a reason why HD = HE. (1)

8.2 Determine, stating reasons, the size of the following angles:

8.2.1
$$\stackrel{\wedge}{\rm D}_1$$
 (2)

$$8.2.2 \quad \stackrel{\wedge}{\mathrm{D}_2}$$
 (2)


$$8.2.3$$
 DEF (2)

$$8.2.4$$
 \hat{G} (2)

$$8.2.5$$
 FDK (2)

8.3 Show, stating reasons, whether EHDF is cyclic. (3) [14]

In \triangle ABC below, AB = 20 cm, BC = 17,34 cm and AC = 7 cm. D and E are points on the sides of the triangle such that DE || AC. AD: DB = 2:3




- 9.1 Determine the lengths of AD and DB. (3)
- 9.2 Calculate, stating reasons, the length of BE. (3)
- 9.3 Prove, stating reasons, that $\triangle BDE \parallel \triangle BAC$. (3)
- 9.4 Hence, determine the length of DE. (3) [12]

A wheel with a diameter 250 mm, has a circumferential velocity of 108 kilometres per hour.


10.1 Convert 108 km/h to m/s. (2) 10.2 Determine the rotational frequency of the wheel in seconds. (5) 10.3 Determine the angular velocity of the wheel in seconds. (3) 10.4 Determine the distance, in km, a point on the wheel will cover in 10 min. (3) 10.5 Determine how long it will take the wheel to make 20 revolutions. (2) [15]

11.1 In diagram below, RSTV is a square with sides 8 cm. RT is an arc of the sector RST.





- 11.1.1 Determine the length of arc RT. (3)
- 11.1.2 Determine the area of sector RST. (3)
- 11.1.3 Hence, calculate the area of the shaded area. (3)
- 11.2 The diagram below is a minor segment of a circle with height 4 cm and chord 10 cm.




Determine the length of the radius of the circle.

(5)

Copyright reserved

11.3 The ordinates in the irregular figure are 9,42; 8,14; 6,42; 7,1; 8 and 7 cm as indicated in the diagram below. The area of the irregular figure is 113,61 cm².



Determine the width of the equal parts on the horizontal axis.

(4)

[18]

TOTAL: 150

INFORMATION SHEET

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$x = -\frac{b}{2a}$$

$$y = \frac{4ac - b^2}{4a}$$

$$a^x = b \Leftrightarrow x = \log_a b$$

$$a > 0$$
, $a \ne 1$ en $b > 0$

$$A = P(1+ni)$$

$$A = P(1-ni)$$

$$A = P(1+i)^n$$

$$A = P(1-i)^n$$

$$i_{eff} = \left(1 + \frac{i^m}{m}\right)^m - 1$$

$$f'(x) = \lim_{h \to 0} \frac{f(x+h) - f(x)}{h}$$

$$\int x^n dx = \frac{x^{n+1}}{n+1} + C \qquad , \quad n \neq -1$$

$$\int kx^n dx = k \cdot \frac{x^{n+1}}{n+1} + C \quad , \quad n \neq -1$$

$$\int \frac{1}{x} dx = \ln(x) + C, \quad x > 0$$

$$\int \frac{k}{x} dx = k \cdot \ln(x) + C, \quad x > 0$$

$$\int a^x dx = \frac{a^x}{\ln a} + C \quad , \quad a > 0$$

$$\int ka^{nx}dx = k \cdot \frac{a^{nx}}{n \ln a} + C \quad , \quad a > 0$$

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

$$\mathbf{M}\left(\frac{x_1+x_2}{2}; \frac{y_1+y_2}{2}\right)$$

$$y = mx + c$$

$$y = mx + c$$
 $y - y_1 = m(x - x_1)$ $m = \frac{y_2 - y_1}{x_2 - x_1}$

$$m = \tan \theta$$

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$$

In ΔABC:

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc \cos A$$

Area =
$$\frac{1}{2}ab.\sin C$$

$$\sin^2\theta + \cos^2\theta = 1$$

$$\tan^2\theta + 1 = \sec^2\theta$$

$$1 + \cot^2 \theta = \csc^2 \theta$$

 $\pi rad = 180^{\circ}$

Angular velocity = $\omega = 2\pi n$ where n = rotation frequency

Angular velocity = $\omega = 360^{\circ}n$ where n = rotation frequency

Circumferential velocity = $v = \pi Dn$ where D = diameter and n = rotation frequency

Circumferential velocity = $v = \omega r$ where ω = Angular velocity and r = radius

Arc length $s = r\theta$ where r = radius and $\theta = \text{central}$ angle in radians

Area of a sector = $\frac{rs}{2}$ where r = radius and s = arc length

Area of a sector = $\frac{r^2\theta}{2}$ where r = radius and θ = central angle in radians

 $4h^2 - 4dh + x^2 = 0$ where h = height of segment, d = diameter of the circle and x = length of chord

 $A_T = a(m_1 + m_2 + m_3 + ... + m_{n-1})$ where a = width of equal parts, $m_1 = \frac{o_1 + o_2}{2}$ and n = number of ordinates

OR

$$\mathbf{A}_{\mathrm{T}} = a \left(\frac{o_1 + o_n}{2} + o_2 + o_3 + o_4 + \ldots + o_{n-1} \right) \qquad \text{where } a = \text{width of equal parts, } o_i = i^{th} \text{ ordinate and } n = \text{number of ordinates}$$