


Source-aligned RNA reconstruction: baseline programs and DFMO

68-hour mouse CD4 T-cell cultures | TMM + limma-voom | animal-blocked models
A: 3 WT animals, Th17p / Th17n / iTreg; B: 4 WT + 3 JMJD3 KO, Th17n / iTreg
Source-aligned reconstruction; exact author gene sets and supplementary-number concordance are unavailable.






How to read this plate

A: genes selected only by vehicle lineage contrasts; no gene z-scoring. Lines connect animal-matched cultures.

B-D: each step is a gene, not an animal. Positive x: higher RNA after DFMO.


Programs use the same study controls. B-study primary program balances WT and KO baseline contrasts equally. No conversion or fate is measured.



Gene selection: BH $q \leq 0.05$ and $|\log_2 FC| \geq \log_2(1.5)$. Curves summarize selected-gene distributions, not independent replicates. A differs from the published 3,414-gene PCA unless exact source selection is recovered. All tested genes and sensitivity partitions are retained.

JMJD3 dependence: direct genotype × DFMO contrasts

GSE162382 | 68-hour bulk RNA | 4 WT + 3 conditional-KO animal labels, paired treatment cultures
Endpoint: (DFMO – vehicle)KO – (DFMO – vehicle)WT on the fitted log₂ RNA-abundance scale.
Dots: estimates; bars: 95% moderated model intervals. Filled dots: BH q ≤ 0.05 across all tested genes in that contrast.



Positive: the KO response is more positive (or less negative); negative: more negative (or less positive).
An interaction is not a mediation test. Open points do not establish equivalence. Intervals are conditional on this model and source labels;
physical specimen independence is unverified. The twelve genes were fixed in the earlier descriptive contract.