Public-data results and biological rationale

Same source lineage; exposed descriptive analyses

MEASURED: A12 candidate nuclei

12 paired patients, LUAD minus normal
IL1RAP detection: +2.45 percentage points
Intensity among detected nuclei: -0.196 log2 units
IL1R1 / IL1RAP co-detection: +1.01 points
After common-depth standardization: +0.16 points

MEASURED: Visium RNA organization

10 paired patients, LUAD minus AAH
IL1B / CLDN4 one-hop statistic: +0.0279
With broad RNA-composition covariates: +0.0095
Lattice adjacency only; spots contain mixtures
Patient-level description, not a spot significance test

WHAT THESE RESULTS CHANGE

The aggregate RNA pattern has detection and spatial components.
Depth, candidate selection and broad composition remain consequential.

ALTERNATIVE EXPLANATIONS RETAINED

Transferred-label selection and epithelial mixtures
RNA depth and background RNA
Tissue geometry and other inflammatory inputs
Sensitivity checks do not identify a unique cause.

FUNCTIONAL HYPOTHESIS STILL UNTESTED

Does recipient configuration alter an IL-1 response?
Needs independently established state identity
and protein / functional response measurements.
RNA co-detection is not receptor assembly or signaling.

Dashed arrows represent interpretation, not demonstrated causation. No accepted mechanism or independent validation.