








A NKX21 · mouse**B NKX21 · human****C Concordance across targets****D Internal source-table check**

Human numeric S5 reproduction remains unresolved; source labels are retained.





A Mouse epithelium

B Human fibroblasts




Marker-panel contrast: mean $\log_2(\text{TMM CPM} + 0.5)$

Target versus same-plate TIGIT + tdTomato · common scale; numbers are effect estimates


A NKX21 · mouse**B NKX21 · human****C AT2 markers and chemokines****D Growth and chemokines**

RNA axes: mean $\log_2[\text{TMM CPM} + 0.5]$ contrasts · 95% intervals in A-B are technical


A Transition and stress markers



B Wnt and AT1 markers




C SLC34A2 marker profile







D Deposited component activities





NKX21	0.19	-0.49	0.52
BECN1	-0.32	0.17	0.04
CSNK2A1	0.53	0.88	0.04
CTNNB1	0.24	0.23	-0.83
ELOVL1	-0.33	-0.04	-0.44
ATP6V0E	-0.27	-0.41	-0.34
SLC34A2	0.84	0.18	-0.03
	ICA 5	ICA 13	ICA 17



A-C: 95% technical intervals · D: deposited S6 values; component signs are arbitrary


A Mouse controls · n = 111**B Human controls · n = 99****C Detection within each compartment****D Epithelial receptor perturbations**

Species were normalized separately · D: 95% technical intervals; day 7 is secondary


A NKX21 sensitivity**B SLC34A2 sensitivity****C NKX21 paired-depth sensitivity****D Cross-target association**

RNA contrasts: mean $\log_2(\text{TMM CPM} + 0.5)$ · A-B ranges are sensitivity ranges, not confidence intervals


A Mouse Hallmark context




B Human Hallmark context




C Intergene-correlation sensitivity




A Species-specific QC retention




B Gene detection · 886 wells



C Species-assignment diagnostic





D Day-14 scaling sensitivity




Paired wells are a subset of species QC passes; wells are not independent preparations.


A Source S7 gene projections




B Budding versus controls




C Budding versus other imaged



A Control score-depth associations



B Target influence



M: mouse marker panel; H: human marker panel · residual correlations do not identify causality