



A





Deposited individual-mouse labels per tissue and assay

**B**

Cell-count concordance with supplementary Tables 1a and 2a




Ageing comparisons require design and release checks. All 73 source records join, but 47 cell totals differ; 24-month females and lung droplet samples are absent. Tissues and brain releases overlap. Two pooled lung libraries are excluded from individual-mouse counts.




To assess tissue remodelling, captured bladder fractions are compared by mouse. From 3 to 24 months, urothelial cells increase (44.0% to 75.9%) and mesenchymal cells decrease (50.3% to 15.7%). These are captured proportions, not absolute abundance. Points: mice; bars: means; triangles: females.


A 3 months · 6 mice · 4,488 cells




B 18 months · 4 mice · 4,375 cells




C 24 months · 4 mice · 4,267 cells




D Deposited clusters 1 and 6









E Deposited clusters 10, 12 and 14







F Marker detection: equal mouse means



Microglial heterogeneity is assessed through deposited clusters: blue, 1/6; orange, 10/12/14; grey, others. From 3 to 24 months, occupancy shifts from 58.7% to 6.9% (blue) and 0.5% to 6.8% (orange). Final-paper cluster mapping remains unresolved; UMAP establishes no trajectory. D/E: mouse points and means; triangles: females.



Separate RNA endpoints assess detection, expression intensity and composition. Lung Cdkn2a detection increases from 0.65% to 1.42% (3 to 24 months). RNA establishes neither senescence nor secretion. Points: mice; bars: means; triangles: females. D compares all captured cells with an equal-type reference restricted to shared types.



To assess sampling effects on apparent clonal expansion, older mice retain higher T-cell clonality after standardizing to 60 reconstructed cells. Tissue sampling and reconstruction remain alternative explanations; clonality does not measure immune function. B-D: mouse points and means; sex is not encoded.



Fixed markers assess whether age-associated RNA changes agree across resident immune populations. Il1b detection directions differ across tissues, and the kidney direction reverses in the male-only subset. Values: 24 minus 3 months, in percentage points; n: young/old mice. Different populations, assays and reused mice preclude independent replication or a conserved-mechanism claim.