
32 protein-coding candidates

Complete union of four reading lists · alphabetical order

	KO−HET CD62L-high	KO−HET CD62L-low	State interaction
Adam8	-0.02	-1.40	-1.37
Adcy6	-0.08	+0.85	+0.93
Ager	-0.24	+0.71	+0.96
Ccr2	-0.45	-1.97	-1.51
Ccr4	-0.59	-1.46	-0.87
Ccr5	-0.16	-1.39	-1.23
Cxcr3	-0.81	-1.35	-0.53
Cxcr6	-0.11	-1.39	-1.29
Dapl1	+0.66	+1.38	+0.72
G0s2	+0.44	+1.23	+0.79
H2-Aa	+0.02	+1.14	+1.12
H2-Ab1	-0.04	+1.03	+1.07
H2-Eb1	-0.08	+1.04	+1.12
Igfbp4	+0.47	+1.51	+1.04
Il10	-0.36	-1.84	-1.48
Il1r2	-0.48	-1.41	-0.93
Il4	-0.07	-1.48	-1.42
Khdc1c	-0.23	+0.71	+0.94
Klk1b22	+0.86	+1.68	+0.82
Klrg1	-0.01	-2.34	-2.32
Lsr	-0.10	-1.43	-1.32
Ly6a	+0.40	-0.79	-1.19
Ly6c2	+0.21	+1.11	+0.91
Ly6d	-0.14	+0.94	+1.09
Myo1f	-0.56	-1.60	-1.04
Ntn4	-0.41	+0.58	+0.99
Prdm1	-0.20	-1.36	-1.15
Rnf130	+0.23	+1.00	+0.77
Rps27	-2.34	+2.50	+4.84
S100a4	-1.58	-2.01	-0.44
S100a6	-1.32	-2.03	-0.71
Tigit	-0.73	-1.78	-1.04



Shared-source exploration, 11 preparations; biological independence is unresolved.
Interaction = low-state contrast − high-state contrast. Values are not ordinary log₂ fold changes.
Colours show measured RNA differences, not significance, pathway activation or target validity.