









Age-associated RNA differences combine captured composition and within-type changes. Cdkn2a detection rises in lung and falls in bladder. Values are 24 minus 3 months, with equal mouse weighting on common cell types. Four terms sum to the total; they are descriptive accounting, not causal effects. Color scales differ between panels.


A Lung FACS · within-type detection




B Bladder droplet · within-type detection




C Lung · Cdkn2a detection







D Bladder · Cdkn2a detection



Within-type profiles locate age-associated RNA differences; fixed young composition asks whether population redistribution accounts for the tissue summary. A/B: 24 minus 3 months. C/D: connected values from each mouse; black bars, means; triangles, females. Sparse types and finer subtype mixtures remain unresolved. RNA detection does not establish senescence.



Local T-cell clone repetition is higher in older thymus and spleen samples; marrow shows a different age pattern. Points are mice and black bars are means. Bottom panels condition on tissue-specific sampling depth, so their magnitudes are not directly comparable across tissues. T-cell subtype, assembly selection and function remain unresolved.



Older local-clone burden persists after fixing tissue representation in the same mice. A-C use mice with thymus, spleen and marrow; D uses all supported mice per tissue. Shared clones indicate observed distribution across tissues, not migration. Only two young mice support the balanced comparison; subtype and reconstruction selection remain alternatives.