










A: numbers are source mouse labels, not independent cross-tissue replicates. Two pooled young lung libraries are excluded.

B: each point is one deposited tissue-mouse record. General and Figure 4 brain releases overlap.




Points: individual mice; triangles: female; circles: male; black bars: arithmetic mouse means. n is shown above each age. Fractions describe captured cells, not absolute tissue abundance. One month is developmental; age is cross-sectional.


A 3 months · 6 mice · 4,488 cells




B 18 months · 4 mice · 4,375 cells




C 24 months · 4 mice · 4,267 cells




D Deposited clusters 1 and 6









E Deposited clusters 10, 12 and 14



F Marker detection: equal mouse means







UMAP and labels are from the author deposit; final-paper cluster correspondence is unresolved. Blue: clusters 1/6; orange: clusters 10/12/14; grey: remaining clusters. D-E: mouse points and means. F: nonzero deposited X; no back-transformation. These panels do not establish a transition or distinct intermediate state.



Mouse points and arithmetic means; triangles denote females. Normalized per-cell RNA is used, not raw counts.


D: equal weighting of cell types present in every retained mouse; see coverage table. RNA detection does not establish senescence or cytokine secretion.




Source Table 9, matched reconstructed T cells only. Points: mouse; bars: equal mouse means; sexes are not encoded in these panels.
 C: exact hypergeometric expectation using within-mouse observed clone labels. Tissue mixture and reconstruction selection remain unresolved.

A

All observed sexes

**B**

Male-only sensitivity



Equal mouse means; n labels are young/old animals. Complete fixed-marker family is retained in the source table; this subset was fixed for display. Resident populations and assays differ. Shared direction is context, not a conserved mechanism or independent organ replication; no interaction test.