

Ezh2-associated RNA differences across Treg contexts

DuPage et al. (2015): KO vs HET | Peeters et al. (2024): Y641F vs WT

A Expression differences across observed states




B Complete 32-gene candidate panel



	CD62L-high	CD62L-low
Adam8	-0.02	-1.40
Adcy6	-0.08	+0.85
Ager	-0.24	+0.71
Ccr2	-0.45	-1.97
Ccr4	-0.59	-1.46
Ccr5	-0.16	-1.39
Cxcr3	-0.81	-1.35
Cxcr6	-0.11	-1.39
Dapl1	+0.66	+1.38
G0s2	+0.44	+1.23
H2-Aa	+0.02	+1.14
H2-Ab1	-0.04	+1.03
H2-Eb1	-0.08	+1.04
Igfbp4	+0.47	+1.51
Il10	-0.36	-1.84
Il1r2	-0.48	-1.41
Il4	-0.07	-1.48
Khdc1c	-0.23	+0.71
Klk1b22	+0.86	+1.68
Klrg1	-0.01	-2.34
Lsr	-0.10	-1.43
Ly6a	+0.40	-0.79
Ly6c2	+0.21	+1.11
Ly6d	-0.14	+0.94
Myo1f	-0.56	-1.60
Ntn4	-0.41	+0.58
Prdm1	-0.20	-1.36
Rnf130	+0.23	+1.00
Rps27	-2.34	+2.50
S100a4	-1.58	-2.01
S100a6	-1.32	-2.03
Tigit	-0.73	-1.78

KO - HET: mean log2(FPKM + 1) difference

C The gain-context comparison changes with culture



Ranges: single-library omission checks, not confidence intervals.

A-B: recovered FOXP3-positive populations; 11 libraries (HET high/low: 2/3; KO high/low: 3/3).
Gray points: all eligible loci; outlined points: the 32 previously selected coding candidates. Selection was exploratory.
CD62L is an observed state, not an activation intervention. C uses a different study and scale; effect sizes are not pooled.