

A
Corrected Compass association
83 variable reactions; 60 pools
4 libraries nested in 2 animals
PGAM
ρ = +0.291
BH q = 0.181
Reaction rank (ascending ρ)
Spearman ρ: consistency vs pathogenicity
B
EGCG vs DMSO in Th17n
Division 1; 5 libraries per arm
IL17F
IL17A
RORC
IL23R
−log₁₀ BH-adjusted P
Library-level log₂ fold change
C
Relative RNA programmes
Dot: observed; gray: 95% random-set range
Th17 effector
Cholesterol / SREBP
Hypoxia / HIF
UPR
OXPHOS
Heat shock
Amino-acid transport
NRF2 / oxidative stress
Glycolysis
Ribosomal proteins
Cell cycle
Serine / one-carbon
Histones
ISR
Centred median log₂ fold change
D
Drug and culture context
Th17 effector +0.54 -1.18 -0.73 -1.12
Cholesterol / SREBP +0.11 +0.10 -0.18 -0.26
Hypoxia / HIF +0.13 +0.25 -0.28 +0.53
UPR +0.06 -0.07 -0.10 -0.09
OXPHOS -0.03 -0.12 -0.01 -0.07
Heat shock +0.02 -0.05 +0.03 -0.06
Amino-acid transport -0.07 -0.04 -0.34 -0.15
NRF2 / oxidative stress -0.07 -0.16 -0.02 +0.03
Glycolysis -0.05 +0.14 -0.32 +0.18
Ribosomal proteins -0.21 +0.01 -0.35 +0.07
Cell cycle -0.26 -0.22 -0.53 -0.15
Serine / one-carbon -0.30 -0.15 -0.50 -0.15
Histones -0.46 -0.26 -0.60 -0.05
ISR -0.49 +0.34 -0.60 -0.15
Th17n EGCG
Th17n DHEA
Th17p EGCG
Th17p DHEA
Centred median log₂ fold change