







Figure 5 | Older reconstructed T-cell repertoires remain more concentrated at a common sampling depth.

Rationale: Higher observed clonality may reflect unequal numbers of reconstructed receptors or different sampled tissues. Source identity correction, mouse-level summaries and equal-depth sampling assess these observational explanations before assigning an immune-function interpretation.

A, cells in within-mouse nonsingleton source clone groups divided by matched reconstructed-cell counts. B, individual-mouse fractions and arithmetic means ($n = 7/4/4$ at 3/18/24 months). C, the exact expected nonsingleton fraction after sampling 60 reconstructed cells per mouse, conditional on the observed clone labels. D, reconstruction denominators on a logarithmic axis. Sex is not encoded. The common depth is the observed minimum, not a power threshold.

Results: The source yields 55/2,076, 479/2,056 and 348/1,868 clonal cells. Numerators match the manuscript, but the young and old denominators differ. Equal-mouse fractions average 2.30%, 22.53% and 18.94%; common-depth expectations are 0.62%, 8.41% and 7.82%. Eight 18-month and three 24-month rows lack metadata matches. The older-versus-younger pattern remains in this conditional sampling comparison, while tissue mixture, reconstruction selection and source-version differences remain unresolved. These results do not establish antigen specificity or immune competence.