

# A Quantum Model for Solving HCP and TSP Graph Problem Based on Quantum $Z_2$ Lattice Gauge Theory

Xiaopeng Cui

(Dated: 2022.6.28)

## Abstract

In this paper, using the condensed closed string characteristics of  $Z_2$  topological quantum phase transition, by mapping a graph to a lattice we develop a novel quantum model and a adiabatic algorithm to solve the shortest path, HCP and TSP graph problems, which only need  $N_e$  qubits for a given graph (the number of vertices and edges is  $N_v, N_e$ ). By the simulation of a number of small random graphs we demonstrate the effectiveness of this quantum algorithm model.

# I. HAMILTONIAN MODEL OF GRAPH PROBLEM BASED ON $Z_2$ GAUGE THEORY

The Hamiltonian of the quantum  $Z_2$  gauge theory[1–3] is defined as

$$\begin{aligned} H &= Z + gX \\ Z &= \sum_{\square} Z_{\square}, \quad Z_{\square} = -\prod_{l \in \square} \sigma_l^z \\ X &= -\sum_l \sigma_l^x, \end{aligned} \tag{1}$$

where the  $\square$  indicates the elementary plaquette of the lattice, which is the mapping of the vertex from the graph.

For graph problem, with the dual mapping from graph to  $Z_2$ -lattice[4], the  $Z_2$  Graph Hamiltonian can be defined as

$$H_{Z_2-G} = H_{problem} + gX \quad \text{or} \quad H_{Z_2-G} = \lambda H_{problem} + X \tag{2}$$

Here are three problems. (1)The Shortest Path of Two Vertex,(2)HCP, (3)TSP

## A. The Shortest Path of Two Vertex

First,we need add a edge  $l_0$  between starting vertex and end vertex with  $w_{l_0} = 0$  and set  $\sigma_{l_0}^z = -1$ . Then the Hamiltonian of the shortest path problem can be defined as

$$\begin{aligned} H_{hcp} &= Z + W \\ Z &= \sum_{\square} Z_{\square}, \quad Z_{\square} = -\prod_{l \in \square} \sigma_l^z \\ W &= -\sum_{l \in \square} w_l \sigma_l^z \end{aligned} \tag{3}$$

Meanings of items:

- 1)  $\sigma_l^z = -1$  means that this edge  $l$  is selected by a path.
- 2)  $Z$  item : the  $\square$  indicates the elementary plaquette of the lattice, or a vertex of its mapping graph.  $d_{\square}$  represents the degree of the vertex  $\square$ .  $Z$  item restricts that the path is a loop in the graph.
- 3)  $W$  item reduce the number of visited vertexs and the weight sum of the path.

## B. HCP

The Hamiltonian of HCP can be defined as

$$\begin{aligned}
H_{hcp} &= Z + V + fF \\
Z &= \sum_{\square} Z_{\square}, \quad Z_{\square} = - \prod_{l \in \square} \sigma_l^z \\
V &= \sum_{\square} V_{\square}, \quad V_{\square} = \left( \sum_{l \in \square} \sigma_l^z - c_{\square} \right)^2, \quad c_{\square} = d_{\square} - 4 \\
F &= ?
\end{aligned} \tag{4}$$

Meanings of items:

- 1)  $\sigma_l^z = -1$  means that this edge  $l$  is selected by a path.
- 2) the  $\square$  indicates the elementary plaquette of the lattice, or a vertex of its mapping graph.  
 $d_{\square}$  represents the degree of the vertex  $\square$ .  $Z$  item restricts that the path is a loop in the graph.
- 3)  $V$  item restricts a single visit for each vertex in the graph (1-visit loop).
- 4)  $F$  facilitate the fusion of split 1-visit loops into one single loop.  $f = 0 - 1$ .  $F$  is in consideration, which should reduce the number of subloop.

## C. TSP

The Hamiltonian of TSP can be defined as

$$\begin{aligned}
H_{tsp} &= H_{hcp} + W \\
W &= - \sum_l w_l \sigma_l^z
\end{aligned} \tag{5}$$

Meanings of items:

- 1)  $W$  represents the minimization of the sum of path weights.  $w_l$  represents the weight of the edge  $l$ .

Simplify Hamiltonian  $H_{hcp}, H_{tsp}$

$$d_{\square} = c_{\square} + 4 \quad (6)$$

$$\begin{aligned}
V_{\square} &= \left( \sum_{l \in \square} \sigma_l^z - c_{\square} \right)^2 \\
&= \left( \sum_{l \in \square} \sigma_l^z \right)^2 - 2c_{\square} \sum_{l \in \square} \sigma_l^z + c_{\square}^2 \\
&= \left[ \left( \sum_{l \in \square} (\sigma_l^z)^2 \right) + \left( \sum_{l, m \in \square, l \neq m} \sigma_l^z \sigma_m^z \right) \right] - 2c_{\square} \sum_{l \in \square} \sigma_l^z + c_{\square}^2, \quad (\sigma_l^z)^2 = +1 \\
&= [d_{\square} + \left( \sum_{l, m \in \square, l \neq m} \sigma_l^z \sigma_m^z \right)] - 2c_{\square} \sum_{l \in \square} \sigma_l^z + c_{\square}^2 \\
&= \left( \sum_{l, m \in \square, l \neq m} \sigma_l^z \sigma_m^z \right) - 2c_{\square} \sum_{l \in \square} \sigma_l^z + [c_{\square}^2 + c_{\square} + 4] \\
V &= \sum_{\square} V_{\square} = \sum_{\square} \left( \sum_{l, m \in \square, l \neq m} \sigma_l^z \sigma_m^z \right) - 2c_{\square} \sum_{l \in \square} \sigma_l^z + [c_{\square}^2 + c_{\square} + 4]
\end{aligned} \quad (7)$$

Remove the constant term, the Hamiltonian of HCP is

$$\begin{aligned}
V_{\square} &= \left( \sum_{l, m \in \square, l \neq m} \sigma_l^z \sigma_m^z \right) - 2c_{\square} \sum_{l \in \square} \sigma_l^z \\
V &= \sum_{\square} V_{\square} = \sum_{\square} \left[ \left( \sum_{l, m \in \square, l \neq m} \sigma_l^z \sigma_m^z \right) - 2c_{\square} \sum_{l \in \square} \sigma_l^z \right] \\
H_{hcp} &= \sum_{\square} - \prod_{l \in \square} \sigma_l^z + \sum_{\square} \left[ \left( \sum_{l, m \in \square, l \neq m} \sigma_l^z \sigma_m^z \right) - 2c_{\square} \sum_{l \in \square} \sigma_l^z \right] + fF
\end{aligned} \quad (8)$$

For TSP, we need add the weight of the edge  $w_l$ . So the Hamiltonian of TSP is

$$H_{tsp} = H_{hcp} - \sum_l w_l \sigma_l^z \quad (9)$$

When  $g$  is equal to 0, the ground state of the system is the optimal solution to the HCP or TSP problem. Obviously, The optimal solution can be obtained by gradually increasing  $\lambda$  from initial Hamiltonian  $-\sum_l \sigma_l^x$  adiabatically. The adiabatic algorithm can refer to the article "Solving Hamiltonian Cycle Problem with Quantum Z2 Lattice Gauge Theory".

## II. QUANTUM SIMULATION OF HCP MODEL IN $3 \times 3$ TORUS LATTICE

Quantum simulations(the  $3 \times 3$  torus lattice) verify that above model works correctly as shown in FIG.1. (1) Loop state condition:  $\langle Z \rangle = -N_v$ ; (2) 1-visit loop state condition:  $\langle V \rangle = 0$ .




FIG. 1. Simulation results. (a) The  $3 \times 3$  torus lattice. (b) The graph corresponding to the  $3 \times 3$  torus lattice. (c)  $f = 0$ , the simulation results of  $H_{Z_2\text{-hcp}}$ . (d)  $f = 0$ , the probability of 1-visit loop state  $P_{\text{loop}}$  vs  $\lambda$ .

### III. CONCLUSION AND DISCUSSION

In this paper, using the condensed close string characteristics of  $\mathbb{Z}_2$  topological quantum phase transition, by mapping a graph to a lattice we develop a novel quantum model and a adiabatic algorithm to solve the shortest path, HCP and TSP graph problem, which only need  $N_e$  qubits (the number of vertices and edges is  $N_v, N_e$ ). Using this algorithm model, by the simulation of a number of small random graphs we demonstrate its effectiveness.

### IV. ACKNOWLEDGE

- 
- [1] X. Cui, Y. Shi, and J. C. Yang, “Circuit-based digital adiabatic quantum simulation and pseudoquantum simulation as new approaches to lattice gauge theory,” J. High Energy Phys **2020**, 160 (2020).
  - [2] S. Sachdev, “Topological order, emergent gauge fields, and fermi surface reconstruction,” Rep. Prog. Phys. **82**, 014001 (2018).
  - [3] F. J. Wegner, “Duality in generalized ising models and phase transitions without local order parameters,” J. Math. Phys. **12**, 2259 (1971).
  - [4] Xiaopeng Cui and Yu Shi, “Solving Hamiltonian Cycle Problem using Quantum  $\mathbb{Z}_2$  Lattice Gauge Theory,” arXiv e-prints , arXiv:2202.08817 (2022), arXiv:2202.08817 [quant-ph].