

APPENDIX X: CHILD MEAL PATTERNS

Child Meal Patterns

Breakfast (Select all three components)			
Food Components and Food Items	Ages 1-2	Ages 3-5	Ages 6-12
Fluid Milk¹	4 fluid ounces	6 fluid ounces	8 fluid ounces
Vegetables, fruits, or portions of both²	¼ cup	½ cup	½ cup
Grains (oz eq)³			
Whole grain-rich or enriched bread	½ slice	½ slice	1 slice
Whole grain-rich or enriched bread product, such as biscuit, roll, or muffin	½ serving	½ serving	1 serving
Whole grain-rich, enriched or fortified cooked breakfast cereal, cereal grain, and/or pasta	¼ cup	¼ cup	½ cup
Whole grain-rich, enriched or fortified ready-to-eat breakfast cereal (dry, cold)			
Flaked or rounds	½ cup	½ cup	1 cup
Puffed cereal	¾ cup	¾ cup	1 ¼ cup
Granola	⅓ cup	⅓ cup	¼ cup

Lunch and Supper (Select all five components)			
Food Components and Food Items	Ages 1-2	Ages 3-5	Ages 6-12
Fluid Milk¹	4 fluid ounces	6 fluid ounces	8 fluid ounces
Meat/meat alternatives			
Lean meat, poultry, or fish	1 ounce	1 ½ ounce	2 ounces
Tofu, soy products, or alternate protein products	1 ounce	1 ½ ounce	2 ounces
Cheese	1 ounce	1 ½ ounce	2 ounces
Large egg	½	¾	1
Cooked dry beans or peas	¼ cup	¾ cup	½ cup
Peanut butter or soy nut butter or other nut or seed butters	2 tbsp	3 tbsp	4 tbsp
Yogurt, plain or flavored, unsweetened or sweetened	4 ounces or ½ cup	6 ounces or ¾ cup	8 ounces or 1 cup
The following may be used to meet no more than 50% of the requirement: Peanuts, soy nuts, tree nuts, or seeds (1 ounce of nuts/seeds = 1 ounce of cooked lean meat, poultry, or fish)	½ ounce = 50%	¾ ounce = 50 %	1 ounce = 50 %
Vegetables²	¼ cup	¼ cup	½ cup
Fruits²	¼ cup	¼ cup	¼ cup
Grains (ounce equivalents)³			
Whole grain-rich or enriched bread	½ slice	½ slice	1 slice
Whole grain-rich or enriched bread product, such as biscuit, roll, or muffin	½ serving	½ serving	1 serving
Whole grain-rich, enriched or fortified cooked breakfast cereal, cereal grain, and/or pasta	¼ cup	¼ cup	½ cup

¹ Must be unflavored whole milk for children age one. Must be unflavored low-fat (1 percent) or unflavored fat-free (skim) milk for children two through five years old. Must be unflavored low-fat (1 percent), unflavored fat-free (skim), or flavored fat-free (skim) milk for children six years old or older.

² Pasteurized full-strength juice may only be used to meet the vegetable or fruit requirement at one meal, including snacks, per day.

³ At least one serving per day, across all eating occasions, must be whole grain-rich. Grain-based desserts do not count toward meeting the grains requirement.

APPENDIX X: CHILD MEAL PATTERNS

Child Meal Patterns

Snack (Select two of the five components)			
Food Components and Food Items	Ages 1-2	Ages 3-5	Ages 6-12
Fluid Milk ¹	4 fluid ounces	4 fluid ounces	8 fluid ounces
Meat/meat alternatives			
Lean meat, poultry, or fish	½ ounce	½ ounce	1 ounces
Tofu, soy products, or alternate protein products	½ ounce	½ ounce	1 ounces
Cheese	½ ounce	½ ounce	1 ounces
Large egg	½	½	½
Cooked dry beans or peas	¼ cup	¼ cup	¼ cup
Peanut butter, soy nut butter, or other nut or seed butters	1 tbsp	1 tbsp	2 tbsp
Yogurt, plain or flavored, unsweetened or sweetened	2 ounces or ¼ cup	2 ounces or ¼ cup	4 ounces or ½ cup
Peanuts, soy nuts, tree nuts, or seed	½ ounce	½ ounce	1 ounce
Vegetables ²	½ cup	½ cup	¾ cup
Fruits ²	½ cup	½ cup	¾ cup
Grains (ounce equivalents) ³			
Whole grain-rich or enriched bread	½ slice	½ slice	1 slice
Whole grain-rich or enriched bread product, such as biscuit, roll, or muffin	½ serving	½ serving	1 serving
Whole grain-rich, enriched or fortified cooked breakfast cereal, cereal grain, and/or pasta	¼ cup	¼ cup	½ cup
Whole grain-rich, enriched or fortified ready-to-eat breakfast cereal (dry, cold)			
Flaked or rounds	½ cup	½ cup	1 cup
Puffed cereal	¾ cup	¾ cup	1 ¼ cup
Granola	¾ cup	¾ cup	¾ cup

¹ Must be unflavored whole milk for children age one. Must be unflavored low-fat (1 percent) or unflavored fat-free (skim) milk for children two through five years old. Must be unflavored low-fat (1 percent), unflavored fat-free (skim), or flavored fat-free (skim) milk for children six years old or older.

² Pasteurized full-strength juice may only be used to meet the vegetable or fruit requirement at one meal, including snacks, per day.

³ At least one serving per day, across all eating occasions, must be whole grain-rich. Grain-based desserts do not count toward meeting the grains requirement.

APPENDIX XI: INFANT MEAL PATTERNS

Infant Meal Patterns

Breakfast, Lunch, and Dinner (Select all three components)	
Birth through 5 months	6 through 11 months
4 -6 fluid ounces breastmilk ¹ or formula ²	6-8 fluid ounces breastmilk ¹ or formula ² ; and 0-4 tablespoons of • Infant cereal² • Meat • Fish • Poultry • Whole egg • Cooked, dry beans or • Cooked dry peas or 0-2 ounces of cheese or 0-4 ounces (volume) of cottage cheese, or • 0-4 ounces or ½ cup of yogurt; or a combination of the above³; and 0-2 tablespoons vegetable or fruit or a combination of both. ^{3,4}
Snacks (Select all three components)	
Birth through 5 months	6 through 11 months
4-6 ounces of breastmilk ¹ or formula ²	2-4 fluid ounces breastmilk or formula ² ; and • 0-1/2 slice bread^{3,5} or • 0-2 crackers ^{3,5} or • 0-4 tablespoons infant cereal ^{2,3} and 0-2 tablespoons vegetable or fruit, or a combination of both. ^{3,4}

¹ Breastmilk or formula, or portions of both, must be served; however, it is recommended that breastmilk be served in place of formula from birth through 11 months. For some breastfed infants who regularly consume less than the amount of breastmilk per feeding, a serving of less than the minimum amount of breastmilk may be offered, with additional breastmilk offered at a later time if the infant will consume more.

² Infant formula and dry infant cereal must be iron fortified.

³ A serving of this component is required when the infant is developmentally ready to accept it.

⁴ Fruit and vegetable juices must not be served.

⁵ A serving of grains must be whole grain-rich, enriched meal, or enriched flour.