

# Conformal prediction and its extensions

## 1 CP: Conformal prediction

Methods for distribution-free prediction.

**Assumption: exchangeability**

- The **data**  $Z_i = (X_i, Y_i)$  are assumed to be exchangeable (for example, i.i.d.).
  - Definition (Vovk, Gammernan, and Shafer 2005; Shafer and Vovk 2008). Suppose that for any collection of  $N$  values, the  $N!$  different orderings are equally likely. Then we say that  $Z_1, \dots, Z_N$  are exchangeable. The exchangeability assumption is slightly weaker than the i.i.d. assumption.
- The **algorithm** which maps data to a fitted model  $\hat{\mu} : \mathcal{X} \rightarrow \mathbb{R}$  is assumed to treat the data points symmetrically.
  - For example, OLS versus WLS.

### 1.1 Split conformal prediction

(inductive conformal prediction)

1. initial training data set: pre-trained model  $\hat{\mu} : \mathcal{X} \rightarrow \mathbb{R}$ .
2. holdout/calibration set: nonconformity scores  $R_i = |Y_i - \hat{\mu}(X_i)|$ ,  $i = 1, \dots, n$ .
3. prediction set:  $\widehat{C}_n(X_{n+1}) = \hat{\mu}(X_{n+1}) \pm Q_{1-\alpha} \left( \sum_{i=1}^n \frac{1}{n+1} \cdot \delta_{R_i} + \frac{1}{n+1} \cdot \delta_{+\infty} \right)$ .  
(the  $\lceil (n+1)(1-\alpha) \rceil$ th smallest of  $R_1, \dots, R_n$ )

Drawback: the loss of accuracy due to sample splitting.

## 1.2 Full conformal prediction

(transductive conformal prediction)

1. training data & a hypothesized test point:  $\hat{\mu}^y = \mathcal{A}((X_1, Y_1), \dots, (X_n, Y_n), (X_{n+1}, y))$  for each  $y \in \mathbb{R}$ .
2. residuals:  $R_i^y = \begin{cases} |Y_i - \hat{\mu}^y(X_i)|, & i = 1, \dots, n \\ |y - \hat{\mu}^y(X_{n+1})|, & i = n + 1 \end{cases}$ .
3. prediction set:  $\widehat{C}_n(X_{n+1}) = \{y \in \mathbb{R} : R_{n+1}^y \leq Q_{1-\alpha}(\sum_{i=1}^{n+1} \frac{1}{n+1} \cdot \delta_{R_i^y})\}$ .

Drawback: a steep computational cost.

*THEOREM:*

$\mathbb{P}\{Y_{n+1} \in \widehat{C}_n(X_{n+1})\} \geq 1 - \alpha$  holds true for both split conformal and full conformal.

## 1.3 Jackknife+

(close to cross-conformal prediction in Vovk (2013), offering a compromise between the computational and statistical costs)

1. training data with  $i$ th point removed:  $\hat{\mu}_{-i} = \mathcal{A}((X_1, Y_1), \dots, (X_{i-1}, Y_{i-1}), (X_{i+1}, Y_{i+1}), \dots, (X_n, Y_n))$ .
2. residuals:  $R_i^{\text{LOO}} = |Y_i - \hat{\mu}_{-i}(X_i)|$ .
3. prediction set:

$$\left[ Q_\alpha \left( \sum_{i=1}^n \frac{1}{n+1} \cdot \delta_{\hat{\mu}_{-i}(X_{n+1}) - R_i^{\text{LOO}}} + \frac{1}{n+1} \cdot \delta_{-\infty} \right), Q_{1-\alpha} \left( \sum_{i=1}^n \frac{1}{n+1} \cdot \delta_{\hat{\mu}_{-i}(X_{n+1}) + R_i^{\text{LOO}}} + \frac{1}{n+1} \cdot \delta_{+\infty} \right) \right]$$

Drawback: while in practice it generally provides coverage close to the target level  $1 - \alpha$ , its theoretical guarantee only ensures  $1 - 2\alpha$  probability of coverage in the worst case.

*THEOREM:*

$\mathbb{P}\{Y_{n+1} \in \widehat{C}_n(X_{n+1})\} \geq 1 - 2\alpha$  holds true for jackknife+.

## 2 Conformal time-series forecasting

Stankeviciute, M Alaa, and Schaar (2021): CF-RNNs

**Multi-horizon time-series forecasting problem**

**Notation:**

- the  $i$ th data point:  $Z_i = (y_{1:t}^{(i)}, y_{t+1:t+H}^{(i)})$ . Note that the label  $y_{t+1:t+H}^{(i)}$  is now an  $H$ -dimensional value, in contrast with the scalar  $y$  value from before.

**Assumption:**

- exchangeable time-series observations

## 2.1 Methodology

(Split conformal prediction)

1. training set: train the underlying (auxiliary) model  $\hat{\mu} : \mathbb{R}^t \rightarrow \mathbb{R}^H$ , which produces multi-horizon forecasts **directly** (conditionally independent predictions).
2. calibration set: obtain the  $H$ -dimensional nonconformity scores

$$R_i = \left[ |y_{t+1}^{(i)} - \hat{y}_{t+1}^{(i)}|, \dots, |y_{t+H}^{(i)} - \hat{y}_{t+H}^{(i)}| \right]^\top.$$

3. prediction set:  $\Gamma_1^\alpha(y_{1:t}^{(n+1)}), \dots, \Gamma_H^\alpha(y_{1:t}^{(n+1)})$ , where  $\Gamma_h^\alpha(y_{1:t}^{(n+1)}) = [\hat{y}_{t+h}^{(n+1)} - \hat{\varepsilon}_h, \hat{y}_{t+h}^{(n+1)} + \hat{\varepsilon}_h]$ ,  $\forall h \in \{1, \dots, H\}$  with the critical nonconformity scores  $\hat{\varepsilon}_1, \dots, \hat{\varepsilon}_H$  become the  $[(n+1)(1 - \alpha/H)]$ -th smallest residuals in the corresponding nonconformity score distributions. (Bonferroni correction)

*THEOREM:*

- $\mathcal{D} = \left\{ (y_{1:t}^{(i)}, y_{t+1:t+H}^{(i)}) \right\}_{i=1}^n$ : **exchangeable** time-series observations.
- $\hat{\mu}$ : model predicting  $H$ -step forecasts using **the direct strategy**.

$$\mathbb{P}(\forall h \in \{1, \dots, H\} \cdot y_{t+h} \in [\hat{y}_{t+h} - \hat{\varepsilon}_h, \hat{y}_{t+h} + \hat{\varepsilon}_h]) \geq 1 - \alpha.$$

## 3 NexCP: Conformal prediction beyond exchangeability

Barber et al. (2023)

1. Nonexchangeable conformal with a **symmetric** algorithm (weights)
2. Nonexchangeable conformal with **nonsymmetric** algorithms (weights & swap)

### 3.1 Notation

- the  $i$ th data point  $Z_i = (X_i, Y_i)$
- the full data sequence  $Z = (Z_1, \dots, Z_{n+1})$
- the sequence after swap  $Z^i = (Z_1, \dots, Z_{i-1}, Z_{n+1}, Z_{i+1}, Z_n, Z_i)$

### 3.2 Methodology

1. Choose **fixed (non-data-dependent)** weights  $w_1, \dots, w_n \in [0, 1]$  with the intuition that a higher weight should be assigned to a data point that is “trusted” more.
2. Normalize weights  $\tilde{w}_i = \frac{w_i}{w_1 + \dots + w_{n+1}}, i = 1, \dots, n$  and  $\tilde{w}_{n+1} = \frac{1}{w_1 + \dots + w_{n+1}}$ .
3. Generate “tagged” data points  $(X_i, Y_i, t_i) \in \mathcal{X} \times \mathbb{R} \times \mathcal{T}$ .
4. Swap data set, resulting in  $Z^K$  with  $K \sim \sum_{i=1}^{n+1} \tilde{w}_i \cdot \delta_i$ , i.e., two data points have swapped tags.
5. Apply algorithm  $\mathcal{A}$  to  $Z^K$  in place of  $Z$ .

### 3.3 Split conformal prediction

- prediction set:  $\widehat{C}_n(X_{n+1}) = \hat{\mu}(X_{n+1}) \pm Q_{1-\alpha}(\sum_{i=1}^n \tilde{w}_i \cdot \delta_{R_i} + \tilde{w}_{n+1} \cdot \delta_{+\infty})$

### 3.4 Full conformal prediction

1. training data & a hypothesized test point:  $\hat{\mu}^{y,k} = \mathcal{A}((X_{\pi_k(i)}, Y_{\pi_k(i)}^y, t_i) : i \in [n+1])$  for any  $y \in \mathbb{R}$  and  $k \in [n+1]$ , where  $\pi_k$  is the permutation on  $[n+1]$  swapping indices  $k$  and  $n+1$ , and  $Y_i^y = \begin{cases} Y_i, & i = 1, \dots, n \\ y, & i = n+1 \end{cases}$ .
2. residuals:  $R_i^{y,k} = \begin{cases} |Y_i - \hat{\mu}^{y,k}(X_i)|, & i = 1, \dots, n \\ |y - \hat{\mu}^{y,k}(X_{n+1})|, & i = n+1 \end{cases}$ .
3. prediction set:  $\widehat{C}_n(X_{n+1}) = \{y : R_{n+1}^{y,K} \leq Q_{1-\alpha}(\sum_{i=1}^{n+1} \tilde{w}_i \cdot \delta_{R_i^{y,K}})\}$ .

*THEOREM:*

- Lower bounds on coverage.

$$\mathbb{P}\{Y_{n+1} \in \widehat{C}_n(X_{n+1})\} \geq 1 - \alpha - \sum_{i=1}^n \tilde{w}_i \cdot d_{TV}(R(Z), R(Z^i))$$

- Upper bounds on coverage.

$$\mathbb{P}\{Y_{n+1} \in \widehat{C}_n(X_{n+1})\} < 1 - \alpha + \tilde{w}_{n+1} + \sum_{i=1}^n \tilde{w}_i \cdot d_{TV}(R(Z), R(Z^i)),$$

if  $R_1^{Y_{n+1},K}, \dots, R_n^{Y_{n+1},K}, R_{n+1}^{Y_{n+1},K}$  are distinct with probability 1.

The results hold true for both nonexchangeable split conformal and full conformal.

So, if  $\tilde{w}_{n+1} = \frac{1}{w_1 + \dots + w_{n+1}}$  is small (the effective sample size is large), then mild violations of exchangeability can only lead to mild undercoverage or to mild overcoverage.

### 3.5 Jackknife+

1. training data with  $i$ th point removed and data tag swapped:

$$\hat{\mu}_{-i}^k = \mathcal{A} \left( (X_{\pi_k(j)}, Y_{\pi_k(j)}, t_j) : j \in [n+1], \pi_k(j) \notin \{i, n+1\} \right).$$

2. residuals:  $R_i^{k, \text{LOO}} = |Y_i - \hat{\mu}_{-i}^k(X_i)|$ .

3. prediction set:

$$\left[ Q_\alpha \left( \sum_{i=1}^n \tilde{w}_i \cdot \delta_{\hat{\mu}_{-i}^K(X_{n+1}) - R_i^{K, \text{LOO}}} + \tilde{w}_{n+1} \cdot \delta_{-\infty} \right), Q_{1-\alpha} \left( \sum_{i=1}^n \tilde{w}_i \cdot \delta_{\hat{\mu}_{-i}^K(X_{n+1}) + R_i^{K, \text{LOO}}} + \tilde{w}_{n+1} \cdot \delta_{+\infty} \right) \right]$$

*THEOREM:*

$$\mathbb{P} \left\{ Y_{n+1} \in \widehat{C}_n(X_{n+1}) \right\} \geq 1 - 2\alpha - \sum_{i=1}^n \tilde{w}_i \cdot d_{\text{TV}}(R_{\text{jack}+}(Z), R_{\text{jack}+}(Z^i))$$

## 4 WCP: Conformal prediction under covariate shift

Tibshirani et al. (2019)

A weighted version of conformal prediction, using a quantile of a suitably weighted empirical distribution of nonconformity scores.

### 4.1 Setup/assumption - Covariate shift

Focus on settings in which the data  $(X_i, Y_i), i = 1, \dots, n+1$  are no longer exchangeable. Specifically,

$$\begin{aligned} (X_i, Y_i) &\stackrel{\text{i.i.d.}}{\sim} P = P_X \times P_{Y|X}, \quad i = 1, \dots, n, \\ (X_{n+1}, Y_{n+1}) &\sim \widetilde{P} = \widetilde{P}_X \times P_{Y|X}, \text{ independently.} \end{aligned} \tag{1}$$

the test and training covariate distributions differ, but the likelihood ratio between the two distributions,  $d\widetilde{P}_X/dP_X$ , must be known exactly or well approximated for correct coverage.

## 4.2 Methodology

**Prediction set:**

- CP:  $\widehat{C}_n(X_{n+1}) = \widehat{\mu}(X_{n+1}) \pm Q_{1-\alpha} \left( \sum_{i=1}^n \frac{1}{n+1} \cdot \delta_{R_i} + \frac{1}{n+1} \cdot \delta_{+\infty} \right)$ .
- NexCP:  $\widehat{C}_n(X_{n+1}) = \widehat{\mu}(X_{n+1}) \pm Q_{1-\alpha} \left( \sum_{i=1}^n \tilde{w}_i \cdot \delta_{R_i} + \tilde{w}_{n+1} \cdot \delta_{+\infty} \right)$ .
  - weights  $w$  are fixed
  - $\tilde{w}_i = \frac{w_i}{w_1 + \dots + w_n + 1}, i = 1, \dots, n$  and  $\tilde{w}_{n+1} = \frac{1}{w_1 + \dots + w_n + 1}$
- WCP:  $\widehat{C}_n(X_{n+1}) = \widehat{\mu}(X_{n+1}) \pm Q_{1-\alpha} \left( \sum_{i=1}^n p_i^w(x) \delta_{R_i} + p_{n+1}^w(x) \delta_{+\infty} \right)$ .
  - $w = d\tilde{P}_X/dP_X$  or  $w \propto d\tilde{P}_X/dP_X$
  - $p_i^w(x) = \frac{w(X_i)}{\sum_{j=1}^n w(X_j) + w(x)}, i = 1, \dots, n$ , and  $p_{n+1}^w(x) = \frac{w(x)}{\sum_{j=1}^n w(X_j) + w(x)}$

The **weight function**  $\hat{w}$  can be estimated using logistic regression, random forests, etc.

*THEOREM:*

Assume data from the model Equation 1. Assume  $\tilde{P}_X$  is absolutely continuous with respect to  $P_X$ , and denote  $w = d\tilde{P}_X/dP_X$ . For any score function  $S$ , and any  $\alpha \in (0, 1)$ , define for  $x \in \mathbb{R}^d$ . Then

$$\mathbb{P} \{ Y_{n+1} \in \widehat{C}_n(X_{n+1}) \} \geq 1 - \alpha.$$

## 4.3 Comparison to NexCP

1. Assumption
  - WCP: covariate shift
  - NexCP: do not make any assumption on the joint distribution of the  $n + 1$  points
2. Weights
  - WCP: a function of the data point  $(X_i, Y_i)$  to compensate for the **known** distribution shift.
  - NexCP: required to be fixed, can compensate for **unknown** violations of the exchangeability assumption, as long as the violations are **small** (to ensure a low coverage gap).
3. Nonsymmetric algorithm
  - WCP: No.
  - NexCP: Yes.
4. For exchangeable data

- WCP: does not have any coverage guarantee.
- NexCP: retains exact coverage.

## 5 ACP: Adaptive conformal inference under distribution shift

Gibbs and Candes (2021)

- No assumptions on the data-generating distribution.
- Modelling the distribution shift as a learning problem in a single parameter whose optimal value is varying over time and must be continuously re-estimated.
- Adjust significance level  $\alpha$  based on rolling coverage of  $Y_t$ .

### 5.1 Methodology

Work with score function  $S(\cdot)$  and quantile function  $\hat{Q}(\cdot)$ .

**Some facts:**

- If the distribution of the data is shifting over time, both functions should be regularly re-estimated to align with the most recent observations.
- The realized miscoverage rate  $M_t(\alpha)$  also varies over time and may not be equal or close to  $\alpha$ .

**Assumptions:**

Assume that there may be an alternative value  $\alpha^* \in [0, 1]$  such that  $M_t(\alpha^*) \cong \alpha$ .

Assume that with probability one,  $\hat{Q}_t(\cdot)$  is continuous, non-decreasing and such that  $\hat{Q}_t(0) = -\infty$  and  $\hat{Q}_t(1) = \infty$ .

**Adaptive conformal inference:**

- Under assumptions,  $M_t(\cdot)$  will be non-decreasing on  $[0, 1]$  with  $M_t(0) = 0$  and  $M_t(1) = 1$ .
- Define  $\alpha_t^* := \sup \{\beta \in [0, 1] : M_t(\beta) \leq \alpha\}$ , then  $M_t(\alpha_t^*) = \alpha$ .
- Use a simple **update process** to perform the calibration.
  - Intuition: after examining the empirical miscoverage frequency of the previous prediction sets, decreasing (increasing) estimate of  $\alpha_t^*$  if the prediction sets were historically under-covering (over-covering)  $Y_t$ .

- Let  $\alpha_1 = \alpha$ , consider the **update**

$$\alpha_{t+1} := \alpha_t + \gamma (\alpha - \text{err}_t)$$

OR

$$\alpha_{t+1} = \alpha_t + \gamma \left( \alpha - \sum_{s=1}^t w_s \text{err}_s \right),$$

where  $\gamma > 0$  is a fixed step size parameter whose choice gives a tradeoff between adaptability and stability.

So the distribution is allowed to shift continuously over time.

*THEOREM:*

With probability one we have that for all  $T \in \mathbb{N}$ ,

$$\left| \frac{1}{T} \sum_{t=1}^T \text{err}_t - \alpha \right| \leq \frac{\max\{\alpha_1, 1 - \alpha_1\} + \gamma}{T\gamma}.$$

In particular,  $\lim_{T \rightarrow \infty} \frac{1}{T} \sum_{t=1}^T \text{err}_t \stackrel{\text{a.s.}}{=} \alpha$ .

## 6 PID: Conformal PID control for time series prediction

Angelopoulos, Candes, and Tibshirani (2023)

- PID: proportional-integral-derivative (control theory).
- reactive + forward-looking.



## 6.1 Methodology

The proposed conformal PID controller is given by

$$q_{t+1} = \underbrace{\eta(\text{err}_t - \alpha)}_{\text{P}} + r_t \underbrace{\left( \sum_{i=1}^t (\text{err}_i - \alpha) \right)}_{\text{I}} + \underbrace{g'_t}_{\text{D}},$$

where  $g_t = \text{err}_t - \alpha$ .

The goal is to achieve **long-run coverage** in time, i.e.,  $\frac{1}{T} \sum_{t=1}^T \text{err}_t = \alpha + o(1)$ , where  $o(1)$  denotes a quantity that tends to zero as  $T \rightarrow \infty$ .

- **P: Quantile tracking.** ACI can be regarded as a special case but it can sometimes output infinite or null prediction sets, while quantile tracking on the scale of the original score sequence does not have this behavior.
- **I: Error integration.**  $r_t$  should be a saturation function to achieve long-run coverage.
- **D: Scorecasting (forward-looking).** It trains a second model to predict the quantile of the next score.

The applications only consider one-step-ahead forecast.

## 7 Others

### 7.1 EnbPI: predictive inference method around ensemble estimators

Xu and Xie ([2021](#))

- For Dynamic time series
- Ensemble point forecasts + update residuals

### 7.2 SPCI: Sequential Predictive Conformal Inference

Xu and Xie ([2023](#))

- Adaptively re-estimate the conditional quantile of non-conformity scores, upon leveraging the temporal dependency among residuals. Random Forest for quantile regression is used.

### 7.3 Kath & Ziel (2021, IJF)

Kath and Ziel (2021)

#### Method 1.

The steps are:

1. Re-arrange time series data as form  $Z = \{(x_1, y_1), \dots, (x_L, y_L)\}$ .
2. Randomly split observations into subsets  $\pi$  (training set) and  $1 - \pi$  (calibration set).
3. Train a random forecast model using the training set and generate out-of-sample forecasts.
4. Calculate non-conformity score.
5. Obtain final prediction interval estimation.

#### Method 2.

The differences compared to Method 1 are:

1. Here an adjusted normalized non-conformity score  $\lambda_{i,h} = \frac{|y_{i,h} - \hat{y}_{i,h}|}{|\hat{\varepsilon}_{i,h}|}$  is used.
2. A second explicit error estimation model is trained to estimate the error associated with the prediction  $\hat{y}_{t,h}$ , i.e.,  $\varepsilon_{t,h}$ .
3. The new interval forecast is now given by  $y_{\alpha,T+1,h} = \hat{y}_{T+1,h} \pm \lambda_{L+1,h}^\alpha |\hat{\varepsilon}_{L+1,h}|$ .

The applications only consider one-step-ahead forecast.

### 7.4 TimeGPT

Garza and Mergenthaler-Canseco (2023)

For uncertainty quantification, they perform rolling forecasts on the latest available data to estimate the model's errors in forecasting the particular target time series.

The  $h$ -step-ahead forecast is generated based on  $h$ -step-ahead non-conformity errors on the calibration set separately.

### 7.5 Weighted quantile estimators

Akinshin (2023)

- The goal is to estimate the distribution at the tail of a time series.

**Desired properties of the weighted quantile estimators:**

- **Requirement R1:** consistency with existing quantile estimators.

$$Q^*(\mathbf{x}, \mathbf{w}, p) = Q(\mathbf{x}, p), \text{ where } \mathbf{w} = \{1, 1, \dots, 1\}.$$

- **Requirement R2:** zero weight support.

$$Q^*(\{x_1, x_2, \dots, x_{n-1}, x_n\}, \{w_1, w_2, \dots, w_{n-1}, 0\}, p) = Q^*(\{x_1, x_2, \dots, x_{n-1}\}, \{w_1, w_2, \dots, w_{n-1}\}, p).$$

- **Requirement R3:** stability (continuity of the quantile estimations with respect to the weight coefficient).

$$\lim_{\varepsilon_i \rightarrow 0} Q^*(\{x_1, x_2, \dots, x_n\}, \{w_1 + \varepsilon_1, w_2 + \varepsilon_2, \dots, w_n + \varepsilon_n\}, p) \rightarrow Q^*(\{x_1, x_2, \dots, x_n\}, \{w_1, w_2, \dots, w_n\}, p).$$

1. Some popular R implementations of weighted quantiles show violations of Requirement R3.
2. In order to satisfy Requirement R2, the sample size needs adjustments. Consider using Kish's effective sample size given by

$$n^*(\mathbf{w}) = \frac{(\sum_{i=1}^n w_i)^2}{\sum_{i=1}^n w_i^2} = \frac{1}{\sum_{i=1}^n \bar{w}_i^2},$$

where  $\mathbf{w}$  is the vector of non-negative weight coefficients,  $\bar{\mathbf{w}}$  is the vector of normalized weights where  $\bar{w}_i = \frac{w_i}{\sum_{i=1}^n w_i}$ .

### Method 1: Weighted Harrell–Davis quantile estimator

- a linear combination of all order statistics.
- pros: provide higher statistical efficiency in the cases of light-tailed distributions.
- cons: not robust (its breakdown point is zero).

The weighted Harrell–Davis quantile estimator is defined as

$$Q_{\text{HD}}^*(\mathbf{x}, \mathbf{w}, p) = \sum_{i=1}^n W_{\text{HD},i}^* \cdot x_{(i)}, \quad W_{\text{HD},i}^* = I_{t_i^*}(\alpha^*, \beta^*) - I_{t_{i-1}^*}(\alpha^*, \beta^*),$$

where  $\alpha^* = (n^* + 1)p$ ,  $\beta^* = (n^* + 1)(1 - p)$ ,  $t_i^* = s_i(\bar{\mathbf{w}})$ ,  $I_{t_i^*}(\alpha^*, \beta^*)$  is the regularized incomplete beta function. For all  $p \in (0; 1)$ ,  $Q_{\text{HD}}^*(\mathbf{x}, \mathbf{w}, p)$  satisfies R1, R2, and R3.

### Method 2: Weighted trimmed Harrell–Davis quantile estimator

- a trimmed modification.
- idea: since most of the linear coefficients  $W_{\text{HD},i}^*$  are pretty small, they do not have a noticeable impact on efficiency, but they significantly reduce the breakdown point.
- pros: allow customizing trade-off between robustness and efficiency

For  $p \in (0; 1)$ , the weighted trimmed Harrell–Davis quantile estimator based on the beta distribution highest density interval  $[L^*; R^*]$  of the given size  $D^*$  (the rule of thumb:  $D^* = 1/\sqrt{n^*}$ ) is defined as

$$Q_{\text{THD}}^*(\mathbf{x}, \mathbf{w}, p) = \sum_{i=1}^n W_{\text{THD},i}^* \cdot x_{(i)}, \quad W_{\text{THD},i}^* = F_{\text{THD}}^*(t_i^*) - F_{\text{THD}}^*(t_{i-1}^*), \quad t_i^* = s_i(\bar{\mathbf{w}}),$$

where

$$F_{\text{THD}}^*(t) = \begin{cases} 0 & \text{for } t < L^*, \\ (I_t(\alpha^*, \beta^*) - I_{L^*}(\alpha^*, \beta^*)) / (I_{R^*}(\alpha^*, \beta^*) - I_{L^*}(\alpha^*, \beta^*)) & \text{for } L^* \leq t \leq R^*, \\ 1 & \text{for } R^* < t. \end{cases}$$

### Method 3: Weighted traditional quantile estimators

- Hyndman and Fan (1996) summarized types 1-9.
  - Types 1–3 have discontinuities, so the corresponding estimators fail to satisfy R3.
  - Here only Types 4-9 are considered (using a linear combination of **two order statistics**).
- Pros: extremely robust, not so efficient.

Table: The Hyndman & Fan (1996) taxonomy of quantile estimators.

Type	h	Equation
1	$np$	$x_{(\lceil h \rceil)}$
2	$np + 1/2$	$(x_{(\lceil h-1/2 \rceil)} + x_{(\lceil h+1/2 \rceil)}) / 2$
3	$np$	$x_{(\lfloor h \rfloor)}$
4	$np$	$x_{(\lfloor h \rfloor)} + (h - \lfloor h \rfloor) (x_{(\lceil h \rceil)} - x_{(\lfloor h \rfloor)})$
5	$np + 1/2$	$x_{(\lfloor h \rfloor)} + (h - \lfloor h \rfloor) (x_{(\lceil h \rceil)} - x_{(\lfloor h \rfloor)})$
6	$(n+1)p$	$x_{(\lfloor h \rfloor)} + (h - \lfloor h \rfloor) (x_{(\lceil h \rceil)} - x_{(\lfloor h \rfloor)})$
7	$(n-1)p + 1$	$x_{(\lfloor h \rfloor)} + (h - \lfloor h \rfloor) (x_{(\lceil h \rceil)} - x_{(\lfloor h \rfloor)})$
8	$(n+1/3)p + 1/3$	$x_{(\lfloor h \rfloor)} + (h - \lfloor h \rfloor) (x_{(\lceil h \rceil)} - x_{(\lfloor h \rfloor)})$
9	$(n+1/4)p + 3/8$	$x_{(\lfloor h \rfloor)} + (h - \lfloor h \rfloor) (x_{(\lceil h \rceil)} - x_{(\lfloor h \rfloor)})$

These estimators can be rewritten in a form that matches the definition of  $Q_{\text{THD}}^*$  given by

$$Q_k^*(\mathbf{x}, \mathbf{w}, p) = \sum_{i=1}^n W_{F_k^*, i}^* \cdot x_{(i)}, \quad W_{F_k^*, i}^* = F_k^*(t_i^*) - F_k^*(t_{i-1}^*), \quad t_i^* = s_i(\bar{\mathbf{w}}),$$

where

$$F_k^*(t) = \begin{cases} 0 & \text{for } t < (h^* - 1)/n^*, \\ tn^* - h^* + 1 & \text{for } (h^* - 1)/n^* \leq t \leq h^*/n^*, \\ 1 & \text{for } t > h^*/n^*. \end{cases}$$

## 7.6 LCP: Localized conformal prediction

Guan (2022)

The **weight** on data point  $i$  is determined as **a function of the distance**  $\|X_i - X_{n+1}\|_2$ , to enable predictive coverage that holds **locally** (in neighborhoods of  $X$  space, that is, an approximation of prediction that holds conditional on the value of  $X_{n+1}$ ).

## 8 Simulation

### 8.1 Setup

Simulate a time series  $y$  with length  $T = 5000$  from an AR(2) model with  $\phi_1 = 0.8$ ,  $\phi_2 = -0.5$ , and  $\sigma^2 = 1$ .

Only consider one-step-ahead forecasting, i.e.,  $h = 1$ .

### 8.2 Split conformal prediction with fixed training set

#### 8.2.1 Details

Let  $n$  be the number of data used to fit an AR(2) model, and  $m$  be the number of data in the calibration set.

- Step 1. Rearrange the time series as  $Z_i = (X_i, Y_i)$ , where  $X_i = y_{i:(i+1)}$  and  $Y_i = y_{i+2}$ ,  $i = 1, 2, \dots, T - 2$ .
- Step 2. Train an AR(2) model  $\hat{\mu}$  based on training set with length  $n$ ,  $Z_{tr} = (Z_1, Z_2, \dots, Z_n)$ .




Figure 1: Simulated time series from an AR(2) model.

- Step 3. Calculate nonconformity scores (absolute residuals) based on calibration set with length  $m$ , i.e.,  $R_i = |Y_i - \hat{\mu}(X_i)|$ ,  $i = n + 1, n + 2, \dots, n + m$ .
- Step 4. Generate PI on test set.  $\hat{C}_m(X_i) = \hat{\mu}(X_i) \pm Q_{1-\alpha} \left( \sum_{j=n+1}^{n+m} w_j \cdot \delta_{R_j} + w_i \cdot \delta_{+\infty} \right)$  for  $i = n + m + 1, \dots, T - 2$ .

### 8.2.2 Results

Let  $n = m = 500$  and fit the AR(2) model using the `lm` function.

Consider methods: CP, WCP.LR, WCP.RF, NexCP with  $\alpha = 0.1$ .

Issues:

- PIs have constant width over the test set. We can generate PIs with varying local width by using a function to perform training on the absolute residuals i.e., to produce an estimator of  $E(R|X)$ .
- When generating weights via GLM or RF in WCP method, we need to use all data from the test set, which is not reasonable.
- Regression-based model.




Figure 2: Local coverage frequencies and width for split conformal prediction methods with fixed training set ( $k=500$ ).

### 8.3 Split conformal prediction with rolling training set

#### 8.3.1 Details

Let  $n$  be the number of observations used to fit an AR(2) model, and  $m$  be the number of observations in the calibration set.

For  $i = n + 1, n + 2, \dots, T$ :

- Step 1. Fit an AR(2) model  $\hat{\mu}_{i-1}$  based on observations  $y_{(i-n):(i-1)}$ , generate one-step-ahead forecast  $\hat{y}_i$ .
- When  $i > n + m$ :
  - Step 2. Calculate weights for the updated calibration set with length  $m$  and the updated test set with length 1 using different methods. Here WCP can not be applied because we only have a test set with length equal to one.
  - Step 3. Generate PI on test set.  $\hat{y}_i \pm Q_{1-\alpha} \left( \sum_{j=i-m}^{i-1} w_j \cdot \delta_{R_j} + w_i \cdot \delta_{+\infty} \right)$ .
  - Step 4 for ACP. Update  $\alpha$  based on the recent empirical miscoverage frequency.
- Step 5. Calculate nonconformity scores (absolute residuals)  $R_i = |y_i - \hat{y}_i|$ .

#### 8.3.2 Results

Let  $n = m = 500$  and fit AR(2) models using the `Arma` function with setting `order = c(2,0,0)`, `include.mean = TRUE`, `method = "CSS"` to make it comparable with the previous result.

Consider methods: AR, CP, NexCP, ACP with  $\alpha = 0.1$ .

Features:

- The fitted model is updating over time.
- PIs have changing width over the test set.
- Other non-regression-based models can be considered.
- Time-consuming because of the rolling window approach.






Figure 3: Local coverage frequencies and width for split conformal prediction methods with rolling training set and calibration set ( $k=500$ ).

## 9 To do list

- Multi-step ahead forecasting
- Other data-dependent weight.
- Papers from other journals such as IJF.
- Theoretical proof.
- Hierarchical time series.

## 10 References

- Akinshin, Andrey. 2023. “Weighted Quantile Estimators.” *arXiv Preprint arXiv:2304.07265*.
- Angelopoulos, Anastasios N, Emmanuel J Candes, and Ryan J Tibshirani. 2023. “Conformal PID Control for Time Series Prediction.” *arXiv Preprint arXiv:2307.16895*.
- Barber, Rina Foygel, Emmanuel J. Candès, Aaditya Ramdas, and Ryan J. Tibshirani. 2023. “Conformal Prediction Beyond Exchangeability.” *The Annals of Statistics* 51 (2). <https://doi.org/10.1214/23-aos2276>.
- Garza, Azul, and Max Mergenthaler-Canseco. 2023. “TimeGPT-1.” *arXiv Preprint arXiv:2310.03589*.
- Gibbs, Isaac, and Emmanuel Candes. 2021. “Adaptive Conformal Inference Under Distribution Shift.” *Advances in Neural Information Processing Systems* 34: 1660–72.
- Guan, Leying. 2022. “Localized Conformal Prediction: A Generalized Inference Framework for Conformal Prediction.” *Biometrika* 110 (1): 33–50. <https://doi.org/10.1093/biomet/asac040>.
- Hyndman, Rob J, and Yanan Fan. 1996. “Sample Quantiles in Statistical Packages.” *The American Statistician* 50 (4): 361–65.
- Kath, Christopher, and Florian Ziel. 2021. “Conformal Prediction Interval Estimation and Applications to Day-Ahead and Intraday Power Markets.” *International Journal of Forecasting* 37 (2): 777–99.
- Shafer, Glenn, and Vladimir Vovk. 2008. “A Tutorial on Conformal Prediction.” *Journal of Machine Learning Research* 9 (3).
- Stankeviciute, Kamile, Ahmed M Alaa, and Mihaela van der Schaar. 2021. “Conformal Time-Series Forecasting.” *Advances in Neural Information Processing Systems* 34: 6216–28.
- Tibshirani, Ryan J, Rina Foygel Barber, Emmanuel Candes, and Aaditya Ramdas. 2019. “Conformal Prediction Under Covariate Shift.” *Advances in Neural Information Processing Systems* 32.
- Vovk, Vladimir. 2013. “Cross-Conformal Predictors.” *Annals of Mathematics and Artificial Intelligence* 74 (1-2): 9–28. <https://doi.org/10.1007/s10472-013-9368-4>.
- Vovk, Vladimir, Alexander Gammerman, and Glenn Shafer. 2005. *Algorithmic Learning in a Random World*. Springer-Verlag. <https://doi.org/10.1007/b106715>.

- Xu, Chen, and Yao Xie. 2021. “Conformal Prediction Interval for Dynamic Time-Series.” In *International Conference on Machine Learning*, 11559–69. PMLR.
- . 2023. “Sequential Predictive Conformal Inference for Time Series.” In *International Conference on Machine Learning*, 38707–27. PMLR.