

# $k$ -contact reduction

B.M. Zawora

(joint work with J. de Lucas, X. Rivas, and S. Vilariño)

**in progress**

Department of Mathematical Methods in Physics,  
Faculty of Physics, University of Warsaw

Universidad de Zaragoza · Zaragoza

19.01.2024

- $k$ -symplectic manifolds
- $k$ -symplectic reduction
- $k$ -contact manifolds
- $k$ -contact reduction of the geometry
- $k$ -contact reduction of the dynamics
- an example of  $k$ -contact reduction

Let  $M$  be an  $n$ -dimensional manifold and let  $TM$  be its tangent bundle.

Let  $M$  be an  $n$ -dimensional manifold and let  $TM$  be its tangent bundle. Consider the Whitney sum of  $k$  copies of its tangent bundle,  $\bigoplus^k TM = TM \oplus_M \cdots \oplus_M^{(k)} TM$ , and the natural projections

$$\mathrm{pr}^\alpha: \bigoplus^k TM \rightarrow TM, \quad \mathrm{pr}_M: \bigoplus^k TM \rightarrow M, \quad \alpha = 1, \dots, k,$$

where  $\mathrm{pr}^\alpha$  denotes the projection onto the  $\alpha$ -th component of the Whitney sum.

Let  $M$  be an  $n$ -dimensional manifold and let  $TM$  be its tangent bundle. Consider the Whitney sum of  $k$  copies of its tangent bundle,  $\bigoplus^k TM = TM \oplus_M \cdots \oplus_M^{(k)} TM$ , and the natural projections

$$\text{pr}^\alpha: \bigoplus^k TM \rightarrow TM, \quad \text{pr}_M: \bigoplus^k TM \rightarrow M, \quad \alpha = 1, \dots, k,$$

where  $\text{pr}^\alpha$  denotes the projection onto the  $\alpha$ -th component of the Whitney sum.

## Definition

A  $k$ -vector field on a manifold  $M$  is a section  $\mathbf{X}: M \rightarrow \bigoplus^k TM$  of the projection  $\text{pr}_M$ . Then,  $\mathfrak{X}^k(M)$  denotes the set of all  $k$ -vector fields on  $M$ .

$$\begin{array}{ccc} TM & \xleftarrow{\text{pr}_\alpha} & \bigoplus^k TM \\ \uparrow x_\alpha & \nearrow \mathbf{x} & \\ M & & \end{array}$$

# $k$ -Contact reduction of the dynamics

Let  $M$  be an  $n$ -dimensional manifold and let  $TM$  be its tangent bundle. Consider the Whitney sum of  $k$  copies of its tangent bundle,  $\bigoplus^k TM = TM \oplus_M \cdots \oplus_M^{(k)} TM$ , and the natural projections

$$\text{pr}^\alpha: \bigoplus^k TM \rightarrow TM, \quad \text{pr}_M: \bigoplus^k TM \rightarrow M, \quad \alpha = 1, \dots, k,$$

where  $\text{pr}^\alpha$  denotes the projection onto the  $\alpha$ -th component of the Whitney sum.

## Definition

A  $k$ -vector field on a manifold  $M$  is a section  $\mathbf{X}: M \rightarrow \bigoplus^k TM$  of the projection  $\text{pr}_M$ . Then,  $\mathfrak{X}^k(M)$  denotes the set of all  $k$ -vector fields on  $M$ .

$$\begin{array}{ccc} TM & \xleftarrow{\text{pr}_\alpha} & \bigoplus^k TM \\ \uparrow X_\alpha & \nearrow \mathbf{X} & \\ M & & \end{array}$$

Taking into account the diagram above, a  $k$ -vector field  $\mathbf{X} \in \mathfrak{X}^k(M)$  amounts to some vector fields  $X_1, \dots, X_k \in \mathfrak{X}(M)$  given by  $X_\alpha = \text{pr}^\alpha \circ \mathbf{X}$  with  $\alpha = 1, \dots, k$ .



## Definition

Let  $h \in C^\infty(M)$  be a Hamiltonian function. A  $k$ -contact Hamiltonian  $k$ -vector field is  $\mathbf{X} = (X_1, \dots, X_k) \in \mathfrak{X}^k(M)$  satisfying

$$\begin{cases} \iota_{X_\alpha} d\eta^\alpha = dh - (R_\alpha h)\eta^\alpha, \\ \iota_{X_\alpha} \eta^\alpha = -h. \end{cases}$$



## Definition

Let  $h \in C^\infty(M)$  be a Hamiltonian function. A  $k$ -contact Hamiltonian  $k$ -vector field is  $\mathbf{X} = (X_1, \dots, X_k) \in \mathfrak{X}^k(M)$  satisfying

$$\begin{cases} \iota_{X_\alpha} d\eta^\alpha = dh - (R_\alpha h)\eta^\alpha, \\ \iota_{X_\alpha} \eta^\alpha = -h. \end{cases}$$

## Theorem

Let  $\mathbf{X}^h = (X_1^h, \dots, X_k^h)$  be a  $k$ -contact Hamiltonian  $k$ -vector field associated with a  $G$ -invariant function  $h \in C^\infty(M)$  relative to the Lie group action  $\Phi$ .

## Definition

Let  $h \in C^\infty(M)$  be a Hamiltonian function. A  $k$ -contact Hamiltonian  $k$ -vector field is  $\mathbf{X} = (X_1, \dots, X_k) \in \mathfrak{X}^k(M)$  satisfying

$$\begin{cases} \iota_{X_\alpha} d\eta^\alpha = dh - (R_\alpha h) \eta^\alpha, \\ \iota_{X_\alpha} \eta^\alpha = -h. \end{cases}$$

## Theorem

Let  $\mathbf{X}^h = (X_1^h, \dots, X_k^h)$  be a  $k$ -contact Hamiltonian  $k$ -vector field associated with a  $G$ -invariant function  $h \in C^\infty(M)$  relative to the Lie group action  $\Phi$ . Assume that  $\Phi_{g*} \mathbf{X}^h = \mathbf{X}^h$  for every  $g \in G$  and  $\mathbf{X}^h$  is tangent to  $\mathbf{J}^{-1}(\mathbb{R}^\times \mu)$ .

## Definition

Let  $h \in C^\infty(M)$  be a Hamiltonian function. A  $k$ -contact Hamiltonian  $k$ -vector field is  $\mathbf{X} = (X_1, \dots, X_k) \in \mathfrak{X}^k(M)$  satisfying

$$\begin{cases} \iota_{X_\alpha} d\eta^\alpha = dh - (R_\alpha h)\eta^\alpha, \\ \iota_{X_\alpha} \eta^\alpha = -h. \end{cases}$$

## Theorem

Let  $\mathbf{X}^h = (X_1^h, \dots, X_k^h)$  be a  $k$ -contact Hamiltonian  $k$ -vector field associated with a  $G$ -invariant function  $h \in C^\infty(M)$  relative to the Lie group action  $\Phi$ . Assume that  $\Phi_{g*} \mathbf{X}^h = \mathbf{X}^h$  for every  $g \in G$  and  $\mathbf{X}^h$  is tangent to  $\mathbf{J}^{-1}(\mathbb{R}^\times \mu)$ . Then, for every  $\alpha = 1, \dots, k$ , the flow  $F_s^\alpha$  of  $X_\alpha^h$  leave  $\mathbf{J}^{-1}(\mathbb{R}^\times \mu)$  invariant and induces a unique flow  $\mathcal{F}_s^\alpha$  on  $\mathbf{J}^{-1}(\mathbb{R}^\times \mu)/K_\mu$  satisfying  $\pi_\mu \circ F_s^\alpha = \mathcal{F}_s^\alpha \circ \pi_\mu$ .



## Example

Consider a manifold  $M = \bigoplus^2 T^*\mathbb{R} \times \mathbb{R}^2$  with coordinates  $(q^1, q^2, p_1^t, p_2^t, p_1^x, p_2^x, s^t, s^x)$ .

## Example

Consider a manifold  $M = \bigoplus^2 T^*\mathbb{R} \times \mathbb{R}^2$  with coordinates  $(q^1, q^2, p_1^t, p_2^t, p_1^x, p_2^x, s^t, s^x)$ . Then,  $(M, \eta)$  is a two-contact manifold, where

$$\eta = \eta^t \otimes e_1 + \eta^x \otimes e_2 = (ds^t - p_1^t dq^1 - p_2^t dq^2) \otimes e_1 + (ds^x - p_1^x dq^1 - p_2^x dq^2) \otimes e_2.$$

## Example

Consider a manifold  $M = \bigoplus^2 T^*\mathbb{R} \times \mathbb{R}^2$  with coordinates  $(q^1, q^2, p_1^t, p_2^t, p_1^x, p_2^x, s^t, s^x)$ . Then,  $(M, \boldsymbol{\eta})$  is a two-contact manifold, where

$$\boldsymbol{\eta} = \eta^t \otimes e_1 + \eta^x \otimes e_2 = (ds^t - p_1^t dq^1 - p_2^t dq^2) \otimes e_1 + (ds^x - p_1^x dq^1 - p_2^x dq^2) \otimes e_2.$$

The Reeb vector fields associated with  $\eta^t$  and  $\eta^x$  are  $R_t = \frac{\partial}{\partial s^t}$  and  $R_x = \frac{\partial}{\partial s^x}$ , respectively.

## Example

Consider a manifold  $M = \bigoplus^2 T^*\mathbb{R} \times \mathbb{R}^2$  with coordinates  $(q^1, q^2, p_1^t, p_2^t, p_1^x, p_2^x, s^t, s^x)$ . Then,  $(M, \eta)$  is a two-contact manifold, where

$$\eta = \eta^t \otimes e_1 + \eta^x \otimes e_2 = (ds^t - p_1^t dq^1 - p_2^t dq^2) \otimes e_1 + (ds^x - p_1^x dq^1 - p_2^x dq^2) \otimes e_2.$$

The Reeb vector fields associated with  $\eta^t$  and  $\eta^x$  are  $R_t = \frac{\partial}{\partial s^t}$  and  $R_x = \frac{\partial}{\partial s^x}$ , respectively. Consider a Lie group action

$$\begin{aligned} \Phi : \mathbb{R}^2 \times M &\ni (\lambda_1, \lambda_2; q^1, q^2, p_1^t, p_2^t, p_1^x, p_2^x, s^t, s^x) \\ &\mapsto (q^1 + \lambda_1, q^2 + \lambda_2, p_1^t, p_2^t, p_1^x, p_2^x, s^t, s^x + \lambda_2) \in M. \end{aligned}$$



## Example

Consider a manifold  $M = \bigoplus^2 T^*\mathbb{R} \times \mathbb{R}^2$  with coordinates  $(q^1, q^2, p_1^t, p_2^t, p_1^x, p_2^x, s^t, s^x)$ . Then,  $(M, \eta)$  is a two-contact manifold, where

$$\eta = \eta^t \otimes e_1 + \eta^x \otimes e_2 = (ds^t - p_1^t dq^1 - p_2^t dq^2) \otimes e_1 + (ds^x - p_1^x dq^1 - p_2^x dq^2) \otimes e_2.$$

The Reeb vector fields associated with  $\eta^t$  and  $\eta^x$  are  $R_t = \frac{\partial}{\partial s^t}$  and  $R_x = \frac{\partial}{\partial s^x}$ , respectively. Consider a Lie group action

$$\begin{aligned} \Phi : \mathbb{R}^2 \times M \ni (\lambda_1, \lambda_2; q^1, q^2, p_1^t, p_2^t, p_1^x, p_2^x, s^t, s^x) \\ \mapsto (q^1 + \lambda_1, q^2 + \lambda_2, p_1^t, p_2^t, p_1^x, p_2^x, s^t, s^x + \lambda_2) \in M. \end{aligned}$$

The fundamental vector fields associated with  $\Phi : \mathbb{R}^2 \times M \rightarrow M$  read

$$\xi_M^1 = \frac{\partial}{\partial s^x}, \quad \xi_M^2 = \frac{\partial}{\partial q^1} + \frac{\partial}{\partial q^2},$$

## Example

Consider a manifold  $M = \bigoplus^2 T^*\mathbb{R} \times \mathbb{R}^2$  with coordinates  $(q^1, q^2, p_1^t, p_2^t, p_1^x, p_2^x, s^t, s^x)$ . Then,  $(M, \eta)$  is a two-contact manifold, where

$$\eta = \eta^t \otimes e_1 + \eta^x \otimes e_2 = (ds^t - p_1^t dq^1 - p_2^t dq^2) \otimes e_1 + (ds^x - p_1^x dq^1 - p_2^x dq^2) \otimes e_2.$$

The Reeb vector fields associated with  $\eta^t$  and  $\eta^x$  are  $R_t = \frac{\partial}{\partial s^t}$  and  $R_x = \frac{\partial}{\partial s^x}$ , respectively. Consider a Lie group action

$$\begin{aligned} \Phi : \mathbb{R}^2 \times M \ni (\lambda_1, \lambda_2; q^1, q^2, p_1^t, p_2^t, p_1^x, p_2^x, s^t, s^x) \\ \mapsto (q^1 + \lambda_1, q^2 + \lambda_2, p_1^t, p_2^t, p_1^x, p_2^x, s^t, s^x + \lambda_2) \in M. \end{aligned}$$

The fundamental vector fields associated with  $\Phi : \mathbb{R}^2 \times M \rightarrow M$  read

$$\xi_M^1 = \frac{\partial}{\partial s^x}, \quad \xi_M^2 = \frac{\partial}{\partial q^1} + \frac{\partial}{\partial q^2},$$

and the contact momentum map  $\mathbf{J}$  is given by

$$\mathbf{J} : M \ni x \mapsto \mu = \mu^1 \otimes e_1 + \mu^2 \otimes e_2 = (0, 1) \otimes e_1 - (p_1^t + p_2^t, p_1^x + p_2^x) \otimes e_2 \in (\mathbb{R}^2)^{*2}.$$

## Example

Consider a manifold  $M = \bigoplus^2 T^*\mathbb{R} \times \mathbb{R}^2$  with coordinates  $(q^1, q^2, p_1^t, p_2^t, p_1^x, p_2^x, s^t, s^x)$ . Then,  $(M, \eta)$  is a two-contact manifold, where

$$\eta = \eta^t \otimes e_1 + \eta^x \otimes e_2 = (ds^t - p_1^t dq^1 - p_2^t dq^2) \otimes e_1 + (ds^x - p_1^x dq^1 - p_2^x dq^2) \otimes e_2.$$

The Reeb vector fields associated with  $\eta^t$  and  $\eta^x$  are  $R_t = \frac{\partial}{\partial s^t}$  and  $R_x = \frac{\partial}{\partial s^x}$ , respectively. Consider a Lie group action

$$\begin{aligned} \Phi : \mathbb{R}^2 \times M \ni (\lambda_1, \lambda_2; q^1, q^2, p_1^t, p_2^t, p_1^x, p_2^x, s^t, s^x) \\ \mapsto (q^1 + \lambda_1, q^2 + \lambda_2, p_1^t, p_2^t, p_1^x, p_2^x, s^t, s^x + \lambda_2) \in M. \end{aligned}$$

The fundamental vector fields associated with  $\Phi : \mathbb{R}^2 \times M \rightarrow M$  read

$$\xi_M^1 = \frac{\partial}{\partial s^x}, \quad \xi_M^2 = \frac{\partial}{\partial q^1} + \frac{\partial}{\partial q^2},$$

and the contact momentum map  $\mathbf{J}$  is given by

$$\mathbf{J} : M \ni x \mapsto \mu = \mu^1 \otimes e_1 + \mu^2 \otimes e_2 = (0, 1) \otimes e_1 - (p_1^t + p_2^t, p_1^x + p_2^x) \otimes e_2 \in (\mathbb{R}^2)^{*2}.$$

Let us fix  $\mu = \mu_1 \otimes e_1$  and recall that

$$T_x \mathbf{J}^{-1}(\mathbb{R}^\times \mu) = \{v_x \in T_x M : T_x \mathbf{J}(v_x) = \lambda \mu, \quad \lambda \in \mathbb{R}^\times\},$$

for every  $x \in \mathbf{J}^{-1}(\mathbb{R}^\times \mu)$ .

## Example

Then,

Then,

$$\begin{aligned}\mathbf{J}^{-1}(\mathbb{R}^\times \boldsymbol{\mu}) &= \{x \in M : p_1^t = -p_2^t, \quad p_1^x = -p_2^x\}, \\ T_x \mathbf{J}^{-1}(\mathbb{R}^\times \boldsymbol{\mu}) &= \left\langle \frac{\partial}{\partial s^x}, \frac{\partial}{\partial s^t}, \frac{\partial}{\partial q^1}, \frac{\partial}{\partial q^2}, \frac{\partial}{\partial p_1^x} - \frac{\partial}{\partial p_2^x}, \frac{\partial}{\partial p_1^t} - \frac{\partial}{\partial p_2^t} \right\rangle. \\ T_x(K_\mu x) &= \left\langle \frac{\partial}{\partial q^1} + \frac{\partial}{\partial q^2} \right\rangle.\end{aligned}$$

Then,

$$\begin{aligned}\mathbf{J}^{-1}(\mathbb{R}^\times \mu) &= \{x \in M : p_1^t = -p_2^t, \quad p_1^x = -p_2^x\}, \\ T_x \mathbf{J}^{-1}(\mathbb{R}^\times \mu) &= \left\langle \frac{\partial}{\partial s^x}, \frac{\partial}{\partial s^t}, \frac{\partial}{\partial q^1}, \frac{\partial}{\partial q^2}, \frac{\partial}{\partial p_1^x} - \frac{\partial}{\partial p_2^x}, \frac{\partial}{\partial p_1^t} - \frac{\partial}{\partial p_2^t} \right\rangle. \\ T_x(K_\mu x) &= \left\langle \frac{\partial}{\partial q^1} + \frac{\partial}{\partial q^2} \right\rangle.\end{aligned}$$

Then, it follows that the reduced manifold  $M_\mu = (\mathbf{J}^{-1}(\mathbb{R}^\times \mu)/K_\mu, \eta_\mu)$  is a 2-contact manifold with

$$\eta_\mu = \eta_\mu^t \otimes e_1 + \eta_\mu^x \otimes e_2 = (ds^t - p^t dq) \otimes e_1 + (ds^x - p^x dq) \otimes e_2,$$

Then,

$$\begin{aligned}\mathbf{J}^{-1}(\mathbb{R}^\times \mu) &= \{x \in M : p_1^t = -p_2^t, \quad p_1^x = -p_2^x\}, \\ T_x \mathbf{J}^{-1}(\mathbb{R}^\times \mu) &= \left\langle \frac{\partial}{\partial s^x}, \frac{\partial}{\partial s^t}, \frac{\partial}{\partial q^1}, \frac{\partial}{\partial q^2}, \frac{\partial}{\partial p_1^x} - \frac{\partial}{\partial p_2^x}, \frac{\partial}{\partial p_1^t} - \frac{\partial}{\partial p_2^t} \right\rangle. \\ T_x(K_\mu x) &= \left\langle \frac{\partial}{\partial q^1} + \frac{\partial}{\partial q^2} \right\rangle.\end{aligned}$$

Then, it follows that the reduced manifold  $M_\mu = (\mathbf{J}^{-1}(\mathbb{R}^\times \mu)/K_\mu, \eta_\mu)$  is a 2-contact manifold with

$$\eta_\mu = \eta_\mu^t \otimes e_1 + \eta_\mu^x \otimes e_2 = (ds^t - p^t dq) \otimes e_1 + (ds^x - p^x dq) \otimes e_2,$$

where  $(q := q^1 - q^2, p^t := p_1^t - p_2^t, p^x := p_1^x - p_2^x, s^t, s^x)$  are local coordinates on  $M_\mu$ .

Then,

$$\begin{aligned}\mathbf{J}^{-1}(\mathbb{R}^\times \mu) &= \{x \in M : p_1^t = -p_2^t, \quad p_1^x = -p_2^x\}, \\ T_x \mathbf{J}^{-1}(\mathbb{R}^\times \mu) &= \left\langle \frac{\partial}{\partial s^x}, \frac{\partial}{\partial s^t}, \frac{\partial}{\partial q^1}, \frac{\partial}{\partial q^2}, \frac{\partial}{\partial p_1^x} - \frac{\partial}{\partial p_2^x}, \frac{\partial}{\partial p_1^t} - \frac{\partial}{\partial p_2^t} \right\rangle. \\ T_x(K_\mu x) &= \left\langle \frac{\partial}{\partial q^1} + \frac{\partial}{\partial q^2} \right\rangle.\end{aligned}$$

Then, it follows that the reduced manifold  $M_\mu = (\mathbf{J}^{-1}(\mathbb{R}^\times \mu)/K_\mu, \eta_\mu)$  is a 2-contact manifold with

$$\eta_\mu = \eta_\mu^t \otimes e_1 + \eta_\mu^x \otimes e_2 = (ds^t - p^t dq) \otimes e_1 + (ds^x - p^x dq) \otimes e_2,$$

where  $(q := q^1 - q^2, p^t := p_1^t - p_2^t, p^x := p_1^x - p_2^x, s^t, s^x)$  are local coordinates on  $M_\mu$ . Let  $h \in C^\infty(M)$  be a Hamiltonian function defined as

$$h(q^1, q^2, p_1^t, p_2^t, p_1^x, p_2^x, s^t, s^x) = \frac{1}{2} \left( (p_1^t)^2 + (p_2^t)^2 - (p_1^x)^2 - (p_2^x)^2 \right) + C(q^1 - q^2) + \gamma s^t,$$



Then,

$$\begin{aligned}\mathbf{J}^{-1}(\mathbb{R}^\times \mu) &= \{x \in M : p_1^t = -p_2^t, \quad p_1^x = -p_2^x\}, \\ T_x \mathbf{J}^{-1}(\mathbb{R}^\times \mu) &= \left\langle \frac{\partial}{\partial s^x}, \frac{\partial}{\partial s^t}, \frac{\partial}{\partial q^1}, \frac{\partial}{\partial q^2}, \frac{\partial}{\partial p_1^x} - \frac{\partial}{\partial p_2^x}, \frac{\partial}{\partial p_1^t} - \frac{\partial}{\partial p_2^t} \right\rangle. \\ T_x(K_\mu x) &= \left\langle \frac{\partial}{\partial q^1} + \frac{\partial}{\partial q^2} \right\rangle.\end{aligned}$$

Then, it follows that the reduced manifold  $M_\mu = (\mathbf{J}^{-1}(\mathbb{R}^\times \mu)/K_\mu, \eta_\mu)$  is a 2-contact manifold with

$$\eta_\mu = \eta_\mu^t \otimes e_1 + \eta_\mu^x \otimes e_2 = (ds^t - p^t dq) \otimes e_1 + (ds^x - p^x dq) \otimes e_2,$$

where  $(q := q^1 - q^2, p^t := p_1^t - p_2^t, p^x := p_1^x - p_2^x, s^t, s^x)$  are local coordinates on  $M_\mu$ . Let  $h \in C^\infty(M)$  be a Hamiltonian function defined as

$$h(q^1, q^2, p_1^t, p_2^t, p_1^x, p_2^x, s^t, s^x) = \frac{1}{2} \left( (p_1^t)^2 + (p_2^t)^2 - (p_1^x)^2 - (p_2^x)^2 \right) + C(q^1 - q^2) + \gamma s^t,$$

where  $C(q^1 - q^2)$  is a coupling function between the two strings.

The integral sections of the two-vector field  $\mathbf{X}$  associated with  $h$  satisfy equations

The integral sections of the two-vector field  $\mathbf{X}$  associated with  $h$  satisfy equations

$$\begin{aligned}\frac{\partial^2 q^1}{\partial t^2} - \frac{\partial^2 q^1}{\partial x^2} &= -\gamma \frac{\partial q^1}{\partial t} - \frac{\partial C}{\partial q}, \\ \frac{\partial^2 q^2}{\partial t^2} - \frac{\partial^2 q^2}{\partial x^2} &= -\gamma \frac{\partial q^2}{\partial t} + \frac{\partial C}{\partial q}.\end{aligned}$$

The integral sections of the two-vector field  $\mathbf{X}$  associated with  $h$  satisfy equations

$$\begin{aligned}\frac{\partial^2 q^1}{\partial t^2} - \frac{\partial^2 q^1}{\partial x^2} &= -\gamma \frac{\partial q^1}{\partial t} - \frac{\partial C}{\partial q}, \\ \frac{\partial^2 q^2}{\partial t^2} - \frac{\partial^2 q^2}{\partial x^2} &= -\gamma \frac{\partial q^2}{\partial t} + \frac{\partial C}{\partial q}.\end{aligned}$$

This system of PDEs corresponds to two coupled damped vibrating strings.

The integral sections of the two-vector field  $\mathbf{X}$  associated with  $h$  satisfy equations

$$\begin{aligned}\frac{\partial^2 q^1}{\partial t^2} - \frac{\partial^2 q^1}{\partial x^2} &= -\gamma \frac{\partial q^1}{\partial t} - \frac{\partial C}{\partial q}, \\ \frac{\partial^2 q^2}{\partial t^2} - \frac{\partial^2 q^2}{\partial x^2} &= -\gamma \frac{\partial q^2}{\partial t} + \frac{\partial C}{\partial q}.\end{aligned}$$

This system of PDEs corresponds to two coupled damped vibrating strings.  
After the  $k$ -contact reduction we get

$$\frac{\partial^2 q}{\partial t^2} - \frac{\partial^2 q}{\partial x^2} = \gamma \frac{\partial q}{\partial t} - 2 \frac{\partial C}{\partial q},$$

which is the equation of a single damped string with an external force acting on it.

Thank you for your attention

# Thank you for your attention

## Literature:

- de Lucas J., Rivas X., Vilariño S., Zawora B.M. *k-Contact Marsden-Weinstein reduction*, work in progress.
- J. C. Marrero, N. Román-Roy, M. Salgado, and S. Vilariño, *Reduction of polysymplectic manifolds*, J. Phys. A **48** (2015) 055206, 43.
- Grabowska K., Grabowski J. *Reductions: precontact versus presymplectic*, Annali di Matematica Pura ed Applicata, **202** (2023), 2803–2839.
- Willett C. *Contact reduction*. Trans. Amer. Math. Soc., **10** (2002), 4245–4260.
- Gaset J., Gràcia X., Muñoz-Lecanda M.C., Rivas X., Román-Roy N. *A contact geometry framework for field theories with dissipation*, Ann. Physics, **414** (2020), 168092.