

# XRONOS: An Open Data Infrastructure for Archaeological Chronology

Joe Roe<sup>a,\*</sup>, Clemens Schmid<sup>b</sup>, Setareh Ebrahimiabareghi<sup>a</sup>, Caroline Heitz<sup>a</sup>,  
Martin Hinz<sup>a,\*</sup>

<sup>a</sup>*University of Bern, Institute of Archaeological Sciences,*

<sup>b</sup>*Max Planck Institute for Evolutionary Anthropology,*

---

## Abstract

XRONOS (<https://xronos.ch>) is an open data infrastructure for the backbone of the archaeological record – chronology. It provides open access to published radiocarbon dates and other chronometric data from any period, anywhere in the world. By collating a large number of existing regional and global compilations of dates, XRONOS offers the most comprehensive radiocarbon database yet published, with over 350,000 radiocarbon and 75,000 site records. It also provides a foundation for expanding the systematic collection of chronometric information beyond radiocarbon, with support for typological and dendrochronological dates and a generalisable data model that can be adapted to other methods of absolute dating. Automated and semi-automated quality control processes ensure that data from diverse sources is continuously integrated and standardised, making it easier to find information of interest and reducing the need for manual data cleaning by end users. In this paper we describe the concept and implementation of XRONOS in relation to the state of the art in chronometric data-sharing, and evaluate its potential as a general-purpose open repository and curation platform for archaeological chronology.

*Keywords:* open data, chronology, chronometry, radiocarbon dating, dendrochronology, typological dating

---

## 1. Introduction

Chronology is the backbone of the archaeological record. As a necessary prerequisite to understanding the context of any past event or process (Lucas, 2004), it has unsurprisingly been at the forefront of methodological development in archaeology for as long as the discipline has existed: from putting finds and events

---

\*Corresponding author

Email addresses: [joeroe@hey.com](mailto:joeroe@hey.com) (Joe Roe), [martin.hinz@unibe.ch](mailto:martin.hinz@unibe.ch) (Martin Hinz)

in sequence (Ford, 1962; Harris, 1979; Petrie, 1899; Thomsen, 1836; Worsaae, 1843) to an increasingly wide array of scientific methods that place them on an absolute timescale (Bada and Helfman, 1975; Daniels et al., 1953; Douglass, 1929; Evernden et al., 1965; Libby, 1955) and an increasingly sophisticated set of statistical tools to build them into chronologies (Buck et al., 1991; Crema, 2024; Levy et al., 2021; Mischka, 2004; Suess, 1967). If archaeology is to be an open science (Lake, 2012), it is therefore critical that effective open access to chronological information be placed front and centre.

Over the last two decades, archaeologists have answered this call by publishing an increasing number of compilations of dates from archaeological contexts as open data. These efforts have facilitated re-evaluations of chronologies themselves (e.g. Higham et al., 2014; Katsianis et al., 2020; Loftus et al., 2019; Prates et al., 2020) but also the development of novel ways of using chronological data (Crema et al., 2024; Crema, 2022; e.g. Grove, 2011; Marom and Wolkowski, 2024; Riris et al., 2024; Silva and Steele, 2014). The focus has been overwhelmingly on radiocarbon dating and most compilations focus on a single region and/or period. The profusion of open radiocarbon data in particular has prompted several initiatives towards a global synthesis (Bird et al., 2022; Bronk Ramsey et al., 2019; e.g. Schmid et al., 2019).

At the same time, the broad range of other types of chronological information used in archaeology—from other radiometric methods to dendrochronology to typological dating and epigraphy—remains relatively difficult to access as open data. Even when it comes to radiocarbon data, the coverage of available compilations is patchy both geographically and in time and of variable quality (see Section 2.1). The publication of many overlapping, non-standardised and mostly static open data resources means that it is still difficult to obtain reliable and up-to-date chronological datasets, especially for applications that crosscut conventional geographic and temporal domains of research. Initiatives towards synthesis have improved this situation, but the goal of a global dataset that is both comprehensive and up-to-date remains elusive.

XRONOS is a new open data infrastructure that aims to provide access to published radiocarbon dates and other chronometric data from any period, anywhere in the world. It is our attempt to move the state of the art in open archaeological chronology beyond the publication of static, one-off resources (‘uploading CSVs,’ Batist, 2023, pp. 188–189), and towards a living digital infrastructure (Kintigh, 2006) embedded in a transparent and sustainable collaborative network. The core of XRONOS is a server application that ingests chronological data from diverse sources, facilitates semi-automated and manual curation of this data, and makes it available via both a web-based graphical user interface (GUI) and machine-readable application programming interface (API). The web frontend can be accessed via <https://xronos.ch> and all components of the software are developed as free and open source software with source code available at <https://github.com/xronos-ch>.

In the remainder of this paper, we describe the concept and implementation

Table 1: Summary of published compilations of radiocarbon dates. For full data, see supplementary materials.

[1] "omitted from manuscript - see attached file `table1.html`"

of XRONOS in relation to the state of the art in open chronometric data in archaeology, and evaluate our progress in achieving these goals as of writing. Since we envisage both XRONOS as a dataset and XRONOS as software to be continually developing resources, the description here should be read as a ‘snapshot’ of the project as of [version 1.1.0](#).

## 2. State of the Art

### 2.1. Compilations of radiocarbon dates

Though an *explicit* emphasis on ‘open data’ is a relatively recent phenomenon in archaeology (Lake, 2012), the open publication of compiled radiocarbon dates has a substantial prehistory. Arnold and Libby (Arnold and Libby, 1951) initiated the tradition of regularly publishing ‘data lists’, a practice was subsequently continued by radiocarbon laboratories as supplements to journals such as *Radiocarbon* and *Archaeometry*. However, as the number of labs and volume of radiocarbon dates being produced grew, this paper-based format became impractical and mostly disappeared (Bronk Ramsey et al., 2019; c.f. e.g. Ndeye et al., 2022), without being replaced by another form of systematic data-sharing or dissemination. Additionally, because date lists were sourced from radiocarbon laboratories directly—not from those who collected the sample—they typically included only very limited contextual information. On the eve of the AMS revolution there was an effort to create a computerised ‘International Radiocarbon Database’ (Kra, 1988)—already by 1989 described as a “much needed, long overdue enterprise” (Kra, 1989, p. 1067)—but it never came to fruition.

Thus, even though radiocarbon data comes from a relatively limited number of sources (some 172 active labs, Radiocarbon, 2024) and has relatively standardised reporting conventions (Bayliss, 2015; Millard, 2014), in practice the only way to produce aggregated datasets in recent decades has been to manually search through relevant literature for dates reported secondarily by the submitter of the sample. This already laborious process is further hampered by a significant inconsistency in how much authors adhere to reporting conventions for measurements and sample metadata, a lack of conventions on the reporting of *contextual* information, weak or nonexistent disciplinary norms regarding the responsibility to publish results openly in a timely fashion, and a range of other issues affecting data reuse (Moody et al., 2021).

Despite these inefficiencies, there have been a profusion of published radiocarbon compilations since the decline of the date list. Our review of the literature identified 61 published since 1994 (Table Table 1 and supplementary materials). This is almost certainly an undercount, because our firsthand knowledge

of regional literature is limited to Europe and West Asia and many resources only ever existed in ‘grey’ formats (e.g. websites that were not indexed and no longer exist). We also restricted ourselves to structured datasets disseminated primarily in a digital format; ‘date lists’ in printed periodicals and gazetteers were excluded.




Figure 1: Cumulative number of radiocarbon compilations published since 1995

The number of available compilations has increased significantly since around 1995 (Figure 1). The first generation came around the turn of the century and consists mostly of online databases with a web frontend. These include some databases operated by radiocarbon labs, for example the Oxford Radiocarbon Lab (ORAU) and the Belgian Royal Institute for Cultural Heritage (KIK-IRPA), and essentially represent a continuation of their date lists in a digital format. The majority, however, were compiled from the literature by individual researchers interested in a particular region and/or period. Notable early examples include ANDES 14C in 1994 (Central Andes, Michczyński et al., 1995), CARD (Canada, Gajewski et al., 2011) and RADON (Europe, Raetzl-Fabian, 1999) in 1999, and CANEW in 2001 (Near East, Reingruber and Thissen, 2005). From 2010, coinciding with broader shifts in scientific publishing (Tenopir et al., 2011), it became more common to publish standalone ‘open data’ products in the form of journal supplements, archives in repositories and/or data papers; the *Journal of Open Archaeology Data*, launched in 2012, has been a prominent venue for this latter category. Most recently there has been a trend towards providing version-controlled plain text data via platforms such as [GitHub](#), reflecting the broader adoption of these tools amongst computational archaeologists over the last decade (Batist and Roe, 2024). The shift from online databases to

wards more static but more preservable open data products is welcome, given how many databases from the first generation have subsequently ceased to be accessible. Version-controlled repositories are particular well-suited to data compilation projects because they allow for continued updates whilst still providing snapshot ‘releases’ that are citeable and can be archived in long-term repositories.




Figure 2: Geographic coverage of published regional radiocarbon compilations according to our survey (binned by UN M49 macroregion).

Although this body of work has greatly improved the accessibility of radiocarbon dates and supported significant methodological advances (Crema et al., 2024; Crema, 2022), some limitations are apparent.

The geographic coverage of regional radiocarbon compilations is markedly uneven (Figure 2). Europe and, especially, North America are over-represented (Alcántara and Pedroza, in press; Chaput and Gajewski, 2016). South America, West Asia, and East Asia are reasonably well-covered, but there practically no systematically compiled dates from East or West Africa, Central or South Asia, or Mainland Southeast Asia. This is probably explained in part by a lower volume of archaeological research and access to radiocarbon dating in these regions, but a lack of attention in compilation work must also be a factor. For example, radiocarbon dating has been an established part of Indian archaeology since at least 1961 (Kusumgar et al., 1963), but we have not able to locate a single systematic compilation of dates from South Asia.<sup>1</sup>

---

<sup>1</sup>We would be very happy to be corrected on this point.

Datasets based on literature review also become out of date almost immediately upon publication, due to the constant production of new dates. Unfortunately this applies to many databases that are in theory continuously updated, as it is common to see them become unmaintained and or unexpectedly become unavailable. Of the 61 published datasets we identified, 33 were intended to be continuously updated, but only 13 have received updates in the last two years. The mean ‘lifespan’ of a dataset from its publication to its last update is around 4 years. Most radiocarbon datasets we reviewed were compiled with a specific goal in mind (e.g. a particular analysis) and, even where there is the intention to keep them updated afterwards, the exigencies of scientific production combined with the labour-intensive nature of the process make that difficult to achieve in practice.

Laboratory databases solve the problem of currency, but tend to have more arbitrary coverage, since the inclusion of data is determined by who submits dates to that lab, not any form of principled curation. There are also comparatively few of them – most active labs no longer directly publish dates that they produce (if they ever did).

Other outstanding problems with existing compilations include various systematic biases in data collection (Clist et al., 2023) and a large degree of overlap and duplication between individual databases. For example, we identified 9 different resources covering Western Europe but none covering South Asia. The quality and accessibility of published compilations is also variable. 50 of the 61 resources we reviewed are not ‘open’ according to the Open Knowledge Foundation’s definition of data openness (“Open data and content can be freely used, modified, and shared by anyone for any purpose,” Open Knowledge Foundation, n.d.), which both limits the access to and reuse potential of these datasets. And even of these, many are not currently available in readily machine-readable formats (e.g. plain text or database files rather than PDFs or hypertext).

The fragmentation of the radiocarbon record into regional datasets also hinders analysis at larger scales. Although the core elements of a radiocarbon date—laboratory identifier, radiocarbon age, measurement error—are more or less standardised, there is no such consistency in contextual information on the sample or site. Such contextual information is important not just for the interpretation of dates, but for filtering out unreliable dates based on sample information (‘chronometric hygiene’ sensu Pettitt et al., 2003) and for correcting for known systematic errors such as the marine reservoir effect (Alves et al., 2018). Most published datasets incorporate all or part of earlier compilations, meaning duplicate records are also very common, but deduplicating them is not a trivial problem due to format variations (see Section 4.2). These issues are by no means impossible to overcome, but adds a significant amount of data-cleaning effort to a process that would otherwise be very amenable to standardisation.




Figure 3: Geographic (quantile kernel density) and temporal (sum calibration) of georeferenced dates in XRONOS and other global radiocarbon compilations

## 2.2. Global radiocarbon compilations

The profusion of radiocarbon compilations over the last decade has naturally prompted many to think globally. Three existing initiatives in particular share similar aims to XRONOS (at least as far as radiocarbon is concerned): c14bazAAR, IntChron, and p3k14c.

The first available synthetic radiocarbon database was c14bazAAR (Schmid et al., 2019), an R package that provides an index of openly published radiocarbon databases and a common interface for retrieving them and performing basic data cleaning. Because c14bazAAR downloads data from its original source repositories, rather than mirroring it, it only includes resources that have been published in a fully open and machine-accessible format. Despite this limitation, it has global coverage and a large number of dates (Figure 3), and was therefore our starting point for data collection for XRONOS.

Another indexical approach is taken by the IntChron project (Bronk Ramsey et al., 2019), which exposes data from multiple sources and exposes them with a common JSON-based web interface. The IntChron specification is open, meaning that radiocarbon labs or compilation projects can implement it independently and thereby allow end users to access their data through a common interface (though to our knowledge it has so far only been adopted by databases associated with the Oxford Radiocarbon Lab). The JSON format also lends itself to the implementation of wrapper libraries, for example the rIntChron package gives direct access to IntChron-indexed databases in R (Roe, 2024).

p3k14c (Bird et al., 2022) instead compiles multiple source databases into a single flat file dataset, with a similar level of coverage to c14bazAAR. The major advantage of this approach is that the data is made internally consistent and has been manually cleaned to an extent, which makes it particularly well-suited to global analyses. The downside is that without the continuous link to the source databases present in the c14bazAAR and IntChron, it can only be kept up to date manually with periodic re-releases. An accompanying package (Bird et al., 2024) provides direct access to the p3k14c dataset in R.

As of December 2024, c14bazAAR had 118,071 radiocarbon dates with unique laboratory identifiers (excluding those sourced from p3k14c and XRONOS), IntChron had 12,388 (excluding those from non-archaeological contexts), and p3k14c had 176,016. The geographic distribution of dates from each is similar (Figure 3), reflecting the large degree in overlap between the sources of each compilation. IntChron, which in practice is currently only used to publish dates associated with the Oxford Radiocarbon Lab, has dates from more diverse contexts, but is an order of magnitude smaller.

## 2.3. Beyond radiocarbon

Radiocarbon has been by far the most active area of open data compilation, but archaeological chronology incorporates a much more diverse range of sources of information (Harding, 1999). In periods beyond the practical limit of radiocarbon dating (c. 55,000 BP), other types of radiometric (K–Ar, U–Pb, etc.),



chemical or luminescence dating offer an alternative (Aitken, 1999). Conversely, in historic periods, or on calibration ‘plateaus’ (Guilderson et al., 2005), radiocarbon is often less important than conventional typological dating (based on artefact characteristics), which in these periods can offer comparable or better temporal resolution, or direct dating based on epigraphy (Heřmánková et al., 2021), numismatics (Kemmers and Myrberg, 2011) or historical sources. In places where it is widely available, dendrochronology (Baillie, 2014) also produces significantly better resolved chronologies and therefore tends to be the main source of chronometric data. Other more application-specific chronological methods include shoreline dating (Brøgger, 1905; Roalkvam, 2023), lichenometry (Benedict, 2009) and rock weathering dating (Andreae et al., 2020; Bednarik, 2020; Whitley, 2012).

Compared to radiocarbon, there are few examples of systematic, open compilations of any of these other types of data. This is most striking when it comes to other radiometric/scientific dating methods, as the data structures and publication modes are very similar to radiocarbon. The ‘Radiocarbon Palaeolithic Europe Database’ (Vermeersch, 2020), despite the name, includes a significant number of thermo- and optically stimulated luminescence, electron spin resonance, uranium–thorium and amino acid dates. Similarly, the AustArch database (Williams et al., 2014) includes luminescence dates alongside radiocarbon, but is limited to Australia and was last updated in 2013. Apart from these and a few other exceptions where other scientific dates are collected alongside radiocarbon, we are not aware of any open compilations of them.

### *2.3.1. Dendrochronology*

With regard to tree-ring data, some databases provide valuable resources for dendrochronological studies in general but are not primarily intended for archaeological contexts. For instance, Dendro4Art specializes in dendrochronological data related to wooden art objects, such as sculptures and panel paintings. While this focus serves art historians and conservationists well, its utility for studying prehistoric datasets is minimal. Similarly, the Dendrochronological Picture Database, maintained by the Swiss Federal Institute for Snow and Avalanche Research (SLF), offers a visual archive of approximately 1,400 images documenting dendrochronological phenomena. Although valuable as an educational resource, it does not provide raw data necessary for chronological or archaeological analyses. Additionally, the OLDLIST and Eastern OLDLIST databases focus on documenting the maximum ages of trees worldwide. Their emphasis on biological longevity, while significant for ecological research, limits their applicability to archaeological or prehistoric investigations.

Among databases that do provide dendrochronological data, the degree to which they support prehistoric research varies substantially. The NOAA International Tree-Ring Data Bank (ITRDB) serves as a global repository of tree-ring measurements. However, its focus remains predominantly on North America, with only 34 datasets representing European prehistoric contexts. This restricts its relevance for studying the European past. Similarly, the ADS

database, maintained by the UK-based Vernacular Architecture Group, compiles dendrochronological data from the UK but is limited to medieval and later periods, making it unsuitable for prehistoric studies.

DendroDB, hosted by the Swiss Federal Institute for Forest, Snow and Landscape Research (WSL), emphasizes ecological and climate studies over archaeological wood material. The CFS-TRenD database, managed by the Canadian Forest Service, compiles over 4,600 datasets from Canadian forests, primarily focusing on boreal ecosystems. Despite its extensive coverage for North America, its geographical specificity and lack of open access restrict its utility for European prehistoric contexts. Similarly, the QUB Dendrochronology Database, managed by Queen’s University Belfast, offers valuable datasets for Ireland and the UK but lacks significant representation of prehistoric material, limiting its application in broader archaeological investigations. The Building Archaeology Research Database (BARD) contains over 24,000 records, including dendrochronological data from more than 2,700 buildings. However, its focus on medieval and post-medieval timber-frame construction further narrows its utility for studies involving prehistoric wood samples.

The Digital Collaboratory for Cultural Dendrochronology (DCCD) presents itself as a potentially valuable international platform for dendrochronology, particularly through its integration with archaeological data services such as ARIADNE. However, it remains heavily biased toward datasets from the Netherlands, which account for more than two-thirds of its entries, while only 0.08% of its *Quercus* data pertain to Switzerland and just 2.5% represent prehistoric datasets. Notably, these estimates date back to 2021, and an updated assessment is currently unattainable following the platform’s migration to DataverseNL, which now charges an annual fee of nearly €6,000 for access. Furthermore, database activity has declined significantly, from 3,846 new project records between 2010 and 2014 to only 83 by the end of 2019. Although 519 additional records have been reported since 2021, it remains unclear whether this figure includes revisions to pre-existing entries, potentially inflating the count. The database’s narrow focus and restrictive access model significantly limit its broader utility for prehistoric research.

Finally, the Strategic Environmental Archaeology Database (SEAD), hosted by Umeå University, integrates multiple environmental proxies, including dendrochronological data. However, the dendrochronological component is largely confined to Swedish data, with limited relevance to prehistoric contexts. While SEAD aims for broader applications, its dendro component has limited utility for studies outside Sweden.

The utility of dendrochronological databases for prehistoric research varies widely. Global resources such as NOAA’s ITRDB and DCCD offer substantial datasets but face significant limitations in practical geographical and temporal scope. Similarly, platforms like DendroDB and BARD primarily cater to historical studies, leaving critical gaps in prehistoric coverage. Specialized resources like OLDLIST, Dendro4Art, and the Dendrochronological Picture Database pro-

vide valuable contributions but lack the direct relevance necessary for archaeological tree-ring analysis. Consequently, researchers focused on prehistoric dendrochronology must navigate a fragmented landscape of databases, each offering distinct strengths and limitations. Addressing these gaps remains crucial for advancing the field.

### *2.3.2. Typological dating*

Typological dates—i.e. relative, expertise- or seriation-based dating based on artefact characteristics—are ubiquitous in archaeological studies but rarely treated as a form of chronometric data in their own right. For example, the majority of the radiocarbon datasets we reviewed (Section 2.1) included some form of typological chronological information in the form of a ‘period’ or ‘culture’ column. This is also typically present in many other forms of systematic compilation work in archaeology, for example site gazetteers. Aggregated typological information from such sources are often used in aoristic analysis and related methods (Crema, 2024; Mischka, 2004). But this presentation of typological dating lacks metadata on how the determination was made and how exactly it is to be understood. Like any archaeological date, a typological date is derived from a physical sample – the object or set of object from which a chronological estimate was derived. Typological dates on one class of object may well clash with other classes of object, or for that matter with scientific dates – does one trust the date on pottery, the date on architecture, or the radiocarbon date? Without additional metadata on e.g. who made the typological determination or what the radiocarbon date was obtained on, such inconsistencies are difficult to resolve. Similarly the absolute date range corresponding to a typological determination (e.g. “Late Neolithic”) can be interpreted in multiple ways depending on the region and intentions of the expert making the determination. PeriodO (Rabinowitz et al., 2016) is a linked open data infrastructure that includes a shared vocabulary of typological periods and corresponding calendar age estimates, and an important step towards addressing the latter problem. However, it remains to be systematically linked to compilations of typological dates (though there are some efforts in this direction e.g. Hannah et al., 2022).

---

What is missing to date is a general-purpose infrastructure for combining all of these types of chronometric information on a global scale. This is the gap that XRONOS aims to fill, starting with three methods: radiocarbon, dendrochronology, and typological dating. These were chosen because they are widely used and relatively advanced in terms of open data, but an important aim of the project is to develop a generalisable data model that can easily scale to any and all types of archaeological chronology (see Section 3.2).

## **3. Concept**

XRONOS inherits its basic structure from RADON (Hinz et al., 2012; Kneisel et al., 2014; Raetzl-Fabian, 1999; Rinne et al., 2024), with a database-backed

web application and a data model that separates radiocarbon dates, contextual information, and sites. Our overall aims in developing XRONOS is to bring this model, which RADON has operated on for more than twenty years, up to date, to generalise it to other types of chronometric information, and to transform it from an online database to a data infrastructure that supports the continuous ingestion, curation, and open dissemination of archaeological chronologies from diverse sources.

### *3.1. Design goals*

XRONOS is our answer to Kintigh’s call (Kintigh, 2006) for digital infrastructures that don’t just provide access to chronological data but enables researchers to “archive, access, integrate, and mine disparate data sets”. It complements several similar open data infrastructures within and outwith archaeology, such as the Global Biodiversity Information Facility (GBIF, Canhos et al., 2004), the Strategic Environmental Archaeology Database (SEAD, Buckland, 2014), IMPACT for mummified human remains (Nelson and Wade, 2015), Neotoma for palaeoecological data (Williams et al., 2018), IsoArch for stable isotope data (Plomp et al., 2022), Poseidon for archaeogenetic data (Schmid et al., 2024), the International Soil Radiocarbon Database (ISRaD, Lawrence et al., 2020), and the ‘Big Interdisciplinary Archaeological Database’ (BIAD), an ambitious new initiative to combine many of these individual domains, including chronology (Reiter et al., 2024). To improve upon existing global syntheses of radiocarbon dates (see Section 2.2), we aimed to develop a living infrastructure that both continually collected data from diverse sources and presented a seamless single database to the user.

Our principal goals for the software were therefore to:

1. Combine all available sources of radiocarbon and other chronometric data in single database
2. Develop robust tools for the continuous ingestion, collation and curation of this data
3. Disseminate the collated and curated data as linked open data within a FAIR framework

Meeting these goals required the development of a) a conceptual data model, including links to other open data resources, that is flexible enough for all forms of chronometric data; and b) a software implementation that supports the main functions of ingesting, curating, and disseminating this data. The individual components of this work are described in more depth below but, briefly, consist of a relational data model implemented in a PostgreSQL database; a Ruby application providing server-based tools for ingestion, curation and dissemination of data; and multiple graphical and programmatic interfaces to the resulting dataset.




Figure 4: Simplified entity relationship diagram showing the XRONOS data model

### 3.2. Data model

At the base of the XRONOS data model (Figure 4) are sets of spatiotemporal coordinates or, as we call them, *chrons*. In an archaeological context, we conceptualise a chron as an assertion linking human activity with a particular point in space and time. Our data model currently encompasses three types of chron: radiocarbon dates, typological dates (e.g. ‘Early Neolithic’) and dendrochronological dates. However we anticipate that the concept will accommodate other types of absolute and relative dating techniques, as the scope of the database expands.

Chrons are conceptually useful because they emphasise that different types of archaeological ‘dates’, drawn from different sources, have essentially the same information content: the location of an event in space and time. We thereby avoid privileging certain sources of chronological data over others (as might be the case if, for example, we treated ‘period’ as a fixed attribute of a site) and can accommodate contradictory (e.g. differences of opinion on typological classification). This is important given that XRONOS aspires to be an authoritative ‘backbone’ with a global scope, so we cannot realistically impose a single chronological scheme or resolve conflicting information provided by specialists. Chrons are useful practically because they expose a common interface for attributes that all types of chronological information share, such as a *terminus post quem* (TPQ), *terminus ante quem* (TAQ), and midpoint estimate. This allows applications to query chronological data from multiple sources, without necessarily having to be aware of the peculiarities of each type of dating.

In order to unify chronological information in the form of a chron, we need a common chronological ‘coordinate system’. The natural choice is a *calendar probability distribution*, which expresses the probability that an event occurred as a function of time on a calendric scale. Most archaeologists are familiar with working with this kind of representation in the form of calibrated radiocarbon dates, but it can be extended and generalised to essentially any kind of chronological information. For example, in aoristic analysis (Mischka, 2004), a periodic time estimate (e.g. the event occurred in the Neolithic) is conceptualised as a uniform probability distribution over the timespan between the known start and end dates of that period. A similar model is used in OxCal (Bronk Ramsey, 2009, a direct inspiration for our approach) to integrate prior chronological information from diverse sources. In practical terms, this model means that the canonical representation the time component of any chron in XRONOS, regardless of source, is a probability distribution over the set of calendar years (arbitrarily measured in years Before Present) in which it could have plausibly occurred. Further statistics, e.g. a midpoint estimate or TPQ/TAQ range, can be derived from this distribution using well-known methods. In this way, we can support many different types of date and much of the implementation of XRONOS can be agnostic to the source of chronological information.

Chrons are located in space through association to a *sample* – the physical object from which a chronological determination was made. The location of samples is represented with geographical coordinates and an associated coordinate reference system (CRS), though since in practice the precise location of single samples is rarely available, this property is usually inherited from the site. We also record relevant metadata on the nature of the sample. For radiocarbon dates, for example, we follow established conventions (Millard, 2014) in recording the type (e.g. charcoal, charred seed) and, where applicable, taxonomic designation (e.g. *Quercus*, *Triticum dicoccum*) of the organic material used for dating. For typological dates, an ideal scenario would be for the sample to represent the particular object from which an inference was made (e.g. ‘Natufian’ might be inferred from ‘lunate-type microlith’). In practice, the best we can glean from most published datasets is the type of material used (e.g. ‘pottery’, ‘lithics’). The same sample can be associated with multiple chronos, including different types of chron. This is useful, for example, for representing replicate radiocarbon dates on the same sample, or radiocarbon dates and dendrochronological made on the same section of wood for wiggle-matching.

Further contextual information is associated with *contexts* and *sites*. The site is the primary geographic container for chronological information. As already mentioned, we typically record the spatial location of chronos using this entity, though it is possible to modify this by providing specific coordinates at the sample level. Sites also have attributes describing their conventional name or names in different languages and are associated with a flexible ‘site type’ typology that combines information on their form and function.

A context represents the specific find-context of a sample, e.g. an architectural

feature, stratigraphic unit, or phase. Since the units and conventions for recording such information vary greatly between different regions and archaeological traditions—and XRONOS is designed with global data in mind—we leave the question of what a context precisely represents open, and only record an unstandardised, free text label for it. Crucially, however, contexts can have a self-referential association to other contexts belonging to the same site. This allows it to encode arbitrary relational structures between contexts, whether they be hierarchical (e.g. phases and sub-phases) or graphical (e.g. stratigraphic). In this way, it can serve as a foundation for chronological modelling.

The series of relations `(chron) > sample > context > site` links the chronological and contextual sides of the XRONOS data model. Each step is a many-to-one association, meaning for example that it is possible to attach multiple chrons to the same sample (e.g. replicated radiocarbon dates on the same material), multiple *types* of chrons to the same sample (e.g. radiocarbon dates on tree-rings for wiggle-matching). Since this kind of information is rarely systematically recorded in our source databases, there are currently few actual records that make use of this feature of the data model. However, we hope it will provide a foundation for more nuanced chronological modelling in the future.

Metadata is incorporated into XRONOS’ data model at the level of the individual records (e.g. all records store their data of creation and last modification) and through two additional types of record: bibliographic *references* and *versions*. Bibliographic references store information on the source of a record in BibTeX format and can be linked through many-to-many associations to sites or chrons. Versions are a special type of record that are associated with all other records (including bibliographic references) and store the previous versions of those records as a series of changesets. In this way all changes to data are recorded and can be reconstructed (or reversed) precisely. This ‘paper trail’ also stores contextual metadata, e.g. who made the change and why. It also means that records that are deleted can be reviewed or restored from their stored version history, which is never discarded. Together these two systems provide a transparent record where the data in XRONOS comes from and how it has been altered, which we view as essential in a scientific data infrastructure.

### 3.3. Linked data

The XRONOS data model presents several opportunities to link to other resources as linked open data (LOD). We use controlled vocabularies such as the GBIF Backbone Taxonomy (GBIF Secretariat, 2023, for taxonomic descriptions of samples) to both standardise fields and to link the two resources based on this shared concept. There are concordances between XRONOS entities and several other specialised data infrastructures in archaeology, such as PeriodO (Rabinowitz et al., 2016, mapping to typological chrons) and gazetters like Pleiades (Bagnall et al., 2006) or Vici (Voorburg, 2012, mapping to site names). Moving beyond archaeology, Wikidata (<https://wikidata.org>) already includes many of the concepts represented in the XRONOS model (e.g. an ‘archaeological site’,

<https://www.wikidata.org/wiki/Q839954>). Linking to Wikidata is especially useful because it disseminates—and thereby preserves—the data compiled in XRONOS beyond the specialist/academic community. It also allows us to enrich the database with contextual information that otherwise be beyond our scope and resources, for example embedding multilingual descriptions of sites from Wikipedia articles on site records where the site has been linked to a Wikidata item.

Conversely, we encourage others to use XRONOS as a linked open data resource by providing stable URLs and a machine-readable interface for every record. Such usages could look like, for example, using a XRONOS URL as a canonical representation of a single radiocarbon date (e.g. <https://xronos.ch/c14s/SMU-257> -> <https://xronos.ch/c14s/23410>). We also plan to implement additional view formats to facilitate this, such as IntChron-format JSON, to allow it to be indexed through the IntChron service, or RDF, which is widely used by many linked open data resources in the digital humanities.

## 4. Implementation

Following a short pilot project in 2019, the first phase of development of XRONOS was completed in 2021–2024. The web interface (<https://xronos.ch>) has been publicly accessible since July 2021. Though we envisage XRONOS as a continuously-developing open source project and ‘living database’, the following offers a snapshot of progress at the end of our first grant-funded implementation phase. We do not aim to be comprehensive, but rather to describe some key elements of XRONOS’ current implementation that illustrate how our concept has been realised in practice.

### 4.1. Software architecture

The XRONOS data model is implemented as a relational database using the free and open source database management system PostgreSQL. However, apart from backups and other routine maintenance procedures, all interaction with the database is via a web application, which thus forms the core of XRONOS’ architecture. The XRONOS web application is written in Ruby and uses CRUD (Create, Read, Update, Delete) and MVC (Model, View, Controller) patterns as implemented in the Ruby on Rails framework.

The XRONOS web application exposes two distinct user interfaces: a graphical user interface accessed through a web browser; and an application programming interface (API). Both interfaces follow a REST (Representational State Transfer) pattern (Verborgh et al., 2015), where each resource (e.g. a single radiocarbon date, a single bibliographic reference, or a single user) is statelessly mapped to a single address. Users can then interact with resources at these addresses using a predictable and uniform interface based on HTTP verbs. For example, the radiocarbon date RTD-8904 is represented by the address <https://xronos.ch/c14s/156205>. Users can view information on this resource by sending a



GET request to that address, regardless of which interface they are using, and authorised users can modify it using POST, PATCH, or DELETE requests. The bibliographic reference associated with this date (Richter et al., 2017) is similarly represented at the address <https://xronos.ch/references/17778>, and can be accessed at that address using the same interface as the radiocarbon date.

This basic REST pattern is augmented by seven ‘actions’ (following the standard pattern in Rails application) that express different ways of interacting with a resource: index, show, destroy, new, create, edit, and update. The ‘show’ action represents interaction with a single resource, as described above. The ‘index’ action, which lists resources of a given type (e.g. <https://xronos.ch/c14s> for radiocarbon dates), is worth special mention because it is through this that the filtering logic at the core of XRONOS’ two interfaces is implemented. By passing a query as HTTP GET parameters to the index action of a resource, the list returned to the user is modified to only include records that match that query. For example, [https://xronos.ch/sites?site%5Bcountry\\_code%5D=CH](https://xronos.ch/sites?site%5Bcountry_code%5D=CH) (the part of the URL after the ? character encodes the SQL WHERE clause `country = 'CH'` as a GET parameter) lists sites in Switzerland. More complex queries can be executed using nested parameters. For example, <https://xronos.ch/c14s?sample%5Bmaterial%5D%5Bname%5D=charcoal> (encoding that the `c14` table should be joined to the `material` table via `sample`, followed by the WHERE clause `material.name = 'charcoal'`) lists radiocarbon dates obtained from charcoal samples. Index actions can also respond with the result in a tabular data format (i.e. `.csv`).

#### 4.2. Data ingestion and curation

The chronological data in XRONOS comes from a variety of sources, including published structured datasets in repositories and journal supplements, other on-line databases, literature review, and direct input from collaborators. Our aim is not just to ‘mirror’ these sources as they are, but integrate them into a single curated and continuously updated database. For the purposes of ingestion, we classify data resources into three categories: static resources, such as supplementary data in published papers, which are imported once; versioned resources, updated on a periodic basis, which we import after each new version; and live resources, which are continuously updated and therefore continuously imported. Records of each type are imported into XRONOS in as close to their original state as possible, i.e. without any corrections or standardisation applied. This ensures that any subsequent changes (even immediate, automatic ones) are entered into the record’s version history, so that the source of any deviations or potential errors can always be reconstructed. The version history also records the direct source of the data for attribution purposes. In addition, a bibliographic reference to the original resource is attached to ensure that the source is clearly attributed even if the record is merged with another one.

Once ingested, we apply a number of automated and semi-automated quality control processes to integrate new data into the existing database. Controlled

Table 2: Automatically-recognised data quality issues currently implemented in XRONOS

Issue	N	Description
Sites		
MISSING_COORDINATES	4452	Missing geographic coordinates
INVALID_COORDINATES	0	Geographic coordinates fall outside the earth’s ellipsoid
MISSING_COUNTRY_CODE	1221	Missing data on what country the site is in
Samples		
MISSING_MATERIAL	138177	Missing data on the sample material
MISSING_TAXON	138182	Missing data on the sample taxon
MISSING_CRS	0	Sample has coordinates but the coordinate reference system is not given
Taxons		
UNKNOWN_TAXON	9260	Sample taxon has not been matched to the GBIF Backbone Taxonomy
LONG_TAXON	509	Description of the sample taxon is implausibly long
Radiocarbon dates		
MISSING_C14_AGE	447	Missing radiocarbon age
VERY_OLD_C14	76	Radiocarbon age older than the effective range of the method (50 ka)
MISSING_C14_ERROR	585	Missing measurement error
MISSING_D14C	238875	Missing <sup>13</sup> C measurement
MISSING_D14C_ERROR	238875	Missing <sup>13</sup> C measurement error
MISSING_C14_METHOD	242233	Missing data on radiocarbon dating method (conventional, AMS, etc.)
MISSING_C14_LAB_ID	1233	Missing laboratory identifier
INVALID_LAB_ID	16053	Laboratory identifier does not match the standard format (e.g. ‘Abc-1234’)
MISSING_C14_LAB	350190	Missing data on the radiocarbon laboratory
Bibliographic references		
MIXED_REFERENCE	21933	Bibliographic reference appears to combine multiple publications
MISSING_BIBTEX	42109	Bibliographic reference without structured data in BibTeX format

vocabularies are used in a number of places in the data model (Figure 4), and we use thesauri to automatically standardise these fields as much as possible. For example, as mentioned above, the taxonomic description of samples is controlled using GBIF’s backbone taxonomy (GBIF Secretariat, 2023), and we also use a thesaurus service provided by GBIF to automatically change variant or obsolete taxonomic names to the canonical version. If the system is not able to standardise a field using the available thesaurus, it is flagged for manual correction. A wide variety of other potential data quality issues (e.g. missing data on what country a site is in) are also flagged for human review by this system (Table 2), which can often be semi-automated (e.g. suggesting close matches in the thesaurus or the country indicated by the record’s coordinates).

A final critical component of XRONOS’ data curation system is duplicate handling. We import data from many overlapping resources (many of which incorporate each other either in whole or in part), so duplicate records are common (as recently discussed by Reiter et al., 2024). The end result of standardising and correcting a record is also often to reveal a duplicate: e.g. the same sample imported from one source as ‘oak’ but another as ‘*Quercus* sp.’ will become a recognisable duplicate record as ‘*Quercus*’, and thus can be automatically merged to a single sample. Such exact duplicates can be merged automatically, with the oldest record becoming the authoritative version, but detecting fuzzier duplicated information (e.g. differences in the spelling of site names) has proved a more difficult problem. As of writing there are therefore still many duplicate records in XRONOS that need to be manually resolved, but we hope to automate much more of this work in the future.

### 4.3. User interfaces

The graphical user interface (GUI) to XRONOS, accessed through a web browser (e.g. at <https://xronos.ch>), uses REST resources and actions as the building blocks for various interfaces through which users can browse, search, retrieve, and analyse chronometric data (Figure 5). Each action on each resource is represented by a page, though not all of these are publicly accessible. The most important of these are the ‘index’ views, which list and summaries all instances of a resource and support filtering and sorting; and ‘show’ views (e.g. Figure 5b for a site), which give a more comprehensive information on an individual record along with visualisations, links to related records, external linked open data resources, and a log of changes made to the data since it was imported into XRONOS (Figure 5c). Pages representing REST resources directly are supplemented by a number of synthetic interfaces, for example the ‘data browser’ (<https://xronos.ch/data>, Figure 5a), which facilitates more complex filtering, or the search interface (<https://xronos.ch/search>). The GUI also includes several resources which are not part of XRONOS’ scientific data model, for example documentation pages, user profiles and news articles; these are not exposed in the API.

Access to various ‘backstage’ interfaces for creating, editing, and deleting data, and monitoring data quality is managed using a user permissions system. Cur-




Figure 5: Views in the XRONOS web graphical user interface

rently only authorised users affiliated with the XRONOS project can access these, but in the future we intend to support open registration and expose editing interfaces to all authenticated users. For this reason, there is no sharp division between a ‘public’ and ‘private’ areas – viewing/querying data and editing/curating data share the same architecture and interface patterns.

The XRONOS API uses the same addresses as the web-based GUI (with the exception of some of the synthetic interfaces mentioned above) but responds with machine-readable data in JSON format, rather than a HTML page. This response can be triggered by appending `.json` to the address or by including a HTTP `content-format` header in the request. Though users can make such requests manually and parse the data with one of several off-the-shelf tools, the primary intended uses of this interface is to provide access for 1) programmatic clients to XRONOS and 2) other web services. The XRONOS R package (Hinz and Roe, 2024) is an example of a programmatic client; it uses the API to facilitate direct querying and retrieval of data from XRONOS in the R statistical programming language (R Core Team, 2024). Similar libraries could be developed in other programming environments used for scientific computing, such as Python or Haskell. The API also provides the foundation for other web services to access XRONOS directly, to embed chronological information in other contexts or otherwise make use of its data resources.

An overarching principle of this software architecture is that all interaction

with XRONOS’ data store, and as much of the data processing and ‘business logic’ of responding to REST requests as possible, is directed through the same server-side routines. First and foremost, this allows us to provide multiple interfaces (i.e. the GUI and API, perhaps more in the future) without duplicating these elements of our codebase. It also improves accessibility for users accessing XRONOS through devices with limited processing capability or through text-only browsers. More broadly, avoiding reliance on client-side processing, e.g. with Javascript or Web Assembly (WASM)—which would be the other option—allows us keep our client interfaces simple (in most cases plain, semantic HTML pages and self-contained stylesheets) and therefore, we hope, sustainable in the face of constantly-evolving client-side technologies and standards. It does have the weakness that, in practical terms, XRONOS is difficult to run and relies on the continued existence of a maintained external server. We have however tried to mitigate this by providing clearly-documented source code and regular data dumps so that, if our instance of XRONOS disappears, or if one simply does not want to use it, it is possible for others to host a XRONOS server of their own. As the sustainability of scientific software and data infrastructures is a pressing problem, in the future it may be desirable to support further decentralisation through, for example, a federated server-server model.

## 5. Evaluation

In this paper we have outlined the conceptual and technical infrastructure developed during the initial, SNF-funded phases of development on XRONOS in 2019 and 2021–2024. These include a generalised data model for radiocarbon and typological dates, extendible to other chronometric information, and associated contextual information; an R- and Ruby-based pipeline for continuous ingestion of data from a variety of sources; continuous, semi-automated data cleaning protocols; a Ruby-on-Rails application providing a web-based frontend to the data and a REST API for programmatic access; and an R package for interfacing with the API. These systems are deployed and publicly accessible at <https://xronos.ch>.

At the current stage of implementation, we argue XRONOS provides a framework for access to chronometric data that is more open, more reliable, and more comprehensive than the previously available global radiocarbon compilations. XRONOS blends aspects of the three existing approaches (c14bazAAR, IntChron, and p3k14c; see Section 2.2) to achieve the same aim of providing access to the global radiocarbon data through a common interface. Like c14bazAAR and IntChron, it is a ‘metadatabase’ that draws from existing data resources and maintains an explicit link to them. But like p3k14c, it integrates these into a single database and applies data curation processes to harmonise them and improve the quality of the information. It has a wider scope than c14bazAAR or IntChron, as it mirrors rather than directly retrieves the source data (allowing us to use resources that aren’t openly published), and does not rely on the authors of these sources to implement a common specification. It

also goes beyond the functionality of p3k14c by providing systems for the continuous ingestion and curation of new data as it is published. With that said, we see the approaches followed by as complementary rather than competing. A parser for XRONOS has been contributed back to c14bazAAR and we also aim to provide an IntChron interface to XRONOS’ data in the near future. New data and corrections from p3k14c are incorporated into XRONOS as they are released.

From 2025, development of XRONOS will continue within the framework of ‘ESTER’, an ERC-funded research project on estimation of prehistoric population development from large, multiproxy datasets. Our immediate development goals include the incorporation of dendrochronology into the database, further refinement of our data curation pipelines, and the public release of the editing interfaces.

Looking beyond the near term, we have endeavoured to create a sustainable infrastructure that can be maintained by a wider scholarly commons – though we must acknowledge that this is a difficult problem, and one that is as much organisational than technical. The source code for all the software components of the system are available online (at <https://github.com/xronos-ch>) under open licenses. The databases of the instance at <https://xronos.ch> are also archived with Zenodo at <https://doi.org/10.5281/zenodo.14282598>. With these two resources we reduce XRONOS’ ‘bus factor’; if we are not able to continue operating XRONOS, somebody else can fully recreate it. Equally importantly, we enable the ‘right to fork’ should others wish to take the software and/or database in another direction. But aside from these extreme scenarios, the long-term sustainability of XRONOS is contingent on the existence of a community of scholars that use and contribute back to it.

## 6. Acknowledgements

The XRONOS project was carried under the direction of Albert Hafner. Individual contributors to the XRONOS database to date include Chiara Huwiler, Rivana Moser, Tomasz Chmielewski, and Stephanie Döppler. A complete and up-to-date list of acknowledgements for the project can be found at <https://xronos.ch/about/acknowledgements>.

We are also grateful to the two anonymous reviewers of our manuscript, whose comments have improved the finished paper.

## 7. Funding Statement

This work was funded by the Swiss National Science Foundation ([SNSF Project #198152](#); [SNSF Project #194326](#)) and the University of Bern (UniBE Initiator Grant 2019).

## 8. Data Accessibility Statement

The data and R code used to produce our analysis of the state of the art in radiocarbon compilation, including all the figures presented here, can be accessed via Zenodo at <https://doi.org/10.5281/zenodo.14282598>.

The database and software described in this paper is open source and can be accessed at <https://github.com/xronos-ch>.

## References

- Aitken, M.J., 1999. *Science-Based Dating in Archaeology*. Routledge, London. <https://doi.org/10.4324/9781315836645>
- Al-Abyadh-Balghelam-Dalma-Jebel, A., Dhanna-Marawah-Marawah, M.R., Yas, R.G.-R.-S.B., n.d. ADIAS Radiocarbon Archive.
- Alcántara, A., 2021. Entre tepalcates. Hacia una comprensión historiográfica de la datación arqueométrica en la arqueología mexicana (1950-2020) (PhD thesis). <https://doi.org/10.13140/RG.2.2.24447.74400/2>
- Alcántara, A., Pedroza, I., in press. Latin American Radiocarbon on the Web Databases and Datasets: A Mexican and Brazilian Perspectives. *Radiocarbon*.
- Alves, E.Q., Macario, K., Ascough, P., Bronk Ramsey, C., 2018. The World-wide Marine Radiocarbon Reservoir Effect: Definitions, Mechanisms, and Prospects. *Reviews of Geophysics* 56, 278–305. <https://doi.org/10.1002/2017RG000588>
- Anderson, D.G., Miller, D.S., Yerka, S.J., Gillam, J.C., Johanson, E.N., Anderson, D.T., Goodyear, A.C., Smallwood, A.M., 2010. *PIDBA (Paleoindian Database of the Americas) 2010: Current Status and Findings*. *Archaeology of Eastern North America* 38, 63–89.
- Andreae, M.O., Al-Amri, A., Andreae, C.M., Guagnin, M., Jochum, K.P., Stoll, B., Weis, U., 2020. Archaeometric studies on the petroglyphs and rock varnish at Kilwa and Sakaka, northern Saudi Arabia. *Arabian Archaeology and Epigraphy* 31, 219–244. <https://doi.org/10.1111/aae.12167>
- Arnold, J.R., Libby, W.F., 1951. Radiocarbon Dates. *Science* 113, 111–120. <https://doi.org/10.1126/science.113.2927.111>
- Bada, J.L., Helfman, P.M., 1975. Amino acid racemization dating of fossil bones. *World Archaeology* 7, 160–173. <https://doi.org/10.1080/00438243.1975.9979630>
- Bagnall, R., Talbert, R.J.A., Tom, E., et al., 2006. *Pleiades: A Community-Built Gazetteer and Graph of Ancient Places*.
- Baillie, M.G.L., 2014. *Tree-ring Dating and Archaeology*. Routledge, London. <https://doi.org/10.4324/9781315748689>
- Banque Nationale de Données Radiocarbones (BANADORA), n.d.
- Barceló Álvarez, J.A., Bogdanović, I., Capuzzo, G., 2013. New developments of technological and archaeological data bases: The example of radiocarbon dating, in: *Arqueología: Actas Del Primer Congreso Internacional de Buenas*

- Prácticas En Patrimonio Mundial, 2013, ISBN 978-84-941030-9-4, págs. 733-749. JAS Arqueología, pp. 733-749.
- Batist, Z., 2023. Archaeological data work as continuous and collaborative practice. <https://doi.org/10.5281/zenodo.8373390>
- Batist, Z., Roe, J., 2024. Open Archaeology, Open Source? Collaborative practices in an emerging community of archaeological software engineers. Internet Archaeol. <https://doi.org/10.11141/ia.67.13>
- Bayliss, A., 2015. Quality in Bayesian chronological models in archaeology. World Archaeol. 47, 677-700. <https://doi.org/10.1080/00438243.2015.1067640>
- Bednarik, R.G., 2020. The use of weathering indices in rock art science and archaeology. Rock Art Research: The Journal of the Australian Rock Art Research Association (AURA) 29, 59-84. <https://doi.org/10.3316/ielapa.275310392544915>
- Benedict, J.B., 2009. A Review of Lichenometric Dating and Its Applications to Archaeology. American Antiquity 74, 143-172. <https://doi.org/10.1017/S0002731600047545>
- Benz, M., 2010. PPND – The Platform for Neolithic Radiocarbon Dates.
- Bird, D., Bocinsky, K., Vander Linden, M., 2024. [p3k14c: P3k14C: A synthetic global database of archaeological radiocarbon dates.](https://doi.org/10.1017/S0002731600047545)
- Bird, D., Miranda, L., Vander Linden, M., Robinson, E., Bocinsky, R.K., Nicholson, C., Capriles, J.M., Finley, J.B., Gayo, E.M., Gil, A., d'Alpoim Guedes, J., Hoggarth, J.A., Kay, A., Loftus, E., Lombardo, U., Mackie, M., Palmisano, A., Solheim, S., Kelly, R.L., Freeman, J., 2022. P3k14c, a synthetic global database of archaeological radiocarbon dates. Scientific Data 9, 27. <https://doi.org/10.1038/s41597-022-01118-7>
- Böhner, U., Schyle, D., 2004. Radiocarbon CONTEXT database.
- Brøgger, W.C., 1905. Strandliniens beliggenhed under stenalderen i det sydøstlige Norge. I kommission hos H. Aschehoug & Company.
- Bronk Ramsey, C., 2009. Bayesian Analysis of Radiocarbon Dates. Radiocarbon 51, 337-360. <https://doi.org/10.1017/S0033822200033865>
- Bronk Ramsey, C., Blaauw, M., Kearney, R., Staff, R.A.S., 2019. The importance of open access to chronological information: The IntChron initiative. Radiocarbon 61, 1-11. <https://doi.org/10.1017/RDC.2019.21>
- Bronk Ramsey, C., Higham, T.F.G., Brock, F., Baker, D., Ditchfield, P., 2009. Radiocarbon Dates from the Oxford Ams System: Archaeometry DaTEList 33. Archaeometry 51, 323-349. <https://doi.org/10.1111/j.1475-4754.2008.00457.x>
- Buck, C.E., Kenworthy, J.B., Litton, C.D., Smith, A.F.M., 1991. Combining archaeological and radiocarbon information: A Bayesian approach to calibration. Antiquity 65, 808-821. <https://doi.org/10.1017/S0003598X00080534>
- Buckland, P.I., 2014. SEAD - The Strategic Environmental Archaeology Database Inter-linking Multiproxy Environmental Data with Archaeological Investigations and Ecology, in: Verhagen, P., Earl, G. (Eds.), Archaeology in the Digital Era: Papers from the 40th Annual Conference of Computer Applications and Quantitative Methods in Archaeology (CAA), Southamp-



- ton, 26-29 March 2012. Amsterdam University Press, pp. 320–331. <https://doi.org/10.1515/9789048519590-035>
- Canhos, V.P., Souza, S. de, Giovanni, R.D., Canhos, D.A.L., 2004. Global Biodiversity Informatics: Setting the scene for a "new world" of ecological forecasting. *Biodiversity Informatics* 1. <https://doi.org/10.17161/bi.v1i0.3>
- Capriles, J.M., 2023. The Bolivian Radiocarbon Database: A Countrywide Compilation of Radiocarbon Dates. *Journal of Open Archaeology Data* 11. <https://doi.org/10.5334/joad.104>
- Capuzzo, G., Boaretto, E., Barceló, J.A., 2014. EUBAR: A Database of  $^{14}\text{C}$  Measurements for the European Bronze Age. A Bayesian Analysis of  $^{14}\text{C}$ -Dated Archaeological Contexts from Northern Italy and Southern France. *Radiocarbon* 56, 851–869. <https://doi.org/10.2458/56.17453>
- Chapple, R.M., 2019. Catalogue of Radiocarbon Determinations & Dendrochronology Dates (August 2019 release).
- Chaput, M.A., Gajewski, K., 2016. Radiocarbon dates as estimates of ancient human population size. *Anthropocene* 15, 3–12. <https://doi.org/10.1016/j.ancene.2015.10.002>
- Clist, B., Denbow, J., Lanfranchi, R., 2023. Using the radiocarbon dates of Central Africa for studying long-term demographic trends of the last 50,000 years: Potential and pitfalls. *Azania: Archaeological Research in Africa* 58, 235–293. <https://doi.org/10.1080/0067270X.2023.2215649>
- Cochrane, E.E., Rieth, T.M., Filimoehala, D., 2021. The first quantitative assessment of radiocarbon chronologies for initial pottery in Island Southeast Asia supports multi-directional Neolithic dispersal. *PLOS ONE* 16, e0251407. <https://doi.org/10.1371/journal.pone.0251407>
- Courtney Mustaphi, C., 2016. Radiocarbon dates from eastern Africa in the CARD2.0 format. <https://doi.org/10.7910/DVN/NJLNRJ>
- Crema, E.R., 2024. A Bayesian alternative for Aoristic analyses in archaeology. <https://doi.org/10.31219/osf.io/98qkx>
- Crema, E.R., 2022. Statistical Inference of Prehistoric Demography from Frequency Distributions of Radiocarbon Dates: A Review and a Guide for the Perplexed. *J Archaeol Method Theory* 29, 1387–1418. <https://doi.org/10.1007/s10816-022-09559-5>
- Crema, E.R., Bloxam, A., Stevens, C.J., Vander Linden, M., 2024. Modelling diffusion of innovation curves using radiocarbon data. *Journal of Archaeological Science* 165, 105962. <https://doi.org/10.1016/j.jas.2024.105962>
- Crema, E.R., Habu, J., Kobayashi, K., Madella, M., 2016. Summed Probability Distribution of  $^{14}\text{C}$  Dates Suggests Regional Divergences in the Population Dynamics of the Jomon Period in Eastern Japan. *PLoS One* 11, e0154809. <https://doi.org/10.1371/journal.pone.0154809>
- d’Errico, F., Banks, W.E., Vanhaeren, M., Laroulandie, V., Langlais, M., 2011. PACEA Geo-Referenced Radiocarbon Database. *PaleoAnthropology* 2011, 1–12.
- Daniels, F., Boyd, C.A., Saunders, D.F., 1953. Thermoluminescence as a Research Tool. *Science* 117, 343–349. <https://doi.org/10.1126/science.117.3040.343>

- de Saulieu, G., Oslisly, R., Nlend, P., Garcin, Y., Chevillotte, H., Kuete, N., 2017. Plateforme des datations archéologiques d’Afrique Centrale. Version 2, 03.
- Dee, M., Wengrow, D., Shortland, A., Stevenson, A., Brock, F., Girdland Flink, L., Bronk Ramsey, C., 2013. An absolute chronology for early Egypt using radiocarbon dating and Bayesian statistical modelling. *Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences* 469, 20130395. <https://doi.org/10.1098/rspa.2013.0395>
- Dee, M.W., Rowland, J.M., Higham, T.F.G., Shortland, A.J., Brock, F., Harris, S.A., Ramsey, C.B., 2012. Synchronising radiocarbon dating and the Egyptian historical chronology by improved sample selection. *Antiquity* 86, 868–883. <https://doi.org/10.1017/S0003598X00047979>
- Díaz-Rodríguez, M., Nielsen, T.K., Maier, A., Riede, F., 2023. An Annotated Compilation of Chronometric Dates for the Middle-Upper Palaeolithic Transition (45–30 ka BP) in Northern Iberia (Spain). *Journal of Open Archaeology Data* 11. <https://doi.org/10.5334/joad.113>
- Douglass, A.E., 1929. The secret of the Southwest solved by talkative tree rings. *National Geographic* 56, 737–770.
- Douglass, K., Hixon, S., Wright, H.T., Godfrey, L.R., Crowley, B.E., Manjakahery, B., Rasolondrainy, T., Crossland, Z., Radimilahy, C., 2019. A critical review of radiocarbon dates clarifies the human settlement of Madagascar. *Quat. Sci. Rev.* 221, 105878. <https://doi.org/10.1016/j.quascirev.2019.105878>
- Evernden, J.F., Curtis, G.H., Bishop, W., Brace, C.L., Clark, J.D., Damon, P.E., Hay, R.L., Hopkins, D.M., Howell, F.C., Knopf, A., Kretzoi, M., Leakey, L.S.B., Maude, H.E., Richards, J.R., Savage, D.E., Wright, H.E., 1965. The Potassium-Argon Dating of Late Cenozoic Rocks in East Africa and Italy [and Comments and Reply]. *Current Anthropology* 6, 342–385. <https://doi.org/10.1086/200619>
- Flohr, P., Fleitmann, D., Matthews, R., Matthews, W., Black, S., 2016. Evidence of resilience to past climate change in Southwest Asia: Early farming communities and the 9.2 and 8.2 ka events. *Quat. Sci. Rev.* 136, 23–39. <https://doi.org/10.1016/j.quascirev.2015.06.022>
- Ford, J.A., 1962. A quantitative method for deriving cultural chronology, *Manuales técnicos*. Pan American Union, Washington.
- Gajewski, K., Munoz, S., Peros, M., Viau, A., Morlan, R., Betts, M., 2011. The Canadian Archaeological Radiocarbon Database (Card): Archaeological 14C Dates in North America and Their Paleoenvironmental Context. *Radiocarbon* 53, 371–394. <https://doi.org/10.1017/S0033822200056630>
- Garnett, M.H., Bryant, C.L., Freeman, S.P.H.T., Gulliver, P., Ascough, P.L., 2023. A DATABASE OF NERC RADIOCARBON MEASUREMENTS DETERMINED BY ACCELERATOR MASS SPECTROMETRY. *Radiocarbon* 1–4. <https://doi.org/10.1017/RDC.2023.18>
- Gayo, E.M., Latorre, C., Santoro, C.M., 2015. Timing of occupation and regional settlement patterns revealed by time-series analyses of an archaeological radiocarbon database for the South-Central Andes (16°–25°S). *Qua-*

- ternary International, Palaeodemography in Southern South America 356, 4–14. <https://doi.org/10.1016/j.quaint.2014.09.076>
- GBIF Secretariat, 2023. GBIF Backbone Taxonomy. <https://doi.org/10.15468/39OMEI>
- Gillespie, R., Gowlett, J.a.J., Hall, E.T., Hedges, R.E.M., 1984. Radiocarbon Measurement by Accelerator Mass Spectrometry: An Early Selection of Dates. *Archaeometry* 26, 15–20. <https://doi.org/10.1111/j.1475-4754.1984.tb00313.x>
- Gregorio de Souza, J., 2020. Rexpand: Radiocarbon dates for the spread of farming and ceramics in tropical South America.
- Großmann, R., Weinelt, M., Müller, J., 2023. Demographic dynamics between 5500 and 3500 calBP (3550–1550 BCE) in selected study regions of Central Europe and the role of regional climate influences. *PLOS ONE* 18, e0291956. <https://doi.org/10.1371/journal.pone.0291956>
- Grove, M., 2011. A Spatio-Temporal Kernel Method for Mapping Changes in Prehistoric Land-Use Patterns. *Archaeometry* 53, 1012–1030. <https://doi.org/10.1111/j.1475-4754.2010.00578.x>
- Guilderson, T.P., Reimer, P.J., Brown, T.A., 2005. The Boon and Bane of Radiocarbon Dating. *Science* 307, 362–364. <https://doi.org/10.1126/science.1104164>
- Hannah, E., Carlin, N., Griffiths, S., Gaydarska, B., Swan, R., Corns, A., 2022. *Project Radiocarbon*. *Archaeology Ireland* 36, 14–17.
- Harding, A., 1999. Establishing Archaeological Chronologies, in: *Companion Encyclopedia of Archaeology*. Routledge.
- Harris, E.C., 1979. The laws of archaeological stratigraphy. *World Archaeology* 11, 111–117. <https://doi.org/10.1080/00438243.1979.9979753>
- Heřmáňková, P., Kaše, V., Sobotková, A., 2021. Inscriptions as data: Digital epigraphy in macro-historical perspective. *Journal of Digital History* 1, 99–141. <https://doi.org/10.1515/jdh-2021-1004>
- Higham, T., Douka, K., Wood, R., Ramsey, C.B., Brock, F., Basell, L., Camps, M., Arrizabalaga, A., Baena, J., Barroso-Ruiz, C., Bergman, C., Boitard, C., Boscatto, P., Caparrós, M., Conard, N.J., Draily, C., Froment, A., Galván, B., Gambassini, P., Garcia-Moreno, A., Grimaldi, S., Haesaerts, P., Holt, B., Iriarte-Chiapusso, M.-J., Jelinek, A., Jordá Pardo, J.F., Maíllo-Fernández, J.-M., Marom, A., Maroto, J., Menéndez, M., Metz, L., Morin, E., Moroni, A., Negrino, F., Panagopoulou, E., Peresani, M., Pirson, S., de la Rasilla, M., Riel-Salvatore, J., Ronchitelli, A., Santamaria, D., Semal, P., Slimak, L., Soler, J., Soler, N., Villaluenga, A., Pinhasi, R., Jacobi, R., 2014. The timing and spatiotemporal patterning of Neanderthal disappearance. *Nature* 512, 306–309. <https://doi.org/10.1038/nature13621>
- Hinz, M., Furrholt, M., Müller, J., Raetzl-Fabian, D., Rinne, C., Sjögren, K.-G., Wotzka, H.-P., 2012. RADON - Radiocarbon dates online 2012. Central European database of 14C dates for the Neolithic and Early Bronze Age. *Jungsteinsite* 14, 1–4.
- Hinz, M., Roe, J., 2024. *Xronos: Client for the 'XRONOS' chronological database*.

- Hoebe, P.W., Peeters, J.H.M., Arnoldussen, S., 2023. Parsing prehistoric patterns: Prospects and limitations of a big radiocarbon dataset for understanding the impact of climate on Late Palaeolithic and Mesolithic populations in northwest Europe (16–7.5 ka calBP). *Journal of Archaeological Science: Reports* 49, 103944. <https://doi.org/10.1016/j.jasrep.2023.103944>
- Hoggarth, J.A., Ebert, C.E., Castelazo-Calva, V.E., 2021. MesoRAD: A New Radiocarbon Data Set for Archaeological Research in Mesoamerica. *Journal of Open Archaeology Data* 9, 10. <https://doi.org/10.5334/joad.83>
- Huet, T., Basílio, A.C., Carvalho, A.F., Cubas, M., Gibaja, J.F., López-Romero, E., Oms, F.X., Mazzucco, N., 2024. NeoNet Atlantic. Radiocarbon Dates for the Late Mesolithic/Early Neolithic Transition in the Southern European Atlantic Coast 12, 2. <https://doi.org/10.5334/joad.120>
- Huet, T., Cubas, M., Gibaja, J.F., Oms, F.X., Mazzucco, N., 2022. NeoNet Dataset. Radiocarbon Dates for the Late Mesolithic/Early Neolithic Transition in the North Central-Western Mediterranean Basin. *Journal of Open Archaeology Data* 10, 3. <https://doi.org/10/gpk67w>
- Katsianis, M., Bevan, A., Styliaras, G., Maniatis, Y., 2020. An Aegean History and Archaeology Written through Radiocarbon Dates. *Journal of Open Archaeology Data* 8. <https://doi.org/10.5334/joad.65>
- Kelly, R.L., Mackie, M.E., Robinson, E., Meyer, J., Berry, M., Boulanger, M., Coddling, B.F., Freeman, J., Garland, C.J., Gingerich, J., Hard, R., Haug, J., Martindale, A., Meeks, S., Miller, M., Miller, S., Perttula, T., Railey, J.A., Reid, K., Scharlotta, I., Spangler, J., Thomas, D.H., Thompson, V., White, A., 2022. A New Radiocarbon Database for the Lower 48 States. *American Antiquity* 87, 581–590. <https://doi.org/10.1017/aaq.2021.157>
- Kemmers, F., Myrberg, N., 2011. Rethinking numismatics. The archaeology of coins. *Archaeological Dialogues* 18, 87–108. <https://doi.org/10.1017/S1380203811000146>
- Kim, H., Lee, G.-A., Crema, E.R., 2021. Bayesian analyses question the role of climate in Chulmun demography. *Scientific Reports* 11, 23797. <https://doi.org/10.1038/s41598-021-03180-4>
- Kintigh, K., 2006. The promise and challenge of archaeological data integration. *Am. Antiq.* 71, 567–578. <https://doi.org/10.1017/S0002731600039810>
- Kneisel, J., Hinz, M., Rinne, C., 2014. RADON-B – Radiocarbon dates online (version 2014). Database for European 14C dates for the Bronze and Early Iron Age.
- Kra, R., 1989. The International Radiocarbon Data Base: A Progress Report. *Radiocarbon* 31, 1067–1075. <https://doi.org/10.1017/S003382220001273X>
- Kra, R., 1988. Updating the past: The Establishment of the International Radiocarbon Data Base. *American Antiquity* 53, 118–125. <https://doi.org/10.2307/281158>
- Kudo, Y., 2018. Approach for Creating Database of the Radiocarbon Dates Published on the Archaeological Research Reports in Japan. *Bulletin of the National Museum of Japanese History* 212, 251.
- Kudo, Y., Sakamoto, M., Hakozaiki, M., Stevens, C.J., Crema, E.R., 2023. An Archaeological Radiocarbon Database of Japan. *Journal of Open Archaeol-*

- ogy Data 11. <https://doi.org/10.5334/joad.115>
- Kusumgar, S., Lal, D., Sarna, R.P., 1963. Tata Institute Radiocarbon Date List I. Radiocarbon 5, 273–282. <https://doi.org/10.1017/S0033822200036894>
- Lake, M., 2012. Open archaeology. World Archaeology 44, 471–478. <https://doi.org/10.1080/00438243.2012.748521>
- Lawrence, C.R., Beem-Miller, J., Hoyt, A.M., Monroe, G., Sierra, C.A., Stoner, S., Heckman, K., Blankinship, J.C., Crow, S.E., McNicol, G., Trumbore, S., Levine, P.A., Vindušková, O., Todd-Brown, K., Rasmussen, C., Hicks Pries, C.E., Schädel, C., McFarlane, K., Doetterl, S., Hatté, C., He, Y., Treat, C., Harden, J.W., Torn, M.S., Estop-Aragonés, C., Asefaw Berhe, A., Keiluweit, M., Della Rosa Kuhnen, Á., Marin-Spiotta, E., Plante, A.F., Thompson, A., Shi, Z., Schimel, J.P., Vaughn, L.J.S., von Fromm, S.F., Wagai, R., 2020. An open-source database for the synthesis of soil radiocarbon data: International Soil Radiocarbon Database (ISRaD) version 1.0. Earth System Science Data 12, 61–76. <https://doi.org/10.5194/essd-12-61-2020>
- Levy, E., Geeraerts, G., Pluquet, F., Piasetzky, E., Fantalkin, A., 2021. Chronological networks in archaeology: A formalised scheme. Journal of Archaeological Science 127, 105225. <https://doi.org/10.1016/j.jas.2020.105225>
- Libby, W.F., 1955. Radiocarbon Dating, Second. ed. University of Chicago Press.
- Linnenlucke, L., Bird, M.I., Petchey, F., Alliston, G., McNiven, I.J., David, B., Ulm, S., 2023. Assessing the Reliability of a Small-Scale Legacy Radiocarbon Dataset Using Chronometric Transparency Approaches: Torres Strait Radiocarbon Database. Journal of Open Archaeology Data 11. <https://doi.org/10.5334/joad.95>
- Lipo, C., 2020. A repository of radiocarbon data for Rapa Nui.
- Loftus, E., Mitchell, P.J., Ramsey, C.B., 2019. An archaeological radiocarbon database for southern Africa. Antiquity 93, 870–885. <https://doi.org/10.15184/aqy.2019.75>
- Lucarini, G., Wilkinson, T., Crema, E.R., Palombini, A., Bevan, A., Broodbank, C., 2020. The MedAfriCarbon Radiocarbon Database and Web Application. Archaeological Dynamics in Mediterranean Africa, ca. 9600–700 BC. Journal of Open Archaeology Data 8. <https://doi.org/10.5334/joad.60>
- Lucas, G., 2004. Beyond chronology, in: The Archaeology of Time. Routledge.
- Manning, K., Colledge, S., Crema, E., Shennan, S., Timpson, A., 2016. The Cultural Evolution of Neolithic Europe. EUROEVOL Dataset 1: Sites, Phases and Radiocarbon Data. Journal of Open Archaeology Data 5, 275. <https://doi.org/10.5334/joad.40>
- Marom, N., Wolkowski, U., 2024. A note on predator-prey dynamics in radiocarbon datasets. Peer Community Journal 4. <https://doi.org/10.24072/pcjournal.395>
- Martínez-Grau, H., Morell-Rovira, B., Antolín, F., 2021. Radiocarbon Dates Associated to Neolithic Contexts (Ca. 5900 – 2000 Cal BC) from the Northwestern Mediterranean Arch to the High Rhine Area. Journal of Open Archaeology Data 9. <https://doi.org/10.5334/joad.72>
- McFadgen, B., Higham, T., Sparks, R.J., 2000. New Zealand radiocarbon

- database for archaeology. *Archaeology in New Zealand* 43, 297–300.
- Michczyński, A., Krzanowski, A., Pazdur, M.F., Ziolkowski, M.S., 1995. A Computer-Based Database for Radiocarbon Dates of Central Andean Archaeology. *Radiocarbon* 37, 337–343. <https://doi.org/10.1017/S0033822200030812>
- Millard, A.R., 2014. Conventions for reporting radiocarbon determinations. *Radiocarbon* 56, 555–559. <https://doi.org/10.2458/56.17455>
- Mischka, D., 2004. Aoristische Analyse in der Archäologie. *Archäologische Informationen* 27, 233–243. <https://doi.org/10.11588/ai.2004.2.12685>
- Moody, B., Dye, T., May, K., Wright, H., Buck, C., 2021. Digital chronological data reuse in archaeology: Three case studies with varying purposes and perspectives. *Journal of Archaeological Science: Reports* 40, 103188. <https://doi.org/10.1016/j.jasrep.2021.103188>
- Ndeye, M., Athie, A.H., Kébé, D., Sène, M., 2022. IFAN CH.DIOP RADIO-CARBON LABORATORY MEASUREMENTS II. *Radiocarbon* 64, 397–414. <https://doi.org/10.1017/RDC.2022.8>
- Nelson, A.J., Wade, A.D., 2015. Impact: Development of a Radiological Mummy Database. *The Anatomical Record* 298, 941–948. <https://doi.org/10.1002/ar.23130>
- Open Knowledge Foundation, n.d. The Open Definition: Defining Open in Open Data, Open Content and Open Knowledge [WWW Document]. URL <https://opendefinition.org/> (accessed 11.25.2024).
- Palmisano, A., Bevan, A., Kabelindde, A., Roberts, N., Shennan, S., 2022a. AIDA: Archive of Italian radiocarbon DAtes.
- Palmisano, A., Bevan, A., Lawrence, D., Shennan, S., 2022b. The NERD Dataset: Near East Radiocarbon Dates between 15,000 and 1,500 cal. Yr. BP 10, 2. <https://doi.org/10.5334/joad.90>
- Palmisano, A., Bevan, A., Shennan, S., 2017. Data and code for demographic trends in the paper "Human responses and non-responses to climatic variations during the Last Glacial-Interglacial transition in the eastern Mediterranean". <https://doi.org/10.14324/000.ds.1570274>
- Pardo-Gordó, S., Vidal-Matutano, P., González-Marrero, M. del C., Chávez-Álvarez, M.E., 2023. The 14Canarias web application. An interactive radiocarbon database for the Canary Islands. *Journal of Open Archaeology Data* 11. <https://doi.org/10.5334/joad.105>
- Perrin, T., 2019. BDA: Une Base de Données Archéologique collaborative en ligne. *Bulletin de la Société préhistorique française* 116, 159–162.
- Petchey, F., Bickler, S., Hughes, L., Bunbury, M., 2022. The Aotearoa/New Zealand Radiocarbon Database Upgrade. *Archaeology in New Zealand* 65, 32–40.
- Petrie, W.M.F., 1899. Sequences in Prehistoric Remains. *The Journal of the Anthropological Institute of Great Britain and Ireland* 29, 295–301. <https://doi.org/10.2307/2843012>
- Pettitt, P.B., Davies, W., Gamble, C.S., Richards, M.B., 2003. Palaeolithic radiocarbon chronology: Quantifying our confidence beyond two half-lives. *J. Archaeol. Sci.* 30, 1685–1693. [https://doi.org/10.1016/S0305-4403\(03](https://doi.org/10.1016/S0305-4403(03)



00070-0

- Plomp, E., Stantis, C., James, H.F., Cheung, C., Snoeck, C., Kootker, L., Kharobi, A., Borges, C., Moreiras Reynaga, D.K., Pospieszny, Ł., Fulminante, F., Stevens, R., Alaica, A.K., Becker, A., de Rochefort, X., Salesse, K., 2022. The IsoArcH initiative: Working towards an open and collaborative isotope data culture in bioarchaeology. *Data in Brief* 45, 108595. <https://doi.org/10.1016/j.dib.2022.108595>
- Prates, L., Politis, G.G., Perez, S.I., 2020. Rapid radiation of humans in South America after the last glacial maximum: A radiocarbon-based study. *PLOS ONE* 15, e0236023. <https://doi.org/10.1371/journal.pone.0236023>
- R Core Team, 2024. *R: A language and environment for statistical computing*. R Foundation for Statistical Computing, Vienna, Austria.
- Rabinowitz, A., Shaw, R., Buchanan, S., Golden, P., Kansa, E., 2016. Making Sense of the Ways We Make Sense of the Past: The Periodo Project. *Bulletin of the Institute of Classical Studies* 59, 42–55. <https://doi.org/10.1111/j.2041-5370.2016.12037.x>
- Rademaker, K., 2024. Updated Peru archaeological radiocarbon database, 20,000–7000 14C BP. *Quaternary International*. <https://doi.org/10.1016/j.quaint.2024.01.012>
- Rademaker, K., Bromley, G.R.M., Sandweiss, D.H., 2013. Peru archaeological radiocarbon database, 13,000–7000 14C B.P. *Quaternary International*, A Late Pleistocene/early Holocene archaeological 14C database for Central and South America: Palaeoenvironmental contexts and demographic interpretations 301, 34–45. <https://doi.org/10.1016/j.quaint.2012.08.2052>
- Radiocarbon, 2024. Radiocarbon Lab Information [WWW Document]. URL <https://radiocarbon.webhost.uits.arizona.edu/laboratories> (accessed 11.25.2024).
- Raetzl-Fabian, D., 1999. Editorial. *Jungsteinsite*.
- Ramsey, C.B., Dee, M.W., Rowland, J.M., Higham, T.F.G., Harris, S.A., Brock, F., Quiles, A., Wild, E.M., Marcus, E.S., Shortland, A.J., 2010. Radiocarbon-Based Chronology for Dynastic Egypt. *Science* 328, 1554–1557. <https://doi.org/10.1126/science.1189395>
- Reingruber, A., Thissen, L., 2017. The 14SEA Project: A 14C database for Southeast Europe and Anatolia (10,000–3000 calBC).
- Reingruber, A., Thissen, L., 2009. Depending on 14C Data: Chronological Frameworks in the Neolithic and Chalcolithic of Southeastern Europe. *Radiocarbon* 51, 751–770. <https://doi.org/10.1017/S0033822200056071>
- Reingruber, A., Thissen, L., 2005. 14C database for the Aegean catchment (Eastern Greece, southern Balkans and western Turkey) 10,000–5500 cal BC, in: Lichter, C. (Ed.), *How Did Farming Reach Europe?*, Byzas. Ege Yayınlar, Istanbul, pp. 295–327.
- Reiter, S.S., Staniuk, R., Kolář, J., Bulatović, J., Rose, H.A., Ryabogina, N.E., Speciale, C., Schjerven, N., Paulsson, B.S., Lee, V.Y.K., Canteri, E., Revill, A., Dahlberg, F., Sabatini, S., Frei, K.M., Racimo, F., Ivanova-Bieg, M., Traylor, W., Kate, E.J., Derenne, E., Frank, L., Woodbridge, J., Fyfe, R., Shennan, S., Kristiansen, K., Thomas, M.G., Timpson, A., 2024. The

- BIAD standards: Recommendations for archaeological data publication and insights from the big interdisciplinary archaeological database. *Open Archaeology* 10, 20240015. <https://doi.org/doi:10.1515/opar-2024-0015>
- Richter, T., Arranz-Otaegui, A., Yeomans, L., Boaretto, E., 2017. High Resolution AMS Dates from Shubayqa 1, northeast Jordan Reveal Complex Origins of Late Epipalaeolithic Natufian in the Levant. *Sci. Rep.* 7, 17025. <https://doi.org/10.1038/s41598-017-17096-5>
- Rinne, C., Kneisel, J., Hinz, M., Furrholt, M., Kruschke, N., Müller, J., Raetzl-Fabian, D., Rodens, M., Sjögren, K.-G., Vandkilde, H., Wotzka, H.-P., 2024. Rado.NB.
- Riris, P., 2021. Carribean-14C: A compilation of 2147 anthropogenic radiocarbon (14C) dates for the Caribbean region from 504 sites across 57 islands.
- Riris, P., Silva, F., Crema, E., Palmisano, A., Robinson, E., Siegel, P.E., French, J.C., Jørgensen, E.K., Maezumi, S.Y., Solheim, S., Bates, J., Davies, B., Oh, Y., Ren, X., 2024. Frequent disturbances enhanced the resilience of past human populations. *Nature* 1–6. <https://doi.org/10.1038/s41586-024-07354-8>
- Roalkvam, I., 2023. Shoredate: An R package for shoreline dating coastal Stone Age sites. *Journal of Open Source Software* 8, 5337. <https://doi.org/10.21105/joss.05337>
- Roe, J., 2024. [Rintchron: Interface to IntChron](#).
- Schmid, C., Ghalichi, A., Lamnidis, T.C., Mudiyansele, D.B.A., Haak, W., Schiffels, S., 2024. Poseidon – A framework for archaeogenetic human genotype data management. *eLife* 13. <https://doi.org/10.7554/eLife.98317.1>
- Schmid, C., Seidensticker, D., Hinz, M., 2019. c14bazAAR: An r package for downloading and preparing C14 dates from different source databases. *Journal of Open Source Software* 4, 1914. <https://doi.org/10.21105/joss.01914>
- Seidensticker, D., Schmid, C., 2021. Dirkseidensticker/aDRAC: V2.0 Pikunda. <https://doi.org/10.5281/zenodo.4530717>
- Silva, F., Steele, J., 2014. New methods for reconstructing geographical effects on dispersal rates and routes from large-scale radiocarbon databases. *Journal of Archaeological Science* 52, 609–620. <https://doi.org/10.1016/j.jas.2014.04.021>
- Suess, H.E., 1967. [Bristlecone pine calibration of the radiocarbon time scale from 4100 B.C. To 1500 B.C.](#)
- Tanindi, O., Erdoğan, B., 2005. Recording the past: Towards an inventory of the archaeological sites of Turkey. *Conservation and Management of Archaeological Sites* 7, 125–130. <https://doi.org/10.1179/135050305793137468>
- Tenopir, C., Allard, S., Douglass, K., Aydinoglu, A.U., Wu, L., Read, E., Manoff, M., Frame, M., 2011. Data Sharing by Scientists: Practices and Perceptions. *PLOS ONE* 6, e21101. <https://doi.org/10.1371/journal.pone.0021101>
- Thomsen, C.J., 1836. *Ledetrad til Nordiske Oldkindighed*. Copenhagen.
- Uriarte González, A., Fernández Freire, C., Fraguas Bravo, A., Castañeda Clemente, N., Capdevila Montes, E., Salas Tovar, E., Gilman, A., Bosque González, I. del, Vicent García, J.M., 2017. IDEArq-C14: Una Infraestructura de Datos Espaciales para la cronología radiocarbónica de la Prehistoria



- Reciente ibérica. <https://doi.org/10.13039/501100003329>
- Van Strydonck, M., De Roock, E., n.d. Radiocarbon and stable isotope measurements. Royal Institute for Cultural Heritage.
- Verborgh, R., van Hooland, S., Cope, A.S., Chan, S., Mannens, E., Van de Walle, R., 2015. The fallacy of the multi-API culture: Conceptual and practical benefits of Representational State Transfer (REST). *Journal of Documentation* 71, 233–252. <https://doi.org/10.1108/JD-07-2013-0098>
- Vermeersch, P.M., 2024. Radiocarbon Palaeolithic Europe Database.
- Vermeersch, P.M., 2020. Radiocarbon Palaeolithic Europe database: A regularly updated dataset of the radiometric data regarding the Palaeolithic of Europe, Siberia included. *Data Brief* 31, 105793. <https://doi.org/10.1016/j.dib.2020.105793>
- Voorburg, R., 2012. Archaeological Atlas of Antiquity - Vici.org [WWW Document]. URL <https://vici.org/> (accessed 11.25.2024).
- Wang, C., Lu, H., Zhang, J., Gu, Z., He, K., 2014. Prehistoric demographic fluctuations in China inferred from radiocarbon data and their linkage with climate change over the past 50,000 years. *Quaternary Science Reviews* 98, 45–59. <https://doi.org/10.1016/j.quascirev.2014.05.015>
- Weninger, B., 2022. CalPal. <https://doi.org/10.5281/ZENODO.7215741>
- Whitley, D.S., 2012. In *Suspect Terrain: Dating Rock Engravings*, in: A Companion to Rock Art. John Wiley & Sons, Ltd, pp. 605–624. <https://doi.org/10.1002/9781118253892.ch34>
- Williams, A.N., Smith, M., 2013. *AustArch3: A database of  $^{14}\text{C}$  and luminescence ages from archaeological sites in southern Australia*. *Australian Archaeology* 102–102.
- Williams, A.N., Smith, M.A., 2012. *AustArch2: A database of  $^{14}\text{C}$  and luminescence ages from archaeological sites in the Top End*. *Australian Archaeology* 146–146.
- Williams, A.N., Smith, M.A., Turney, C.S.M., Cupper, M.L., 2008. *AustArch1: A Database of  $^{14}\text{C}$  and Luminescence Ages from Archaeological Sites in the Australian Arid Zone*. *Australian Archaeology* 99–99.
- Williams, A., Ulm, S., Smith, M., Reid, J., 2014. *AustArch: A Database of  $^{14}\text{C}$  and Non- $^{14}\text{C}$  Ages from Archaeological Sites in Australia - Composition, Compilation and Review (Data Paper)*. *Internet Archaeology*. <https://doi.org/10.11141/ia.36.6>
- Williams, J.W., Grimm, E.C., Blois, J.L., Charles, D.F., Davis, E.B., Goring, S.J., Graham, R.W., Smith, A.J., Anderson, M., Arroyo-Cabrales, J., Ashworth, A.C., Betancourt, J.L., Bills, B.W., Booth, R.K., Buckland, P.I., Curry, B.B., Giesecke, T., Jackson, S.T., Latorre, C., Nichols, J., Purdum, T., Roth, R.E., Stryker, M., Takahara, H., 2018. The Neotoma Paleoecology Database, a multiproxy, international, community-curated data resource. *Quaternary Research* 89, 156–177. <https://doi.org/10.1017/qua.2017.105>
- Worsaae, J.J.A., 1843. *Danmarks Oldtid oplyst ved Oldsager og Gravhøie*. Selskabet for Trykkefrihedens rette Brug, Copenhagen.