

**Cool summers are stormy summers: precession reorganizes extratropical
cyclone activity through the quiet season**

Xiaoxu Shi^a

^a *Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research, Bremerhaven,
Germany*

Corresponding author: Xiaoxu Shi, shixiaoxu@sml-zhuhai.cn

7 ABSTRACT: Extratropical cyclones peak in winter, while orbital precession reshapes the seasonal
8 insolation cycle mainly in summer. Thus storm tracks might be expected to ignore the precession
9 cycle. This is examined with twelve equilibrium simulations using a coupled climate model AWI-
10 ESM2, spanning a full precession cycle at 30° longitude increments under fixed eccentricity and
11 obliquity. Extratropical cyclones are explicitly tracked based on 6-hourly atmospheric pressure
12 fields for both hemispheres. While the annual-mean cyclone response to precession is small,
13 substantial seasonal response emerges. In the North Atlantic, the North Pacific, and the Southern
14 Ocean the response focuses on local summer. We also find summers spent near perihelion feature
15 suppressed storminess, whereas summers near aphelion experience intensified cyclone activity,
16 yielding antiphase mid-latitude storm variability between the two hemispheres over the precession
17 cycle. This signal is carried by cyclone number and position, and it follows the pronounced
18 summer response of the Eady growth rate driven primarily by vertical wind shear in all three
19 basins. Precession therefore shapes extratropical storm tracks mostly through their quiet season
20 (i.e., summer). Summer-sensitive proxy archives should record a strong and hemispherically
21 antiphased storminess signal at the precession period that annually integrating archives would
22 miss. Moreover, monthly storm responses are phase-locked to the calendar timing of perihelion,
23 lagged by 1-2 months due to surface thermal inertia. This fingerprint also appears when we apply
24 the same cyclone tracking algorithm to pre-industrial (PI, 0 ka, perihelion falling in winter), Last
25 Interglacial (LIG, 127 ka, perihelion falling in summer) and mid-Holocene (MH, 6 ka, perihelion
26 falling in autumn), with weaker amplitudes matching their weaker eccentricity.

27 SIGNIFICANCE STATEMENT: Long orbital cycles drive global climate variability, and the
28 strongest of these, precession, reshapes sunlight mainly in summer, while mid-latitude storms
29 happen mainly in winter. This creates an apparent disconnect between precessional forcing and
30 storm track variability. Climate model simulations covering a full precession cycle show that
31 the orbit nevertheless controls storminess by acting through summer. When perihelion falls in a
32 hemisphere's summer, that summer has fewer mid-latitude storms and the storm belt sits farther
33 poleward, and the two hemispheres evolve in antiphase through the precession cycle. Simulations
34 of the Last Interglacial, when perihelion falls in northern summer, reproduce this pattern. This
35 tells researchers where to look for orbital storm signals in geological records, namely in archives
36 sensitive to summer, not to the yearly average.

37 1. Introduction

38 Extratropical cyclones govern mid-latitude weather systems. They transport the majority of
39 poleward heat and moisture outside the tropics, supply a substantial fraction of mid-latitude pre-
40 cipitation, and aggregate into the canonical storm-track pathways of the North Atlantic, North
41 Pacific, and Southern Ocean (Hoskins and Valdes 1990; Chang et al. 2002; Hoskins and Hodges
42 2002, 2005). These cyclones develop via baroclinic instability driven by the meridional tempera-
43 ture gradient (Eady 1949), a gradient that reaches its maximum during winter. Cyclone activities
44 consequently peak in the cold season, with consistent wintertime maxima (e.g., frequency) evi-
45 dent in both the Northern Hemisphere (Hoskins and Hodges 2002) and the Southern Hemisphere
46 (Simmonds and Keay 2000). Mid-latitude cyclogenesis is therefore most favored during winter.

47 Precession, by contrast, exerts its primary forcing on warm seasons. Slow variations in the longi-
48 tude of perihelion represent the leading orbital control on seasonal insolation cycles. It redistributes
49 incoming solar radiation, and the amplitude of the anomalies scales with a season's background
50 solar radiation (Berger 1978; Laskar et al. 2004). Mid-latitude summertime insolation fluctuates
51 by over 100 W m^{-2} across a full precession cycle, whereas midwinter insolation experiences only
52 minor orbital-driven changes. This stark contrast explains why summer solar forcing serves as
53 the most impactful component of climate changes (Huybers 2006). Since extratropical cyclone
54 activity is inherently winter-favored, this invites a simple expectation that precession should exert

negligible control over mid-latitude storm tracks. But this expectation has never been put to a clean test.

Previous studies have looked closely at precessional signals, and they find seasonal shifts in climate. This is well established in monsoon research, from the classic orbital experiments of Kutzbach and Guetter (1986) to single-forcing designs that isolate precession explicitly (Bosmans et al. 2015; Merlis et al. 2013), and cave records show the northern and southern monsoons swinging in antiphase as perihelion migrates through the year (Wang et al. 2008). The extratropics have received far less attention. Kaspar et al. (2007) simulated a stronger and poleward-shifted North Atlantic winter storm track for the Eemian from two orbital time slices. Brayshaw et al. (2010) linked Holocene orbital shifts to North Atlantic and Mediterranean winter storm tracks via changes in baroclinicity. And Varma et al. (2012) found the Southern Hemisphere westerlies shift under Holocene orbital forcing. Explicit cyclone tracking has entered paleoclimate work mainly at the Last Glacial Maximum, where ice sheets dominate the forcing (Pinto and Ludwig 2020; Raible et al. 2021), and in transient simulations of the last millennium (Hess and Chemke 2025). For tropical cyclones the precessional response has recently been examined in high-resolution paleoclimate simulations, where it is governed by atmospheric moisture (Raavi et al. 2023). Here our study provides the extratropical counterpart, where cyclone growth is baroclinic. To our knowledge, no previous study has applied explicit cyclone tracking across a full precession cycle to isolate the orbital response of the extratropical storm tracks, and none has asked which season carries the response.

Many paleoclimate archives that record past storm activity, such as wind-blown dust and sediment, or rainfall linked to westerly winds, only capture signals from specific seasons instead of annual averages. If a proxy record mainly reflects one single season, it can either hide or falsely create an orbital signal, depending on which time of year carries the strongest response (Laepfle et al. 2011). Fully understanding which season holds the clearest precessional storm signal is therefore essential for properly interpreting these geological records. Moreover, different metrics for cyclone behavior, e.g., total storm counts, how much rain each storm produces, and storm intensity, react differently to orbital forcing. We must evaluate these quantities separately before drawing conclusions from any single geological proxy.

84 To fill this gap, we run 12 equilibrium climate simulations covering a full precession cycle.
85 Every extratropical cyclone in both hemispheres is tracked individually using 6-hour sea level
86 pressure. Our experimental design isolates precession at amplified eccentricity with all other
87 boundary conditions fixed, so that the seasonal storm response to precession forcing alone can be
88 examined. The results are then tested in simulations of real orbital states, namely PI, MH, and
89 LIG. Section 2 describes our model and simulation, and the method we use for storm tracking and
90 analysis. Section 3 presents the results. We discuss and conclude in section 4 and 5 respectively.

91 **2. Methodology**

92 *a. Model and experiments*

93 The simulations are performed with the coupled Earth system model AWI-ESM2 (Sidorenko
94 et al. 2019). The atmosphere component is ECHAM6 at T63 resolution with 47 vertical levels
95 (Stevens et al. 2013), which includes the land surface module JSBACH (Reick et al. 2021). The
96 ocean and sea ice component is FESOM2, a finite-volume model formulated on an unstructured
97 mesh (Danilov et al. 2017), with a spatially varying resolution of from 25 km over high latitudes
98 and coastal regions, to 150 km over open oceans.

99 We analyze 12 equilibrium simulations, taken from the idealized precession ensemble introduced
100 by Shi et al. (2025). The longitude of perihelion ω is prescribed at values from 0° to 330° in steps of
101 30° (labeled as p000, p030..., and p330), while eccentricity is fixed at 0.058 and obliquity at 24.5° .
102 The eccentricity value sits near the upper bound reached during the Quaternary and amplifies the
103 precessional effects on insolation. All remaining boundary conditions follow the pre-industrial
104 setup, and each phase is branched from a long pre-industrial control. We analyze 30 years of
105 6-hourly output from the equilibrated part of each simulation. The calendar month of perihelion
106 for each phase is computed from orbital geometry (Berger 1978), which provides the insolation
107 fields shown in Fig. 1c-d.

108 To test our results under realistic orbital configurations, we additionally conducted three equilib-
109 rium simulations including a PI (0 ka) control run, a MH (6 ka) experiment, and a LIG (127 ka)
110 experiment. Boundary conditions follow the protocols of PMIP4 (Otto-Bliesner et al. 2017). Or-
111 bital parameters are calculated following Berger (1977), and the greenhouse gas concentrations are
112 taken from multi-archive reconstructions from the Greenland and the Antarctic ice cores (Flückiger

et al. 2002; Monnin et al. 2004; Schneider et al. 2013; Schilt et al. 2010; Buiron et al. 2011). Each simulation is integrated for 1,000 model years with the trend of simulated global mean surface temperature in the final 100 model years not exceeding ± 0.05 K/century. We preserved 6-hourly sea level pressure from each simulation. At 127 ka, perihelion occurs close to the June solstice with an eccentricity of 0.039378; at 6 ka, perihelion falls within September, while the pre-industrial control features perihelion in early January.

b. Converting fixed-day calendar into angular calendar

Precession moves perihelion through the seasonal cycle, and Kepler’s second law (Kepler 1953) then changes the lengths of the seasons (Fig. 1f-g). This is the calendar effect long known in paleoclimate modeling (Joussaume and Braconnot 1997; Bartlein and Shafer 2019; Shi et al. 2021). Monthly and seasonal averages at different orbital configurations are therefore only comparable when the averaging windows follow the same position of the Earth along its orbit.

Following Shi et al. (2021), we compute the length of a month (season) by calculating how much time the Earth needs to move from the respective starting point to the endpoint. We first define the mean anomaly M as the angle between the perihelion and Earth’s position based on the assumption that the orbit describes a perfect circle with the sun at the center by:

$$M = \frac{2\pi}{T} \cdot t_p \quad (1)$$

Here, t_p denotes the time elapsed since Earth passes the perihelion and T is the Earth’s revolution period, namely 1 year or 365 days. Taking into account the orbit’s eccentricity ϵ , we define the eccentric anomaly E via:

$$E - \epsilon \cdot \sin(E) = M \quad (2)$$

Equation 2 can be solved with the application of Newton’s method (Danby and Burkardt 1983), and E follows from the true anomaly θ , the angle at the Sun between perihelion and the actual position of the Earth (see Eq. 3.13b of Curtis 2014):

$$E = 2 \cdot \arctan\left(\sqrt{\frac{1-\epsilon}{1+\epsilon}} \cdot \tan\left(\frac{\theta}{2}\right)\right) \quad (3)$$

The above equations relate t_p to θ by:

$$t_p(\theta) = \frac{MT}{2\pi} = \frac{(E - \epsilon \cdot \sin(E))T}{2\pi} \quad (4)$$

This allows us to define seasons with respect to Earth's position on the orbit rather than relying on a fixed number of days as used in our modern calendar. The insolation panels of Fig. 1 are computed on this angular calendar. Re-binning the 6-hourly cyclone statistics onto the same calendar changes the local-summer response amplitudes by at most 1.5 percentage points and the phase-locking lags by less than half a month, while the autumn response strengthens slightly. All storm diagnostics are therefore reported on the model's own fixed calendar.

c. Cyclone detection

Extratropical cyclones are detected and tracked with the algorithm of Crawford et al. (2021), and Fig. 2 summarizes the procedure. 6-hourly sea level pressure is first interpolated onto 100-km equal-area grids centered on each pole, and the two hemispheres are processed separately. A grid cell qualifies as a cyclone center when its pressure p_c lies below that of all 8 neighbors and the surrounding field keeps a sufficient inward gradient. The gradient test compares p_c with the mean pressure $\bar{p}(d)$ on a one-cell ring at distance d from the candidate,

$$\frac{\bar{p}(d) - p_c}{d} \geq \gamma \quad (5)$$

where $d = 1000$ km and $\gamma = 750 \text{ Pa} (1000 \text{ km})^{-1}$, and candidates over terrain above 1500 m are discarded. Centers closer than 1000 km are treated as one multicenter system. The cyclone area A is obtained by growing closed isobars around each center at a 200-Pa interval until a contour fails to close, and the last closed isobar defines the edge pressure p_e (Fig. 2b). Each cyclone carries the depth and the area-equivalent radius

$$D = p_e - p_c \quad (6)$$

$$R = \sqrt{A/\pi} \quad (7)$$

Tracking links the centers of consecutive 6-hourly fields (Fig. 2c). For every active cyclone a first-guess position is projected from its previous displacement, damped to three quarters,

$$\mathbf{x}_q = \mathbf{x}_t + 0.75 (\mathbf{x}_t - \mathbf{x}_{t-1}) \quad (8)$$

and a center of the next field qualifies as the continuation when it lies within $v_{\max}\Delta t$ of both the current position $\mathbf{x}_t$ and the projection $\mathbf{x}_q$, with $v_{\max} = 150 \text{ km h}^{-1}$ and $\Delta t = 6 \text{ h}$, a searching radius of 900 km per step. Among the qualifying candidates the one nearest to the projection continues the track. Centers left unmatched at the new time start tracks as genesis events, and tracks left unmatched end as lysis. Only systems poleward of 30° latitude enter the analysis.

Several track-based diagnostics are derived. Track density counts every 6-hourly cyclone position, system density counts each cyclone once at its mean position, and genesis density counts the first point of each track. For every cyclone the lifespan, the maximum depth relative to the outermost closed isobar, the minimum central pressure, the mean cyclone area, and the propagation speed are recorded. We further define a "storm day" field, representing the number of days per month during which a grid point lies inside a detected cyclone. It therefore measures the storm-influence time of each grid point, and it is affected by most cyclone metrics, such as the number of cyclones, their area, lifespan, and propagation speed.

d. Baroclinicity diagnostics

The maximum Eady growth rate serves as the measure of baroclinicity (Eady 1949; Lindzen and Farrell 1980; Hoskins and Valdes 1990). It is computed from wind, temperature, and geopotential height in the 925-700 hPa layer as

$$\sigma = 0.31 |f| \frac{|\partial U / \partial z|}{N}, \quad (9)$$

where f is the Coriolis parameter, $|\partial U / \partial z|$ the vertical wind shear, and N the Brunt-Väisälä frequency. The response to precession is quantified between the end members p090 and p270, which place perihelion at the June and December solstices.

To separate the roles of vertical shear and stratification we follow the substitution approach of Camargo et al. (2007). Writing $\sigma = 0.31 |f| S / N$ with $S = |\partial U / \partial z|$, the difference is split by

182 changing one factor at a time while holding the other at its p270 value,

$$\Delta\sigma_S = 0.31 |f| \frac{S_{90} - S_{270}}{N_{270}}, \quad (10)$$

$$\Delta\sigma_N = 0.31 |f| S_{270} \left(\frac{1}{N_{90}} - \frac{1}{N_{270}} \right), \quad (11)$$

184 so that $\Delta\sigma = \Delta\sigma_S + \Delta\sigma_N + \varepsilon$, with a small cross term ε .

185 *e. Basins and statistics*

186 Three storm-track basins are analyzed, the North Atlantic (30–75°N, 80°W–20°E), the North
 187 Pacific (30–75°N, 120°E–120°W), and the Southern Ocean (40–75°S, all longitudes). Basin means
 188 are area weighted. Uncertainty estimates use the interannual spread of the 30 individual years in
 189 each phase, and the significance of the end-member difference maps is assessed with a two-sided
 190 Welch t test (Welch 1947) at each grid point.

191 **3. Results**

192 *a. Summer sensitivity*

193 The 12 precession-phase annual mid-latitude storm climatologies exhibit only minor differences,
 194 and phase anomalies relative to the multi-phase ensemble mean manifest as weak, slowly evolving
 195 dipoles aligned along major storm track pathways (Fig. 3). The basin-averaged time series quantify
 196 the muted amplitude of this annual-mean signal, with across-phase ranges of 10%, 17% and 4% in
 197 the North Atlantic, the North Pacific and the Southern Ocean, varying smoothly with ω . We also
 198 observe an interhemispheric antiphase.

199 There is also a pronounced seasonal cycle. Extratropical cyclone activities peak in DJF in North-
 200 ern Hemisphere basins and JJA across the Southern Ocean, with this seasonal timing unchanged
 201 across all orbital setups (Fig. 4a–c). Cold-season cyclones also carry greater intensity, with winter
 202 peak cyclone depths exceeding summertime values by 70-100 % within the Northern Hemisphere
 203 storm basins (not shown). Overall, winter dominates mid-latitude cyclone activity uniformly across
 204 hemispheres and all orbital phases.

205 Precession exerts negligible control over annual-mean storm conditions but profoundly reshapes
 206 seasonal cyclone variability, and its strongest modulation targets the low-activity warm season.
 207 Though each precession phase retains the same wintertime peak in basin-averaged storm track
 208 density, cross-phase deviations emerge almost entirely within each basin's local summer (Fig. 4a–c).
 209 Anomalies between end-members (p090 and p270) quantify this stark seasonal contrast (Fig. 4d–f).
 210 For the North Pacific, track density drops by $0.10 \text{ cell}^{-1} \text{ month}^{-1}$ in boreal summer (JJA), about
 211 44% of the phase mean, against an annual-mean decrease of only 7%. The North Atlantic follows
 212 the same pattern, with a summertime decrease near 23% versus 5% annually. The Southern Ocean
 213 mirrors this behavior in its local summer (DJF), where track density rises by 19%, against a 2%
 214 annual increase. All three basins identify local summer as the most orbitally sensitive season.
 215 Moreover, summer and winter vary in antiphase across the precession cycle, with perihelion
 216 suppressing the summer storms while slightly enhancing the winter ones. Therefore the annual
 217 signal is muted.

218 This summertime storm response is locked to orbital geometry. When basin-averaged track-
 219 density anomalies are plotted against both calendar month and precession angle ω , the band of
 220 peak anomalies traces a diagonal ridge aligned with the calendar timing of perihelion across the
 221 full phase space (Fig. 4g–i), with a delay of 1–2 months. Summers spent near perihelion exhibit
 222 suppressed storm activity, whereas aphelion summers feature intensified storminess. It should be
 223 noted that northern summer perihelion aligns with southern summer aphelion, precessional forcing
 224 thus drives antiphased mid-latitude storm signals in the two hemispheres.

225 The pronounced seasonal dependence of orbital storm signals originates from insolation forcing.
 226 Averaged across the Northern Hemisphere storm-track band ($40\text{--}70^\circ\text{N}$), precessional insolation
 227 variability peaks at 105 W m^{-2} during boreal summer (JJA) and falls to merely 22 W m^{-2} in
 228 boreal winter (DJF). The Southern Hemisphere storm band ($40\text{--}70^\circ\text{S}$) exhibits mirror-symmetric
 229 behavior, with a maximum seasonal forcing amplitude of 105 W m^{-2} in austral summer (DJF)
 230 versus just 22 W m^{-2} in austral winter (JJA) (Fig. 1e). The underlying mechanism lies in the
 231 multiplicative nature of precessional insolation modulation. Shifting perihelion across the annual
 232 cycle alters the Sun–Earth distance for each month, which rescales monthly incoming solar flux by
 233 a factor proportional to eccentricity e (approximately $4e$, equaling 23% for $e = 0.058$, Fig. 1a–b).
 234 Midwinter at mid-latitudes features minimal climatological insolation, leaving little baseline flux

to be modulated by precessional shifts. Precession forcing therefore exerts its strongest modulation on local summer in each hemisphere, and barely perturbs deep winter conditions (Fig. 1a-e).

Orbital insolation anomalies are translated into extratropical storm activity via shifts in mid-latitude baroclinicity. Differences in maximum Eady growth rate between the extreme precession phases $\omega = 90^\circ$ and $\omega = 270^\circ$ reach their peak magnitude within each basin's local summer (Fig. 5). Normalized against seasonal climatological means, summertime Eady growth rate responses hit 31 % in the North Atlantic, a pronounced 92 % in the North Pacific, and 21 % in the Southern Ocean. A decomposition analysis further reveals that vertical wind shear dominates the precessional Eady growth rate anomalies, while static stability contributes negligible signal strength. This is consistent across all storm tracks.

b. Meridional shift

Spatial difference maps between the two orbital end-members (p090 minus p270) yields weak responses for the annual means (Fig. 6a). By contrast, boreal summer (JJA) exhibits a prominent dipole pattern across Northern Hemisphere basins, with fewer storm days through the mid-latitude core and more along the Arctic flank at perihelion summer, the signature of a poleward shift joined to an overall weakening. In DJF the signal migrates to the Southern Ocean and forms a circumpolar banded structure of opposite sign, marking enhanced mid-latitude storm activity and an equatorward shift of the southern storm belt.

The latitudinal displacement of storm tracks driving these dipole anomalies is quantified in Fig. 7. For the North Pacific boreal summer, core storm latitude shifts from 53°N when perihelion falls in northern winter to 61°N when it falls in northern summer, yielding an 8° total poleward migration that peaks precisely at $\omega = 90^\circ$. The North Atlantic summertime storm track undergoes a 4.4° poleward shift following identical orbital behavior, while the Southern Ocean mirrors this response with a 3.8° poleward displacement when perihelion coincides with austral summer (near p270). A perihelion summer therefore weakens storm activity and shifts the storm population poleward, consistent with the poleward shift of baroclinic activity evident in the Eady growth rate dipole (Fig. 5a).

By contrast, the annual-mean track latitude varies by no more than 1.3° in any basin. Winter storm-track latitudes only fluctuate by $1.5\text{--}2.8^\circ$ across the full precession cycle, and their latitudinal

extrema do not align with solstitial orbital phases. This further confirms our earlier conclusion that mid-latitude winter storms are orbitally insensitive. Spring and autumn display intermediate latitudinal variability with a coherent phase dependence. For example, autumn poleward displacement maximizes near $\omega = 150^\circ$, the orbital configuration that places perihelion in late summer. Latitudinal poleward shifts therefore track the calendar timing of perihelion identically to Fig. 4g–i, with a uniform lag of 1–2 months, consistent with surface thermal inertia. Figure 8 quantifies this basin-dependent lag. All basin-scale storm extrema cluster along basin-specific dashed lines offset from the perihelion reference. The response lags the perihelion calendar by distinct intervals across regions, near 1 month in the North Pacific, 1-2 months in the North Atlantic, and slightly over 2 months in the Southern Ocean.

c. Cyclone precipitation

The hydrological signature of precessional forcing evolves in antiphase with storm dynamical responses (Fig. 9). During boreal summer over the North Pacific, mean precipitation per individual storm rises by as much as 15 % at the orbital phase where total storm frequency reaches its minimum, yielding nearly perfectly opposing variability between the two metrics. Perihelion summers feature warmer, more humid mid-latitude atmospheres, which amplifies precipitation intensity within the remaining cyclones. Nevertheless, the sharp reduction in storm number dominates the total signal, such that basin-wide summertime precipitation drops by 17 % between the two extreme orbital phases. The fractional share of precipitation originating from cyclones also declines markedly, with a full-cycle range of 9.5 % in North Pacific summer compared to just 3 % for the annual mean. Precessional forcing thus exerts two counteracting controls on extratropical hydroclimate simultaneously. It reduces total cyclone counts while boosting precipitation intensity per storm.

d. Realistic paleosimulations (MH and LIG)

Our results based on the 12 idealized simulations can be validated against the realistic orbital-state simulations (PI, MH, and LIG) introduced in Section 2. In LIG, perihelion occurs near June solstice, and the tracked cyclone responses fully reproduce our core theoretical pattern. Relative to PI, summer storm track density decreases by 8% over the North Atlantic and 11% over the North Pacific; winter cyclone changes remain within approximately 3%. Meanwhile, austral summer

storm track density across the Southern Ocean rises by 8%. Northern Hemisphere summer storm tracks shift poleward by 1–2 degrees, while the Southern Hemisphere summer storm belt shifts equatorward (Fig. 10). The summer difference composites show identical weakening and poleward displacement of Northern Hemisphere storm tracks as seen in our amplified idealized end-member simulations (Fig. 11). In MH, perihelion happens in September, leaving seasonal forcing dominated by warm September conditions with cool March conditions. Consistent with this seasonal signal, MH storm variability features fewer, poleward-shifted cyclones in autumn and the opposite trend in spring. The magnitude of storm responses in these paleosimulations is much weaker than that of our amplified end-member contrasts (p090 vs. p270, Fig. 6), consistent with their smaller eccentricity values of the real orbits. Notably, these experiments combine realistic precession with their own obliquity and greenhouse gases, so they test our theoretical prediction under multi-factor natural orbital forcing rather than isolating precessional effects.

4. Discussion

The seasonal preference seen in storm responses originates purely from orbital geometry. Precession redistributes insolation in proportion to the insolation a season already has, meaning winter hemispheres have very little background insolation for orbital changes to modify (Berger 1978). Fig. 1 clearly illustrates this stark seasonal contrast in forcing strength. Winter insolation varies by only 22 W m^{-2} across the precession cycle, while summer values reach 105 W m^{-2} within mid-latitude storm-track latitudes. This is model independent and will hold in any climate models as it simply reflects the orbital geometry behind the well-known summer-dominated orbital forcing framework (Huybers 2006). Existing monsoon studies has long confirmed that summer bears the strongest precessional signal (Kutzbach and Guetter 1986; Bosmans et al. 2015). But the new insight of our work lies elsewhere. This summer-focused orbital forcing acts on a winter-dominated weather system, reshaping storm activity exclusively during its normally quiet warm season. Note that the term “quiet” here refers to the weaker overall intensity of the summer storm track, not its population. Both hemisphere still hosts abundant storm systems in local summer, and they are merely weaker, shallower and move more slowly compared to winter cyclones.

The mechanism that converts summer insolation into summer storms is the baroclinic instability. Our factor decomposition analysis confirms that the shear term carries essentially the whole Eady

321 growth rate response in all three ocean basins, while static stability plays a negligible role. We
322 see similar physical mechanism operating in the modern Northern Hemisphere, where greenhouse
323 warming takes the place of perihelion-driven heating. Stronger summer warming at high latitudes
324 has weakened summertime large-scale circulation, eddy kinetic energy, and the number of strong
325 summer cyclones (Coumou et al. 2015; Chang et al. 2016; Gertler and O’Gorman 2019). Future
326 climate projections show the same shear-based decline of Northern Hemisphere summer cyclone
327 activity and baroclinic energy as globe warms (Lehmann et al. 2014; Priestley and Catto 2022).
328 But a difference should be noted here, as greenhouse warming is a transient forcing that both
329 hemispheres change in phase, whereas our simulations are equilibrated and precession swings both
330 hemispheres in antiphase. Even so, both forcings follow one unified rule that is the core theme of
331 this study: cooler summers host more storm activity.

332 Compared to summer, storm systems in winter are much more stable. Across all three ocean
333 basins, winter storm activity changes by less than 10% over a full precession cycle, even though
334 winter Eady growth rates shift by 16–18% between the two orbital end-members. Weak winter
335 insolation forcing cannot fully explain this stability. The winter storm track appears saturated
336 in the sense that ample baroclinicity is available in every phase, so that moderate changes in the
337 growth rate do not translate into changes in occurrence. Transient simulation of the last millennium
338 likewise found winter cyclone nearly insensitive to external forcing (Raible et al. 2018; Hess and
339 Chemke 2025), and those forcings acted to affect storm track in summer (Hess and Chemke 2025).
340 Our findings expand this conclusion to the far larger swings in solar radiation driven by precession,
341 and this insensitivity only holds for wintertime conditions.

342 Geological records sensitive to summertime conditions within mid-latitude storm belts should
343 preserve prominent precession-scale storm signals, whereas proxies that average climate signals
344 across winter or the full year will show only minor orbital imprint. Just as observed for tem-
345 perature reconstructions, the seasonal bias of a given archive can either generate artificial orbital
346 storm signals or fully obscure true precessional variations (Laepple et al. 2011). Furthermore,
347 summer-sensitive storm records from both hemispheres will fluctuate oppositely over precessional
348 cycles, forming an extratropical equivalent of the well-documented monsoon seesaw (Wang et al.
349 2008). Current published sediment records already support the feasibility of testing this frame-
350 work. Distinct precession-period signals have been detected in Northern Hemisphere mid-latitude

351 westerly proxies (Li et al. 2019), as well as North Pacific dust records. Precessional variation
352 appears specifically in moisture-sensitive dust fractions (Zhong et al. 2024). Meanwhile, dust
353 intensity archives that capture winter-only signals display very weak precessional power (Chen
354 et al. 2014). Regarding the Southern Hemisphere, subtropical margins of the southern westerly
355 belt exhibit clear precession cycles (Lamy et al. 2019), consistent with our results. Records of the
356 southern westerlies also show orbital-band migrations of the belt itself (Lamy et al. 2010; Varma
357 et al. 2012), and glacial-state cyclone modeling has shown how strongly reconstructed storminess
358 depends on which cyclone property a proxy senses (Pinto and Ludwig 2020).

359 Our study is based on idealized simulations designed to isolate pure precessional forcing. The
360 experimental setup holds eccentricity, obliquity, ice sheets, and greenhouse gas concentrations
361 fixed and does not incorporate transient climate evolution or orbital cross-interactions. As such,
362 our results represent dynamical responses to idealized orbital conditions rather than reconstructions
363 of specific geological epochs. Future work could incorporate time-varying boundary conditions
364 and full orbital combinations to further generalize the storm seasonal response across broader
365 paleoclimate backgrounds.

366 5. Conclusions

367 In our study, we carry out a set of equilibrium climate simulations covering a full precession cycle,
368 with explicit tracking of all extratropical cyclones across both hemispheres to isolate precessional
369 forcing. We find that precession regulates extratropical cyclone primarily through local summer,
370 the season with naturally low storm activity, while winter storm tracks has only minor change
371 across all precession phases. Storm activity calms down when perihelion falls in local summer
372 and intensifies when aphelion coincides with local summer. And, as perihelion summer in one
373 hemisphere matches aphelion summer in the other, mid-latitude storminess of the two hemispheres
374 varies in antiphase during a precession cycle. This orbital signal propagates through shear-driven
375 baroclinicity and expresses itself in cyclones numbers and position, both of which are phase locked
376 to the calendar month of perihelion, with a delay of 1-2 months. Our result also bring clear
377 implications for paleoclimate proxy interpretation. Orbital storm signals should be extracted from
378 mid-latitude archives sensitive to summer climate, and the recorded signals will display opposite
379 variations between the two hemispheres.




FIG. 1. (a-b) Range of monthly mean insolation across the 12 precession phases for the 40–70°N and 40–70°S bands, computed on the classical fixed-day calendar (red) and on the angular calendar (blue). The dashed line marks the distance factor $4e/(1-e^2)^2$ multiplied by the monthly mean insolation. (c-d) Insolation anomaly as a function of calendar month and ω on the angular calendar, with the perihelion (red) and aphelion (blue) months overlaid. (e) Insolation range in the two bands for each season on the angular calendar, computed as maximum minus minimum insolation across the 12 phases. Units: W m^{-2} . (f),(g) The orbital geometry behind the calendar drift, shown for two precession end-members (i.e., p090 and p270). The equinox and solstice lines through the Sun divide the orbit into the four seasons. Seasons on the perihelion side last about 85 days and seasons on the aphelion side 98 days, and this nearly two-week contrast swaps sides between the two phases. Day counts hold for $e = 0.058$ with a 365-day year, while the drawn eccentricity is exaggerated for clarity.




FIG. 2. Cyclone detection and tracking procedure. (a,c) One 6-hourly sea level pressure field of phase p000 (contours, 8-hPa interval) with all detected cyclone centers poleward of 30°N (dots) and their area-equivalent radii $R = \sqrt{A/\pi}$ (circles), 15 January of one model year. (b,e) The deepest system of that field, with the outermost closed isobar p_e (dashed), the cyclone center (cross), the central and edge pressures and depth annotated, and the binary mask cells (shading) defined by $\text{SLP} < p_e$ inside the $\sqrt{A}$ search window; contour interval 4 hPa. (c,f) All cyclone tracks of the same month (lines) with genesis points (dots). (d) Data processing workflow.




FIG. 3. Annual-mean storm days across the precession cycle. (a)–(l) Annual storm-day anomaly of each phase relative to the cross-phase mean (shading), with the phase-mean climatology outlined at 5 and 8 d month⁻¹ (contours). (m)–(o) Basin-mean annual storm days against ω with interannual standard errors; the across-phase range as a percentage of the phase mean is annotated in each panel.




FIG. 4. Seasonal concentration of the precessional storm response, from the Lagrangian tracks. (a)–(c) Basin-mean track density (cell⁻¹ month⁻¹) by calendar month for the North Atlantic, the North Pacific, and the Southern Ocean, one line per phase colored by ω , with local summer shaded. (d)–(f) Track-density difference between the end members p090 and p270 (cell⁻¹ month⁻¹) for the annual mean and the four seasons, with the local-summer bar highlighted. (g)–(i) Basin-mean track-density anomaly relative to the phase mean as a function of calendar month and precession phase (unsmoothed), with the perihelion (red) and aphelion (blue) calendar months of each phase overlaid.




FIG. 5. Precessional response of the maximum Eady growth rate (925–700 hPa). (a) p090 minus p270 difference for JJA and (b) for DJF (day^{-1}). (c) Seasonal response amplitude in each basin, the RMS of the p090 minus p270 difference over the storm box as a percentage of the seasonal climatology, with stars marking local summer. (d) Projection shares of the vertical-shear and static-stability factors in the local-summer difference pattern.




FIG. 6. Difference in storm days (d month^{-1}) between p090 and p270 for (a) the annual mean, (b) JJA, and (c) DJF. Stippling marks grid points where the difference is significant at the 95% level based on a two-sided Welch t test.




FIG. 7. Storm-track latitude across the precession cycle, the storm-day-weighted mean latitude over each basin sector shown as the poleward anomaly from each season's phase mean, for (a) the North Atlantic, (b) the North Pacific, and (c) the Southern Ocean. Lines give the annual mean and the four seasons, with the local summer of each basin drawn heavy with markers. Vertical dotted lines mark the phase with perihelion in the local summer of each basin.




FIG. 8. Phase locking of the storm response to the perihelion calendar. For each calendar month and basin, the phase of the first harmonic across the 12 precession phases of (a) basin-mean storm days, shown as the ω of the activity minimum, and (b) the storm-day-weighted track latitude, shown as the ω of the poleward maximum. Marker size scales with the harmonic amplitude. The red line gives the ω that places perihelion in that calendar month, and dashed lines repeat it delayed by each basin's lag over the amplitude-strong months (annotated in months).




FIG. 9. Storm-attributed precipitation across the precession cycle. (a) North Pacific JJA percent anomalies from the phase mean of basin-mean storm days, per-storm precipitation, and total precipitation; the dotted line marks the phase with perihelion in NH summer. (b) Across-phase range of the storm-attributed share of precipitation, accumulated precipitation inside storm footprints divided by total accumulated precipitation (percentage points), for the annual mean and local summer in each basin.




FIG. 10. Seasonal storm response of the real-orbit experiments, tracked with the same pipeline as the main ensemble and referenced to the preindustrial control. (a)–(c) Anomalies of basin-mean track density for the mid-Holocene (gold) and the Last Interglacial (red), as percentages of the preindustrial mean, for the annual mean and the four seasons, with error bars giving twice the standard error of the 30-year difference and the local summer shaded. Track density is normalized per T63 grid cell, and the annotated preindustrial values give the absolute scale in the units of Fig. 4. (d)–(f) The corresponding change in the track-point mean latitude, positive poleward.




FIG. 11. Storm-day anomalies of the real-orbit experiments relative to the preindustrial control, from the same storm-day product as the main ensemble. (a)–(d) Mid-Holocene minus preindustrial for DJF, MAM, JJA, and SON. (e)–(h) Last Interglacial minus preindustrial. Contours outline the preindustrial climatology at 5 and 8 d month^{-1} . The isolated warm-season maxima over subtropical continental interiors reflect stationary surface heat lows rather than traveling cyclones.

505 *Acknowledgments.* The simulations were performed at the German Climate Computing Center
506 (DKRZ). [TODO funding and grant numbers before submission.]

507 *Data availability statement.* The AWI-ESM simulations of the idealized precession ensemble
508 are described in Shi et al. (2025). The cyclone-track database, the gridded storm statistics, and
509 the analysis scripts underlying all figures will be archived and assigned a DOI upon acceptance.
510 [TODO repository DOI.]

References

- Bartlein, P. J., and S. L. Shafer, 2019: Paleo calendar-effect adjustments in time-slice and transient climate-model simulations (paleocaladjust v1. 0): Impact and strategies for data analysis. *Geoscientific Model Development*, **12** (9), 3889–3913.
- Berger, A., 1977: Long-term variations of the earth’s orbital elements. *Celestial Mechanics*, **15** (1), 53–74.
- Berger, A., 1978: Long-term variations of daily insolation and quaternary climatic changes. *Journal of Atmospheric Sciences*, **35** (12), 2362–2367.
- Bosmans, J., S. Drijfhout, E. Tuenter, F. Hilgen, and L. Lourens, 2015: Response of the north african summer monsoon to precession and obliquity forcings in the ec-earth gcm. *Climate dynamics*, **44** (1), 279–297.
- Brayshaw, D. J., B. Hoskins, and E. Black, 2010: Some physical drivers of changes in the winter storm tracks over the north atlantic and mediterranean during the holocene. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, **368** (1931), 5185–5223.
- Buiron, D., and Coauthors, 2011: Taldice-1 age scale of the talos dome deep ice core, east antarctica. *Climate of the Past*, **7** (1), 1–16.
- Camargo, S. J., K. A. Emanuel, and A. H. Sobel, 2007: Use of a genesis potential index to diagnose enso effects on tropical cyclone genesis. *Journal of Climate*, **20** (19), 4819–4834.
- Chang, E. K., S. Lee, and K. L. Swanson, 2002: Storm track dynamics. *Journal of climate*, **15** (16), 2163–2183.
- Chang, E. K., C.-G. Ma, C. Zheng, and A. M. Yau, 2016: Observed and projected decrease in northern hemisphere extratropical cyclone activity in summer and its impacts on maximum temperature. *Geophysical Research Letters*, **43** (5), 2200–2208.
- Chen, J., X. Liu, and V. A. Kravchinsky, 2014: Response of the high-resolution chinese loess grain size record to the 50° n integrated winter insolation during the last 500,000 years. *Geophysical Research Letters*, **41** (17), 6244–6251.

538 Coumou, D., J. Lehmann, and J. Beckmann, 2015: The weakening summer circulation in the
539 northern hemisphere mid-latitudes. *Science*, **348** (6232), 324–327.

540 Crawford, A. D., E. A. Schreiber, N. Sommer, M. C. Serreze, J. C. Stroeve, and D. G. Barber,
541 2021: Sensitivity of northern hemisphere cyclone detection and tracking results to fine spatial
542 and temporal resolution using era5. *Monthly Weather Review*, **149** (8), 2581–2598.

543 Curtis, H., 2014: Orbital position as a function of time. *Orbital Mechanics for Engineering*
544 *Students, 3rd edition, Elsevier, Amsterdam, the Netherlands*, 145–186.

545 Danby, J., and T. Burkardt, 1983: The solution of kepler’s equation, i. *Celestial Mechanics*, **31** (2),
546 95–107.

547 Danilov, S., D. Sidorenko, Q. Wang, and T. Jung, 2017: The finite-volume sea ice–ocean model
548 (fesom2). *Geoscientific Model Development*, **10** (2), 765–789.

549 Eady, E. T., 1949: Long waves and cyclone waves. *Tellus*, **1** (3), 33–52.

550 Flückiger, J., E. Monnin, B. Stauffer, J. Schwander, T. F. Stocker, J. Chappellaz, D. Raynaud, and
551 J.-M. Barnola, 2002: High-resolution holocene n2o ice core record and its relationship with ch4
552 and co2. *Global Biogeochemical Cycles*, **16** (1), 10–1.

553 Gertler, C. G., and P. A. O’Gorman, 2019: Changing available energy for extratropical cyclones and
554 associated convection in northern hemisphere summer. *Proceedings of the National Academy of*
555 *Sciences*, **116** (10), 4105–4110.

556 Hess, O., and R. Chemke, 2025: Forced changes in the north atlantic storm track over the last
557 millennium. *Geophysical Research Letters*, **52** (15), e2025GL115 894.

558 Hoskins, B. J., and K. I. Hodges, 2002: New perspectives on the northern hemisphere winter storm
559 tracks. *Journal of the Atmospheric Sciences*, **59** (6), 1041–1061.

560 Hoskins, B. J., and K. I. Hodges, 2005: A new perspective on southern hemisphere storm tracks.
561 *Journal of climate*, **18** (20), 4108–4129.

562 Hoskins, B. J., and P. J. Valdes, 1990: On the existence of storm-tracks. *Journal of Atmospheric*
563 *Sciences*, **47** (15), 1854–1864.

- Huybers, P., 2006: Early pleistocene glacial cycles and the integrated summer insolation forcing. *Science*, **313** (5786), 508–511.
- Joussaume, S., and P. Braconnot, 1997: Sensitivity of paleoclimate simulation results to season definitions. *Journal of Geophysical Research: Atmospheres*, **102** (D2), 1943–1956.
- Kaspar, F., T. Spanghel, and U. Cubasch, 2007: Northern hemisphere winter storm tracks of the eemian interglacial and the last glacial inception. *Climate of the Past*, **3** (2), 181–192.
- Kepler, J., 1953: *Epitome astronomiae copernicanae*. Jo. Plancus.
- Kutzbach, J. E., and P. J. Guetter, 1986: The influence of changing orbital parameters and surface boundary conditions on climate simulations for the past 18 000 years. *Journal of atmospheric sciences*, **43** (16), 1726–1759.
- Laepple, T., M. Werner, and G. Lohmann, 2011: Synchronicity of antarctic temperatures and local solar insolation on orbital timescales. *Nature*, **471** (7336), 91–94.
- Lamy, F., R. Kilian, H. W. Arz, J.-P. Francois, J. Kaiser, M. Prange, and T. Steinke, 2010: Holocene changes in the position and intensity of the southern westerly wind belt. *Nature Geoscience*, **3** (10), 695–699.
- Lamy, F., and Coauthors, 2019: Precession modulation of the south pacific westerly wind belt over the past million years. *Proceedings of the National Academy of Sciences*, **116** (47), 23 455–23 460.
- Laskar, J., P. Robutel, F. Joutel, M. Gastineau, A. C. Correia, and B. Levrard, 2004: A long-term numerical solution for the insolation quantities of the earth. *Astronomy & Astrophysics*, **428** (1), 261–285.
- Lehmann, J., D. Coumou, K. Frieler, A. V. Eliseev, and A. Levermann, 2014: Future changes in extratropical storm tracks and baroclinicity under climate change. *Environmental Research Letters*, **9** (8), 084 002.
- Li, Y., Y. Song, Q. Yin, L. Han, and Y. Wang, 2019: Orbital and millennial northern mid-latitude westerlies over the last glacial period. *Climate Dynamics*, **53** (5), 3315–3324.

- 590 Lindzen, R., and B. Farrell, 1980: A simple approximate result for the maximum growth rate of
591 baroclinic instabilities. *Journal of Atmospheric Sciences*, **37** (7), 1648–1654.
- 592 Merlis, T. M., T. Schneider, S. Bordoni, and I. Eisenman, 2013: Hadley circulation response to
593 orbital precession. part i: Aquaplanets. *Journal of Climate*, **26** (3), 740–753.
- 594 Monnin, E., and Coauthors, 2004: Evidence for substantial accumulation rate variability in antarctica
595 during the holocene, through synchronization of co₂ in the taylor dome, dome c and dml ice
596 cores. *Earth and Planetary Science Letters*, **224** (1-2), 45–54.
- 597 Otto-Bliesner, B., and Coauthors, 2017: The pmip4 contribution to cmip6–part 2: Two interglacials,
598 scientific objective and experimental design for holocene and last interglacial simulations.
599
- 600 Pinto, J. G., and P. Ludwig, 2020: Extratropical cyclones over the north atlantic and western europe
601 during the last glacial maximum and implications for proxy interpretation. *Climate of the Past*,
602 **16** (2), 611–626.
- 603 Priestley, M. D., and J. L. Catto, 2022: Future changes in the extratropical storm tracks and cyclone
604 intensity, wind speed, and structure. *Weather and Climate Dynamics*, **3**, 337–360.
- 605 Raavi, P. H., J.-E. Chu, A. Timmermann, S.-S. Lee, and K. J. Walsh, 2023: Moisture control
606 of tropical cyclones in high-resolution simulations of paleoclimate and future climate. *Nature
607 Communications*, **14** (1), 6426.
- 608 Raible, C. C., M. Messmer, F. Lehner, T. F. Stocker, and R. Blender, 2018: Extratropical cyclone
609 statistics during the last millennium and the 21st century. *Climate of the Past*, **14** (10), 1499–1514.
- 610 Raible, C. C., J. G. Pinto, P. Ludwig, and M. Messmer, 2021: A review of past changes in
611 extratropical cyclones in the northern hemisphere and what can be learned for the future. *Wiley
612 Interdisciplinary Reviews: Climate Change*, **12** (1), e680.
- 613 Reick, C. H., and Coauthors, 2021: Jsbach 3-the land component of the mpi earth system model:
614 documentation of version 3.2.

Schilt, A., M. Baumgartner, T. Blunier, J. Schwander, R. Spahni, H. Fischer, and T. F. Stocker, 2010: Glacial–interglacial and millennial-scale variations in the atmospheric nitrous oxide concentration during the last 800,000 years. *Quaternary Science Reviews*, **29** (1-2), 182–192.

Schneider, R., J. Schmitt, P. Köhler, F. Joos, and H. Fischer, 2013: A reconstruction of atmospheric carbon dioxide and its stable carbon isotopic composition from the penultimate glacial maximum to the last glacial inception. *Climate of the Past*, **9** (6), 2507–2523.

Shi, X., M. Werner, H. Yang, Q. Gao, J. Liu, and G. Lohmann, 2025: Precession controls on climate and water isotope signals in northern africa. *Paleoceanography and Paleoclimatology*, **40** (1), e2024PA004 999.

Shi, X., and Coauthors, 2021: Calendar effects on surface air temperature and precipitation based on model-ensemble equilibrium and transient simulations from pmip4 and pacmedy. *Climate of the Past Discussions*, **2021**, 1–29.

Sidorenko, D., and Coauthors, 2019: Evaluation of fesom2. 0 coupled to echam6. 3: preindustrial and highresmpip simulations. *Journal of Advances in Modeling Earth Systems*, **11** (11), 3794–3815.

Simmonds, I., and K. Keay, 2000: Mean southern hemisphere extratropical cyclone behavior in the 40-year ncep–ncar reanalysis. *Journal of Climate*, **13** (5), 873–885.

Stevens, B., and Coauthors, 2013: Atmospheric component of the mpi-m earth system model: Echam6. *Journal of Advances in Modeling Earth Systems*, **5** (2), 146–172.

Varma, V., and Coauthors, 2012: Holocene evolution of the southern hemisphere westerly winds in transient simulations with global climate models. *Climate of the Past*, **8** (2), 391–402.

Wang, Y., and Coauthors, 2008: Millennial-and orbital-scale changes in the east asian monsoon over the past 224,000 years. *nature*, **451** (7182), 1090–1093.

Welch, B. L., 1947: The generalization of ‘student’s’ problem when several different population variances are involved. *Biometrika*, **34** (1-2), 28–35.

Zhong, Y., and Coauthors, 2024: Orbital controls on north pacific dust flux during the late quaternary. *Geophysical Research Letters*, **51** (4), e2023GL106 631.