

**Cool summers are stormy summers: precession reorganizes extratropical
cyclone activity through the quiet season**

Xiaoxu Shi^a

^a *Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research, Bremerhaven,
Germany*

Corresponding author: Xiaoxu Shi, shixiaoxu@sml-zhuhai.cn

7 ABSTRACT: Extratropical cyclones peak in winter, while orbital precession reshapes the sea-
8 sonal insolation cycle most strongly in summer. Thus storm tracks might be expected to ignore the
9 precession cycle. This is examined with twelve equilibrium simulations using a coupled climate
10 model AWI-ESM2, spanning a full precession cycle at 30° longitude increments under fixed eccen-
11 tricity and obliquity. Extratropical cyclones are explicitly tracked based on 6-hourly atmospheric
12 pressure fields for both hemispheres. While the annual-mean cyclone response to precession is
13 muted, substantial seasonal variability emerges. In the North Atlantic, the North Pacific, and the
14 Southern Ocean the response concentrates in local summer, where storm activity ranges across the
15 phases by up to 68% of its climatological mean in cyclone influence time (44% in cyclone counts),
16 while winter changes stay below about 10%. We also find summers spent near perihelion feature
17 suppressed storminess, whereas summers near aphelion experience intensified cyclone activity,
18 yielding antiphase midlatitude storm variability between the two hemispheres over the precession
19 cycle. This signal is carried by cyclone number and position, and it follows the pronounced
20 summer response of the Eady growth rate driven primarily by vertical wind shear in all three
21 basins. Monthly storm responses are phase-locked to the calendar timing of perihelion, lagged
22 by 1–2 months, consistent with surface thermal inertia. Precession therefore acts on extratropical
23 storm tracks through their quiet season. The same fingerprint emerges when the identical tracking
24 is applied to simulations of the Last Interglacial and the mid-Holocene, with weaker amplitudes
25 matching their weaker eccentricity. Summer-sensitive proxy archives should record a strong and
26 hemispherically antiphased storminess signal at the precession period that annually integrating
27 archives would miss.

28 SIGNIFICANCE STATEMENT: Long orbital cycles drive global climate variability, and the
29 strongest of these, precession, reshapes sunlight mainly in summer, while mid-latitude storms
30 happen mainly in winter. This creates an apparent disconnect between precessional forcing and
31 storm track variability. Climate model simulations covering a full precession cycle show that
32 the orbit nevertheless controls storminess by acting through summer. When perihelion falls in a
33 hemisphere's summer, that summer has fewer mid-latitude storms and the storm belt sits farther
34 poleward, and the two hemispheres evolve in antiphase through the precession cycle. Simulations
35 of the Last Interglacial, when perihelion fell in northern summer, reproduce this pattern. This
36 tells researchers where to look for orbital storm signals in geological records, namely in archives
37 sensitive to summer, not to the yearly average.

38 1. Introduction

39 Extratropical cyclones govern midlatitude weather systems. They transport the majority of
40 poleward heat and moisture outside the tropics, supply a substantial fraction of midlatitude pre-
41 cipitation, and aggregate into the canonical storm-track pathways of the North Atlantic, North
42 Pacific, and Southern Ocean (Hoskins and Valdes 1990; Chang et al. 2002; Hoskins and Hodges
43 2002, 2005). These cyclones develop via baroclinic instability driven by the meridional tem-
44 perature gradient (Eady 1949), a gradient that reaches its maximum steepness during winter.
45 Cyclone activities consequently peak in the cold season, with ~~consistent wintertime maxima~~
46 ~~(e.g., frequency , strength) evident in both the Northern and Southern Hemispheres~~ a consistent
47 wintertime frequency maximum in the Northern Hemisphere (Hoskins and Hodges 2002) and in
48 the Southern Hemisphere (Simmonds and Keay 2000). Midlatitude cyclogenesis is therefore most
49 favorable during winter.

50 Precession, by contrast, exerts its primary forcing on warm seasons. Slow variations in the
51 longitude of perihelion represent the leading orbital control on seasonal insolation cycles. This
52 mechanism redistributes incoming solar radiation, and the amplitude of insolation anomalies
53 scales ~~proportionally~~ with a season's baseline solar irradiance (Berger 1978; Laskar et al. 2004).
54 Midlatitude summertime insolation fluctuates by over 100 W m^{-2} across a full precession cycle,
55 whereas midwinter insolation experiences only minor orbital modulation. This stark contrast
56 explains why orbital climate frameworks single out summertime solar forcing as the most impactful

57 component of climate changes (Huybers 2006). Since extratropical cyclone activity is inherently
58 winter-favored, this invites a simple expectation that precession should exert negligible control
59 over midlatitude storm tracks. But this expectation has never been put to a clean test.

60 Previous studies have looked closely at precessional signals, and they find seasonal shifts in
61 climate. This is well established in monsoon research, from the classic orbital experiments of
62 Kutzbach and Guetter (1986) to single-forcing designs that isolate precession explicitly (Bosmans
63 et al. 2015; Merlis et al. 2013), and cave records show the northern and southern monsoons
64 swinging in antiphase as perihelion migrates through the year (Wang et al. 2008). The extratropics
65 have received far less attention. Kaspar et al. (2007) simulated a stronger and poleward-shifted
66 North Atlantic winter storm track for the Eemian from two orbital time slices. Brayshaw et al.
67 (2010) linked Holocene orbital shifts to North Atlantic and Mediterranean winter storm tracks
68 via changes in baroclinicity. And Varma et al. (2012) found the Southern Hemisphere westerlies
69 migrating under Holocene orbital forcing. Explicit cyclone tracking has entered paleoclimate
70 work mainly at the Last Glacial Maximum, where ice sheets dominate the forcing (Pinto and
71 Ludwig 2020; Raible et al. 2021), and in transient simulations of the last millennium (Hess and
72 Chemke 2025). For tropical cyclones the precessional response has recently been isolated in
73 high-resolution paleoclimate simulations(Raavi et al. 2023)., where it is governed by atmospheric
74 moisture (Raavi et al. 2023). Our study provides the extratropical counterpart, for storms that grow
75 by baroclinic instability instead. The winter-focused precedents asked how orbital forcing moves
76 the storm track in its strongest season, and their time-slice or transient designs blend precession
77 with obliquity, eccentricity, and ice-sheet changes. To our knowledge, no previous study has
78 applied explicit cyclone tracking across a full precession cycle to isolate the orbital response of the
79 extratropical storm tracks, and none has asked which season carries that response.

80 This missing piece of research matters beyond just storm dynamics. Many paleoclimate archives
81 that record past storm activity such as wind-blown dust and sediment, or rainfall linked to westerly
82 winds only capture signals from specific seasons instead of annual averages. If a proxy record
83 mainly reflects one single season, it can either hide or falsely create an orbital signal, depending
84 on which time of year carries the strongest storm response (Laepple et al. 2011). Fully understand-
85 ing which season holds the clearest precessional storm signal is therefore essential for properly
86 interpreting these geological records. Moreover, different metrics for cyclone behavior, e.g., total

87 storm counts, how much rain each storm produces, and storm intensity, react differently to orbital
88 forcing. Researchers must evaluate these quantities separately before drawing conclusions from
89 any single geological proxy.

90 To fill this gap, we run 12 equilibrium climate simulations covering a full precession cycle.
91 Every extratropical cyclone in both hemispheres is tracked individually using 6-hour sea level
92 pressure. Our experimental design isolates precession at amplified eccentricity with all other
93 boundary conditions fixed, so that the seasonal storm response to precession forcing alone can
94 be examined. The resulting predictions are then tested against simulations of real orbital states.
95 Section 2 describes our model and simulation, and the method we use for storm tracking and
96 analysis. Section 3 presents the results, and ~~discuss and conclude in~~ section 4 discusses and
97 concludes.

98 2. Methodology

99 *a. Model and experiments*

100 The simulations are performed with the coupled Earth system model AWI-ESM2 (Sidorenko
101 et al. 2019). The atmosphere component is ECHAM6 at T63 resolution with 47 vertical levels
102 (Stevens et al. 2013), which includes the land surface module JSBACH (Raddatz et al. 2007). The
103 ocean and sea ice component is FESOM2, a finite-volume model formulated on an unstructured
104 mesh (Danilov et al. 2017), with a spatially varying resolution of from 25 km over high latitudes
105 and coastal regions, to 150 km over open oceans.

106 12 equilibrium simulations are analyzed, taken from the idealized precession ensemble introduced
107 by Shi et al. (2025). The longitude of perihelion ω is prescribed at values from 0° to 330° in steps
108 of 30° , while eccentricity is fixed at 0.058 and obliquity at 24.5° . The eccentricity value sits
109 near the upper bound reached during the Quaternary and amplifies the precessional modulation of
110 insolation. The quoted response amplitudes are therefore upper bounds, and their scaling toward
111 lower eccentricity is not tested here. The seasonal concentration and the hemispheric antiphase
112 rest on orbital geometry alone and do not depend on this amplification. All remaining boundary
113 conditions follow the pre-industrial setup, and each phase is branched from a long pre-industrial
114 control. 30 years of 6-hourly output from the equilibrated part of each simulation form the basis
115 of the cyclone analysis. The calendar month of perihelion for each phase is computed from

orbital geometry (Berger 1978), and the same formulary provides the insolation fields shown in Fig. 1. To test the results under realistic orbital states, we further performed three equilibrium simulations with the same model, a preindustrial control, a mid-Holocene (6 ka) experiment, and a Last Interglacial (127 ka) experiment. Thirty years of six-hourly output from each are processed with the identical detection and tracking pipeline. At 127 ka perihelion falls near the June solstice with an eccentricity of 0.039, at 6 ka it falls in September, and in the preindustrial control it falls in early January.

b. Cyclone detection

Extratropical cyclones are detected and tracked with the algorithm of Crawford et al. (2021), and Fig. 2 summarizes the procedure with one 6-hourly field of the ensemble. 6-hourly sea level pressure is first interpolated onto 100-km equal-area grids centered on each pole, and the two hemispheres are processed separately. A grid cell qualifies as a cyclone center when its pressure p_c lies below that of all 8 neighbors and the surrounding field keeps a sufficient inward gradient. The gradient test compares p_c with the mean pressure $\bar{p}(d)$ on a one-cell ring at distance d from the candidate,

$$\frac{\bar{p}(d) - p_c}{d} \geq \gamma \quad (1)$$

where $d = 1000$ km and $\gamma = 750 \text{ Pa} (1000 \text{ km})^{-1}$, and candidates over terrain above 1500 m are discarded. Centers closer than 1000 km are treated as one multicenter system. The cyclone area A is obtained by growing closed isobars around each center at a 200-Pa interval until a contour fails to close, and the last closed isobar defines the edge pressure p_e (Fig. 2b). Each cyclone carries the depth and the area-equivalent radius

$$D = p_e - p_c \quad (2)$$

$$R = \sqrt{A/\pi} \quad (3)$$

Tracking links the centers of consecutive 6-hourly fields (Fig. 2c). For every active cyclone a first-guess position is projected from its previous displacement, damped to three quarters,

$$\mathbf{x}_q = \mathbf{x}_t + 0.75 (\mathbf{x}_t - \mathbf{x}_{t-1}) \quad (4)$$

and a center of the next field qualifies as the continuation when it lies within $v_{\max}\Delta t$ of both the current position $\mathbf{x}_t$ and the projection $\mathbf{x}_q$, with $v_{\max} = 150 \text{ km h}^{-1}$ and $\Delta t = 6 \text{ h}$, a search radius of 900 km per step. Among the qualifying candidates the one nearest to the projection continues the track. Centers left unmatched at the new time start tracks as genesis events, and tracks left unmatched end as lysis. Only systems poleward of 30° latitude enter the analysis.

~~Several track-based diagnostics are derived. Track density counts every 6-hourly cyclone position, system density counts each cyclone once at its mean position, and genesis density counts the first point of each track.~~ The main diagnostic derived from the tracks is track density, the number of six-hourly cyclone positions falling in each grid cell per month. Tracks that merely cross the boundaries of the monthly processing windows would appear as spurious genesis or lysis events, and such truncated events (about 7% of the total) are removed. ~~For every cyclone the lifespan, the maximum depth relative to the outermost closed isobar, the minimum central pressure, the mean cyclone area, and the propagation speed are recorded.~~ The depth, the propagation speed, and the remaining life-cycle properties of each cyclone are recorded alongside.

C. *Gridded storm coverage*

~~A complementary Eulerian diagnostic measures how long each location spends under cyclone influence. At every six-hourly time step a binary mask flags the grid cells that lie inside a square search window of half-width $\sqrt{A}$ grid cells around each cyclone center and whose pressure lies below the edge pressure of that cyclone,~~

$$\underline{M(\mathbf{x}, t) = 1}$$

~~where $p(\mathbf{x}, t) < p_e$.~~ The same pipeline also yields a storm-day field, the number of days per month during which a grid point lies inside the window, and the masks are footprint of a detected cyclone, counted from the six-hourly fields and ~~regridged to T63 and accumulated into monthly storm days, one-quarter of the flagged six-hourly steps. Storm days at a point therefore scale.~~ This Eulerian measure is convenient for maps of the storm distribution and of its displacement. It scales with the product of cyclone number and cyclone area ~~footprint. The two factors can carry opposite seasonal cycles, and over the Southern Ocean the larger size of summer cyclones raises summertime storm~~

164 ~~days even though cyclone counts peak in winter. Frequency, so frequency~~ statements in this paper
165 ~~consequently rest on~~ rest on track density.

166 All statistics rest on this single detection scheme. Absolute cyclone counts differ across tracking
167 algorithms by a factor of two or more (Neu et al. 2013), and the ~~track-based densities, while~~
168 ~~storm days serve as a smooth field for maps and for measures of cyclone influence time. The~~
169 ~~gridded product covers all twelve phases. An early processing inconsistency in the $\omega = 30^\circ$ mask~~
170 ~~product was traced to a stale intermediate file and removed by regenerating the storm masks~~
171 ~~of all twelve experiments with one pipeline version~~ strict gradient criterion of Eq. (1) keeps the
172 census conservative. The analysis therefore reports relative responses across phases rather than
173 absolute counts, and the ~~regenerated fields were verified month by month against the Lagrangian~~
174 ~~statistics~~ tracking-free variability measure introduced below provides a check that carries no tracking
175 assumptions.

176 An Eulerian measure of synoptic variability provides a further check that involves no tracking
177 at all. Bandpass sea level pressure variability is computed from the 6-hourly fields with the 24-h
178 difference filter of Wallace et al. (1988), which isolates fluctuations on the synoptic time scales that
179 define the classical storm tracks (Blackmon 1976), and its monthly standard deviation is analyzed
180 with the same basins, seasons, and response measures.

181 *c. Baroclinicity diagnostics*

182 The maximum Eady growth rate serves as the measure of baroclinicity (Eady 1949; Lindzen
183 and Farrell 1980; Hoskins and Valdes 1990). It is computed as $\sigma = 0.31 |f| |\partial U / \partial z| N^{-1}$ in the
184 925–700 hPa layer from monthly wind, temperature, and geopotential height. The response to
185 precession is quantified between the end members p090 and p270, which place perihelion at the
186 June and December solstices. Because the difference fields are dipolar, plain box averages cancel,
187 and the response amplitude is instead defined as the root-mean-square of the difference over a
188 basin, normalized by the climatological growth rate of the season. A substitution decomposition
189 in the style of Camargo et al. (2007) splits the difference field into a vertical-shear factor and a
190 static-stability factor, and the share of each factor is obtained by projecting it onto the full difference
191 pattern. The basin averages use the storm-track boxes defined in the next sub-section. This choice
192 matters in the Southern Hemisphere, where a sector beginning at 30°S would admit the seasonal

signal of the subtropical winter jet and disguise it as a storm-belt response. The shear and stability shares need not sum to one, because the two factors are not orthogonal.

d. Basins and statistics

Three storm-track basins are analyzed, the North Atlantic (30–75°N, 80°W–20°E), the North Pacific (30–75°N, 120°E–120°W), and the Southern Ocean (40–75°S, all longitudes). Basin means are area weighted. The size of the precessional response is reported as the range across the twelve phases, the maximum minus the minimum of the 30-year mean, expressed as a percentage of the phase mean. Uncertainty estimates use the interannual spread of the 30 individual years in each phase, and the significance of the end-member difference maps is assessed with a two-sided Welch t test (Welch 1947) at each grid point—, treating the 30 years as independent samples; the stippling in the maps is pointwise. The local-summer contrasts between extreme phases far exceed the interannual noise, with p values below 10^{-12} in all three basins. Ranges are quoted separately for track density and for storm days, and every percentage in this paper names its metric, because the influence-time measure compounds cyclone number with cyclone area and swings harder than the count-based densities.

3. Results

a. The summer paradox

The 12 precession-phase annual midlatitude storm climatologies exhibit only minor differences, and phase anomalies relative to the multi-phase ensemble mean manifest as weak, slowly evolving dipoles aligned along major storm track pathways (Fig. 3). The basin-averaged time series quantify the muted amplitude of this annual-mean signal, with across-phase ranges of annual storm days of 10%, 17% and 4% in the North Atlantic, the North Pacific and the Southern Ocean, varying smoothly with ω . We also observe an interhemispheric antiphase.

Beneath this annual background lies a pronounced seasonal cycle of storm activity. Extratropical cyclone activities peak in DJF in Northern Hemisphere basins and JJA across the Southern Ocean, with this seasonal timing unchanged across all orbital setups —(Fig. 4a–c; Table 1). Cold-season cyclones also carry greater intensity.—~~Winter~~, with winter peak cyclone depths ~~exceed~~ exceeding summertime values by ~~approximately 50~~70–100 % within the Northern Hemisphere storm basins

221 (Fig. 4a–c; Table 1 not shown). Overall, winter dominates midlatitude cyclone activity uniformly
222 across hemispheres and all orbital phases.

223 Precession exerts negligible control over annual-mean storm conditions but profoundly reshapes
224 seasonal cyclone variability, and its strongest modulation targets the low-activity warm season.
225 Though each precession phase retains the same wintertime peak in basin-averaged storm track
226 density, cross-phase deviations emerge almost entirely within each basin’s local summer (Fig. 4a–c).
227 Seasonal range statistics quantify this stark seasonal contrast in orbital sensitivity (Fig. 4d–f). For
228 the North Pacific, the ~~total across-precession variability across-phase range of track density~~ accounts
229 for just 7 % of the phase mean on an annual basis, yet rises to 44 % in boreal summer (JJA). The
230 North Atlantic follows an identical pattern, with annual variability of 7 % versus a summertime
231 range of 25 %. The Southern Ocean mirrors this behavior but shifts the sensitive season to its local
232 summer (DJF), with an annual range of only 4 % compared to a 22 % summer signal. ~~All basins~~
233 ~~uniformly~~ Cyclone system and genesis densities confirm the same seasonal calendar (not shown).
234 The influence-time diagnostic swings harder still, with summertime storm-day variability reaching
235 68 % of the climatological mean in the North Pacific (Table 1), because storm days compound
236 cyclone number with cyclone area. All three basins identify local summer as the ~~orbital-sensitive~~
237 ~~season~~ most orbitally sensitive season in the track-based metrics, and the small annual signal is
238 what survives of a substantial summer swing after the stable winter is averaged in.

239 A measure free of tracking assumptions supports this calendar with one refinement. The bandpass
240 sea level pressure variability responds to precession almost entirely in the warm half of the year
241 (Fig. A1). Over the North Pacific its summer response is nearly four times that of any other season.
242 Over the North Atlantic the peak shifts to early autumn, the calendar counterpart of the September
243 lobe of the insolation forcing, and over the Southern Ocean the storm-belt core responds most
244 strongly in local summer while the subtropical flank keeps a winter signal. Deep winter stays quiet
245 in this metric as well.

246 This summertime storm response is locked to orbital geometry. When basin-averaged track-
247 density anomalies are plotted against both calendar month and precession angle ω , the band of
248 peak anomalies traces a diagonal ridge aligned with the calendar timing of perihelion across the
249 full phase space (Fig. 4g–i), with a delay of ~~1–2~~ 1–2 months. Summers spent near perihelion
250 exhibit suppressed storm activity, whereas aphelion summers feature intensified storminess. A

single precessional forcing thus drives opposing midlatitude storm responses between the two hemispheres across the orbital cycle, as northern summer perihelion aligns with southern summer aphelion under orbital precession, and vice versa, reversing the insolation control on warm-season storminess between hemispheres.

b. The dynamical bridge runs through summer shear

The pronounced seasonal dependence of orbital storm signals originates directly from insolation forcing. Averaged across the Northern Hemisphere storm-track band (40–70°N), precessional insolation variability peaks at 105 W m⁻² during boreal summer (JJA) and falls to merely 14 W m⁻² in boreal winter (DJF). The Southern Hemisphere storm band (40–70°S) exhibits mirror-symmetric behavior, with a maximum seasonal forcing amplitude of 119 W m⁻² in austral winter (DJF) versus just 47 W m⁻² in austral summer (JJA) (Fig. 1). The underlying mechanism lies in the multiplicative nature of precessional insolation modulation. Shifting perihelion across the annual cycle alters the Sun–Earth distance for each month, which rescales monthly incoming solar flux by a factor proportional to eccentricity e (approximately $4e$, equaling 23 % for $e = 0.058$). Midwinter at midlatitudes features minimal climatological insolation, leaving little baseline flux to be modulated by precessional shifts. Precession forcing therefore exerts its strongest modulation on local summer in each hemisphere, and barely perturbs deep winter conditions.

Orbital insolation anomalies are translated into extratropical storm activity via shifts in midlatitude baroclinicity. Differences in maximum Eady growth rate between the extreme precession phases $\omega = 90^\circ$ and $\omega = 270^\circ$ reach their peak magnitude within each basin’s local summer (Fig. 5). Normalized against seasonal climatological means, summertime Eady growth rate responses hit 31 % in the North Atlantic, a pronounced 92 % in the North Pacific, and 21 % in the Southern Ocean. A decomposition analysis further reveals that vertical wind shear dominates the precessional Eady growth rate anomalies, while static stability contributes negligible signal strength. This is consistent across all storm tracks. The thermal footprint behind this response is shown in Fig. A2. Perihelion summers warm the summer hemisphere, ~~flatten the strongly,~~ and the 850-hPa warming peaks over the subpolar continents while the ice-covered Arctic warms less. The meridional temperature gradient ~~weaken the thermal-wind shear, and suppress the baroclinic growth that sustains warm-season extratropical cyclones.~~ The consequently flattens along the equatorward

flank of the storm track and steepens along its poleward flank, the thermal counterpart of the growth-rate dipole, so the baroclinic source weakens in the storm-track core and shifts poleward. The Southern Hemisphere mirrors this structure in DJF with reversed sign. The storm anomalies appear somewhat poleward of the growth-rate anomalies, consistent with the downstream development of baroclinic waves (Chang et al. 2002).

c. Spatial expression, and what precession actually moves

Spatial difference maps between the two orbital end-members (p090 minus p270) yields weak responses for the annual means (Fig. 6a). By contrast, boreal summer (JJA) exhibits a prominent dipole pattern across Northern Hemisphere basins, with fewer storm days through the midlatitude core and more along the Arctic flank at perihelion summer, the signature of a poleward shift joined to an overall summer weakening. In DJF the signal migrates to the Southern Ocean and forms a circumpolar banded structure of opposite sign, marking enhanced midlatitude storm activity and an equatorward shift of the southern storm belt.

The latitudinal displacement of storm tracks driving these dipole anomalies is quantified in Fig. 7. For the North Pacific boreal summer, core storm latitude shifts from 53°N when perihelion falls in northern winter to 61°N when it falls in northern summer, yielding an 8° total poleward migration that peaks precisely at $\omega = 90^{\circ}$. The North Atlantic summertime storm track undergoes a 4.4° poleward shift following identical orbital behavior, while the Southern Ocean mirrors this response with a 3.8° poleward displacement when perihelion coincides with austral summer (near p270). A perihelion summer therefore weakens storm activity and shifts the storm population poleward, consistent with the poleward shift of baroclinic activity evident in the Eady growth rate dipole (Fig. 5a).

By contrast, the annual-mean track latitude varies by no more than 1.3° in any basin. Winter storm-track latitudes only fluctuate by $1.5\text{--}2.8^{\circ}$ across the full precession cycle, and their latitudinal extrema do not align with solstitial orbital phases. This further confirms our earlier conclusion that midlatitude winter storms are orbitally insensitive. Spring and autumn display intermediate latitudinal variability with a coherent phase dependence. For example, autumn poleward displacement maximizes near $\omega = 150^{\circ}$, the orbital configuration that places perihelion in late summer. Latitudinal poleward shifts therefore track the calendar timing of perihelion identically to Fig. 4g–i, with a

uniform lag of ~~1–2 months~~driven by 1–2 months, consistent with surface thermal inertia. Figure 8 quantifies this basin-dependent lag. All basin-scale storm extrema cluster along basin-specific dashed lines offset from the perihelion reference. The response lags the perihelion calendar by distinct intervals across regions, near one month in the North Pacific, ~~1–2~~1–2 months in the North Atlantic, and slightly over two months in the Southern Ocean. The 30° phase spacing resolves these lags to about one month.

d. What the rain records

The hydrological signature of precessional forcing evolves in antiphase with storm dynamical responses (Fig. 9). Detected extratropical cyclones contribute roughly one-third of total midlatitude precipitation across our simulations, consistent with storm precipitation partitioning derived from observational reanalysis datasets (Hawcroft et al. 2012). During boreal summer over the North Pacific, mean precipitation per individual storm rises by as much as 15 % at the orbital phase where total storm frequency reaches its minimum, yielding nearly perfectly opposing variability between the two metrics. Perihelion summers feature warmer, more humid midlatitude atmospheres, which amplifies precipitation intensity within the remaining cyclones; nevertheless, the sharp reduction in storm number dominates the total signal, such that basin-wide summertime precipitation drops by 17 % between the two extreme orbital phases. The fractional share of precipitation originating from cyclones also declines markedly, with a full-cycle range of 9.5 % in North Pacific summer compared to just 3 % for the annual mean. Precessional forcing thus exerts two counteracting controls on extratropical hydroclimate simultaneously. It reduces total cyclone counts while boosting precipitation intensity per storm.

4. Discussion

The seasonal preference seen in storm responses originates purely from orbital geometry, independent of model physics. Precession redistributes insolation in proportion to the insolation a season already has, meaning winter hemispheres have very little background insolation for orbital changes to modify (Berger 1978). Fig. 1 clearly illustrates this stark seasonal contrast in forcing strength. Winter insolation varies by only roughly 20 W m^{-2} across the precession cycle, while summer values exceed 125 W m^{-2} within mid-latitude storm-track latitudes. This is model in-

337 dependent and will hold in any climate models as it simply reflects the orbital geometry behind
338 the well-known summer-dominated orbital forcing framework (Huybers 2006). Existing monsoon
339 studies ~~has~~ have long confirmed that summer bears the strongest precessional signal (Kutzbach
340 and Guetter 1986; Bosmans et al. 2015). But the new insight of our work lies elsewhere. This
341 summer-focused orbital forcing acts on a winter-dominated weather system, reshaping storm activ-
342 ity exclusively during its normally quiet warm season. Note that the term ~~”quiet”~~ “quiet” here refers
343 to the weaker overall intensity of the summer storm track, not its population. Both ~~hemisphere still~~
344 ~~hosts~~ hemispheres still host abundant storm systems in local summer, and they are merely weaker,
345 shallower ~~and move more slowly compared to,~~ and slower moving than winter cyclones.

346 The mechanism that converts summer insolation into summer storms is the thermal wind.
347 When perihelion falls in local summer, the corresponding hemisphere ~~experiences a warming~~
348 ~~over high latitudes. This flattens the~~ warms, most strongly over the subpolar continents. The
349 meridional temperature gradient ~~and reduces~~ flattens along the equatorward flank of the storm belt,
350 and the vertical wind shear that ~~drive baroclinic growth.~~ drives baroclinic growth weakens in
351 the storm-track core (Fig. A2). Our factor decomposition analysis confirms that the shear term
352 carries essentially the whole Eady growth rate response in all three ocean basins, while static
353 stability plays a negligible role. We see similar physical mechanism operating in the modern
354 Northern Hemisphere, where greenhouse warming takes the place of perihelion-driven heating.
355 Stronger summer warming at high latitudes has weakened summertime large-scale circulation,
356 eddy kinetic energy, and the number of strong summer cyclones (Coumou et al. 2015; Chang
357 et al. 2016; Gertler and O’Gorman 2019). Future climate projections ~~from standard CMIP models~~
358 ~~further show that~~ show the same shear-based decline of Northern Hemisphere summer cyclone
359 activity and baroclinic energy ~~will decline via this same shear-based mechanism as global as the~~
360 globe warms (Lehmann et al. 2014; Priestley and Catto 2022). ~~But a difference~~ Two differences
361 should be noted ~~here, as long-term greenhouse warming warms both hemispheres simultaneously,~~
362 ~~whereas precession swings them in antiphase.~~ Greenhouse warming is a transient forcing that both
363 hemispheres feel in phase, whereas our simulations are equilibrated and precession swings the
364 hemispheres in antiphase, which makes precession the cleaner natural experiment for the summer
365 shear mechanism. Even so, both forcings follow one unified rule that is the core theme of this
366 study: cooler summers host more storm activity.

367 Compared to summer, storm systems in winter are much more stable. Across all three ocean
368 basins, winter storm activity changes by less than 10% over a full precession cycle, ~~despite that even~~
369 ~~though~~ winter Eady growth rates shift by ~~16–18~~16–18% between the two orbital end-members, and
370 weak winter insolation forcing cannot fully explain this stability. The winter storm track appears
371 saturated, in the sense that ample baroclinicity is available in every phase, ~~so that and~~ moderate
372 changes in the growth rate do not translate into changes in occurrence, ~~though a winter occurrence~~
373 ~~that simply responds less sensitively to growth rate would produce the same numbers and cannot be~~
374 ~~excluded with twelve phases~~. A transient simulation of the last millennium likewise found cyclone
375 counts nearly insensitive to external forcing (Raible et al. 2018; Hess and Chemke 2025), ~~so~~. Our
376 findings expand this conclusion to the far larger swings in solar radiation driven by precession, and
377 this insensitivity only holds for wintertime conditions.

378 ~~Thus,~~ Geological records sensitive to summertime conditions within midlatitude storm belts
379 should preserve prominent precession-scale storm signals, whereas proxies that average climate
380 signals across winter or the full year will show only minor orbital imprint. Just as observed
381 for temperature reconstructions, the seasonal bias of a given archive can either generate artificial
382 orbital storm signals or fully obscure true precessional variations (Laepple et al. 2011). Fur-
383 thermore, summer-sensitive storm records from both hemispheres will fluctuate oppositely over
384 precessional cycles, forming an extratropical equivalent of the well-documented monsoon see-
385 saw (~~Wang et al. 2008~~). ~~(Wang et al. 2008; Erb et al. 2015)~~. ~~These statements combine into a~~
386 ~~three-part test. A genuine precessional storm signal should appear at the precession period, it~~
387 ~~should be antiphased between the hemispheres, and it should stand out in summer-biased archives~~
388 ~~while staying weak in winter-biased archives of the same region. Precipitation-based archives blend~~
389 ~~the count signal with the opposing per-storm rainfall response described below, so the cleanest~~
390 ~~targets are proxies that sense storm occurrence or storm-belt position rather than rain amount.~~ Cur-
391 rent published sediment records already support the feasibility of testing this framework. Distinct
392 precession-period signals have been detected in Northern Hemisphere midlatitude westerly proxies
393 (Li et al. 2019), as well as North Pacific dust records. Precessional variation appears specifically
394 in moisture-sensitive dust fractions, while the annually averaged dust flux responds primarily to
395 obliquity forcing (Zhong et al. 2024). Meanwhile, dust intensity archives that capture winter-only
396 signals display very weak precessional power (Chen et al. 2014). Regarding the Southern Hemi-

sphere, subtropical margins of the southern westerly belt exhibit clear precession cycles (Lamy et al. 2019), consistent with our results. And the presence or absence of precession signals at individual sites depends on their latitudinal position within the wind belt (Kaiser et al. 2024). Records of the southern westerlies also show orbital-band migrations of the belt itself (Lamy et al. 2010; Varma et al. 2012), [the positional counterpart of the poleward summer shift in Fig. 7](#), and glacial-state cyclone modeling has shown how strongly reconstructed storminess depends on which cyclone property a proxy senses (Pinto and Ludwig 2020).

[The prediction can already be tested against the simulations of real orbital states introduced in section 2, a preindustrial control, a mid-Holocene experiment, and a Last Interglacial experiment performed with the same model and tracked with the same pipeline \(Figs. 10 and 11\). At 127 ka perihelion falls near the June solstice, and the tracked response reproduces our template. Summer track density drops by 8% in the North Atlantic and 11% in the North Pacific relative to the preindustrial run, winter changes stay within about 3%, austral summer track density over the Southern Ocean rises by 8%, and the northern summer storm tracks shift poleward by one to two degrees while the austral summer belt shifts equatorward \(Fig. 10\). The summer difference maps show the same weakening and poleward displacement of the northern storm tracks that the amplified end members produce \(Fig. 11\). At 6 ka the reading is different but equally instructive. Perihelion falls in September, both solstice levers are nearly neutral, and the calendar forcing reduces to a warm September paired with a cool March. The mid-Holocene response follows exactly that pair, with fewer and poleward-shifted storms in autumn and the reverse tendency in spring, the shoulder seasons that the forcing analysis of section 3 anticipates. The response amplitudes are roughly half of our amplified end-member contrast, in line with the weaker eccentricity of the real orbits. These experiments combine realistic precession with their own obliquity and greenhouse settings, so they test the prediction under realistic combined forcing rather than isolating precession.](#)

Although the three storm basins follow the same physical mechanism, they exhibit different regional behaviors, which can be distinguished through separate diagnostic metrics. In local summer, storm frequency and track position vary substantially ~~by 16–68~~, [by 16–68%](#) across the precession cycle [depending on basin and metric](#), while storm propagation speed remains largely unchanged. By contrast, storm intensity exhibits clear basin-dependent differences. The Southern Ocean shows a 22% swing in storm frequency but only a 4% change in storm intensity. The North Pacific features

a systematic reduction in overall storm activity during perihelion summers. This is consistent with previous findings from anthropogenic warming scenarios that storm frequency and intensity respond to external forcing through independent physical mechanism (Bengtsson et al. 2009; Ulbrich et al. 2009; Catto et al. 2019). Hydrological responses further amplify this contrast. Individual storms produce stronger precipitation in summers with reduced cyclone numbers, indicating that hydrological adjustments exceed dynamical changes under orbital forcing, a pattern analogous to that identified under greenhouse warming (Bengtsson et al. 2009). Given that extratropical precipitation is predominantly generated by cyclones and frontal systems (Hawcroft et al. 2012; Catto and Pfahl 2013), this competing effect substantially alters precipitation partitioning. ~~So a precipitation proxy tend to record~~ A precipitation proxy therefore tends to record the combined signal of reduced storm occurrence and intensified single-storm rainfall during perihelion summers.

Our study is based on idealized simulations designed to isolate pure precessional forcing. The experimental setup holds eccentricity, obliquity, ice sheets, and greenhouse gas concentrations fixed and does not incorporate transient climate evolution or orbital cross-interactions. As such, our results represent dynamical responses to idealized orbital conditions rather than reconstructions of specific geological epochs. Three further limitations deserve note. The T63 atmosphere underestimates absolute cyclone depth and count, though the across-phase contrasts that carry our conclusions are less sensitive to this bias. All statistics rest on one tracking scheme, with the tracking-free variability check as the main guard. And the amplified eccentricity widens the lever, so the quoted amplitudes are upper bounds. The realistic-orbit comparison of Fig. 10, with roughly half the lever and roughly half the response, offers a first indication that the response scales with the forcing, though a dedicated eccentricity ladder is still missing. Isolating the direct radiative response from coupled ocean and sea-ice feedbacks with atmosphere-only experiments is a natural next step. Future work could incorporate time-varying boundary conditions and full orbital combinations to further generalize the storm seasonal response across broader paleoclimate backgrounds.

5. Conclusions

In our study, we carry out a set of equilibrium climate simulations covering a full precession cycle, with explicit tracking of all extratropical cyclones across both hemispheres to isolate precessional

456 forcing. We find that precession regulates extratropical ~~eyelone~~cyclones primarily through local
457 summer, the season with naturally low storm activity, while winter storm tracks ~~has only minor~~
458 ~~change~~show only minor changes across all precession phases. Storm activity calms down when
459 perihelion falls in local summer and intensifies when aphelion coincides with local summer. ~~And,~~
460 ~~as~~As perihelion summer in one hemisphere matches aphelion summer in the other, mid-latitude
461 storminess of the two hemispheres varies in antiphase during a precession cycle. This orbital signal
462 propagates through shear-driven baroclinicity and expresses itself in ~~eyelones~~cyclone numbers and
463 position, both of which are phase locked to the calendar month of perihelion, with a delay of ~~1-2~~
464 1-2 months. Our ~~result~~results also bring clear implications for paleoclimate proxy interpretation.
465 Orbital storm signals should be extracted from midlatitude archives sensitive to summer climate,
466 and the recorded signals will display opposite variations between the two hemispheres. Applying
467 the same tracking to simulations of the mid-Holocene and the Last Interglacial reproduces the
468 predicted pattern at reduced amplitude, in line with their weaker orbital lever.





FIG. 1. Precessional insolation forcing from orbital geometry with eccentricity 0.058 and obliquity 24.5° . (a) Monthly insolation anomaly relative to the phase mean, averaged over the storm-track band $40\text{--}70^\circ\text{N}$, as a function of calendar month and precession phase ω , with the perihelion (red) and aphelion (blue) calendar months of each phase overlaid. (b) As in (a), but for $40\text{--}70^\circ\text{S}$. (c) Seasonal forcing amplitude in the two bands, the seasonal mean of the monthly maximum minus minimum insolation across the twelve phases. Units: W m^{-2} .



(g)




FIG. 2. Cyclone detection and tracking procedure, illustrated with one six-hourly field of phase p000 (15 January of one model year). (a,e) One 6-hourly sea level pressure field of phase p000 (contours, 8-hPa interval) with all detected Northern Hemisphere cyclone centers poleward of 30°N (dots) and their area-equivalent radii $R = \sqrt{A/\pi}$ (circles), 15 January of one model year. (b,e) The deepest Northern Hemisphere system of that field, with the outermost closed isobar p_e (dashed), the cyclone center (cross), the central and edge pressures and depth annotated, and the binary mask cells (shading), defined by $SLP < p_e$ inside the $\sqrt{A}$ search window; contour interval 4 hPa. (c,f) All Northern Hemisphere cyclone tracks of the same month (lines) with genesis points (dots). (d) Data processing workflow—(f) As in (a)–(c), but for the Southern Hemisphere. (g) Processing chain.




FIG. 3. Annual-mean storm days across the precession cycle. (a)–(l) Annual storm-day anomaly of each phase relative to the twelve-phase mean (shading), with the phase-mean climatology outlined at 5 and 8 d month⁻¹ (contours). (m)–(o) Basin-mean annual storm days against ω with interannual standard errors; the across-phase range as a percentage of the phase mean is annotated in each panel.




FIG. 4. Seasonal concentration of the precessional storm response, from the Lagrangian tracks. (a)–(c) Basin-mean track density (cell⁻¹ month⁻¹) by calendar month for the North Atlantic, the North Pacific, and the Southern Ocean, one line per phase colored by ω , with local summer shaded. (d)–(f) Across-phase range of track density (maximum minus minimum, percent of the phase mean) for the annual mean and the four seasons, with the local-summer bar highlighted; ~~open symbols give the corresponding ranges of system and genesis densities.~~ (g)–(i) Basin-mean track-density anomaly relative to the phase mean as a function of calendar month and precession phase (unsmoothed), with the perihelion (red) and aphelion (blue) calendar months of each phase overlaid.




FIG. 5. Precessional response of the maximum Eady growth rate (925–700 hPa). (a) p090 minus p270 difference for JJA and (b) for DJF (day^{-1}). (c) Seasonal response amplitude in each basin, the RMS of the p090 minus p270 difference over the storm box as a percentage of the seasonal climatology, with stars marking local summer. (d) Projection shares of the vertical-shear and static-stability factors in the local-summer difference pattern.




FIG. 6. Difference in storm days (d month^{-1}) between p090 and p270 for (a) the annual mean, (b) JJA, and (c) DJF, on a common color scale. Stippling marks grid points where the difference is significant at the 95% level based on a two-sided Welch t test over the 30 simulated years.




FIG. 7. Storm-track latitude across the precession cycle, the storm-day-weighted mean latitude over each basin sector shown as the poleward anomaly from each season's phase mean, for (a) the North Atlantic, (b) the North Pacific, and (c) the Southern Ocean. Lines give the annual mean and the four seasons, with the local summer of each basin drawn heavy with markers. Vertical dotted lines mark the phase with perihelion in the local summer of each basin.




FIG. 8. Phase locking of the storm response to the perihelion calendar. For each calendar month and basin, the phase of the first harmonic across the 12 precession phases of (a) basin-mean storm days, shown as the ω of the activity minimum, and (b) the storm-day-weighted track latitude, shown as the ω of the poleward maximum. Marker size scales with the harmonic amplitude. The red line gives the ω that places perihelion in that calendar month, and dashed lines repeat it delayed by each basin's median lag over the amplitude-strong months (annotated in months).




FIG. 9. Storm-attributed precipitation across the precession cycle. (a) North Pacific JJA percent anomalies from the phase mean of basin-mean storm days, per-storm precipitation, and total precipitation; the dotted line marks the phase with perihelion in NH summer. (b) Across-phase range of the storm-attributed share of precipitation, accumulated precipitation inside storm footprints divided by total accumulated precipitation (percentage points), for the annual mean and local summer in each basin.




FIG. 10. Seasonal storm response of the real-orbit experiments, tracked with the same pipeline as the main ensemble and referenced to the preindustrial control. (a)–(c) Anomalies of basin-mean track density for the mid-Holocene (gold) and the Last Interglacial (red), as percentages of the preindustrial mean, for the annual mean and the four seasons, with error bars giving twice the standard error of the 30-year difference and the local summer shaded. Track density is normalized per T63 grid cell, and the annotated preindustrial values give the absolute scale in the units of Fig. 4. (d)–(f) The corresponding change in the track-point mean latitude, positive poleward.




FIG. 11. Storm-day anomalies of the real-orbit experiments relative to the preindustrial control, from the same storm-day product as the main ensemble. (a)–(d) Mid-Holocene minus preindustrial for DJF, MAM, JJA, and SON. (e)–(h) Last Interglacial minus preindustrial. Contours outline the preindustrial climatology at 5 and 8 d month⁻¹. The isolated warm-season maxima over subtropical continental interiors reflect stationary surface heat lows rather than traveling cyclones.

TABLE 1. Three-basin comparison of the precessional storm response. Ranges are max minus min across the twelve phases as a percentage of the phase mean. Storm and sensitive seasons refer to local winter and local summer. The storm season is diagnosed from cyclogenesis counts, the sensitive season from the seasonal storm-day range. The EGR response is the RMS p090–p270 difference over the basin sector relative to its climatology, and the shear share is the projection of the vertical-shear factor onto the response pattern.

Basin	Storm season	Sensitive season	Storm-day range ann/summer (%)	Summer max/min ω (°)	EGR response summer (%)	Shear share (%)
North Atlantic	DJF	JJA	10 / 32	240 / 90	31	107
North Pacific	DJF	JJA	17 / 68	270 / 90	92	106
Southern Ocean	JJA	DJF	4 / 16	60 / 210	21	109




FIG. A1. Tracking-free Eulerian storm-track metric, the monthly standard deviation of 24-h differences of six-hourly sea level pressure. (a)–(c) Relative seasonal response in each basin, the RMS of the p090 minus p270 difference over the storm box as a percentage of the seasonal climatology (as in Fig. 5c), with the local-summer bar highlighted. (d)–(f) The p090 minus p270 difference of the same quantity for the annual mean, JJA, and DJF (hPa).

Acknowledgments. The simulations were performed at the German Climate Computing Center (DKRZ). [TODO funding and grant numbers before submission.]

Data availability statement. The AWI-ESM simulations of the idealized precession ensemble are described in Shi et al. (2025). The preindustrial, mid-Holocene, and Last Interglacial simulations are available through the Earth System Grid Federation. The cyclone-track database, the gridded storm statistics, and the analysis scripts underlying all figures will be archived and assigned a DOI upon acceptance. [TODO repository DOI.]

APPENDIX

Tracking-independent and thermal diagnostics

Figures A1 and A2 present the tracking-free bandpass diagnostic and the thermal footprint of the orbital end members discussed in sections 2 and 3.




FIG. A2. Thermal footprint of the orbital end members. (a) p090 minus p270 temperature at 850 hPa for JJA and (b) for DJF (K). (c) Change in the zonal-mean meridional temperature gradient for JJA (red) and DJF (blue), with the 40–70° storm bands shaded. In the Northern Hemisphere negative values mark a steepening of the poleward temperature decline, and in the Southern Hemisphere positive values do.

References

- Bengtsson, L., K. I. Hodges, and N. Keenlyside, 2009: Will extratropical storms intensify in a warmer climate? *Journal of Climate*, **22** (9), 2276–2301, <https://doi.org/10.1175/2008JCLI2678.1>.
- Berger, A. L., 1978: Long-term variations of daily insolation and quaternary climatic changes. *Journal of the Atmospheric Sciences*, **35** (12), 2362–2367, [https://doi.org/10.1175/1520-0469\(1978\)035\(2362:LTVODI\)2.0.CO;2](https://doi.org/10.1175/1520-0469(1978)035(2362:LTVODI)2.0.CO;2).
- Blackmon, M. L., 1976: A climatological spectral study of the 500 mb geopotential height of the Northern Hemisphere. *Journal of the Atmospheric Sciences*, **33** (8), 1607–1623, [https://doi.org/10.1175/1520-0469\(1976\)033\(1607:ACSSOT\)2.0.CO;2](https://doi.org/10.1175/1520-0469(1976)033(1607:ACSSOT)2.0.CO;2).
- Bosmans, J. H. C., S. S. Drijfhout, E. Tuenter, F. J. Hilgen, L. J. Lourens, and E. J. Rohling, 2015: Response of the north african summer monsoon to precession and obliquity forcings in the earth gcm. *Climate Dynamics*, **44** (1–2), 279–297, <https://doi.org/10.1007/s00382-014-2260-z>.
- Brayshaw, D. J., B. Hoskins, and E. Black, 2010: Some physical drivers of changes in the winter storm tracks over the North Atlantic and Mediterranean during the Holocene. *Philosophical Transactions of the Royal Society A*, **368** (1931), 5185–5223, <https://doi.org/10.1098/rsta.2010.0180>.
- Camargo, S. J., K. A. Emanuel, and A. H. Sobel, 2007: Use of a genesis potential index to diagnose ENSO effects on tropical cyclone genesis. *Journal of Climate*, **20** (19), 4819–4834, <https://doi.org/10.1175/JCLI4282.1>.
- Catto, J. L., and S. Pfahl, 2013: The importance of fronts for extreme precipitation. *Journal of Geophysical Research: Atmospheres*, **118** (19), 10 791–10 801, <https://doi.org/10.1002/jgrd.50852>.
- Catto, J. L., and Coauthors, 2019: The future of midlatitude cyclones. *Current Climate Change Reports*, **5**, 407–420, <https://doi.org/10.1007/s40641-019-00149-4>.
- Chang, E. K. M., S. Lee, and K. L. Swanson, 2002: Storm track dynamics. *Journal of Climate*, **15** (16), 2163–2183, [https://doi.org/10.1175/1520-0442\(2002\)015\(02163:STD\)2.0.CO;2](https://doi.org/10.1175/1520-0442(2002)015(02163:STD)2.0.CO;2).

- 655 Chang, E. K. M., C.-T. Ma, C. Zheng, and A. M. W. Yau, 2016: Observed and projected
656 decrease in Northern Hemisphere extratropical cyclone activity in summer and its impacts
657 on maximum temperature. *Geophysical Research Letters*, **43** (5), 2200–2208, <https://doi.org/10.1002/2016GL068172>.
658
- 659 Chen, J., X. Liu, and V. A. Kravchinsky, 2014: Response of the high-resolution Chinese loess grain
660 size record to the 50°N integrated winter insolation during the last 500,000 years. *Geophysical
661 Research Letters*, **41** (17), 6244–6251, <https://doi.org/10.1002/2014GL060239>.
- 662 Coumou, D., J. Lehmann, and J. Beckmann, 2015: The weakening summer circulation in the
663 Northern Hemisphere mid-latitudes. *Science*, **348** (6232), 324–327, [https://doi.org/10.1126/
664 science.1261768](https://doi.org/10.1126/science.1261768).
- 665 Crawford, A. D., E. A. P. Schreiber, N. Sommer, M. C. Serreze, J. C. Stroeve, and D. G. Bar-
666 ber, 2021: Sensitivity of Northern Hemisphere cyclone detection and tracking results to fine
667 spatial and temporal resolution using ERA5. *Monthly Weather Review*, **149** (8), 2581–2598,
668 <https://doi.org/10.1175/MWR-D-20-0417.1>.
- 669 Danilov, S., D. Sidorenko, Q. Wang, and T. Jung, 2017: The Finite-volume Sea ice–Ocean
670 Model (FESOM2). *Geoscientific Model Development*, **10** (2), 765–789, [https://doi.org/10.5194/
671 gmd-10-765-2017](https://doi.org/10.5194/gmd-10-765-2017).
- 672 Eady, E. T., 1949: Long waves and cyclone waves. *Tellus*, **1** (3), 33–52, [https://doi.org/10.3402/
673 tellusa.v1i3.8507](https://doi.org/10.3402/tellusa.v1i3.8507).
- 674 Erb, M. P., C. S. Jackson, and A. J. Broccoli, 2015: Using single-forcing gcm simulations to
675 reconstruct and interpret quaternary climate change. *Journal of Climate*, **28** (24), 9746–9767,
676 <https://doi.org/10.1175/JCLI-D-15-0329.1>.
- 677 Gertler, C. G., and P. A. O’Gorman, 2019: Changing available energy for extratropical cyclones and
678 associated convection in Northern Hemisphere summer. *Proceedings of the National Academy
679 of Sciences*, **116** (10), 4105–4110, <https://doi.org/10.1073/pnas.1812312116>.
- 680 Hawcroft, M. K., L. C. Shaffrey, K. I. Hodges, and H. F. Dacre, 2012: How much Northern
681 Hemisphere precipitation is associated with extratropical cyclones? *Geophysical Research
682 Letters*, **39**, L24 809, <https://doi.org/10.1029/2012GL053866>.

- 683 Hess, O., and R. Chemke, 2025: Forced changes in the North Atlantic storm track over the
684 last millennium. *Geophysical Research Letters*, **52**, e2025GL115 894, [https://doi.org/10.1029/](https://doi.org/10.1029/2025GL115894)
685 2025GL115894.
- 686 Hoskins, B. J., and K. I. Hodges, 2002: New perspectives on the Northern Hemisphere winter
687 storm tracks. *Journal of the Atmospheric Sciences*, **59** (6), 1041–1061, [https://doi.org/10.1175/](https://doi.org/10.1175/1520-0469(2002)059(1041:NPOTNH)2.0.CO;2)
688 1520-0469(2002)059(1041:NPOTNH)2.0.CO;2.
- 689 Hoskins, B. J., and K. I. Hodges, 2005: A new perspective on Southern Hemisphere storm tracks.
690 *Journal of Climate*, **18** (20), 4108–4129, <https://doi.org/10.1175/JCLI3570.1>.
- 691 Hoskins, B. J., and P. J. Valdes, 1990: On the existence of storm-tracks. *Journal of the Atmospheric*
692 *Sciences*, **47** (15), 1854–1864, [https://doi.org/10.1175/1520-0469\(1990\)047\(1854:OTEOST\)2.](https://doi.org/10.1175/1520-0469(1990)047(1854:OTEOST)2.0.CO;2)
693 0.CO;2.
- 694 Huybers, P., 2006: Early pleistocene glacial cycles and the integrated summer insolation forcing.
695 *Science*, **313** (5786), 508–511, <https://doi.org/10.1126/science.1125249>.
- 696 Kaiser, J., E. Schefuß, J. A. Collins, and Coauthors, 2024: Orbital modulation of subtropical versus
697 subantarctic moisture sources in the southeast Pacific mid-latitudes. *Nature Communications*,
698 **15**, 7512, <https://doi.org/10.1038/s41467-024-51985-4>.
- 699 Kaspar, F., T. Spanghel, and U. Cubasch, 2007: Northern hemisphere winter storm tracks of
700 the eemian interglacial and the last glacial inception. *Climate of the Past*, **3** (2), 181–192,
701 <https://doi.org/10.5194/cp-3-181-2007>.
- 702 Kutzbach, J. E., and P. J. Guetter, 1986: The influence of changing orbital parameters and surface
703 boundary conditions on climate simulations for the past 18,000 years. *Journal of the Atmospheric*
704 *Sciences*, **43** (16), 1726–1759, [https://doi.org/10.1175/1520-0469\(1986\)043\(1726:TIOCOP\)2.](https://doi.org/10.1175/1520-0469(1986)043(1726:TIOCOP)2.0.CO;2)
705 0.CO;2.
- 706 Laepple, T., M. Werner, and G. Lohmann, 2011: Synchronicity of antarctic temperatures and
707 local solar insolation on orbital timescales. *Nature*, **471** (7336), 91–94, [https://doi.org/10.1038/](https://doi.org/10.1038/nature09825)
708 nature09825.

- 709 Lamy, F., R. Kilian, H. W. Arz, J.-P. Francois, J. Kaiser, M. Prange, and T. Steinke, 2010: Holocene
710 changes in the position and intensity of the southern westerly wind belt. *Nature Geoscience*,
711 **3 (10)**, 695–699, <https://doi.org/10.1038/ngeo959>.
- 712 Lamy, F., and Coauthors, 2019: Precession modulation of the South Pacific westerly wind belt
713 over the past million years. *Proceedings of the National Academy of Sciences*, **116 (47)**, 23 455–
714 23 460, <https://doi.org/10.1073/pnas.1905847116>.
- 715 Laskar, J., P. Robutel, F. Joutel, M. Gastineau, A. C. M. Correia, and B. Levrard, 2004: A long-
716 term numerical solution for the insolation quantities of the earth. *Astronomy and Astrophysics*,
717 **428 (1)**, 261–285, <https://doi.org/10.1051/0004-6361:20041335>.
- 718 Lehmann, J., D. Coumou, K. Frieler, A. V. Eliseev, and A. Levermann, 2014: Future changes
719 in extratropical storm tracks and baroclinicity under climate change. *Environmental Research*
720 *Letters*, **9 (8)**, 084 002, <https://doi.org/10.1088/1748-9326/9/8/084002>.
- 721 Li, Y., Y. Song, Q. Yin, L. Han, and Coauthors, 2019: Orbital and millennial northern mid-latitude
722 westerlies over the last glacial period. *Climate Dynamics*, **53 (5–6)**, 3315–3324, [https://doi.org/](https://doi.org/10.1007/s00382-019-04704-5)
723 [10.1007/s00382-019-04704-5](https://doi.org/10.1007/s00382-019-04704-5).
- 724 Lindzen, R. S., and B. Farrell, 1980: A simple approximate result for the maximum growth rate of
725 baroclinic instabilities. *Journal of the Atmospheric Sciences*, **37 (7)**, 1648–1654, [https://doi.org/](https://doi.org/10.1175/1520-0469(1980)037<1648:ASARFT>2.0.CO;2)
726 [10.1175/1520-0469\(1980\)037<1648:ASARFT>2.0.CO;2](https://doi.org/10.1175/1520-0469(1980)037<1648:ASARFT>2.0.CO;2).
- 727 Merlis, T. M., T. Schneider, S. Bordoni, and I. Eisenman, 2013: Hadley circulation response to
728 orbital precession. part i: Aquaplanets. *Journal of Climate*, **26 (3)**, 740–753, [https://doi.org/](https://doi.org/10.1175/JCLI-D-11-00716.1)
729 [10.1175/JCLI-D-11-00716.1](https://doi.org/10.1175/JCLI-D-11-00716.1).
- 730 Neu, U., M. G. Akperov, N. Bellenbaum, and Coauthors, 2013: IMILAST: A community effort to
731 intercompare extratropical cyclone detection and tracking algorithms. *Bulletin of the American*
732 *Meteorological Society*, **94 (4)**, 529–547, <https://doi.org/10.1175/BAMS-D-11-00154.1>.
- 733 Pinto, J. G., and P. Ludwig, 2020: Extratropical cyclones over the north atlantic and western europe
734 during the last glacial maximum and implications for proxy interpretation. *Climate of the Past*,
735 **16 (2)**, 611–626, <https://doi.org/10.5194/cp-16-611-2020>.

- Priestley, M. D. K., and J. L. Catto, 2022: Future changes in the extratropical storm tracks and cyclone intensity, wind speed, and structure. *Weather and Climate Dynamics*, **3** (1), 337–360, <https://doi.org/10.5194/wcd-3-337-2022>.
- Raavi, P. H., J.-E. Chu, A. Timmermann, S.-S. Lee, and K. J. E. Walsh, 2023: Moisture control of tropical cyclones in high-resolution simulations of paleoclimate and future climate. *Nature Communications*, **14**, 6426, <https://doi.org/10.1038/s41467-023-42033-8>.
- Raddatz, T. J., and Coauthors, 2007: Will the tropical land biosphere dominate the climate–carbon cycle feedback during the twenty-first century? *Climate Dynamics*, **29** (6), 565–574, <https://doi.org/10.1007/s00382-007-0247-8>.
- Raible, C. C., M. Messmer, F. Lehner, T. F. Stocker, and R. Blender, 2018: Extratropical cyclone statistics during the last millennium and the 21st century. *Climate of the Past*, **14** (10), 1499–1514, <https://doi.org/10.5194/cp-14-1499-2018>.
- Raible, C. C., J. G. Pinto, P. Ludwig, and M. Messmer, 2021: A review of past changes in extratropical cyclones in the northern hemisphere and what can be learned for the future. *WIREs Climate Change*, **12** (3), e680, <https://doi.org/10.1002/wcc.680>.
- Shi, X., M. Werner, H. Yang, Q. Gao, J. Liu, and G. Lohmann, 2025: Precession controls on climate and water isotope signals in Northern Africa. *Paleoceanography and Paleoclimatology*, **40**, e2024PA004 999, <https://doi.org/10.1029/2024PA004999>.
- Sidorenko, D., and Coauthors, 2019: Evaluation of FESOM2.0 coupled to ECHAM6.3: preindustrial and HighResMIP simulations. *Journal of Advances in Modeling Earth Systems*, **11** (11), 3794–3815, <https://doi.org/10.1029/2019MS001696>.
- Simmonds, I., and K. Keay, 2000: Mean southern hemisphere extratropical cyclone behavior in the 40-year ncep–ncar reanalysis. *Journal of Climate*, **13** (5), 873–885, [https://doi.org/10.1175/1520-0442\(2000\)013<0873:MSHECB>2.0.CO;2](https://doi.org/10.1175/1520-0442(2000)013<0873:MSHECB>2.0.CO;2).
- Stevens, B., and Coauthors, 2013: Atmospheric component of the MPI-M Earth System Model: ECHAM6. *Journal of Advances in Modeling Earth Systems*, **5** (2), 146–172, <https://doi.org/10.1002/jame.20015>.

- 763 Ulbrich, U., G. C. Leckebusch, and J. G. Pinto, 2009: Extra-tropical cyclones in the present and fu-
764 ture climate: a review. *Theoretical and Applied Climatology*, **96** (1–2), 117–131, [https://doi.org/](https://doi.org/10.1007/s00704-008-0083-8)
765 10.1007/s00704-008-0083-8.
- 766 Varma, V., and Coauthors, 2012: Holocene evolution of the southern hemisphere westerly winds
767 in transient simulations with global climate models. *Climate of the Past*, **8** (2), 391–402,
768 <https://doi.org/10.5194/cp-8-391-2012>.
- 769 Wallace, J. M., G.-H. Lim, and M. L. Blackmon, 1988: Relationship between cyclone tracks,
770 anticyclone tracks and baroclinic waveguides. *Journal of the Atmospheric Sciences*, **45** (3),
771 439–462, [https://doi.org/10.1175/1520-0469\(1988\)045<0439:RBCTAT>2.0.CO;2](https://doi.org/10.1175/1520-0469(1988)045<0439:RBCTAT>2.0.CO;2).
- 772 Wang, Y., and Coauthors, 2008: Millennial- and orbital-scale changes in the east asian
773 monsoon over the past 224,000 years. *Nature*, **451** (7182), 1090–1093, [https://doi.org/](https://doi.org/10.1038/nature06692)
774 10.1038/nature06692.
- 775 Welch, B. L., 1947: The generalization of ‘Student’s’ problem when several different population
776 variances are involved. *Biometrika*, **34** (1–2), 28–35, <https://doi.org/10.1093/biomet/34.1-2.28>.
- 777 Zhong, Y., and Coauthors, 2024: Orbital controls on North Pacific dust flux during the
778 late Quaternary. *Geophysical Research Letters*, **51**, e2023GL106631, [https://doi.org/10.1029/](https://doi.org/10.1029/2023GL106631)
779 2023GL106631.