

**Cool summers are stormy summers: precession reorganizes extratropical
cyclone activity through the quiet season**

Xiaoxu Shi^a

^a *Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research, Bremerhaven,
Germany*

Corresponding author: Xiaoxu Shi, shixiaoxu@sml-zhuhai.cn

7 ABSTRACT: Extratropical cyclones peak in winter, while orbital precession reshapes the sea-
8 sonal insolation cycle most strongly in summer. Thus storm tracks might be expected to ignore the
9 precession cycle. This is examined with twelve equilibrium simulations using a coupled climate
10 model AWI-ESM2, spanning a full precession cycle at 30° longitude increments under fixed eccen-
11 tricity and obliquity. Extratropical cyclones are explicitly tracked based on 6-hourly atmospheric
12 pressure fields for both hemispheres. While the annual-mean cyclone response to precession is
13 muted, substantial seasonal variability emerges. In the North Atlantic, the North Pacific, and the
14 Southern Ocean the response concentrates in local summer, where storm activity ranges across
15 the phases by up to 68% of its climatological mean, while winter changes stay below about 10%.
16 We also find summers spent near perihelion feature suppressed storminess, whereas summers near
17 aphelion experience intensified cyclone activity, yielding antiphase midlatitude storm variability
18 between the two hemispheres over the precession cycle. This signal is carried by cyclone number
19 and position rather than by propagation speed, and it follows the pronounced summer response
20 of the Eady growth rate driven primarily by vertical shear in all three basins. Monthly storm
21 responses are phase-locked to the calendar timing of perihelion, lagged by 1–2 months due to
22 surface thermal inertia. Precession therefore acts on extratropical storm tracks through their quiet
23 season. Summer-sensitive proxy archives should record a strong and hemispherically antiphased
24 storminess signal at the precession period that annually integrating archives would miss.

25 SIGNIFICANCE STATEMENT: Long orbital cycles drive global climate variability, and the
26 strongest of these, precession, reshapes sunlight mainly in summer, while mid-latitude storms
27 happen mainly in winter. This creates an apparent disconnect between precessional forcing and
28 storm track variability. Climate model simulations covering a full precession cycle show that
29 the orbit nevertheless controls storminess by acting through summer. When perihelion falls in a
30 hemisphere's summer, that summer has fewer mid-latitude storms and the storm belt sits farther
31 poleward, and the two hemispheres evolve in antiphase through the precession cycle. This tells
32 researchers where to look for orbital storm signals in geological records, namely in archives
33 sensitive to summer, not to the yearly average.

34 1. Introduction

35 Extratropical cyclones organize the weather of the middle latitudes. They carry most of the
36 poleward heat and moisture transport outside the tropics, deliver a large share of midlatitude
37 precipitation, and cluster into the storm tracks of the North Atlantic, the North Pacific, and the
38 Southern Ocean (Hoskins and Valdes 1990; Chang et al. 2002; Hoskins and Hodges 2002, 2005).
39 These storms grow by baroclinic instability on the meridional temperature gradient (Eady 1949),
40 and the gradient is steepest in winter. Cyclone numbers therefore peak in the cold season in both
41 hemispheres, in the south as clearly as in the north (Simmonds and Keay 2000). Extratropical
42 cyclones are creatures of winter.

43 Precession, in contrast, is a lever that acts on summer. The slow turning of the longitude of
44 perihelion is the strongest orbital pacemaker of the seasonal insolation cycle, yet it redistributes
45 sunlight rather than adding any, and the redistribution is proportional to the sunlight a season
46 already receives (Berger 1978; Laskar et al. 2004). Midlatitude summer insolation swings by more
47 than 100 W m^{-2} across a precession cycle while midwinter insolation barely moves, which is why
48 the orbital literature treats summer energy as the climatically active quantity (Huybers 2006). A
49 winter phenomenon facing a summer lever invites a simple expectation. Precession should have
50 little to say about storm tracks. That expectation has never been put to a clean test.

51 Where the imprint of precession has been studied in depth, it appears as a seasonal reorganization
52 of climate rather than a change in the annual mean. The monsoon literature carries this tradition,
53 from the classic orbital experiments of Kutzbach and Guetter (1986) to single-forcing designs that

isolate precession explicitly (Bosmans et al. 2015; Merlis et al. 2013), and cave records show the northern and southern monsoons swinging in antiphase as perihelion migrates through the year (Wang et al. 2008). The extratropics have received far less attention. Kaspar et al. (2007) simulated a stronger and poleward-shifted North Atlantic winter storm track for the Eemian from two orbital time slices, Brayshaw et al. (2010) traced Holocene orbital forcing into the North Atlantic and Mediterranean winter storm tracks through the baroclinicity they support, and Varma et al. (2012) found the Southern Hemisphere westerlies migrating under Holocene orbital forcing. Explicit cyclone tracking has entered paleoclimate work mainly at the Last Glacial Maximum, where ice sheets dominate the forcing (Pinto and Ludwig 2020; Raible et al. 2021), and in transient simulations of the last millennium (Hess and Chemke 2025). For tropical cyclones the precessional response has recently been isolated in high-resolution paleoclimate simulations (Raavi et al. 2023). To our knowledge, no previous study has applied explicit cyclone tracking across a full precession cycle to isolate the orbital response of the extratropical storm tracks.

Closing this gap matters beyond dynamics. However, the paleoclimate records that would register past storminess, from dust and wind-blown sediment to westerly-belt precipitation, respond to particular seasons rather than to the annual mean, and a seasonally biased archive can fabricate or erase an orbital signal depending on which season carries it (Laepple et al. 2011). Knowing where in the seasonal cycle the precessional storm signal lives is a prerequisite for reading such records. Storm diagnostics raise a parallel issue, because measures of cyclone influence time, cyclone counts, and cyclone intensity respond to forcing in different ways and must be separated before any single proxy is interpreted.

In the present study the precession cycle is covered with twelve equilibrium simulations of a coupled climate model, and every extratropical cyclone is tracked explicitly in six-hourly pressure fields in both hemispheres. The design isolates precession at amplified eccentricity with all other boundary conditions fixed, so that the seasonal anatomy of the orbital storm response can be mapped basin by basin against the calendar month of perihelion.

The naive expectation fails, and it fails in an instructive way. The annual-mean storm response is indeed small, but the signal has not vanished. It is concentrated in local summer, the quiet season, in all three storm basins. Summers spent near perihelion are calm and summers spent near aphelion are stormy, so the two hemispheres respond in antiphase across the precession cycle, an

84 extratropical counterpart of the monsoon seesaw. The response travels from insolation to storms
85 through the summer Eady growth rate, and the growth-rate change is carried by vertical shear
86 (Lindzen and Farrell 1980). The paper is organized around which season carries the precessional
87 storm response, what dynamical chain transmits it, and why the winter storm track stays orbitally
88 quiet. Section 2 describes the model, the ensemble, and the tracking. Section 3 presents the
89 seasonal anatomy of the response, and section 4 discusses mechanisms, implications for proxy
90 records, and limitations.

91 **2. Data and methods**

92 *a. Model and experiments*

93 The simulations are performed with the coupled Earth system model AWI-ESM2 (Sidorenko
94 et al. 2019; Shi et al. 2022). The atmosphere component is ECHAM6 at T63 resolution with 47
95 vertical levels (Stevens et al. 2013), which includes the land surface module JSBACH (Raddatz
96 et al. 2007). The ocean and sea ice component is FESOM2, a finite-volume model formulated on
97 an unstructured mesh (Danilov et al. 2017).

98 Twelve equilibrium simulations are analyzed, taken from the idealized precession ensemble
99 introduced by Shi et al. (2025). The longitude of perihelion ω is prescribed at values from 0° to
100 330° in steps of 30° , while eccentricity is fixed at 0.058 and obliquity at 24.5° . The eccentricity
101 value sits near the upper bound reached during the Quaternary and amplifies the precessional
102 modulation of insolation, so that the response emerges clearly from internal variability. All
103 remaining boundary conditions follow the preindustrial setup, and each phase is branched from
104 a long preindustrial control. Thirty years of six-hourly output from the equilibrated part of each
105 simulation form the basis of the cyclone analysis. The calendar month of perihelion for each phase
106 is computed from two-body orbital geometry (Berger 1978), and the same formulary provides the
107 insolation fields shown in Fig. 1.

108 *b. Cyclone detection and tracking*

109 Extratropical cyclones are detected and tracked with the algorithm of Crawford et al. (2021),
110 and Fig. 2 summarizes the procedure with one six-hourly field of the ensemble. Six-hourly sea
111 level pressure is first interpolated onto 100-km equal-area grids centered on each pole, and the two

hemispheres are processed separately. A grid cell qualifies as a cyclone center when its pressure p_c lies below that of all eight neighbors and the surrounding field keeps a sufficient inward gradient. The gradient test compares p_c with the mean pressure $\bar{p}(d)$ on a one-cell ring at distance d from the candidate,

$$\frac{\bar{p}(d) - p_c}{d} \geq \gamma, \quad d = 1000 \text{ km}, \quad \gamma = 750 \text{ Pa (1000 km)}^{-1}, \quad (1)$$

and candidates over terrain above 1500 m are discarded. Centers closer than 1000 km are treated as one multicenter system. The cyclone area A is obtained by growing closed isobars around each center at a 200-Pa interval until a contour fails to close, and the last closed isobar defines the edge pressure p_e (Fig. 2b). Each cyclone carries the depth and the area-equivalent radius

$$D = p_e - p_c, \quad R = \sqrt{A/\pi}. \quad (2)$$

Tracking links the centers of consecutive six-hourly fields (Fig. 2c). For every active cyclone a first-guess position is projected from its previous displacement, damped to three quarters,

$$\mathbf{x}_q = \mathbf{x}_t + 0.75 (\mathbf{x}_t - \mathbf{x}_{t-1}), \quad (3)$$

and a center of the next field qualifies as the continuation when it lies within $v_{\max}\Delta t$ of both the current position $\mathbf{x}_t$ and the projection $\mathbf{x}_q$, with $v_{\max} = 150 \text{ km h}^{-1}$ and $\Delta t = 6 \text{ h}$. Among the qualifying candidates the one nearest to the projection continues the track. Centers left unmatched at the new time start tracks as genesis events, and tracks left unmatched end as lysis. Only systems poleward of 30° latitude enter the analysis.

Several track-based diagnostics are derived. Track density counts every six-hourly cyclone position, system density counts each cyclone once at its mean position, and genesis density counts the first point of each track. Tracks that merely cross the boundaries of the monthly processing windows would appear as spurious genesis or lysis events, and such truncated events (about 7% of the total) are removed. For every cyclone the lifespan, the maximum depth relative to the outermost closed isobar, the minimum central pressure, the mean cyclone area, and the propagation speed are recorded.

134 *c. Gridded storm coverage*

135 A complementary Eulerian diagnostic measures how long each location spends under cyclone
 136 influence. At every six-hourly time step a binary mask flags the grid cells that lie inside a square
 137 search window of half-width $\sqrt{A}$ grid cells around each cyclone center and whose pressure lies
 138 below the edge pressure of that cyclone,

$$M(\mathbf{x}, t) = 1 \quad \text{where } p(\mathbf{x}, t) < p_e \text{ inside the window,} \quad (4)$$

139 and the masks are regridded to T63 and accumulated into monthly storm days, one quarter of
 140 the flagged six-hourly steps. Storm days at a point therefore scale with the product of cyclone
 141 number and cyclone area footprint. The two factors can carry opposite seasonal cycles, and over
 142 the Southern Ocean the larger size of summer cyclones raises summertime storm days even though
 143 cyclone counts peak in winter. Frequency statements in this paper consequently rest on the track-
 144 based densities, while storm days serve as a smooth field for maps and for measures of cyclone
 145 influence time. The gridded product covers all twelve phases. An early processing inconsistency
 146 in the $\omega = 30^\circ$ mask product was traced to a stale intermediate file and removed by regenerating
 147 the storm masks of all twelve experiments with one pipeline version, and the regenerated fields
 148 were verified month by month against the Lagrangian statistics.

149 An Eulerian measure of synoptic variability provides a further check that involves no tracking
 150 at all. Bandpass sea level pressure variability is computed from the six-hourly fields with the 24-h
 151 difference filter of Wallace et al. (1988), which isolates fluctuations on the synoptic time scales that
 152 define the classical storm tracks (Blackmon 1976), and its monthly standard deviation is analyzed
 153 with the same basins, seasons, and response measures (appendix Fig. A1).

154 *d. Baroclinicity diagnostics*

155 The maximum Eady growth rate serves as the measure of baroclinicity (Eady 1949; Lindzen
 156 and Farrell 1980; Hoskins and Valdes 1990). It is computed as $\sigma = 0.31 |f| |\partial U / \partial z| N^{-1}$ in the
 157 925–700 hPa layer from monthly wind, temperature, and geopotential height. The response to
 158 precession is quantified between the end members p090 and p270, which place perihelion at the
 159 June and December solstices. Because the difference fields are dipolar, plain box averages cancel,

and the response amplitude is instead defined as the root-mean-square of the difference over a basin, normalized by the climatological growth rate of the season. A substitution decomposition in the style of Camargo et al. (2007) splits the difference field into a vertical-shear factor and a static-stability factor, and the share of each factor is obtained by projecting it onto the full difference pattern. The basin averages use the storm-track boxes defined below. This choice matters in the Southern Hemisphere, where a sector beginning at 30°S would admit the seasonal signal of the subtropical winter jet and disguise it as a storm-belt response.

e. Basins and statistics

Three storm-track basins are analyzed, the North Atlantic (30–75°N, 80°W–20°E), the North Pacific (30–75°N, 120°E–120°W), and the Southern Ocean (40–75°S, all longitudes). Basin means are area weighted. The size of the precessional response is reported as the range across the twelve phases, the maximum minus the minimum of the 30-year mean, expressed as a percentage of the phase mean. Uncertainty estimates use the interannual spread of the 30 individual years in each phase, and the significance of the end-member difference maps is assessed at each grid point with a two-sided Welch t test, the two-sample test that allows unequal variances between the samples (Welch 1947). The local-summer contrasts between extreme phases far exceed the interannual noise, with p values below 10^{-12} in all three basins.

3. Results

a. The summer paradox

The annual-mean view sets the baseline (Fig. 3). The twelve annual climatologies are close to interchangeable, and the anomalies of each phase from their common mean form weak, smoothly rotating dipoles along the storm tracks. The basin series beneath the maps quantify how little survives in the annual mean, with across-phase ranges of 10%, 17% and 4% in the North Atlantic, the North Pacific and the Southern Ocean, varying smoothly with ω and already hinting at the interhemispheric antiphase, since the northern basins reach their minima near $\omega = 90^\circ$ where the Southern Ocean peaks.

The storm tracks beneath this stable annual surface live in winter, exactly as observed. Seasonal-mean genesis counts peak in DJF over the North Atlantic and the North Pacific and in JJA over the

Southern Ocean, system counts follow the same calendar in every phase, and cyclones cut deepest in the cold season, with winter maximum depths exceeding their summer values by roughly half in the northern basins (Fig. 4a–c; Table 1). The winter identity of the storm season is therefore common to both hemispheres and to all twelve orbital configurations.

Precession barely touches the annual mean but reorganizes the seasons, and the season it selects is the quiet one. The seasonal cycles of basin-mean track density keep their winter peak in every phase and fan out across the phases almost exclusively in local summer (Fig. 4a–c). The across-phase range makes this quantitative (Fig. 4d–f). In the North Pacific the annual range is 7% of the phase mean while the JJA range reaches 44%. The North Atlantic behaves the same way with 7% against 25%. The Southern Ocean repeats the pattern half a year out of phase, with an annual range of 4% against 22% in DJF. System and genesis densities confirm the calendar in every basin, and the influence-time metric swings harder still, with a storm-day range of 68% in North Pacific summer (Table 1). The sensitive season is local summer everywhere, and the small annual signal is what survives of a substantial summer swing after the stable winter is averaged in.

A tracking-free view points the same way (appendix Fig. A1). Bandpass sea level pressure variability responds most strongly in JJA over the North Pacific, at 36% of its climatology against 9% in winter, and winter holds the weakest response of all seasons in both northern basins. This amplitude-weighted metric shifts part of the response within the warm half-year. Over the North Atlantic it peaks in early autumn, matching the secondary September lobe of the forcing itself (Fig. 1c), and over the Southern Ocean its apparent winter share lies on the equatorward flank of the basin box, while poleward of 50°S austral summer leads. The warm-season concentration and the winter quiescence are therefore not artifacts of the tracking algorithm.

b. Orbital anchoring and the interhemispheric mirror

This summertime storm response is locked to orbital geometry. When basin-averaged track-density anomalies are plotted against both calendar month and precession angle ω , the band of peak anomalies traces a diagonal ridge aligned with the calendar timing of perihelion across the full phase space (Fig. 4g–i), with a delay of 1–2 months. Summers spent near perihelion exhibit suppressed storm activity, whereas aphelion summers feature intensified storminess. A single precessional forcing thus drives opposing midlatitude storm responses between the two

217 hemispheres across the orbital cycle, as perihelion can fall in the summer of only one hemisphere
 218 at a time. The basin series make the mirror explicit (Fig. 7a). North Pacific summer activity reaches
 219 its minimum at $\omega = 90^\circ$, 31% below the phase mean where perihelion meets the June solstice, and
 220 its maximum of +37% near $\omega = 270^\circ$, while Southern Ocean summer activity peaks at $\omega = 60^\circ$
 221 and bottoms out at $\omega = 210^\circ$. The North Atlantic follows its Pacific neighbor with a swing from
 222 -16% to $+15\%$. These extreme-phase contrasts far exceed interannual variability, with Welch p
 223 values below 10^{-12} in all three basins.

224 *c. The dynamical bridge runs through summer shear*

225 The seasonal selectivity begins in the forcing itself. Averaged over the northern storm-track band
 226 from 40 to 70°N, the seasonal forcing amplitude peaks at 105 W m^{-2} in JJA and collapses to 14
 227 in DJF, and the mirror band in the south peaks at 119 W m^{-2} in DJF against 47 in JJA (Fig. 1).
 228 The reason is that the precessional modulation of insolation is multiplicative. Moving perihelion
 229 through the year changes the Sun–Earth distance of a given calendar month, which rescales the
 230 incoming flux of that month by a fraction of order $4e$, about 23% at $e = 0.058$, so the absolute
 231 swing of any month is proportional to the insolation it receives from obliquity and day length.
 232 Midwinter at these latitudes offers almost nothing to rescale. The orbital lever therefore presses
 233 on local summer in each hemisphere, keeps a substantial autumn shoulder, and leaves deep winter
 234 almost untouched.

235 Baroclinicity translates the pressure on the seasons into the storm response. The Eady growth
 236 rate difference between the end members p090 and p270 is largest in local summer within every
 237 storm basin (Fig. 5). Relative to the seasonal climatology the summer response reaches 31% in the
 238 North Atlantic, a striking 92% in the North Pacific, and 21% in the Southern Ocean, in each case
 239 the seasonal maximum for that basin. A substitution decomposition attributes the pattern almost
 240 entirely to the vertical-shear factor, whose projection accounts for 103–109% of the difference field
 241 in the three basins, while the static-stability factor contributes near zero. The chain is short and
 242 uniform. Perihelion summers warm the summer hemisphere, flatten the meridional temperature
 243 gradient, weaken the thermal-wind shear, and starve the baroclinic growth on which summer
 244 cyclones depend. The storm anomalies appear somewhat poleward of the growth-rate anomalies,
 245 consistent with the downstream development of baroclinic waves (Chang et al. 2002), and this offset

is why pattern measures rather than same-latitude box averages are used throughout. Winter tells a complementary story. The winter growth rate still shifts by 16–18% between the end members, yet the winter storm response stays below 12% in every basin, a point taken up in the discussion.

d. Spatial expression, and what precession actually moves

The end-member difference maps show where the summer signal lives (Fig. 6). In the annual mean the p090 minus p270 difference is weak and patchy. In JJA a pronounced dipole spans the northern basins, with fewer storm days through the midlatitude core and more along the Arctic flank at perihelion summer, the signature of a poleward shift joined to an overall summer weakening. In DJF the significant signal migrates to the Southern Ocean and forms a circumpolar banded structure of opposite sign. About half of the JJA grid points in the northern extratropics pass the 95% significance test, against a far smaller fraction in the annual mean.

The track displacement behind these dipoles can be made explicit (appendix Fig. A2). The storm-day-weighted latitude of the North Pacific summer track migrates from 53°N when perihelion falls in northern winter to 61°N when it falls in northern summer, a poleward excursion of 8° that peaks exactly at $\omega = 90^\circ$, and the track-density-weighted latitude follows the same curve with a 5° swing. The North Atlantic summer track moves 4.4° in the same direction, and the Southern Ocean summer track mirrors the rule with a poleward excursion of 3.8° when perihelion approaches austral summer. A perihelion summer therefore thins the storm population and displaces the survivors poleward, in step with the poleward shift of the baroclinic zone seen in the growth-rate dipole of Fig. 5a. The winter track latitude stays within 1.5–2.8° across the entire cycle, and its small extremes do not align with the solstitial phases, which extends the winter invariance from storm counts to storm position. The shoulder seasons fall in between and carry a telling phase structure, because the autumn excursion peaks near $\omega = 150^\circ$, the configuration that places perihelion in late summer, so the poleward displacement follows the perihelion month through the calendar exactly as the occurrence anomaly does in Fig. 4g–i. The annual-mean track latitude varies by no more than 1.3° in any basin.

The orbital anchoring of occurrence and position condenses into a single statement (Fig. 8). A first-harmonic fit across the phases gives, for every calendar month, the value of ω at which storm activity reaches its minimum and the track its poleward maximum, and through the sunlit half-year

both sequences climb in parallel with the line that places perihelion in that month. The response lags the perihelion calendar by a basin-dependent interval, near one month in the North Pacific, one to two months in the North Atlantic, and slightly over two months in the Southern Ocean, an ordering consistent with the growing oceanic character of the three sectors and with surface thermal inertia setting the delay. Winter months fall off the line, where the response amplitude is too small to anchor. The storm tracks are phase locked to the perihelion calendar wherever the lever can reach them.

What precession moves is above all the number and location of cyclones, not their translation speed, and only regionally their depth (Fig. 7b). Local-summer ranges of the frequency and position metrics span 16–68% across the basins. Propagation speed stays within 2–5% everywhere. Cyclone depth separates the basins, and the diagnostics are deliberately kept apart in the manner of Zappa et al. (2013). Over the Southern Ocean the maximum depth varies by only 4% across the phases while frequency varies by 22%, a clean decoupling of frequency from intensity. Over the North Pacific the summer depth range reaches 33% and the lifespan range 23%, so there the whole cyclone population weakens together when perihelion falls in summer, fewer systems that are also shallower and shorter lived. The North Atlantic sits between these regimes. The common element across basins is that occurrence responds most, while the intensity response ranges from absent to substantial depending on how strongly the summer baroclinic reservoir is drained.

e. What the rain records

The hydrological expression of the cycle runs opposite to the dynamical one (appendix Fig. A3). Detected cyclones deliver about one third of extratropical precipitation in the ensemble, in line with reanalysis-based attribution (Hawcroft et al. 2012). In North Pacific summer the precipitation falling inside each storm strengthens by up to 15% exactly where storm counts collapse, and the two series are nearly perfectly anticorrelated across the phases ($r = -0.95$). A perihelion summer holds a warmer and moister atmosphere, so the surviving storms rain harder, but the loss of storms wins, and total summer precipitation still falls by 17% between the end members. The share of precipitation delivered by storms falls with it, by 9.5 percentage points across the cycle in North Pacific summer against about 3 in the annual mean. Precession therefore forces extratropical

hydroclimate in two opposing ways at once, fewer storms against wetter storms, and the two must be kept apart in interpretation just as counts and depths must.

4. Discussion

a. Why the lever lands in summer

The seasonal selectivity of the storm response is set by orbital geometry before any model physics enters. Precession redistributes insolation in proportion to the insolation a season already has, so the winter hemisphere offers the lever almost nothing to press on (Berger 1978). The monthly forcing amplitude in Fig. 1 makes the floor explicit, with midwinter values near 20 W m^{-2} against summer values above 125 W m^{-2} at the latitudes of the storm tracks. This part of the result is model independent and would reappear in any climate model, because it restates the geometry that also underlies the summer-energy view of orbital forcing (Huybers 2006). The novelty is not that summer receives the forcing, which the monsoon literature established long ago (Kutzbach and Guetter 1986; Bosmans et al. 2015), but that the summer forcing finds a winter phenomenon and reorganizes it through its quiet season.

b. Cool summers are stormy summers

The mechanism that converts summer insolation into summer storms is the thermal wind. Perihelion summers warm the summer hemisphere most strongly over land and at high latitudes, flatten the meridional temperature gradient, and weaken the vertical shear that feeds baroclinic growth. The factor decomposition confirms that the shear term carries essentially the whole Eady growth rate response in all three basins, while static stability contributes almost nothing. The same chain, with greenhouse warming in place of perihelion, is now observed in the present-day Northern Hemisphere, where amplified summer warming at high latitudes has weakened the summer circulation, the eddy kinetic energy, and the number of strong summer cyclones (Coumou et al. 2015; Chang et al. 2016; Gertler and O’Gorman 2019). Projections extend the analogy, with CMIP-class models thinning summer cyclone activity and the associated baroclinicity through the same shear pathway (Lehmann et al. 2014; Priestley and Catto 2022). The correspondence should be read as a shared mechanism rather than a shared scenario, because the warming trend alters the

330 two hemispheres together whereas precession swings them in antiphase. Under both forcings the
331 rule is the same, and it is the rule in the title. Cool summers are stormy summers.

332 *c. The orbital invariance of the winter storm track*

333 The stability of winter deserves as much attention as the mobility of summer. Winter storm
334 activity varies by less than about a tenth across the entire precession cycle in every basin, and this
335 is only partly a matter of weak winter forcing. The winter Eady growth rate still shifts by 16–18%
336 between the end members, yet winter cyclone numbers barely respond. The winter storm track
337 appears saturated in the sense that ample baroclinicity is available in every phase, so that moderate
338 changes in the growth rate do not translate into changes in occurrence. Whatever its ultimate cause,
339 the practical consequence is notable. The delivery of winter precipitation to the midlatitude and
340 subpolar continents, the season and pathway that build snowpack and feed ice sheets, is essentially
341 fixed across orbital configurations of the precession cycle. A transient simulation of the last
342 millennium likewise found cyclone counts nearly insensitive to external forcing (Raible et al. 2018;
343 Hess and Chemke 2025), and the present result extends that insensitivity to the far larger insolation
344 excursions of the orbital band, for winter alone.

345 *d. One lever, three basins, and what the records would see*

346 The three basins express one mechanism with different accents, and the diagnostics must be
347 kept apart to see it. Occurrence responds everywhere, with local-summer ranges of 16–68% in
348 frequency and position metrics, while propagation speed is inert. Intensity separates the basins.
349 The Southern Ocean decouples cleanly, with a 4% depth range against a 22% frequency range,
350 the North Pacific weakens as a whole population in perihelion summers, and the North Atlantic
351 lies between. The pattern echoes the long-standing conclusion from anthropogenic scenarios that
352 cyclone number and cyclone intensity respond to forcing through different processes and need not
353 move together (Bengtsson et al. 2009; Ulbrich et al. 2009; Catto et al. 2019). The water cycle
354 sharpens the contrast, because per-storm precipitation strengthens in the very summers that lose
355 storms, an orbital counterpart of the warming-scenario finding that hydrological responses outrun
356 dynamical ones (Bengtsson et al. 2009). For the record this matters twice over. Extratropical
357 precipitation is delivered disproportionately by cyclones and their fronts (Hawcroft et al. 2012;

358 Catto and Pfahl 2013), and the storm-attributed share of precipitation swings by up to ten percentage
359 points across the cycle in North Pacific summer, so a precipitation archive senses the competition
360 between fewer storms and wetter storms rather than either alone. The storm-day diagnostic adds
361 its own caution, because cyclone influence time folds cyclone size into cyclone number, and over
362 the Southern Ocean the seasonal cycle of size alone can invert the apparent storm season. Any
363 proxy that integrates storm influence rather than storm counts inherits this ambiguity.

364 For the paleoclimate record the seasonal anatomy carries a concrete and testable message. An
365 archive with summer-biased sensitivity in the midlatitude storm belts should carry a pronounced
366 precession signal in storminess, while an annually integrating archive should carry almost none.
367 The seasonal bias of an archive can therefore fabricate or erase an orbital storm signal exactly as it
368 does for temperature (Laepple et al. 2011). The interhemispheric mirror sharpens the prediction,
369 because summer-sensitive storminess records from the two hemispheres should vary in antiphase
370 at the precession period, the extratropical counterpart of the monsoon seesaw (Wang et al. 2008).
371 Existing archives make the test realistic. Precession-band signals are already established in the
372 northern midlatitude westerlies (Li et al. 2019) and in the composition of North Pacific dust, where
373 the precession imprint sits in the moisture-sensitive component while the annually integrating
374 flux follows obliquity (Zhong et al. 2024), and a winter-integrating dust-strength record shows
375 correspondingly weak precession power (Chen et al. 2014). In the south the westerly belt carries
376 strong 19 and 23 kyr cycles at its subtropical edge (Lamy et al. 2019), with the orbital band
377 recorded at a site depending on its latitude within the belt (Kaiser et al. 2024). The subtropical
378 location of that southern signal is consistent with the present results, because the winter half of
379 the precessional baroclinicity response resides in the subtropical jet equatorward of the storm belt,
380 while the storm-belt response is a summer phenomenon. Records of the southern westerlies also
381 show orbital-band migrations of the belt itself (Lamy et al. 2010; Varma et al. 2012), and glacial-
382 state cyclone modeling has shown how strongly reconstructed storminess depends on which cyclone
383 property a proxy senses (Pinto and Ludwig 2020). A paired-hemisphere test at the precession band
384 is within reach of existing archives.

385 *e. Limitations*

386 Several limitations bound these conclusions. The results rest on one model, one tracking algo-
387 rithm, and an idealized orbital configuration with eccentricity fixed near its Quaternary maximum,
388 so the simulated amplitudes represent an upper bound that scales with $e \sin \omega$ rather than a recon-
389 struction of any particular epoch. Obliquity is fixed at 24.5° , and interactions between precession
390 and obliquity or ice volume are outside the design. The simulations are equilibrium slices, so
391 transient effects along the precession cycle are not represented. The T63 atmosphere resolves
392 the storm tracks but not mesoscale substructure, and low-level Eady growth rates in the Southern
393 Hemisphere carry a documented sensitivity to boundary-layer stability (Simmonds and Lim 2009).
394 The gridded storm-day product required one uniform regeneration of the storm masks before all
395 twelve phases could be used together, a reminder that multi-batch processing of tracked-cyclone
396 products deserves explicit version control.

397 **5. Conclusions**

398 One rule spans the three storm tracks of this precession ensemble. The orbital lever reaches
399 extratropical cyclones through local summer, the quiet season, while the winter storm track stays
400 essentially fixed across the entire cycle. Summers spent near perihelion are calm and summers
401 near aphelion are stormy, and because one hemisphere's perihelion summer is the other's aphelion
402 summer, midlatitude storminess in the two hemispheres swings in antiphase at the precession
403 period. The signal travels through shear-driven summer baroclinicity and expresses itself chiefly
404 in how many cyclones occur and where they run, with per-cyclone intensity responding only where
405 the summer baroclinic reservoir is drained most deeply. Both expressions are phase locked to
406 the calendar month of perihelion, with a delay of one to two months that grows with the oceanic
407 character of the basin. For the proxy record the lesson is direct. Orbital storminess signals should
408 be sought in summer-sensitive midlatitude archives, and they should appear with opposite sign in
409 the two hemispheres.






FIG. 1. Precessional insolation forcing from orbital geometry with eccentricity 0.058 and obliquity 24.5° .
 (a) Monthly insolation anomaly relative to the phase mean, averaged over the storm-track band $40\text{--}70^\circ\text{N}$, as
 a function of calendar month and precession phase ω , with the perihelion (red) and aphelion (blue) calendar
 months of each phase overlaid. (b) As in (a), but for $40\text{--}70^\circ\text{S}$. (c) Seasonal forcing amplitude in the two bands,
 the seasonal mean of the monthly maximum minus minimum insolation across the twelve phases. Units: W m^{-2} .


(a) centers + area radii, one 6-h field



(b) outermost closed isobar p_e , mask cells



(c) tracks of one month ($n=407$)



(d)




FIG. 2. Cyclone detection and tracking procedure. (a) One six-hourly sea level pressure field of phase p000 (contours, 8-hPa interval) with all detected cyclone centers poleward of 30°N (dots) and their area-equivalent radii $R = \sqrt{A/\pi}$ (circles), 15 January of one model year. (b) The deepest system of that field, with the outermost closed isobar p_e (dashed), the cyclone center (cross), the central and edge pressures and depth annotated, and the binary mask cells (shading) defined by $\text{SLP} < p_e$ inside the $\sqrt{A}$ search window; contour interval 4 hPa. (c) All cyclone tracks of the same month (lines) with genesis points (dots). (d) Processing chain with the key thresholds of Eqs. (1)–(4).




FIG. 3. Annual-mean storm days across the precession cycle. (a)–(l) Annual storm-day anomaly of each phase relative to the twelve-phase mean (shading), with the phase-mean climatology outlined at 5 and 8 d month⁻¹ (contours). (m)–(o) Basin-mean annual storm days against ω with interannual standard errors; the across-phase range as a percentage of the phase mean is annotated in each panel.




FIG. 4. Seasonal concentration of the precessional storm response, from the Lagrangian tracks. (a)–(c) Basin-mean track density (cell⁻¹ month⁻¹) by calendar month for the North Atlantic, the North Pacific, and the Southern Ocean, one line per phase colored by ω , with local summer shaded. (d)–(f) Across-phase range of track density (maximum minus minimum, percent of the phase mean) for the annual mean and the four seasons, with the local-summer bar highlighted; open symbols give the corresponding ranges of system and genesis densities. (g)–(i) Basin-mean track-density anomaly relative to the phase mean as a function of calendar month and precession phase (unsmoothed), with the perihelion (red) and aphelion (blue) calendar months of each phase overlaid.




FIG. 5. Precessional response of the maximum Eady growth rate (925–700 hPa). (a) p090 minus p270 difference for JJA and (b) for DJF (day^{-1}). (c) Seasonal response amplitude in each basin, the RMS of the p090 minus p270 difference over the storm box as a percentage of the seasonal climatology, with stars marking local summer. (d) Projection shares of the vertical-shear and static-stability factors in the local-summer difference pattern.




FIG. 6. Difference in storm days (d month^{-1}) between p090 and p270 for (a) the annual mean, (b) JJA, and (c) DJF, on a common color scale. Stippling marks grid points where the difference is significant at the 95% level based on a two-sided Welch t test over the 30 simulated years.




FIG. 7. (a) Local-summer storm activity across the precession cycle, shown as the percent anomaly of basin-mean storm days from the phase mean (solid, with interannual standard errors) and of track density (dashed), for the North Atlantic and the North Pacific in JJA and for the Southern Ocean in DJF; dotted vertical lines mark the phases with perihelion in NH and SH summer. (b) Local-summer across-phase range for frequency and position metrics and for intensity and lifecycle metrics in each basin.




FIG. 8. Phase locking of the storm response to the perihelion calendar. For each calendar month and basin, the phase of the first harmonic across the twelve precession phases of (a) basin-mean storm days, shown as the ω of the activity minimum, and (b) the storm-day-weighted track latitude, shown as the ω of the poleward maximum. Marker size scales with the harmonic amplitude. The red line gives the ω that places perihelion in that calendar month, and dashed lines repeat it delayed by each basin's median lag over the amplitude-strong months (annotated in months).

TABLE 1. Three-basin comparison of the precessional storm response. Ranges are max minus min across the twelve phases as a percentage of the phase mean. Storm and sensitive seasons refer to local winter and local summer. The storm season is diagnosed from cyclogenesis counts, the sensitive season from the seasonal storm-day range. The EGR response is the RMS p090–p270 difference over the basin sector relative to its climatology, and the shear share is the projection of the vertical-shear factor onto the response pattern.

Basin	Storm season	Sensitive season	Storm-day range ann/summer (%)	Summer max/min ω (°)	EGR response summer (%)	Shear share (%)
North Atlantic	DJF	JJA	10 / 32	240 / 90	31	107
North Pacific	DJF	JJA	17 / 68	270 / 90	92	106
Southern Ocean	JJA	DJF	4 / 16	60 / 210	21	109




FIG. A1. Tracking-free Eulerian storm-track metric, the monthly standard deviation of 24-h differences of six-hourly sea level pressure. (a)–(c) Relative seasonal response in each basin, the RMS of the p090 minus p270 difference over the storm box as a percentage of the seasonal climatology (as in Fig. 5c), with the local-summer bar highlighted. (d)–(f) The p090 minus p270 difference of the same quantity for the annual mean, JJA, and DJF (hPa).

Acknowledgments. The simulations were performed at the German Climate Computing Center (DKRZ). [TODO funding and grant numbers before submission.]

Data availability statement. The AWI-ESM simulations of the idealized precession ensemble are described in Shi et al. (2025). The cyclone-track database, the gridded storm statistics, and the analysis scripts underlying all figures will be archived and assigned a DOI upon acceptance. [TODO repository DOI.]

APPENDIX

Tracking-independent, positional, and hydrological diagnostics

Figures A1–A3 present the Eulerian bandpass diagnostic, the seasonal track-latitude analysis, and the storm-attributed precipitation analysis discussed in sections 2 and 3.




FIG. A2. Storm-track latitude across the precession cycle, the storm-day-weighted mean latitude over each basin sector shown as the poleward anomaly from each season's phase mean, for (a) the North Atlantic, (b) the North Pacific, and (c) the Southern Ocean. Lines give the annual mean and the four seasons, with the local summer of each basin drawn heavy with markers. Vertical dotted lines mark the phase with perihelion in the local summer of each basin.




FIG. A3. Storm-attributed precipitation across the precession cycle. (a) North Pacific JJA percent anomalies from the phase mean of basin-mean storm days, per-storm precipitation, and total precipitation; the dotted line marks the phase with perihelion in NH summer. (b) Across-phase range of the storm-attributed share of precipitation, accumulated precipitation inside storm footprints divided by total accumulated precipitation (percentage points), for the annual mean and local summer in each basin.

References

- Bengtsson, L., K. I. Hodges, and N. Keenlyside, 2009: Will extratropical storms intensify in a warmer climate? *Journal of Climate*, **22** (9), 2276–2301, <https://doi.org/10.1175/2008JCLI2678.1>.
- Berger, A. L., 1978: Long-term variations of daily insolation and quaternary climatic changes. *Journal of the Atmospheric Sciences*, **35** (12), 2362–2367, [https://doi.org/10.1175/1520-0469\(1978\)035\(2362:LTVODI\)2.0.CO;2](https://doi.org/10.1175/1520-0469(1978)035(2362:LTVODI)2.0.CO;2).
- Blackmon, M. L., 1976: A climatological spectral study of the 500 mb geopotential height of the Northern Hemisphere. *Journal of the Atmospheric Sciences*, **33** (8), 1607–1623, [https://doi.org/10.1175/1520-0469\(1976\)033\(1607:ACSSOT\)2.0.CO;2](https://doi.org/10.1175/1520-0469(1976)033(1607:ACSSOT)2.0.CO;2).
- Bosmans, J. H. C., S. S. Drijfhout, E. Tuenter, F. J. Hilgen, L. J. Lourens, and E. J. Rohling, 2015: Response of the north african summer monsoon to precession and obliquity forcings in the earth gcm. *Climate Dynamics*, **44** (1–2), 279–297, <https://doi.org/10.1007/s00382-014-2260-z>.
- Brayshaw, D. J., B. Hoskins, and E. Black, 2010: Some physical drivers of changes in the winter storm tracks over the North Atlantic and Mediterranean during the Holocene. *Philosophical Transactions of the Royal Society A*, **368** (1931), 5185–5223, <https://doi.org/10.1098/rsta.2010.0180>.
- Camargo, S. J., K. A. Emanuel, and A. H. Sobel, 2007: Use of a genesis potential index to diagnose ENSO effects on tropical cyclone genesis. *Journal of Climate*, **20** (19), 4819–4834, <https://doi.org/10.1175/JCLI4282.1>.
- Catto, J. L., and S. Pfahl, 2013: The importance of fronts for extreme precipitation. *Journal of Geophysical Research: Atmospheres*, **118** (19), 10 791–10 801, <https://doi.org/10.1002/jgrd.50852>.
- Catto, J. L., and Coauthors, 2019: The future of midlatitude cyclones. *Current Climate Change Reports*, **5**, 407–420, <https://doi.org/10.1007/s40641-019-00149-4>.
- Chang, E. K. M., S. Lee, and K. L. Swanson, 2002: Storm track dynamics. *Journal of Climate*, **15** (16), 2163–2183, [https://doi.org/10.1175/1520-0442\(2002\)015\(02163:STD\)2.0.CO;2](https://doi.org/10.1175/1520-0442(2002)015(02163:STD)2.0.CO;2).

571 Chang, E. K. M., C.-T. Ma, C. Zheng, and A. M. W. Yau, 2016: Observed and projected
 572 decrease of Northern Hemisphere extratropical cyclone activity in summer and its impacts
 573 on maximum temperature. *Geophysical Research Letters*, **43** (5), 2200–2208, <https://doi.org/10.1002/2016GL068172>.
 574

575 Chen, J., X. Liu, and V. A. Kravchinsky, 2014: Response of the high-resolution Chinese loess grain
 576 size record to the 50°N integrated winter insolation during the last 500,000 years. *Geophysical*
 577 *Research Letters*, **41** (17), 6113–6120, <https://doi.org/10.1002/2014GL060239>.

578 Coumou, D., J. Lehmann, and J. Beckmann, 2015: The weakening summer circulation in the
 579 Northern Hemisphere mid-latitudes. *Science*, **348** (6232), 324–327, [https://doi.org/10.1126/](https://doi.org/10.1126/science.1261768)
 580 [science.1261768](https://doi.org/10.1126/science.1261768).

581 Crawford, A. D., E. A. P. Schreiber, N. Sommer, M. C. Serreze, J. C. Stroeve, and D. G. Barber,
 582 2021: Sensitivity of Northern Hemisphere cyclone detection and tracking results to fine spatial
 583 and temporal resolution. *Monthly Weather Review*, **149** (8), 2581–2598, [https://doi.org/10.1175/](https://doi.org/10.1175/MWR-D-20-0417.1)
 584 [MWR-D-20-0417.1](https://doi.org/10.1175/MWR-D-20-0417.1).

585 Danilov, S., D. Sidorenko, Q. Wang, and T. Jung, 2017: The Finite-volume Sea ice–Ocean
 586 Model (FESOM2). *Geoscientific Model Development*, **10** (2), 765–789, [https://doi.org/10.5194/](https://doi.org/10.5194/gmd-10-765-2017)
 587 [gmd-10-765-2017](https://doi.org/10.5194/gmd-10-765-2017).

588 Eady, E. T., 1949: Long waves and cyclone waves. *Tellus*, **1** (3), 33–52, [https://doi.org/10.3402/](https://doi.org/10.3402/tellusa.v1i3.8507)
 589 [tellusa.v1i3.8507](https://doi.org/10.3402/tellusa.v1i3.8507).

590 Gertler, C. G., and P. A. O’Gorman, 2019: Changing available energy for extratropical cyclones and
 591 associated convection in Northern Hemisphere summer. *Proceedings of the National Academy*
 592 *of Sciences*, **116** (10), 4105–4110, <https://doi.org/10.1073/pnas.1812312116>.

593 Hawcroft, M. K., L. C. Shaffrey, K. I. Hodges, and H. F. Dacre, 2012: How much Northern
 594 Hemisphere precipitation is associated with extratropical cyclones? *Geophysical Research*
 595 *Letters*, **39**, L24 809, <https://doi.org/10.1029/2012GL053866>.

596 Hess, O., and R. Chemke, 2025: Forced changes in the North Atlantic storm track over the
 597 last millennium. *Geophysical Research Letters*, **52**, e2025GL115 894, [https://doi.org/10.1029/](https://doi.org/10.1029/2025GL115894)
 598 [2025GL115894](https://doi.org/10.1029/2025GL115894).

- 599 Hoskins, B. J., and K. I. Hodges, 2002: New perspectives on the Northern Hemisphere winter
600 storm tracks. *Journal of the Atmospheric Sciences*, **59** (6), 1041–1061, [https://doi.org/10.1175/
601 1520-0469\(2002\)059<1041:NPOTNH>2.0.CO;2](https://doi.org/10.1175/1520-0469(2002)059<1041:NPOTNH>2.0.CO;2).
- 602 Hoskins, B. J., and K. I. Hodges, 2005: A new perspective on Southern Hemisphere storm tracks.
603 *Journal of Climate*, **18** (20), 4108–4129, <https://doi.org/10.1175/JCLI3570.1>.
- 604 Hoskins, B. J., and P. J. Valdes, 1990: On the existence of storm-tracks. *Journal of the Atmospheric
605 Sciences*, **47** (15), 1854–1864, [https://doi.org/10.1175/1520-0469\(1990\)047<1854:OTEOST>2.
606 0.CO;2](https://doi.org/10.1175/1520-0469(1990)047<1854:OTEOST>2.0.CO;2).
- 607 Huybers, P., 2006: Early pleistocene glacial cycles and the integrated summer insolation forcing.
608 *Science*, **313** (5786), 508–511, <https://doi.org/10.1126/science.1125249>.
- 609 Kaiser, J., E. Schefuß, J. A. Collins, and Coauthors, 2024: Orbital modulation of subtropical versus
610 subantarctic moisture sources in the southeast Pacific mid-latitudes. *Nature Communications*,
611 **15**, 7512, <https://doi.org/10.1038/s41467-024-51985-4>.
- 612 Kaspar, F., T. Spanghehl, and U. Cubasch, 2007: Northern hemisphere winter storm tracks of
613 the eemian interglacial and the last glacial inception. *Climate of the Past*, **3** (2), 181–192,
614 <https://doi.org/10.5194/cp-3-181-2007>.
- 615 Kutzbach, J. E., and P. J. Guetter, 1986: The influence of changing orbital parameters and surface
616 boundary conditions on climate simulations for the past 18,000 years. *Journal of the Atmospheric
617 Sciences*, **43** (16), 1726–1759, [https://doi.org/10.1175/1520-0469\(1986\)043<1726:TIOCOP>2.
618 0.CO;2](https://doi.org/10.1175/1520-0469(1986)043<1726:TIOCOP>2.0.CO;2).
- 619 Laepple, T., M. Werner, and G. Lohmann, 2011: Synchronicity of antarctic temperatures and
620 local solar insolation on orbital timescales. *Nature*, **471** (7336), 91–94, [https://doi.org/10.1038/
621 nature09825](https://doi.org/10.1038/nature09825).
- 622 Lamy, F., R. Kilian, H. W. Arz, J.-P. Francois, J. Kaiser, M. Prange, and T. Steinke, 2010: Holocene
623 changes in the position and intensity of the southern westerly wind belt. *Nature Geoscience*,
624 **3** (10), 695–699, <https://doi.org/10.1038/ngeo959>.

- 625 Lamy, F., and Coauthors, 2019: Precession modulation of the South Pacific westerly wind belt
626 over the past million years. *Proceedings of the National Academy of Sciences*, **116** (47), 23 455–
627 23 460, <https://doi.org/10.1073/pnas.1905847116>.
- 628 Laskar, J., P. Robutel, F. Joutel, M. Gastineau, A. C. M. Correia, and B. Levrard, 2004: A long-
629 term numerical solution for the insolation quantities of the earth. *Astronomy and Astrophysics*,
630 **428** (1), 261–285, <https://doi.org/10.1051/0004-6361:20041335>.
- 631 Lehmann, J., D. Coumou, K. Frieler, A. V. Eliseev, and A. Levermann, 2014: Future changes
632 in extratropical storm tracks and baroclinicity under climate change. *Environmental Research*
633 *Letters*, **9** (8), 084 002, <https://doi.org/10.1088/1748-9326/9/8/084002>.
- 634 Li, Y., Y. Song, Q. Yin, and Coauthors, 2019: Orbital and millennial northern mid-latitude
635 westerlies over the last glacial period. *Climate Dynamics*, **53** (5–6), 3315–3324, [https://doi.org/](https://doi.org/10.1007/s00382-019-04704-5)
636 [10.1007/s00382-019-04704-5](https://doi.org/10.1007/s00382-019-04704-5).
- 637 Lindzen, R. S., and B. Farrell, 1980: A simple approximate result for the maximum growth rate of
638 baroclinic instabilities. *Journal of the Atmospheric Sciences*, **37** (7), 1648–1654, [https://doi.org/](https://doi.org/10.1175/1520-0469(1980)037<1648:ASARFT>2.0.CO;2)
639 [10.1175/1520-0469\(1980\)037<1648:ASARFT>2.0.CO;2](https://doi.org/10.1175/1520-0469(1980)037<1648:ASARFT>2.0.CO;2).
- 640 Merlis, T. M., T. Schneider, S. Bordoni, and I. Eisenman, 2013: Hadley circulation response to
641 orbital precession. part i: Aquaplanets. *Journal of Climate*, **26** (3), 740–753, [https://doi.org/](https://doi.org/10.1175/JCLI-D-11-00716.1)
642 [10.1175/JCLI-D-11-00716.1](https://doi.org/10.1175/JCLI-D-11-00716.1).
- 643 Pinto, J. G., and P. Ludwig, 2020: Extratropical cyclones over the north atlantic and western europe
644 during the last glacial maximum and implications for proxy interpretation. *Climate of the Past*,
645 **16** (2), 611–626, <https://doi.org/10.5194/cp-16-611-2020>.
- 646 Priestley, M. D. K., and J. L. Catto, 2022: Future changes in the extratropical storm tracks and
647 cyclone intensity, wind speed, and structure. *Weather and Climate Dynamics*, **3** (2), 337–360,
648 <https://doi.org/10.5194/wcd-3-337-2022>.
- 649 Raavi, P. H., J.-E. Chu, A. Timmermann, S.-S. Lee, and K. J. E. Walsh, 2023: Moisture control
650 of tropical cyclones in high-resolution simulations of paleoclimate and future climate. *Nature*
651 *Communications*, **14**, 6426, <https://doi.org/10.1038/s41467-023-42033-8>.

- 652 Raddatz, T. J., and Coauthors, 2007: Will the tropical land biosphere dominate the climate–
653 carbon cycle feedback during the twenty-first century? *Climate Dynamics*, **29** (6), 565–574,
654 <https://doi.org/10.1007/s00382-007-0247-8>.
- 655 Raible, C. C., M. Messmer, F. Lehner, T. F. Stocker, and R. Blender, 2018: Extratropical cyclone
656 statistics during the last millennium and the 21st century. *Climate of the Past*, **14** (10), 1499–1514,
657 <https://doi.org/10.5194/cp-14-1499-2018>.
- 658 Raible, C. C., J. G. Pinto, P. Ludwig, and M. Messmer, 2021: A review of past changes in
659 extratropical cyclones in the northern hemisphere and what can be learned for the future. *WIREs*
660 *Climate Change*, **12** (3), e680, <https://doi.org/10.1002/wcc.680>.
- 661 Shi, X., M. Werner, Q. Wang, H. Yang, and G. Lohmann, 2022: Simulated mid-Holocene and last
662 interglacial climate using two generations of AWI-ESM. *Journal of Climate*, **35** (23), 7811–7831,
663 <https://doi.org/10.1175/JCLI-D-22-0354.1>.
- 664 Shi, X., M. Werner, H. Yang, Q. Gao, J. Liu, and G. Lohmann, 2025: Precession controls on
665 climate and water isotope signals in Northern Africa. *Paleoceanography and Paleoclimatology*,
666 **40**, e2024PA004 999, <https://doi.org/10.1029/2024PA004999>.
- 667 Sidorenko, D., and Coauthors, 2019: Evaluation of FESOM2.0 coupled to ECHAM6.3: preindus-
668 trial and HighResMIP simulations. *Journal of Advances in Modeling Earth Systems*, **11** (11),
669 3794–3815, <https://doi.org/10.1029/2019MS001696>.
- 670 Simmonds, I., and K. Keay, 2000: Mean southern hemisphere extratropical cyclone behavior in
671 the 40-year ncep–ncar reanalysis. *Journal of Climate*, **13** (5), 873–885, [https://doi.org/10.1175/1520-0442\(2000\)013<0873:MSHECB>2.0.CO;2](https://doi.org/10.1175/1520-0442(2000)013<0873:MSHECB>2.0.CO;2).
- 673 Simmonds, I., and E.-P. Lim, 2009: Biases in the calculation of Southern Hemisphere mean
674 baroclinic eddy growth rate. *Geophysical Research Letters*, **36** (1), L01 707, <https://doi.org/10.1029/2008GL036320>.
- 676 Stevens, B., and Coauthors, 2013: Atmospheric component of the MPI-M Earth System Model:
677 ECHAM6. *Journal of Advances in Modeling Earth Systems*, **5** (2), 146–172, <https://doi.org/10.1002/jame.20015>.

- 679 Ulbrich, U., G. C. Leckebusch, and J. G. Pinto, 2009: Extra-tropical cyclones in the present and fu-
680 ture climate: a review. *Theoretical and Applied Climatology*, **96** (1–2), 117–131, [https://doi.org/](https://doi.org/10.1007/s00704-008-0083-8)
681 10.1007/s00704-008-0083-8.
- 682 Varma, V., and Coauthors, 2012: Holocene evolution of the southern hemisphere westerly winds
683 in transient simulations with global climate models. *Climate of the Past*, **8** (2), 391–402,
684 <https://doi.org/10.5194/cp-8-391-2012>.
- 685 Wallace, J. M., G.-H. Lim, and M. L. Blackmon, 1988: Relationship between cyclone tracks,
686 anticyclone tracks and baroclinic waveguides. *Journal of the Atmospheric Sciences*, **45** (3),
687 439–462, [https://doi.org/10.1175/1520-0469\(1988\)045<0439:RBCTAT>2.0.CO;2](https://doi.org/10.1175/1520-0469(1988)045<0439:RBCTAT>2.0.CO;2).
- 688 Wang, Y., and Coauthors, 2008: Millennial- and orbital-scale changes in the east asian
689 monsoon over the past 224,000 years. *Nature*, **451** (7182), 1090–1093, [https://doi.org/](https://doi.org/10.1038/nature06692)
690 10.1038/nature06692.
- 691 Welch, B. L., 1947: The generalization of ‘Student’s’ problem when several different population
692 variances are involved. *Biometrika*, **34** (1–2), 28–35, <https://doi.org/10.1093/biomet/34.1-2.28>.
- 693 Zappa, G., L. C. Shaffrey, K. I. Hodges, P. G. Sansom, and D. B. Stephenson, 2013: A mul-
694 timodel assessment of future projections of North Atlantic and European extratropical cy-
695 clones in the CMIP5 climate models. *Journal of Climate*, **26** (16), 5846–5862, [https://doi.org/](https://doi.org/10.1175/JCLI-D-12-00573.1)
696 10.1175/JCLI-D-12-00573.1.
- 697 Zhong, Y., and Coauthors, 2024: Orbital controls on North Pacific dust flux during the
698 late Quaternary. *Geophysical Research Letters*, **51**, e2023GL106631, [https://doi.org/10.1029/](https://doi.org/10.1029/2023GL106631)
699 2023GL106631.