

**Cool summers are stormy summers: precession reorganizes extratropical  
cyclone activity through the quiet season**

Xiaoxu Shi<sup>a</sup>

<sup>a</sup> *Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research, Bremerhaven,  
Germany*

*Corresponding author: Xiaoxu Shi, shixiaoxu@sml-zhuhai.cn*

7 ABSTRACT: Extratropical cyclones cluster in winter, while precession reshapes the seasonal  
8 insolation cycle most strongly in summer, so storm tracks might be expected to ignore the precession  
9 cycle. This expectation is tested with twelve equilibrium simulations of the coupled climate  
10 model AWI-ESM-2.1 that cover the precession cycle at intervals of thirty degrees under amplified  
11 eccentricity and fixed obliquity, with extratropical cyclones tracked explicitly in six-hourly pressure  
12 fields in both hemispheres. The annual-mean storm response is small. The seasonal response is  
13 not. In the North Atlantic, the North Pacific, and the Southern Ocean the response concentrates in  
14 local summer, where storm activity ranges across the phases by up to two thirds of its mean while  
15 winter changes stay below about a tenth. We find that summers spent near perihelion are calm  
16 and summers spent near aphelion are stormy, so midlatitude storminess in the two hemispheres  
17 varies in antiphase across the precession cycle. The signal is carried by cyclone number and  
18 position rather than by propagation speed, and it follows the pronounced summer response of the  
19 Eady growth rate, which is dominated by vertical shear in all three basins. Month by month,  
20 the response is phase locked to the calendar date of perihelion, delayed by one to two months of  
21 surface thermal inertia. Precession therefore acts on extratropical storm tracks through their quiet  
22 season. Summer-sensitive proxy archives should record a strong and hemispherically antiphased  
23 storminess signal at the precession period that annually integrating archives would miss.

24 SIGNIFICANCE STATEMENT: Ice-age cycles are paced by slow changes in Earth's orbit, and  
25 the strongest of these, precession, reshapes sunlight mainly in summer, while midlatitude storms  
26 happen mainly in winter. Climate model simulations covering a full precession cycle show that  
27 the orbit nevertheless controls storminess, by acting through summer. When perihelion falls in a  
28 hemisphere's summer, that summer has fewer storms and the storm belt sits farther poleward, and  
29 the two hemispheres respond in opposite phase. This tells researchers where to look for orbital  
30 storm signals in geological records, namely in archives sensitive to summer, not to the yearly  
31 average.

## 32 1. Introduction

33 Extratropical cyclones organize the weather of the middle latitudes. They carry most of the  
34 poleward heat and moisture transport outside the tropics, deliver a large share of midlatitude  
35 precipitation, and cluster into the storm tracks of the North Atlantic, the North Pacific, and the  
36 Southern Ocean (Hoskins and Valdes 1990; Chang et al. 2002; Hoskins and Hodges 2002, 2005).  
37 These storms grow by baroclinic instability on the meridional temperature gradient (Eady 1949),  
38 and the gradient is steepest in winter. Cyclone numbers therefore peak in the cold season in both  
39 hemispheres, in the south as clearly as in the north (Simmonds and Keay 2000). Extratropical  
40 cyclones are creatures of winter.

41 Precession, in contrast, is a lever that acts on summer. The slow turning of the longitude of  
42 perihelion is the strongest orbital pacemaker of the seasonal insolation cycle, yet it redistributes  
43 sunlight rather than adding any, and the redistribution is proportional to the sunlight a season  
44 already receives (Berger 1978; Laskar et al. 2004). Midlatitude summer insolation swings by more  
45 than  $100 \text{ W m}^{-2}$  across a precession cycle while midwinter insolation barely moves, which is why  
46 the orbital literature treats summer energy as the climatically active quantity (Huybers 2006). A  
47 winter phenomenon facing a summer lever invites a simple expectation. Precession should have  
48 little to say about storm tracks. That expectation has never been put to a clean test.

49 Where the imprint of precession has been studied in depth, it appears as a seasonal reorganization  
50 of climate rather than a change in the annual mean. The monsoon literature carries this tradition,  
51 from the classic orbital experiments of Kutzbach and Guetter (1986) to single-forcing designs that  
52 isolate precession explicitly (Bosmans et al. 2015; Merlis et al. 2013), and cave records show

53 the northern and southern monsoons swinging in antiphase as perihelion migrates through the  
54 year (Wang et al. 2008). The extratropics have received far less attention. Kaspar et al. (2007)  
55 simulated a stronger and poleward-shifted North Atlantic winter storm track for the Eemian from  
56 two orbital time slices, Brayshaw et al. (2010) traced Holocene orbital forcing into the North  
57 Atlantic and Mediterranean winter storm tracks through the baroclinicity they support, and Varma  
58 et al. (2012) found the Southern Hemisphere westerlies migrating under Holocene orbital forcing.  
59 Explicit cyclone tracking has entered paleoclimate work mainly at the Last Glacial Maximum,  
60 where ice sheets dominate the forcing (Pinto and Ludwig 2020; Raible et al. 2021), and in transient  
61 simulations of the last millennium (Hess and Chemke 2025). For tropical cyclones the precessional  
62 response has recently been isolated in high-resolution paleoclimate simulations (Raavi et al. 2023).  
63 To our knowledge, no previous study has applied explicit cyclone tracking across a full precession  
64 cycle to isolate the orbital response of the extratropical storm tracks.

65 Closing this gap matters beyond dynamics. However, the paleoclimate records that would register  
66 past storminess, from dust and wind-blown sediment to westerly-belt precipitation, respond to  
67 particular seasons rather than to the annual mean, and a seasonally biased archive can fabricate  
68 or erase an orbital signal depending on which season carries it (Laepple et al. 2011). Knowing  
69 where in the seasonal cycle the precessional storm signal lives is a prerequisite for reading such  
70 records. Storm diagnostics raise a parallel issue, because measures of cyclone influence time,  
71 cyclone counts, and cyclone intensity respond to forcing in different ways and must be separated  
72 before any single proxy is interpreted.

73 In the present study the precession cycle is covered with twelve equilibrium simulations of a  
74 coupled climate model, and every extratropical cyclone is tracked explicitly in six-hourly pressure  
75 fields in both hemispheres. The design isolates precession at amplified eccentricity with all other  
76 boundary conditions fixed, so that the seasonal anatomy of the orbital storm response can be  
77 mapped basin by basin against the calendar month of perihelion.

78 The naive expectation fails, and it fails in an instructive way. The annual-mean storm response  
79 is indeed small, but the signal has not vanished. It is concentrated in local summer, the quiet  
80 season, in all three storm basins. Summers spent near perihelion are calm and summers spent near  
81 aphelion are stormy, so the two hemispheres respond in antiphase across the precession cycle, an  
82 extratropical counterpart of the monsoon seesaw. The response travels from insolation to storms

83 through the summer Eady growth rate, and the growth-rate change is carried by vertical shear  
84 (Lindzen and Farrell 1980). The paper is organized around which season carries the precessional  
85 storm response, what dynamical chain transmits it, and why the winter storm track stays orbitally  
86 quiet. Section 2 describes the model, the ensemble, and the tracking. Section 3 presents the  
87 seasonal anatomy of the response, and section 4 discusses mechanisms, implications for proxy  
88 records, and limitations.

## 89 **2. Data and methods**

### 90 *a. Model and experiments*

91 The simulations are performed with AWI-ESM-2.1-wiso, an isotope-enabled version of the AWI  
92 Earth System Model (Shi et al. 2023). The atmosphere component is ECHAM6 at T63 resolution  
93 with 47 vertical levels (Stevens et al. 2013), which includes the land surface module JSBACH  
94 (Raddatz et al. 2007). The ocean and sea ice component is FESOM2, a finite-volume model  
95 formulated on an unstructured mesh (Danilov et al. 2017; Sidorenko et al. 2019). The water  
96 isotope tracers carried by the model are not used in the present study.

97 Twelve equilibrium simulations are analyzed, taken from the idealized precession ensemble  
98 introduced by Shi et al. (2025). The longitude of perihelion  $\omega$  is prescribed at values from  $0^\circ$  to  
99  $330^\circ$  in steps of  $30^\circ$ , while eccentricity is fixed at 0.058 and obliquity at  $24.5^\circ$ . The eccentricity  
100 value sits near the upper bound reached during the Quaternary and amplifies the precessional  
101 modulation of insolation, so that the response emerges clearly from internal variability. All  
102 remaining boundary conditions follow the preindustrial setup, and each phase is branched from  
103 a long preindustrial control. Thirty years of six-hourly output from the equilibrated part of each  
104 simulation form the basis of the cyclone analysis. The calendar month of perihelion for each phase  
105 is computed from two-body orbital geometry (Berger 1978), and the same formulary provides the  
106 insolation fields shown in Fig. 1.

### 107 *b. Cyclone detection and tracking*

108 Extratropical cyclones are detected and tracked with the algorithm of Crawford et al. (2021),  
109 and Fig. 2 summarizes the procedure with one six-hourly field of the ensemble. Six-hourly sea  
110 level pressure is first interpolated onto 100-km equal-area grids centered on each pole, and the two

hemispheres are processed separately. A grid cell qualifies as a cyclone center when its pressure  $p_c$  lies below that of all eight neighbors and the surrounding field keeps a sufficient inward gradient. The gradient test compares  $p_c$  with the mean pressure  $\bar{p}(d)$  on a one-cell ring at distance  $d$  from the candidate,

$$\frac{\bar{p}(d) - p_c}{d} \geq \gamma, \quad d = 1000 \text{ km}, \quad \gamma = 750 \text{ Pa (1000 km)}^{-1}, \quad (1)$$

and candidates over terrain above 1500 m are discarded. Centers closer than 1000 km are treated as one multicenter system. The cyclone area  $A$  is obtained by growing closed isobars around each center at a 200-Pa interval until a contour fails to close, and the last closed isobar defines the edge pressure  $p_e$  (Fig. 2b). Each cyclone carries the depth and the area-equivalent radius

$$D = p_e - p_c, \quad R = \sqrt{A/\pi}. \quad (2)$$

Tracking links the centers of consecutive six-hourly fields (Fig. 2c). For every active cyclone a first-guess position is projected from its previous displacement, damped to three quarters,

$$\mathbf{x}_q = \mathbf{x}_t + 0.75 (\mathbf{x}_t - \mathbf{x}_{t-1}), \quad (3)$$

and a center of the next field qualifies as the continuation when it lies within  $v_{\max}\Delta t$  of both the current position  $\mathbf{x}_t$  and the projection  $\mathbf{x}_q$ , with  $v_{\max} = 150 \text{ km h}^{-1}$  and  $\Delta t = 6 \text{ h}$ . Among the qualifying candidates the one nearest to the projection continues the track. Centers left unmatched at the new time start tracks as genesis events, and tracks left unmatched end as lysis. Only systems poleward of  $30^\circ$  latitude enter the analysis.

Several track-based diagnostics are derived. Track density counts every six-hourly cyclone position, system density counts each cyclone once at its mean position, and genesis density counts the first point of each track. Tracks that merely cross the boundaries of the monthly processing windows would appear as spurious genesis or lysis events, and such truncated events (about 7% of the total) are removed. For every cyclone the lifespan, the maximum depth relative to the outermost closed isobar, the minimum central pressure, the mean cyclone area, and the propagation speed are recorded.

### 133 *c. Gridded storm coverage*

134 A complementary Eulerian diagnostic measures how long each location spends under cyclone  
 135 influence. At every six-hourly time step a binary mask flags the grid cells that lie inside a square  
 136 search window of half-width  $\sqrt{A}$  grid cells around each cyclone center and whose pressure lies  
 137 below the edge pressure of that cyclone,

$$M(\mathbf{x}, t) = 1 \quad \text{where } p(\mathbf{x}, t) < p_e \text{ inside the window,} \quad (4)$$

138 and the masks are regridded to T63 and accumulated into monthly storm days, one quarter of  
 139 the flagged six-hourly steps. Storm days at a point therefore scale with the product of cyclone  
 140 number and cyclone area footprint. The two factors can carry opposite seasonal cycles, and over  
 141 the Southern Ocean the larger size of summer cyclones raises summertime storm days even though  
 142 cyclone counts peak in winter. Frequency statements in this paper consequently rest on the track-  
 143 based densities, while storm days serve as a smooth field for maps and for measures of cyclone  
 144 influence time. The gridded product covers all twelve phases. An early processing inconsistency  
 145 in the  $\omega = 30^\circ$  mask product was traced to a stale intermediate file and removed by regenerating  
 146 the storm masks of all twelve experiments with one pipeline version, and the regenerated fields  
 147 were verified month by month against the Lagrangian statistics.

148 An Eulerian measure of synoptic variability provides a further check that involves no tracking  
 149 at all. Bandpass sea level pressure variability is computed from the six-hourly fields with the 24-h  
 150 difference filter of Wallace et al. (1988), which isolates fluctuations on the synoptic time scales that  
 151 define the classical storm tracks (Blackmon 1976), and its monthly standard deviation is analyzed  
 152 with the same basins, seasons, and response measures (appendix Fig. A1).

### 153 *d. Baroclinicity diagnostics*

154 The maximum Eady growth rate serves as the measure of baroclinicity (Eady 1949; Lindzen  
 155 and Farrell 1980; Hoskins and Valdes 1990). It is computed as  $\sigma = 0.31 |f| |\partial U / \partial z| N^{-1}$  in the  
 156 925–700 hPa layer from monthly wind, temperature, and geopotential height. The response to  
 157 precession is quantified between the end members p090 and p270, which place perihelion at the  
 158 June and December solstices. Because the difference fields are dipolar, plain box averages cancel,

and the response amplitude is instead defined as the root-mean-square of the difference over a basin, normalized by the climatological growth rate of the season. A substitution decomposition in the style of Camargo et al. (2007) splits the difference field into a vertical-shear factor and a static-stability factor, and the share of each factor is obtained by projecting it onto the full difference pattern. The basin averages use the storm-track boxes defined below. This choice matters in the Southern Hemisphere, where a sector beginning at 30°S would admit the seasonal signal of the subtropical winter jet and disguise it as a storm-belt response.

#### *e. Basins and statistics*

Three storm-track basins are analyzed, the North Atlantic (30–75°N, 80°W–20°E), the North Pacific (30–75°N, 120°E–120°W), and the Southern Ocean (40–75°S, all longitudes). Basin means are area weighted. The size of the precessional response is reported as the range across the twelve phases, the maximum minus the minimum of the 30-year mean, expressed as a percentage of the phase mean. Uncertainty estimates use the interannual spread of the 30 individual years in each phase, and the significance of the end-member difference maps is assessed with a two-sided Welch  $t$  test at each grid point. The local-summer contrasts between extreme phases far exceed the interannual noise, with  $p$  values below  $10^{-12}$  in all three basins.

### **3. Results**

#### *a. Winter storm tracks and what storm days measure*

The three storm tracks of the ensemble live in winter, exactly as observed. Seasonal-mean genesis counts peak in DJF over the North Atlantic and the North Pacific and in JJA over the Southern Ocean, and system counts follow the same calendar in every phase. Cyclones also cut deepest in the cold season, with winter maximum depths exceeding their summer values by roughly half in the northern basins. The winter identity of the storm season is therefore common to both hemispheres and to all twelve orbital configurations (Fig. 4a–c; Table 1).

The gridded storm-day field tells a partly different story, and the difference is instructive. Basin-mean storm days over the Southern Ocean stay high through austral summer although summer cyclones are fewer. The elevation reflects cyclone size rather than cyclone number. Summer systems there are about twenty percent larger in area than their winter counterparts, and storm days



count the time a grid cell spends inside any cyclone, so the product of number and footprint hides the winter peak in counts. A diagnostic of cyclone influence time is not a diagnostic of cyclone frequency, and the two must be read together. Frequency statements below therefore rest on the track-based densities, while storm days provide the smooth fields used for maps.

## *b. The summer paradox*

The annual-mean view sets the baseline (Fig. 3). The twelve annual climatologies are close to interchangeable, and the anomalies of each phase from their common mean form weak, smoothly rotating dipoles along the storm tracks. The basin series beneath the maps quantify how little survives in the annual mean, with across-phase ranges of 10%, 17% and 4% in the North Atlantic, the North Pacific and the Southern Ocean, varying smoothly with  $\omega$  and already hinting at the interhemispheric antiphase, since the northern basins reach their minima near  $\omega = 90^\circ$  where the Southern Ocean peaks.

Precession barely touches the annual mean but reorganizes the seasons, and the season it selects is the quiet one. The seasonal cycles of basin-mean track density keep their winter peak in every phase and fan out across the phases almost exclusively in local summer (Fig. 4a–c). The across-phase range makes this quantitative (Fig. 4d–f). In the North Pacific the annual range is 7% of the phase mean while the JJA range reaches 44%. The North Atlantic behaves the same way with 7% against 25%. The Southern Ocean repeats the pattern half a year out of phase, with an annual range of 4% against 22% in DJF. System and genesis densities confirm the calendar in every basin, and the influence-time metric swings harder still, with a storm-day range of 68% in North Pacific summer (Table 1). The sensitive season is local summer everywhere, and the small annual signal is what survives of a substantial summer swing after the stable winter is averaged in.

A tracking-free view points the same way (appendix Fig. A1). Bandpass sea level pressure variability responds most strongly in JJA over the North Pacific, at 36% of its climatology against 9% in winter, and winter holds the weakest response of all seasons in both northern basins. This amplitude-weighted metric shifts part of the response within the warm half-year. Over the North Atlantic it peaks in early autumn, matching the secondary September lobe of the forcing itself (Fig. 1c), and over the Southern Ocean its apparent winter share lies on the equatorward flank of

215 the basin box, while poleward of 50°S austral summer leads. The warm-season concentration and  
216 the winter quiescence are therefore not artifacts of the tracking algorithm.

### 217 *c. Orbital anchoring and the interhemispheric mirror*

218 The summer response is anchored to the orbit, not to the calendar. When the basin-mean storm-  
219 day anomaly is drawn as a function of calendar month and precession phase, the anomaly ridge  
220 climbs through the diagram along the calendar month of perihelion (Fig. 5). The sign is fixed by  
221 the orbital geometry. Summers spent near perihelion are quiet and summers spent near aphelion  
222 are stormy. North Pacific summer activity reaches its minimum at  $\omega = 90^\circ$ , the configuration  
223 with perihelion at the June solstice, where it falls 31% below the phase mean, and its maximum  
224 of +37% near  $\omega = 270^\circ$  (Fig. 8a). The North Atlantic swings between  $-16\%$  and  $+15\%$  at the  
225 same phases. The Southern Ocean mirrors this behavior, with a DJF maximum at  $\omega = 60^\circ$  and a  
226 minimum at  $\omega = 210^\circ$ , because a perihelion that falls in northern summer is an aphelion summer  
227 for the south. One lever thus drives the two hemispheres in antiphase across the precession cycle.  
228 The extreme-phase contrasts far exceed interannual variability, with Welch  $p$  values below  $10^{-12}$   
229 in all three basins.

### 230 *d. The dynamical bridge runs through summer shear*

231 The seasonal selectivity begins in the forcing itself. Averaged over the northern storm-track band  
232 from 40 to 70°N, the seasonal forcing amplitude peaks at  $105 \text{ W m}^{-2}$  in JJA and collapses to 14  
233 in DJF, and the mirror band in the south peaks at  $119 \text{ W m}^{-2}$  in DJF against 47 in JJA (Fig. 1).  
234 Perihelion can only amplify a season that already receives sunlight, so the orbital lever presses on  
235 local summer in each hemisphere, keeps a substantial autumn shoulder, and leaves deep winter  
236 almost untouched.

237 Baroclinicity translates the pressure on the seasons into the storm response. The Eady growth  
238 rate difference between the end members p090 and p270 is largest in local summer within every  
239 storm basin (Fig. 6). Relative to the seasonal climatology the summer response reaches 31% in the  
240 North Atlantic, a striking 92% in the North Pacific, and 21% in the Southern Ocean, in each case  
241 the seasonal maximum for that basin. A substitution decomposition attributes the pattern almost  
242 entirely to the vertical-shear factor, whose projection accounts for 103–109% of the difference field

in the three basins, while the static-stability factor contributes near zero. The chain is short and uniform. Perihelion summers warm the summer hemisphere, flatten the meridional temperature gradient, weaken the thermal-wind shear, and starve the baroclinic growth on which summer cyclones depend. The storm anomalies appear somewhat poleward of the growth-rate anomalies, consistent with the downstream development of baroclinic waves (Chang et al. 2002), and this offset is why pattern measures rather than same-latitude box averages are used throughout. Winter tells a complementary story. The winter growth rate still shifts by 16–18% between the end members, yet the winter storm response stays below 12% in every basin, a point taken up in the discussion.

#### *e. Spatial expression, and what precession actually moves*

The end-member difference maps show where the summer signal lives (Fig. 7). In the annual mean the p090 minus p270 difference is weak and patchy. In JJA a pronounced dipole spans the northern basins, with fewer storm days through the midlatitude core and more along the Arctic flank at perihelion summer, the signature of a poleward shift joined to an overall summer weakening. In DJF the significant signal migrates to the Southern Ocean and forms a circumpolar banded structure of opposite sign. About half of the JJA grid points in the northern extratropics pass the 95% significance test, against a far smaller fraction in the annual mean.

The track displacement behind these dipoles can be made explicit (appendix Fig. A2). The storm-day-weighted latitude of the North Pacific summer track migrates from 53°N when perihelion falls in northern winter to 61°N when it falls in northern summer, a poleward excursion of 8° that peaks exactly at  $\omega = 90^\circ$ , and the track-density-weighted latitude follows the same curve with a 5° swing. The North Atlantic summer track moves 4.4° in the same direction, and the Southern Ocean summer track mirrors the rule with a poleward excursion of 3.8° when perihelion approaches austral summer. A perihelion summer therefore thins the storm population and displaces the survivors poleward, in step with the poleward shift of the baroclinic zone seen in the growth-rate dipole of Fig. 6a. The winter track latitude stays within 1.5–2.8° across the entire cycle, and its small extremes do not align with the solstitial phases, which extends the winter invariance from storm counts to storm position. The shoulder seasons fall in between and carry a telling phase structure, because the autumn excursion peaks near  $\omega = 150^\circ$ , the configuration that places perihelion in late summer, so the poleward displacement follows the perihelion month through the calendar exactly

272 as the occurrence anomaly does in Fig. 5. The annual-mean track latitude varies by no more than  
273  $1.3^\circ$  in any basin.

274 The orbital anchoring of occurrence and position condenses into a single statement (Fig. 9). A  
275 first-harmonic fit across the phases gives, for every calendar month, the value of  $\omega$  at which storm  
276 activity reaches its minimum and the track its poleward maximum, and through the sunlit half-year  
277 both sequences climb in parallel with the line that places perihelion in that month. The response  
278 lags the perihelion calendar by a basin-dependent interval, near one month in the North Pacific,  
279 one to two months in the North Atlantic, and slightly over two months in the Southern Ocean,  
280 an ordering consistent with the growing oceanic character of the three sectors and with surface  
281 thermal inertia setting the delay. Winter months fall off the line, where the response amplitude is  
282 too small to anchor. The storm tracks are phase locked to the perihelion calendar wherever the  
283 lever can reach them.

284 What precession moves is above all the number and location of cyclones, not their translation  
285 speed, and only regionally their depth (Fig. 8b). Local-summer ranges of the frequency and position  
286 metrics span 16–68% across the basins. Propagation speed stays within 2–5% everywhere. Cyclone  
287 depth separates the basins, and the diagnostics are deliberately kept apart in the manner of Zappa  
288 et al. (2013). Over the Southern Ocean the maximum depth varies by only 4% across the phases  
289 while frequency varies by 22%, a clean decoupling of frequency from intensity. Over the North  
290 Pacific the summer depth range reaches 33% and the lifespan range 23%, so there the whole  
291 cyclone population weakens together when perihelion falls in summer, fewer systems that are also  
292 shallower and shorter lived. The North Atlantic sits between these regimes. The common element  
293 across basins is that occurrence responds most, while the intensity response ranges from absent to  
294 substantial depending on how strongly the summer baroclinic reservoir is drained.

#### 295 *f. What the rain records*

296 The hydrological expression of the cycle runs opposite to the dynamical one (appendix Fig. A3).  
297 Detected cyclones deliver about one third of extratropical precipitation in the ensemble, in line  
298 with reanalysis-based attribution (Hawcroft et al. 2012). In North Pacific summer the precipitation  
299 falling inside each storm strengthens by up to 15% exactly where storm counts collapse, and the  
300 two series are nearly perfectly anticorrelated across the phases ( $r = -0.95$ ). A perihelion summer

holds a warmer and moister atmosphere, so the surviving storms rain harder, but the loss of storms wins, and total summer precipitation still falls by 17% between the end members. The share of precipitation delivered by storms falls with it, by 9.5 percentage points across the cycle in North Pacific summer against about 3 in the annual mean. Precession therefore forces extratropical hydroclimate in two opposing ways at once, fewer storms against wetter storms, and the two must be kept apart in interpretation just as counts and depths must.

## 4. Discussion

### *a. Why the lever lands in summer*

The seasonal selectivity of the storm response is set by orbital geometry before any model physics enters. Precession redistributes insolation in proportion to the insolation a season already has, so the winter hemisphere offers the lever almost nothing to press on (Berger 1978). The monthly forcing amplitude in Fig. 1 makes the floor explicit, with midwinter values near  $20 \text{ W m}^{-2}$  against summer values above  $125 \text{ W m}^{-2}$  at the latitudes of the storm tracks. This part of the result is model independent and would reappear in any climate model, because it restates the geometry that also underlies the summer-energy view of orbital forcing (Huybers 2006). The novelty is not that summer receives the forcing, which the monsoon literature established long ago (Kutzbach and Guetter 1986; Bosmans et al. 2015), but that the summer forcing finds a winter phenomenon and reorganizes it through its quiet season.

### *b. Cool summers are stormy summers*

The mechanism that converts summer insolation into summer storms is the thermal wind. Perihelion summers warm the summer hemisphere most strongly over land and at high latitudes, flatten the meridional temperature gradient, and weaken the vertical shear that feeds baroclinic growth. The factor decomposition confirms that the shear term carries essentially the whole Eady growth rate response in all three basins, while static stability contributes almost nothing. The same chain, with greenhouse warming in place of perihelion, is now observed in the present-day Northern Hemisphere, where amplified summer warming at high latitudes has weakened the summer circulation, the eddy kinetic energy, and the number of strong summer cyclones (Coumou et al. 2015; Chang et al. 2016; Gertler and O’Gorman 2019). Projections extend the analogy, with

CMIP-class models thinning summer cyclone activity and the associated baroclinicity through the same shear pathway (Lehmann et al. 2014; Priestley and Catto 2022). The correspondence should be read as a shared mechanism rather than a shared scenario, because the warming trend alters the two hemispheres together whereas precession swings them in antiphase. Under both forcings the rule is the same, and it is the rule in the title. Cool summers are stormy summers.

*c. The orbital invariance of the winter storm track*

The stability of winter deserves as much attention as the mobility of summer. Winter storm activity varies by less than about a tenth across the entire precession cycle in every basin, and this is only partly a matter of weak winter forcing. The winter Eady growth rate still shifts by 16–18% between the end members, yet winter cyclone numbers barely respond. The winter storm track appears saturated in the sense that ample baroclinicity is available in every phase, so that moderate changes in the growth rate do not translate into changes in occurrence. Whatever its ultimate cause, the practical consequence is notable. The delivery of winter precipitation to the midlatitude and subpolar continents, the season and pathway that build snowpack and feed ice sheets, is essentially fixed across orbital configurations of the precession cycle. A transient simulation of the last millennium likewise found cyclone counts nearly insensitive to external forcing (Raible et al. 2018; Hess and Chemke 2025), and the present result extends that insensitivity to the far larger insolation excursions of the orbital band, for winter alone.

*d. One lever, three basins, and what the records would see*

The three basins express one mechanism with different accents, and the diagnostics must be kept apart to see it. Occurrence responds everywhere, with local-summer ranges of 16–68% in frequency and position metrics, while propagation speed is inert. Intensity separates the basins. The Southern Ocean decouples cleanly, with a 4% depth range against a 22% frequency range, the North Pacific weakens as a whole population in perihelion summers, and the North Atlantic lies between. The pattern echoes the long-standing conclusion from anthropogenic scenarios that cyclone number and cyclone intensity respond to forcing through different processes and need not move together (Bengtsson et al. 2009; Ulbrich et al. 2009; Catto et al. 2019). The water cycle sharpens the contrast, because per-storm precipitation strengthens in the very summers that lose

357 storms, an orbital counterpart of the warming-scenario finding that hydrological responses outrun  
358 dynamical ones (Bengtsson et al. 2009). For the record this matters twice over. Extratropical  
359 precipitation is delivered disproportionately by cyclones and their fronts (Hawcroft et al. 2012;  
360 Catto and Pfahl 2013), and the storm-attributed share of precipitation swings by up to ten percentage  
361 points across the cycle in North Pacific summer, so a precipitation archive senses the competition  
362 between fewer storms and wetter storms rather than either alone. The storm-day diagnostic adds  
363 its own caution, because cyclone influence time folds cyclone size into cyclone number, and over  
364 the Southern Ocean the seasonal cycle of size alone can invert the apparent storm season. Any  
365 proxy that integrates storm influence rather than storm counts inherits this ambiguity.

366 For the paleoclimate record the seasonal anatomy carries a concrete and testable message. An  
367 archive with summer-biased sensitivity in the midlatitude storm belts should carry a pronounced  
368 precession signal in storminess, while an annually integrating archive should carry almost none.  
369 The seasonal bias of an archive can therefore fabricate or erase an orbital storm signal exactly as it  
370 does for temperature (Laepple et al. 2011). The interhemispheric mirror sharpens the prediction,  
371 because summer-sensitive storminess records from the two hemispheres should vary in antiphase  
372 at the precession period, the extratropical counterpart of the monsoon seesaw (Wang et al. 2008).  
373 Existing archives make the test realistic. Precession-band signals are already established in the  
374 northern midlatitude westerlies (Li et al. 2019) and in the composition of North Pacific dust, where  
375 the precession imprint sits in the moisture-sensitive component while the annually integrating  
376 flux follows obliquity (Zhong et al. 2024), and a winter-integrating dust-strength record shows  
377 correspondingly weak precession power (Chen et al. 2014). In the south the westerly belt carries  
378 strong 19 and 23 kyr cycles at its subtropical edge (Lamy et al. 2019), with the orbital band  
379 recorded at a site depending on its latitude within the belt (Kaiser et al. 2024). The subtropical  
380 location of that southern signal is consistent with the present results, because the winter half of  
381 the precessional baroclinicity response resides in the subtropical jet equatorward of the storm belt,  
382 while the storm-belt response is a summer phenomenon. Records of the southern westerlies also  
383 show orbital-band migrations of the belt itself (Lamy et al. 2010; Varma et al. 2012), and glacial-  
384 state cyclone modeling has shown how strongly reconstructed storminess depends on which cyclone  
385 property a proxy senses (Pinto and Ludwig 2020). A paired-hemisphere test at the precession band  
386 is within reach of existing archives.

## 387 *e. Limitations*

388 Several limitations bound these conclusions. The results rest on one model, one tracking algo-  
389 rithm, and an idealized orbital configuration with eccentricity fixed near its Quaternary maximum,  
390 so the simulated amplitudes represent an upper bound that scales with  $e \sin \omega$  rather than a recon-  
391 struction of any particular epoch. Obliquity is fixed at  $24.5^\circ$ , and interactions between precession  
392 and obliquity or ice volume are outside the design. The simulations are equilibrium slices, so  
393 transient effects along the precession cycle are not represented. The T63 atmosphere resolves  
394 the storm tracks but not mesoscale substructure, and low-level Eady growth rates in the Southern  
395 Hemisphere carry a documented sensitivity to boundary-layer stability (Simmonds and Lim 2009).  
396 The gridded storm-day product required one uniform regeneration of the storm masks before all  
397 twelve phases could be used together, a reminder that multi-batch processing of tracked-cyclone  
398 products deserves explicit version control.

## 399 **5. Conclusions**

400 One rule spans the three storm tracks of this precession ensemble. The orbital lever reaches  
401 extratropical cyclones through local summer, the quiet season, while the winter storm track stays  
402 essentially fixed across the entire cycle. Summers spent near perihelion are calm and summers  
403 near aphelion are stormy, and because one hemisphere's perihelion summer is the other's aphelion  
404 summer, midlatitude storminess in the two hemispheres swings in antiphase at the precession  
405 period. The signal travels through shear-driven summer baroclinicity and expresses itself chiefly  
406 in how many cyclones occur and where they run, with per-cyclone intensity responding only where  
407 the summer baroclinic reservoir is drained most deeply. Both expressions are phase locked to  
408 the calendar month of perihelion, with a delay of one to two months that grows with the oceanic  
409 character of the basin. For the proxy record the lesson is direct. Orbital storminess signals should  
410 be sought in summer-sensitive midlatitude archives, and they should appear with opposite sign in  
411 the two hemispheres.








FIG. 1. Precessional insolation forcing from orbital geometry with eccentricity 0.058 and obliquity  $24.5^\circ$ .  
 (a) Monthly insolation anomaly relative to the phase mean, averaged over the storm-track band  $40\text{--}70^\circ\text{N}$ , as  
 a function of calendar month and precession phase  $\omega$ , with the perihelion (red) and aphelion (blue) calendar  
 months of each phase overlaid. (b) As in (a), but for  $40\text{--}70^\circ\text{S}$ . (c) Seasonal forcing amplitude in the two bands,  
 the seasonal mean of the monthly maximum minus minimum insolation across the twelve phases. Units:  $\text{W m}^{-2}$ .


(a) centers + area radii, one 6-h field



(b) outermost closed isobar  $p_e$ , mask cells



(c) tracks of one month ( $n=407$ )



(d)




FIG. 2. Cyclone detection and tracking procedure. (a) One six-hourly sea level pressure field of phase p000 (contours, 8-hPa interval) with all detected cyclone centers poleward of 30°N (dots) and their area-equivalent radii  $R = \sqrt{A/\pi}$  (circles), 15 January of one model year. (b) The deepest system of that field, with the outermost closed isobar  $p_e$  (dashed), the cyclone center (cross), the central and edge pressures and depth annotated, and the binary mask cells (shading) defined by  $SLP < p_e$  inside the  $\sqrt{A}$  search window; contour interval 4 hPa. (c) All cyclone tracks of the same month (lines) with genesis points (dots). (d) Processing chain with the key thresholds of Eqs. (1)–(4).




FIG. 3. Annual-mean storm days across the precession cycle. (a)–(l) Annual storm-day anomaly of each phase relative to the twelve-phase mean (shading), with the phase-mean climatology outlined at 5 and 8 d month<sup>-1</sup> (contours). (m)–(o) Basin-mean annual storm days against  $\omega$  with interannual standard errors; the across-phase range as a percentage of the phase mean is annotated in each panel.




FIG. 4. Seasonal concentration of the precessional storm response, from the Lagrangian tracks. (a)–(c) Basin-mean track density (cell<sup>-1</sup> month<sup>-1</sup>) by calendar month for the North Atlantic, the North Pacific, and the Southern Ocean, one line per phase colored by  $\omega$ , with local summer shaded. (d)–(f) Across-phase range of track density (maximum minus minimum, percent of the phase mean) for the annual mean and the four seasons, with the local-summer bar highlighted; open symbols give the corresponding ranges of system and genesis densities.




FIG. 5. Basin-mean storm-day anomaly (d month<sup>-1</sup>) relative to the phase mean as a function of calendar month and precession phase for (a) the North Atlantic, (b) the North Pacific, and (c) the Southern Ocean, with the perihelion (red) and aphelion (blue) calendar months of each phase overlaid.




FIG. 6. Precessional response of the maximum Eady growth rate (925–700 hPa). (a) p090 minus p270 difference for JJA and (b) for DJF ( $\text{day}^{-1}$ ). (c) Seasonal response amplitude in each basin, the RMS of the p090 minus p270 difference over the storm box as a percentage of the seasonal climatology, with stars marking local summer. (d) Projection shares of the vertical-shear and static-stability factors in the local-summer difference pattern.




FIG. 7. Difference in storm days ( $\text{d month}^{-1}$ ) between p090 and p270 for (a) the annual mean, (b) JJA, and (c) DJF, on a common color scale. Stippling marks grid points where the difference is significant at the 95% level based on a two-sided Welch  $t$  test over the 30 simulated years.




FIG. 8. (a) Local-summer storm activity across the precession cycle, shown as the percent anomaly of basin-mean storm days from the phase mean (solid, with interannual standard errors) and of track density (dashed), for the North Atlantic and the North Pacific in JJA and for the Southern Ocean in DJF; dotted vertical lines mark the phases with perihelion in NH and SH summer. (b) Local-summer across-phase range for frequency and position metrics and for intensity and lifecycle metrics in each basin.




FIG. 9. Phase locking of the storm response to the perihelion calendar. For each calendar month and basin, the phase of the first harmonic across the twelve precession phases of (a) basin-mean storm days, shown as the  $\omega$  of the activity minimum, and (b) the storm-day-weighted track latitude, shown as the  $\omega$  of the poleward maximum. Marker size scales with the harmonic amplitude. The red line gives the  $\omega$  that places perihelion in that calendar month, and dashed lines repeat it delayed by each basin's median lag over the amplitude-strong months (annotated in months).

501     TABLE 1. Three-basin comparison of the precessional storm response. Ranges are max minus min across the  
502     twelve phases as a percentage of the phase mean. The storm season is diagnosed from cyclogenesis counts, the  
503     sensitive season from the seasonal storm-day range. The EGR response is the RMS p090–p270 difference over  
504     the basin sector relative to its climatology; the dominant factor is its projection share.

| Basin          | Storm season | Sensitive season | Storm-day range ann/summer (%) | Track-density range summer (%) | Summer max/min phase       |
|----------------|--------------|------------------|--------------------------------|--------------------------------|----------------------------|
| North Atlantic | DJF (winter) | JJA              | 10 / 32                        | 24                             | $\omega=240$ / $\omega=90$ |
| North Pacific  | DJF (winter) | JJA              | 17 / 68                        | 44                             | $\omega=270$ / $\omega=90$ |
| Southern Ocean | JJA (winter) | DJF              | 4 / 16                         | 22                             | $\omega=60$ / $\omega=210$ |




FIG. A1. Tracking-free Eulerian storm-track metric, the monthly standard deviation of 24-h differences of six-hourly sea level pressure. (a)–(c) Relative seasonal response in each basin, the RMS of the p090 minus p270 difference over the storm box as a percentage of the seasonal climatology (as in Fig. 6c), with the local-summer bar highlighted. (d)–(f) The p090 minus p270 difference of the same quantity for the annual mean, JJA, and DJF (hPa).

**Acknowledgments.** The simulations were performed at the German Climate Computing Center (DKRZ). [TODO funding and grant numbers before submission.]

**Data availability statement.** The AWI-ESM simulations of the idealized precession ensemble are described in Shi et al. (2025). The cyclone-track database, the gridded storm statistics, and the analysis scripts underlying all figures will be archived and assigned a DOI upon acceptance. [TODO repository DOI.]

## APPENDIX

### Tracking-independent, positional, and hydrological diagnostics

Figures A1–A3 present the Eulerian bandpass diagnostic, the seasonal track-latitude analysis, and the storm-attributed precipitation analysis discussed in sections 2 and 3.






FIG. A2. Storm-track latitude across the precession cycle, the storm-day-weighted mean latitude over each basin sector shown as the poleward anomaly from each season's phase mean, for (a) the North Atlantic, (b) the North Pacific, and (c) the Southern Ocean. Lines give the annual mean and the four seasons, with the local summer of each basin drawn heavy with markers. Vertical dotted lines mark the phase with perihelion in the local summer of each basin.




FIG. A3. Storm-attributed precipitation across the precession cycle. (a) North Pacific JJA percent anomalies from the phase mean of basin-mean storm days, per-storm precipitation, and total precipitation; the dotted line marks the phase with perihelion in NH summer. (b) Across-phase range of the storm-attributed share of precipitation, accumulated precipitation inside storm footprints divided by total accumulated precipitation (percentage points), for the annual mean and local summer in each basin.

## References

- Bengtsson, L., K. I. Hodges, and N. Keenlyside, 2009: Will extratropical storms intensify in a warmer climate? *Journal of Climate*, **22** (9), 2276–2301, <https://doi.org/10.1175/2008JCLI2678.1>.
- Berger, A. L., 1978: Long-term variations of daily insolation and quaternary climatic changes. *Journal of the Atmospheric Sciences*, **35** (12), 2362–2367, [https://doi.org/10.1175/1520-0469\(1978\)035\(2362:LTVODI\)2.0.CO;2](https://doi.org/10.1175/1520-0469(1978)035(2362:LTVODI)2.0.CO;2).
- Blackmon, M. L., 1976: A climatological spectral study of the 500 mb geopotential height of the Northern Hemisphere. *Journal of the Atmospheric Sciences*, **33** (8), 1607–1623, [https://doi.org/10.1175/1520-0469\(1976\)033\(1607:ACSSOT\)2.0.CO;2](https://doi.org/10.1175/1520-0469(1976)033(1607:ACSSOT)2.0.CO;2).
- Bosmans, J. H. C., S. S. Drijfhout, E. Tuenter, F. J. Hilgen, L. J. Lourens, and E. J. Rohling, 2015: Response of the north african summer monsoon to precession and obliquity forcings in the earth gcm. *Climate Dynamics*, **44** (1–2), 279–297, <https://doi.org/10.1007/s00382-014-2260-z>.
- Brayshaw, D. J., B. Hoskins, and E. Black, 2010: Some physical drivers of changes in the winter storm tracks over the North Atlantic and Mediterranean during the Holocene. *Philosophical Transactions of the Royal Society A*, **368** (1931), 5185–5223, <https://doi.org/10.1098/rsta.2010.0180>.
- Camargo, S. J., K. A. Emanuel, and A. H. Sobel, 2007: Use of a genesis potential index to diagnose ENSO effects on tropical cyclone genesis. *Journal of Climate*, **20** (19), 4819–4834, <https://doi.org/10.1175/JCLI4282.1>.
- Catto, J. L., and S. Pfahl, 2013: The importance of fronts for extreme precipitation. *Journal of Geophysical Research: Atmospheres*, **118** (19), 10 791–10 801, <https://doi.org/10.1002/jgrd.50852>.
- Catto, J. L., and Coauthors, 2019: The future of midlatitude cyclones. *Current Climate Change Reports*, **5**, 407–420, <https://doi.org/10.1007/s40641-019-00149-4>.
- Chang, E. K. M., S. Lee, and K. L. Swanson, 2002: Storm track dynamics. *Journal of Climate*, **15** (16), 2163–2183, [https://doi.org/10.1175/1520-0442\(2002\)015\(02163:STD\)2.0.CO;2](https://doi.org/10.1175/1520-0442(2002)015(02163:STD)2.0.CO;2).

572 Chang, E. K. M., C.-T. Ma, C. Zheng, and A. M. W. Yau, 2016: Observed and projected  
 573 decrease of Northern Hemisphere extratropical cyclone activity in summer and its impacts  
 574 on maximum temperature. *Geophysical Research Letters*, **43** (5), 2200–2208, <https://doi.org/10.1002/2016GL068172>.  
 575

576 Chen, J., X. Liu, and V. A. Kravchinsky, 2014: Response of the high-resolution Chinese loess grain  
 577 size record to the 50°N integrated winter insolation during the last 500,000 years. *Geophysical*  
 578 *Research Letters*, **41** (17), 6113–6120, <https://doi.org/10.1002/2014GL060239>.

579 Coumou, D., J. Lehmann, and J. Beckmann, 2015: The weakening summer circulation in the  
 580 Northern Hemisphere mid-latitudes. *Science*, **348** (6232), 324–327, [https://doi.org/10.1126/](https://doi.org/10.1126/science.1261768)  
 581 [science.1261768](https://doi.org/10.1126/science.1261768).

582 Crawford, A. D., E. A. P. Schreiber, N. Sommer, M. C. Serreze, J. C. Stroeve, and D. G. Barber,  
 583 2021: Sensitivity of Northern Hemisphere cyclone detection and tracking results to fine spatial  
 584 and temporal resolution. *Monthly Weather Review*, **149** (8), 2581–2598, [https://doi.org/10.1175/](https://doi.org/10.1175/MWR-D-20-0417.1)  
 585 [MWR-D-20-0417.1](https://doi.org/10.1175/MWR-D-20-0417.1).

586 Danilov, S., D. Sidorenko, Q. Wang, and T. Jung, 2017: The Finite-volume Sea ice–Ocean  
 587 Model (FESOM2). *Geoscientific Model Development*, **10** (2), 765–789, [https://doi.org/10.5194/](https://doi.org/10.5194/gmd-10-765-2017)  
 588 [gmd-10-765-2017](https://doi.org/10.5194/gmd-10-765-2017).

589 Eady, E. T., 1949: Long waves and cyclone waves. *Tellus*, **1** (3), 33–52, [https://doi.org/10.3402/](https://doi.org/10.3402/tellusa.v1i3.8507)  
 590 [tellusa.v1i3.8507](https://doi.org/10.3402/tellusa.v1i3.8507).

591 Gertler, C. G., and P. A. O’Gorman, 2019: Changing available energy for extratropical cyclones and  
 592 associated convection in Northern Hemisphere summer. *Proceedings of the National Academy*  
 593 *of Sciences*, **116** (10), 4105–4110, <https://doi.org/10.1073/pnas.1812312116>.

594 Hawcroft, M. K., L. C. Shaffrey, K. I. Hodges, and H. F. Dacre, 2012: How much Northern  
 595 Hemisphere precipitation is associated with extratropical cyclones? *Geophysical Research*  
 596 *Letters*, **39**, L24 809, <https://doi.org/10.1029/2012GL053866>.

597 Hess, O., and R. Chemke, 2025: Forced changes in the North Atlantic storm track over the  
 598 last millennium. *Geophysical Research Letters*, **52**, e2025GL115 894, [https://doi.org/10.1029/](https://doi.org/10.1029/2025GL115894)  
 599 [2025GL115894](https://doi.org/10.1029/2025GL115894).

600 Hoskins, B. J., and K. I. Hodges, 2002: New perspectives on the Northern Hemisphere winter  
 601 storm tracks. *Journal of the Atmospheric Sciences*, **59** (6), 1041–1061, [https://doi.org/10.1175/  
 602 1520-0469\(2002\)059<1041:NPOTNH>2.0.CO;2](https://doi.org/10.1175/1520-0469(2002)059<1041:NPOTNH>2.0.CO;2).

603 Hoskins, B. J., and K. I. Hodges, 2005: A new perspective on Southern Hemisphere storm tracks.  
 604 *Journal of Climate*, **18** (20), 4108–4129, <https://doi.org/10.1175/JCLI3570.1>.

605 Hoskins, B. J., and P. J. Valdes, 1990: On the existence of storm-tracks. *Journal of the Atmospheric  
 606 Sciences*, **47** (15), 1854–1864, [https://doi.org/10.1175/1520-0469\(1990\)047<1854:OTEOST>2.  
 607 0.CO;2](https://doi.org/10.1175/1520-0469(1990)047<1854:OTEOST>2.0.CO;2).

608 Huybers, P., 2006: Early pleistocene glacial cycles and the integrated summer insolation forcing.  
 609 *Science*, **313** (5786), 508–511, <https://doi.org/10.1126/science.1125249>.

610 Kaiser, J., E. Schefuß, J. A. Collins, and Coauthors, 2024: Orbital modulation of subtropical versus  
 611 subantarctic moisture sources in the southeast Pacific mid-latitudes. *Nature Communications*,  
 612 **15**, 7512, <https://doi.org/10.1038/s41467-024-51985-4>.

613 Kaspar, F., T. Spanghehl, and U. Cubasch, 2007: Northern hemisphere winter storm tracks of  
 614 the eemian interglacial and the last glacial inception. *Climate of the Past*, **3** (2), 181–192,  
 615 <https://doi.org/10.5194/cp-3-181-2007>.

616 Kutzbach, J. E., and P. J. Guetter, 1986: The influence of changing orbital parameters and surface  
 617 boundary conditions on climate simulations for the past 18,000 years. *Journal of the Atmospheric  
 618 Sciences*, **43** (16), 1726–1759, [https://doi.org/10.1175/1520-0469\(1986\)043<1726:TIOCOP>2.  
 619 0.CO;2](https://doi.org/10.1175/1520-0469(1986)043<1726:TIOCOP>2.0.CO;2).

620 Laepple, T., M. Werner, and G. Lohmann, 2011: Synchronicity of antarctic temperatures and  
 621 local solar insolation on orbital timescales. *Nature*, **471** (7336), 91–94, [https://doi.org/10.1038/  
 622 nature09825](https://doi.org/10.1038/nature09825).

623 Lamy, F., R. Kilian, H. W. Arz, J.-P. Francois, J. Kaiser, M. Prange, and T. Steinke, 2010: Holocene  
 624 changes in the position and intensity of the southern westerly wind belt. *Nature Geoscience*,  
 625 **3** (10), 695–699, <https://doi.org/10.1038/ngeo959>.

- 626 Lamy, F., and Coauthors, 2019: Precession modulation of the South Pacific westerly wind belt  
627 over the past million years. *Proceedings of the National Academy of Sciences*, **116** (47), 23 455–  
628 23 460, <https://doi.org/10.1073/pnas.1905847116>.
- 629 Laskar, J., P. Robutel, F. Joutel, M. Gastineau, A. C. M. Correia, and B. Levrard, 2004: A long-  
630 term numerical solution for the insolation quantities of the earth. *Astronomy and Astrophysics*,  
631 **428** (1), 261–285, <https://doi.org/10.1051/0004-6361:20041335>.
- 632 Lehmann, J., D. Coumou, K. Frieler, A. V. Eliseev, and A. Levermann, 2014: Future changes  
633 in extratropical storm tracks and baroclinicity under climate change. *Environmental Research*  
634 *Letters*, **9** (8), 084 002, <https://doi.org/10.1088/1748-9326/9/8/084002>.
- 635 Li, Y., Y. Song, Q. Yin, and Coauthors, 2019: Orbital and millennial northern mid-latitude  
636 westerlies over the last glacial period. *Climate Dynamics*, **53** (5–6), 3315–3324, [https://doi.org/](https://doi.org/10.1007/s00382-019-04704-5)  
637 [10.1007/s00382-019-04704-5](https://doi.org/10.1007/s00382-019-04704-5).
- 638 Lindzen, R. S., and B. Farrell, 1980: A simple approximate result for the maximum growth rate of  
639 baroclinic instabilities. *Journal of the Atmospheric Sciences*, **37** (7), 1648–1654, [https://doi.org/](https://doi.org/10.1175/1520-0469(1980)037<1648:ASARFT>2.0.CO;2)  
640 [10.1175/1520-0469\(1980\)037<1648:ASARFT>2.0.CO;2](https://doi.org/10.1175/1520-0469(1980)037<1648:ASARFT>2.0.CO;2).
- 641 Merlis, T. M., T. Schneider, S. Bordoni, and I. Eisenman, 2013: Hadley circulation response to  
642 orbital precession. part i: Aquaplanets. *Journal of Climate*, **26** (3), 740–753, [https://doi.org/](https://doi.org/10.1175/JCLI-D-11-00716.1)  
643 [10.1175/JCLI-D-11-00716.1](https://doi.org/10.1175/JCLI-D-11-00716.1).
- 644 Pinto, J. G., and P. Ludwig, 2020: Extratropical cyclones over the north atlantic and western europe  
645 during the last glacial maximum and implications for proxy interpretation. *Climate of the Past*,  
646 **16** (2), 611–626, <https://doi.org/10.5194/cp-16-611-2020>.
- 647 Priestley, M. D. K., and J. L. Catto, 2022: Future changes in the extratropical storm tracks and  
648 cyclone intensity, wind speed, and structure. *Weather and Climate Dynamics*, **3** (2), 337–360,  
649 <https://doi.org/10.5194/wcd-3-337-2022>.
- 650 Raavi, P. H., J.-E. Chu, A. Timmermann, S.-S. Lee, and K. J. E. Walsh, 2023: Moisture control  
651 of tropical cyclones in high-resolution simulations of paleoclimate and future climate. *Nature*  
652 *Communications*, **14**, 6426, <https://doi.org/10.1038/s41467-023-42033-8>.

653 Raddatz, T. J., and Coauthors, 2007: Will the tropical land biosphere dominate the climate–  
654 carbon cycle feedback during the twenty-first century? *Climate Dynamics*, **29** (6), 565–574,  
655 <https://doi.org/10.1007/s00382-007-0247-8>.

656 Raible, C. C., M. Messmer, F. Lehner, T. F. Stocker, and R. Blender, 2018: Extratropical cyclone  
657 statistics during the last millennium and the 21st century. *Climate of the Past*, **14** (10), 1499–1514,  
658 <https://doi.org/10.5194/cp-14-1499-2018>.

659 Raible, C. C., J. G. Pinto, P. Ludwig, and M. Messmer, 2021: A review of past changes in  
660 extratropical cyclones in the northern hemisphere and what can be learned for the future. *WIREs*  
661 *Climate Change*, **12** (3), e680, <https://doi.org/10.1002/wcc.680>.

662 Shi, X., A. Cauquoin, G. Lohmann, L. Jonkers, Q. Wang, H. Yang, Y. Sun, and M. Werner,  
663 2023: Simulated stable water isotopes during the mid-Holocene and pre-industrial periods  
664 using AWI-ESM-2.1-wiso. *Geoscientific Model Development*, **16**, 5153–5178, [https://doi.org/](https://doi.org/10.5194/gmd-16-5153-2023)  
665 [10.5194/gmd-16-5153-2023](https://doi.org/10.5194/gmd-16-5153-2023).

666 Shi, X., M. Werner, H. Yang, Q. Gao, J. Liu, and G. Lohmann, 2025: Precession controls on  
667 climate and water isotope signals in Northern Africa. *Paleoceanography and Paleoclimatology*,  
668 **40**, e2024PA004 999, <https://doi.org/10.1029/2024PA004999>.

669 Sidorenko, D., and Coauthors, 2019: Evaluation of FESOM2.0 coupled to ECHAM6.3: preindus-  
670 trial and HighResMIP simulations. *Journal of Advances in Modeling Earth Systems*, **11** (11),  
671 3794–3815, <https://doi.org/10.1029/2019MS001696>.

672 Simmonds, I., and K. Keay, 2000: Mean southern hemisphere extratropical cyclone behavior in  
673 the 40-year ncep–ncar reanalysis. *Journal of Climate*, **13** (5), 873–885, [https://doi.org/10.1175/](https://doi.org/10.1175/1520-0442(2000)013<0873:MSHECB>2.0.CO;2)  
674 [1520-0442\(2000\)013<0873:MSHECB>2.0.CO;2](https://doi.org/10.1175/1520-0442(2000)013<0873:MSHECB>2.0.CO;2).

675 Simmonds, I., and E.-P. Lim, 2009: Biases in the calculation of Southern Hemisphere mean  
676 baroclinic eddy growth rate. *Geophysical Research Letters*, **36** (1), L01 707, [https://doi.org/](https://doi.org/10.1029/2008GL036320)  
677 [10.1029/2008GL036320](https://doi.org/10.1029/2008GL036320).

678 Stevens, B., and Coauthors, 2013: Atmospheric component of the MPI-M Earth System Model:  
679 ECHAM6. *Journal of Advances in Modeling Earth Systems*, **5** (2), 146–172, [https://doi.org/](https://doi.org/10.1002/jame.20015)  
680 [10.1002/jame.20015](https://doi.org/10.1002/jame.20015).

- 681 Ulbrich, U., G. C. Leckebusch, and J. G. Pinto, 2009: Extra-tropical cyclones in the present and fu-  
682 ture climate: a review. *Theoretical and Applied Climatology*, **96** (1–2), 117–131, [https://doi.org/](https://doi.org/10.1007/s00704-008-0083-8)  
683 10.1007/s00704-008-0083-8.
- 684 Varma, V., and Coauthors, 2012: Holocene evolution of the southern hemisphere westerly winds  
685 in transient simulations with global climate models. *Climate of the Past*, **8** (2), 391–402,  
686 <https://doi.org/10.5194/cp-8-391-2012>.
- 687 Wallace, J. M., G.-H. Lim, and M. L. Blackmon, 1988: Relationship between cyclone tracks,  
688 anticyclone tracks and baroclinic waveguides. *Journal of the Atmospheric Sciences*, **45** (3),  
689 439–462, [https://doi.org/10.1175/1520-0469\(1988\)045<0439:RBCTAT>2.0.CO;2](https://doi.org/10.1175/1520-0469(1988)045<0439:RBCTAT>2.0.CO;2).
- 690 Wang, Y., and Coauthors, 2008: Millennial- and orbital-scale changes in the east asian  
691 monsoon over the past 224,000 years. *Nature*, **451** (7182), 1090–1093, [https://doi.org/](https://doi.org/10.1038/nature06692)  
692 10.1038/nature06692.
- 693 Zappa, G., L. C. Shaffrey, K. I. Hodges, P. G. Sansom, and D. B. Stephenson, 2013: A mul-  
694 timodel assessment of future projections of North Atlantic and European extratropical cy-  
695 clones in the CMIP5 climate models. *Journal of Climate*, **26** (16), 5846–5862, [https://doi.org/](https://doi.org/10.1175/JCLI-D-12-00573.1)  
696 10.1175/JCLI-D-12-00573.1.
- 697 Zhong, Y., and Coauthors, 2024: Orbital controls on North Pacific dust flux during the  
698 late Quaternary. *Geophysical Research Letters*, **51**, e2023GL106631, [https://doi.org/10.1029/](https://doi.org/10.1029/2023GL106631)  
699 2023GL106631.