

**Cool summers are stormy summers: precession reorganizes extratropical
cyclone activity through the quiet season**

Xiaoxu Shi^a

^a *Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research, Bremerhaven,
Germany*

Corresponding author: Xiaoxu Shi, shixiaoxu@sml-zhuhai.cn

7 ABSTRACT: Extratropical cyclones peak in winter, while orbital precession reshapes the seasonal
8 insolation cycle mainly in summer. Thus storm tracks might be expected to ignore the precession
9 cycle. This is examined with twelve equilibrium simulations using a coupled climate model AWI-
10 ESM2, spanning a full precession cycle at 30° longitude increments under fixed eccentricity and
11 obliquity. Extratropical cyclones are explicitly tracked based on 6-hourly atmospheric pressure
12 fields for both hemispheres. While the annual-mean cyclone response to precession is small,
13 substantial seasonal response emerges. In the North Atlantic, the North Pacific, and the Southern
14 Ocean the response focuses on local summer. We also find summers spent near perihelion feature
15 suppressed storminess, whereas summers near aphelion experience intensified cyclone activity,
16 yielding antiphase mid-latitude storm variability between the two hemispheres over the precession
17 cycle. This signal is carried by cyclone number and position, and it follows the pronounced
18 summer response of the Eady growth rate driven primarily by vertical wind shear in all three
19 basins. Precession therefore shapes extratropical storm tracks mostly through their quiet season
20 (i.e., summer). Summer-sensitive proxy archives should record a strong and hemispherically
21 antiphased storminess signal at the precession period that annually integrating archives would
22 miss. Moreover, monthly storm responses are phase-locked to the calendar timing of perihelion,
23 lagged by 1-2 months due to surface thermal inertia. This fingerprint also appears when we apply
24 the same cyclone tracking algorithm to pre-industrial (PI, 0 ka, perihelion falling in winter), Last
25 Interglacial (LIG, 127 ka, perihelion falling in summer) and mid-Holocene (MH, 6 ka, perihelion
26 falling in autumn), with weaker amplitudes matching their weaker eccentricity.

27 SIGNIFICANCE STATEMENT: Long orbital cycles drive global climate variability, and the
28 strongest of these, precession, reshapes sunlight mainly in summer, while mid-latitude storms
29 happen mainly in winter. This creates an apparent disconnect between precessional forcing and
30 storm track variability. Climate model simulations covering a full precession cycle show that
31 the orbit nevertheless controls storminess by acting through summer. When perihelion falls in a
32 hemisphere's summer, that summer has fewer mid-latitude storms and the storm belt sits farther
33 poleward, and the two hemispheres evolve in antiphase through the precession cycle. Simulations
34 of the Last Interglacial, when perihelion falls in northern summer, reproduce this pattern. This
35 tells researchers where to look for orbital storm signals in geological records, namely in archives
36 sensitive to summer, not to the yearly average.

37 1. Introduction

38 Extratropical cyclones govern mid-latitude weather systems. They transport the majority of
39 poleward heat and moisture outside the tropics, supply a substantial fraction of mid-latitude pre-
40 cipitation, and aggregate into the canonical storm-track pathways of the North Atlantic, North
41 Pacific, and Southern Ocean (Hoskins and Valdes 1990; Chang et al. 2002; Hoskins and Hodges
42 2002, 2005). These cyclones develop via baroclinic instability driven by the meridional tempera-
43 ture gradient (Eady 1949), a gradient that reaches its maximum during winter. Cyclone activities
44 consequently peak in the cold season, with consistent wintertime maxima (e.g., frequency) evi-
45 dent in both the Northern Hemisphere (Hoskins and Hodges 2002) and the Southern Hemisphere
46 (Simmonds and Keay 2000). Mid-latitude ~~cyclones are therefore most favorable~~ cyclogenesis is
47 therefore most favored during winter.

48 Precession, by contrast, exerts its primary forcing on warm seasons. Slow variations in the longi-
49 tude of perihelion represent the leading orbital control on seasonal insolation cycles. It redistributes
50 incoming solar radiation, and the amplitude of the anomalies scales with a season's background
51 solar radiation (Berger 1978; Laskar et al. 2004). Mid-latitude summertime insolation fluctuates
52 by over 100 W m^{-2} across a full precession cycle, whereas midwinter insolation experiences only
53 minor orbital-driven changes. This stark contrast explains why summer solar forcing serves as
54 the most impactful component of climate changes (Huybers 2006). Since extratropical cyclone
55 activity is inherently winter-favored, this invites a simple expectation that precession should exert

negligible control over mid-latitude storm tracks. But this expectation has never been put to a clean test.

Previous studies have looked closely at precessional signals, and they find seasonal shifts in climate. This is well established in monsoon research, from the classic orbital experiments of Kutzbach and Guetter (1986) to single-forcing designs that isolate precession explicitly (Bosmans et al. 2015; Merlis et al. 2013), and cave records show the northern and southern monsoons swinging in antiphase as perihelion migrates through the year (Wang et al. 2008). The extratropics have received far less attention. Kaspar et al. (2007) simulated a stronger and poleward-shifted North Atlantic winter storm track for the Eemian from two orbital time slices. Brayshaw et al. (2010) linked Holocene orbital shifts to North Atlantic and Mediterranean winter storm tracks via changes in baroclinicity. And Varma et al. (2012) found the Southern Hemisphere westerlies shift under Holocene orbital forcing. Explicit cyclone tracking has entered paleoclimate work mainly at the Last Glacial Maximum, where ice sheets dominate the forcing (Pinto and Ludwig 2020; Raible et al. 2021), and in transient simulations of the last millennium (Hess and Chemke 2025). For tropical cyclones the precessional response has recently been examined in high-resolution paleoclimate simulations, where it is governed by atmospheric moisture (Raavi et al. 2023). Here our study provides the extratropical counterpart, where cyclone growth is baroclinic. To our knowledge, no previous study has applied explicit cyclone tracking across a full precession cycle to isolate the orbital response of the extratropical storm tracks, and none has asked which season carries the response.

Many paleoclimate archives that record past storm activity, such as wind-blown dust and sediment, or rainfall linked to westerly winds, only capture signals from specific seasons instead of annual averages. If a proxy record mainly reflects one single season, it can either hide or falsely create an orbital signal, depending on which time of year carries the strongest response (Laepfle et al. 2011). Fully understanding which season holds the clearest precessional storm signal is therefore essential for properly interpreting these geological records. Moreover, different metrics for cyclone behavior, e.g., total storm counts, how much rain each storm produces, and storm intensity, react differently to orbital forcing. We must evaluate these quantities separately before drawing conclusions from any single geological proxy.

85 To fill this gap, we run 12 equilibrium climate simulations covering a full precession cycle. Every
86 extratropical cyclone in both hemispheres is tracked individually using 6-hour sea level pressure.
87 Our experimental design isolates precession at amplified eccentricity with all other boundary
88 conditions fixed, so that the seasonal storm response to precession forcing alone can be examined.
89 The results are then tested in ~~simulationso~~simulations of real orbital states, ~~such as namely~~ PI,
90 MH, and LIG. Section 2 describes our model and simulation, and the method we use for storm
91 tracking and analysis. Section 3 presents the results. We discuss and conclude in section 4 and 5
92 respectively.

93 2. Methodology

94 *a. Model and experiments*

95 The simulations are performed with the coupled Earth system model AWI-ESM2 (Sidorenko
96 et al. 2019). The atmosphere component is ECHAM6 at T63 resolution with 47 vertical levels
97 (Stevens et al. 2013), which includes the land surface module JSBACH (Reick et al. 2021). The
98 ocean and sea ice component is FESOM2, a finite-volume model formulated on an unstructured
99 mesh (Danilov et al. 2017), with a spatially varying resolution of from 25 km over high latitudes
100 and coastal regions, to 150 km over open oceans.

101 We analyze 12 equilibrium simulations, taken from the idealized precession ensemble introduced
102 by Shi et al. (2025). The longitude of perihelion ω is prescribed at values from 0° to 330° in steps of
103 30° (labeled as p000, p030..., and p330), while eccentricity is fixed at 0.058 and obliquity at 24.5° .
104 The eccentricity value sits near the upper bound reached during the Quaternary and amplifies the
105 precessional effects on insolation. All remaining boundary conditions follow the pre-industrial
106 setup, and each phase is branched from a long pre-industrial control. We analyze 30 years of
107 6-hourly output from the equilibrated part of each simulation. The calendar month of perihelion
108 for each phase is computed from orbital geometry (Berger 1978), which provides the insolation
109 fields shown in Fig. 1c-d.

110 To test our results under realistic orbital configurations, we additionally conducted three equilib-
111 rium simulations including a PI (0 ka) control run, a MH (6 ka) experiment, and a LIG (127 ka)
112 experiment. Boundary conditions follow the protocols of PMIP4 (Otto-Bliesner et al. 2017). Or-
113 bital parameters are calculated following Berger (1977), and the greenhouse gas concentrations are

114 taken from multi-archive reconstructions from the Greenland and the Antarctic ice cores (Flückiger
 115 et al. 2002; Monnin et al. 2004; Schneider et al. 2013; Schilt et al. 2010; Buiron et al. 2011). Each
 116 simulation is integrated for 1,000 model years with the trend of simulated global mean surface
 117 temperature in the final 100 model years not exceeding ± 0.05 K/century. We preserved 6-hourly
 118 sea level pressure from each simulation. At 127 ka, perihelion occurs close to the June solstice with
 119 an eccentricity of 0.039378; at 6 ka, perihelion falls within September, while the pre-industrial
 120 control features perihelion in early January.

121 *b. Converting ~~fix-day~~ fixed-day calendar into angular calendar*

122 Precession moves perihelion through the seasonal cycle, and Kepler’s second law (Kepler 1953)
 123 then changes the lengths of the seasons (Fig. 1f-g). This is the calendar effect long known in
 124 paleoclimate modeling (Joussaume and Braconnot 1997; Bartlein and Shafer 2019; Shi et al.
 125 2021). ~~And-month/~~Monthly and seasonal averages at different orbital configurations are therefore
 126 only comparable when the ~~months/seasons~~ averaging windows follow the same position of the
 127 Earth along its orbit.

128 Following Shi et al. (2021), we compute the length of a month (season) by calculating how much
 129 time the Earth needs to move from the respective starting point to the endpoint. We first define the
 130 mean anomaly M as the angle between the perihelion and Earth’s position based on the assumption
 131 that the orbit describes a perfect circle with the sun at the center by:

$$132 \quad M = \frac{2\pi}{T} \cdot t_p \quad (1)$$

133 Here, t_p denotes the time elapsed since Earth passes the perihelion and T is the Earth’s revolution
 134 period, namely 1 year or 365 days. Taking into account the orbit’s eccentricity ϵ , we define the
 135 eccentric anomaly E via:

$$136 \quad E - \epsilon \cdot \sin(E) = M \quad (2)$$

137 Equation 2 can be solved with the application of Newton’s method (Danby and Burkardt 1983),
 138 and ~~E can be found using the following expression~~ E follows from the true anomaly θ , the angle at
 139 the Sun between perihelion and the actual position of the Earth (see Eq. 3.13b of Curtis 2014):

$$E = 2 \cdot \arctan\left(\sqrt{\frac{1-\epsilon}{1+\epsilon}} \cdot \tan\left(\frac{\theta}{2}\right)\right) \quad (3)$$

The above equations ~~implicitly relates~~ relate t_p to θ by:

$$t_p(\theta) = \frac{MT}{2\pi} = \frac{(E - \epsilon \cdot \sin(E))T}{2\pi} \quad (4)$$

This allows us to define seasons with respect to Earth's position on the orbit rather than relying on a fixed number of days as used in our modern calendar. The insolation panels of Fig. 1 are computed on this angular calendar. Re-binning the 6-hourly cyclone statistics onto the same calendar changes the local-summer response amplitudes by at most 1.5 percentage points and the phase-locking lags by less than half a month, while the autumn response strengthens slightly. All storm diagnostics are therefore reported on the model's own fixed calendar.

c. Cyclone detection

Extratropical cyclones are detected and tracked with the algorithm of Crawford et al. (2021), and Fig. 2 summarizes the procedure. 6-hourly sea level pressure is first interpolated onto 100-km equal-area grids centered on each pole, and the two hemispheres are processed separately. A grid cell qualifies as a cyclone center when its pressure p_c lies below that of all 8 neighbors and the surrounding field keeps a sufficient inward gradient. The gradient test compares p_c with the mean pressure $\bar{p}(d)$ on a one-cell ring at distance d from the candidate,

$$\frac{\bar{p}(d) - p_c}{d} \geq \gamma \quad (5)$$

where $d = 1000$ km and $\gamma = 750 \text{ Pa} (1000 \text{ km})^{-1}$, and candidates over terrain above 1500 m are discarded. Centers closer than 1000 km are treated as one multicenter system. The cyclone area A is obtained by growing closed isobars around each center at a 200-Pa interval until a contour fails to close, and the last closed isobar defines the edge pressure p_e (Fig. 2b). Each cyclone carries the depth and the area-equivalent radius

$$D = p_e - p_c \quad (6)$$

$$R = \sqrt{A/\pi} \quad (7)$$

Tracking links the centers of consecutive 6-hourly fields (Fig. 2c). For every active cyclone a first-guess position is projected from its previous displacement, damped to three quarters,

$$\mathbf{x}_q = \mathbf{x}_t + 0.75 (\mathbf{x}_t - \mathbf{x}_{t-1}) \quad (8)$$

and a center of the next field qualifies as the continuation when it lies within $v_{\max}\Delta t$ of both the current position $\mathbf{x}_t$ and the projection $\mathbf{x}_q$, with $v_{\max} = 150 \text{ km h}^{-1}$ and $\Delta t = 6 \text{ h}$, a searching radius of 900 km per step. Among the qualifying candidates the one nearest to the projection continues the track. Centers left unmatched at the new time start tracks as genesis events, and tracks left unmatched end as lysis. Only systems poleward of 30° latitude enter the analysis.

Several track-based diagnostics are derived. Track density counts every 6-hourly cyclone position, system density counts each cyclone once at its mean position, and genesis density counts the first point of each track. For every cyclone the lifespan, the maximum depth relative to the outermost closed isobar, the minimum central pressure, the mean cyclone area, and the propagation speed are recorded. We further define a "storm day" field, representing the number of days per month during which a grid point lies inside a detected cyclone. It therefore measures the storm-influence time of each grid point, and it is affected by most cyclone metrics, such as the number of cyclones, their area, lifespan, and propagation speed.

d. Baroclinicity diagnostics

The maximum Eady growth rate serves as the measure of baroclinicity (Eady 1949; Lindzen and Farrell 1980; Hoskins and Valdes 1990). It is computed from wind, temperature, and geopotential height in the 925-700 hPa layer as

$$\sigma = 0.31 |f| \frac{|\partial U / \partial z|}{N}, \quad (9)$$

180 where f is the Coriolis parameter, $|\partial U/\partial z|$ the vertical wind shear, and N the Brunt-Väisälä
 181 frequency. The response to precession is quantified between the end members p090 and p270,
 182 which place perihelion at the June and December solstices.

183 To separate the roles of vertical shear and stratification we follow the substitution approach of
 184 Camargo et al. (2007). Writing $\sigma = 0.31 |f| S/N$ with $S = |\partial U/\partial z|$, the difference is split by
 185 changing one factor at a time while holding the other at its p270 value,

$$\Delta\sigma_S = 0.31 |f| \frac{S_{90} - S_{270}}{N_{270}}, \quad (10)$$

$$\Delta\sigma_N = 0.31 |f| S_{270} \left(\frac{1}{N_{90}} - \frac{1}{N_{270}} \right), \quad (11)$$

187 so that $\Delta\sigma = \Delta\sigma_S + \Delta\sigma_N + \varepsilon$, with a small cross term ε .

188 *e. Basins and statistics*

189 Three storm-track basins are analyzed, the North Atlantic (30–75°N, 80°W–20°E), the North
 190 Pacific (30–75°N, 120°E–120°W), and the Southern Ocean (40–75°S, all longitudes). Basin means
 191 are area weighted. Uncertainty estimates use the interannual spread of the 30 individual years in
 192 each phase, and the significance of the end-member difference maps is assessed with a two-sided
 193 Welch t test (Welch 1947) at each grid point.

194 **3. Results**

195 *a. Summer sensitivity*

196 The 12 precession-phase annual mid-latitude storm climatologies exhibit only minor differences,
 197 and phase anomalies relative to the multi-phase ensemble mean manifest as weak, slowly evolving
 198 dipoles aligned along major storm track pathways (Fig. 3). The basin-averaged time series quantify
 199 the muted amplitude of this annual-mean signal, with across-phase ranges of 10%, 17% and 4% in
 200 the North Atlantic, the North Pacific and the Southern Ocean, varying smoothly with ω . We also
 201 observe an interhemispheric antiphase.

202 There is also a pronounced seasonal cycle. Extratropical cyclone activities peak in DJF in North-
 203 ern Hemisphere basins and JJA across the Southern Ocean, with this seasonal timing unchanged

across all orbital setups (Fig. 4a–c). Cold-season cyclones also carry greater intensity, with winter peak cyclone depths exceeding summertime values by 70–100 % within the Northern Hemisphere storm basins (not shown). Overall, winter dominates mid-latitude cyclone activity uniformly across hemispheres and all orbital phases.

Precession exerts negligible control over annual-mean storm conditions but profoundly reshapes seasonal cyclone variability, and its strongest modulation targets the low-activity warm season. Though each precession phase retains the same wintertime peak in basin-averaged storm track density, cross-phase deviations emerge almost entirely within each basin’s local summer (Fig. 4a–c). Anomalies between end-members (p090 and p270) quantify this stark seasonal contrast (Fig. 4d–f). For the North Pacific, ~~the anomaly of track density decreases by 7.1~~ track density drops by 0.10 cell⁻¹ month⁻¹ in boreal summer (JJA), about 44% of the phase mean ~~on an annual basis, and by -23% in boreal summer (JJA), against an annual-mean decrease of only 7%.~~ The North Atlantic follows ~~an identical the same~~ pattern, with ~~annual-mean decreased by -10% versus~~ a summertime decrease by ~~roughly 40% near 23% versus 5% annually.~~ The Southern Ocean mirrors this behavior ~~but shifts the sensitive season to in~~ its local summer (DJF), ~~with an annual increase by 3.7% compared to a 20% summer signal~~ where track density rises by 19%, against a 2% annual increase. All three basins identify local summer as the most orbitally sensitive season. Moreover, summer and winter vary in antiphase across the precession cycle, with perihelion suppressing the summer storms while slightly enhancing the winter ones. Therefore the annual signal is muted.

This summertime storm response is locked to orbital geometry. When basin-averaged track-density anomalies are plotted against both calendar month and precession angle ω , the band of peak anomalies traces a diagonal ridge aligned with the calendar timing of perihelion across the full phase space (Fig. 4g–i), with a delay of 1–2 months. Summers spent near perihelion exhibit suppressed storm activity, whereas aphelion summers feature intensified storminess. It should be noted that northern summer perihelion aligns with southern summer aphelion, precessional forcing thus drives antiphased mid-latitude storm signals in the two hemispheres.

The pronounced seasonal dependence of orbital storm signals originates from insolation forcing. Averaged across the Northern Hemisphere storm-track band (40–70°N), precessional insolation variability peaks at 105 W m⁻² during boreal summer (JJA) and falls to merely 22 W m⁻² in boreal winter (DJF). The Southern Hemisphere storm band (40–70°S) exhibits mirror-symmetric

behavior, with a maximum seasonal forcing amplitude of 105 W m^{-2} in austral summer (DJF) versus just 22 W m^{-2} in austral winter (JJA) (Fig. 1e). The underlying mechanism lies in the multiplicative nature of precessional insolation modulation. Shifting perihelion across the annual cycle alters the Sun–Earth distance for each month, which rescales monthly incoming solar flux by a factor proportional to eccentricity e (approximately $4e$, equaling 23% for $e = 0.058$, Fig. 1a-b). Midwinter at mid-latitudes features minimal climatological insolation, leaving little baseline flux to be modulated by precessional shifts. Precession forcing therefore exerts its strongest modulation on local summer in each hemisphere, and barely perturbs deep winter conditions (Fig. 1a-e).

Orbital insolation anomalies are translated into extratropical storm activity via shifts in mid-latitude baroclinicity. Differences in maximum Eady growth rate between the extreme precession phases $\omega = 90^\circ$ and $\omega = 270^\circ$ reach their peak magnitude within each basin’s local summer (Fig. 5). Normalized against seasonal climatological means, summertime Eady growth rate responses hit 31 % in the North Atlantic, a pronounced 92 % in the North Pacific, and 21 % in the Southern Ocean. A decomposition analysis further reveals that vertical wind shear dominates the precessional Eady growth rate anomalies, while static stability contributes negligible signal strength. This is consistent across all storm tracks.

b. Meridional shift

Spatial difference maps between the two orbital end-members (p090 minus p270) yields weak responses for the annual means (Fig. 6a). By contrast, boreal summer (JJA) exhibits a prominent dipole pattern across Northern Hemisphere basins, with fewer storm days through the mid-latitude core and more along the Arctic flank at perihelion summer, the signature of a poleward shift joined to an overall weakening. In DJF the signal migrates to the Southern Ocean and forms a circumpolar banded structure of opposite sign, marking enhanced mid-latitude storm activity and an equatorward shift of the southern storm belt.

The latitudinal displacement of storm tracks driving these dipole anomalies is quantified in Fig. 7. For the North Pacific boreal summer, core storm latitude shifts from 53°N when perihelion falls in northern winter to 61°N when it falls in northern summer, yielding an 8° total poleward migration that peaks precisely at $\omega = 90^\circ$. The North Atlantic summertime storm track undergoes a 4.4° poleward shift following identical orbital behavior, while the Southern Ocean mirrors this response

with a 3.8° poleward displacement when perihelion coincides with austral summer (near p270). A perihelion summer therefore weakens storm activity and shifts the storm population poleward, consistent with the poleward shift of baroclinic activity evident in the Eady growth rate dipole (Fig. 5a).

By contrast, the annual-mean track latitude varies by no more than 1.3° in any basin. Winter storm-track latitudes only fluctuate by $1.5\text{--}2.8^\circ$ across the full precession cycle, and their latitudinal extrema do not align with solstitial orbital phases. This further confirms our earlier conclusion that mid-latitude winter storms are orbitally insensitive. Spring and autumn display intermediate latitudinal variability with a coherent phase dependence. For example, autumn poleward displacement maximizes near $\omega = 150^\circ$, the orbital configuration that places perihelion in late summer. Latitudinal poleward shifts therefore track the calendar timing of perihelion identically to Fig. 4g–i, with a uniform lag of 1–2 months, consistent with surface thermal inertia. Figure 8 quantifies this basin-dependent lag. All basin-scale storm extrema cluster along basin-specific dashed lines offset from the perihelion reference. The response lags the perihelion calendar by distinct intervals across regions, near 1 month in the North Pacific, 1–2 months in the North Atlantic, and slightly over 2 months in the Southern Ocean.

c. Cyclone precipitation

The hydrological signature of precessional forcing evolves in antiphase with storm dynamical responses (Fig. 9). During boreal summer over the North Pacific, mean precipitation per individual storm rises by as much as 15 % at the orbital phase where total storm frequency reaches its minimum, yielding nearly perfectly opposing variability between the two metrics. Perihelion summers feature warmer, more humid mid-latitude atmospheres, which amplifies precipitation intensity within the remaining cyclones. Nevertheless, the sharp reduction in storm number dominates the total signal, such that basin-wide summertime precipitation drops by 17 % between the two extreme orbital phases. The fractional share of precipitation originating from cyclones also declines markedly, with a full-cycle range of 9.5 % in North Pacific summer compared to just 3 % for the annual mean. Precessional forcing thus exerts two counteracting controls on extratropical hydroclimate simultaneously. It reduces total cyclone counts while boosting precipitation intensity per storm.

291 *d. Realistic paleosimulations (MH and LIG)*

292 Our results based on the 12 idealized simulations can be validated against the realistic orbital-
293 state simulations (PI, MH, and LIG) introduced in Section 2. In LIG, perihelion occurs near June
294 solstice, and the tracked cyclone responses fully reproduce our core theoretical pattern. Relative to
295 PI, summer storm track density decreases by 8% over the North Atlantic and 11% over the North
296 Pacific; winter cyclone changes remain within approximately 3%. Meanwhile, austral summer
297 storm track density across the Southern Ocean rises by 8%. Northern Hemisphere summer storm
298 tracks shift poleward by 1–2 degrees, while the Southern Hemisphere summer storm belt shifts
299 equatorward (Fig. 10). The summer difference composites show identical weakening and poleward
300 displacement of Northern Hemisphere storm tracks as seen in our amplified idealized end-member
301 simulations (Fig. 11). In MH, perihelion happens in September, leaving seasonal forcing dominated
302 by warm September conditions with cool March conditions. Consistent with this seasonal signal,
303 MH storm variability features fewer, poleward-shifted cyclones in autumn and the opposite trend
304 in spring. The magnitude of storm responses in these paleosimulations is much weaker than that
305 of our amplified end-member contrasts (p090 vs. p270, Fig. 6), consistent with their smaller
306 eccentricity values of the real orbits. Notably, these experiments combine realistic precession with
307 their own obliquity and greenhouse gases, so they test our theoretical prediction under multi-factor
308 natural orbital forcing rather than isolating precessional effects.

309 **4. Discussion**

310 The seasonal preference seen in storm responses originates purely from orbital geometry. Pre-
311 cession redistributes insolation in proportion to the insolation a season already has, meaning winter
312 hemispheres have very little background insolation for orbital changes to modify (Berger 1978).
313 Fig. 1 clearly illustrates this stark seasonal contrast in forcing strength. Winter insolation varies
314 by only 22 W m^{-2} across the precession cycle, while summer values reach 105 W m^{-2} within
315 mid-latitude storm-track latitudes. This is model independent and will hold in any climate models
316 as it simply reflects the orbital geometry behind the well-known summer-dominated orbital forcing
317 framework (Huybers 2006). Existing monsoon studies has long confirmed that summer bears
318 the strongest precessional signal (Kutzbach and Guetter 1986; Bosmans et al. 2015). But the new
319 insight of our work lies elsewhere. This summer-focused orbital forcing acts on a winter-dominated

320 weather system, reshaping storm activity exclusively during its normally quiet warm season. Note
321 that the term "quiet" here refers to the weaker overall intensity of the summer storm track, not its
322 population. Both hemisphere still hosts abundant storm systems in local summer, and they are
323 merely weaker, shallower and move more slowly compared to winter cyclones.

324 The mechanism that converts summer insolation into summer storms is the baroclinic instability.
325 Our factor decomposition analysis confirms that the shear term carries essentially the whole Eady
326 growth rate response in all three ocean basins, while static stability plays a negligible role. We
327 see similar physical mechanism operating in the modern Northern Hemisphere, where greenhouse
328 warming takes the place of perihelion-driven heating. Stronger summer warming at high latitudes
329 has weakened summertime large-scale circulation, eddy kinetic energy, and the number of strong
330 summer cyclones (Coumou et al. 2015; Chang et al. 2016; Gertler and O’Gorman 2019). Future
331 climate projections show the same shear-based decline of Northern Hemisphere summer cyclone
332 activity and baroclinic energy as globe warms (Lehmann et al. 2014; Priestley and Catto 2022).
333 But a difference should be noted here, as greenhouse warming is a transient forcing that both
334 hemispheres change in phase, whereas our simulations are equilibrated and precession swings both
335 hemispheres in antiphase. Even so, both forcings follow one unified rule that is the core theme of
336 this study: cooler summers host more storm activity.

337 Compared to summer, storm systems in winter are much more stable. Across all three ocean
338 basins, winter storm activity changes by less than 10% over a full precession cycle, even though
339 winter Eady growth rates shift by 16–18% between the two orbital end-members. Weak winter
340 insolation forcing cannot fully explain this stability. The winter storm track appears saturated
341 in the sense that ample baroclinicity is available in every phase, so that moderate changes in the
342 growth rate do not translate into changes in occurrence. Transient simulation of the last millennium
343 likewise found winter cyclone nearly insensitive to external forcing (Raible et al. 2018; Hess and
344 Chemke 2025), and those forcings acted to affect storm track in summer (Hess and Chemke 2025).
345 Our findings expand this conclusion to the far larger swings in solar radiation driven by precession,
346 and this insensitivity only holds for wintertime conditions.

347 Geological records sensitive to summertime conditions within mid-latitude storm belts should
348 preserve prominent precession-scale storm signals, whereas proxies that average climate signals
349 across winter or the full year will show only minor orbital imprint. Just as observed for tem-

perature reconstructions, the seasonal bias of a given archive can either generate artificial orbital storm signals or fully obscure true precessional variations (Laepple et al. 2011). Furthermore, summer-sensitive storm records from both hemispheres will fluctuate oppositely over precessional cycles, forming an extratropical equivalent of the well-documented monsoon seesaw (Wang et al. 2008). Current published sediment records already support the feasibility of testing this framework. Distinct precession-period signals have been detected in Northern Hemisphere mid-latitude westerly proxies (Li et al. 2019), as well as North Pacific dust records. Precessional variation appears specifically in moisture-sensitive dust fractions (Zhong et al. 2024). Meanwhile, dust intensity archives that capture winter-only signals display very weak precessional power (Chen et al. 2014). Regarding the Southern Hemisphere, subtropical margins of the southern westerly belt exhibit clear precession cycles (Lamy et al. 2019), consistent with our results. Records of the southern westerlies also show orbital-band migrations of the belt itself (Lamy et al. 2010; Varma et al. 2012), and glacial-state cyclone modeling has shown how strongly reconstructed storminess depends on which cyclone property a proxy senses (Pinto and Ludwig 2020).

Our study is based on idealized simulations designed to isolate pure precessional forcing. The experimental setup holds eccentricity, obliquity, ice sheets, and greenhouse gas concentrations fixed and does not incorporate transient climate evolution or orbital cross-interactions. As such, our results represent dynamical responses to idealized orbital conditions rather than reconstructions of specific geological epochs. Future work could incorporate time-varying boundary conditions and full orbital combinations to further generalize the storm seasonal response across broader paleoclimate backgrounds.

5. Conclusions

In our study, we carry out a set of equilibrium climate simulations covering a full precession cycle, with explicit tracking of all extratropical cyclones across both hemispheres to isolate precessional forcing. We find that precession regulates extratropical cyclone primarily through local summer, the season with naturally low storm activity, while winter storm tracks has only minor change across all precession phases. Storm activity calms down when perihelion falls in local summer and intensifies when aphelion coincides with local summer. And, as perihelion summer in one hemisphere matches aphelion summer in the other, mid-latitude storminess of the two hemispheres

379 varies in antiphase during a precession cycle. This orbital signal propagates through shear-driven
380 baroclinicity and expresses itself in cyclones numbers and position, both of which are phase locked
381 to the calendar month of perihelion, with a delay of 1-2 months. Our result also bring clear
382 implications for paleoclimate proxy interpretation. Orbital storm signals should be extracted from
383 mid-latitude archives sensitive to summer climate, and the recorded signals will display opposite
384 variations between the two hemispheres.




FIG. 1. (a-b) Range of monthly mean insolation across the 12 precession phases for the 40–70°N and 40–70°S bands, computed on the classical fixed-day calendar (red) and on the angular calendar (blue). The dashed line marks the distance factor $4e/(1-e^2)^2$ multiplied by the monthly mean insolation. (c-d) Insolation anomaly as a function of calendar month and ω on the angular calendar, with the perihelion (red) and aphelion (blue) months overlaid. (e) Insolation range in the two bands for each season on the angular calendar, computed as maximum minus minimum insolation across the 12 phases. Units: W m^{-2} . (f),(g) The orbital geometry behind the calendar drift, shown for two precession end-members (i.e., p090 and p270). The equinox and solstice lines through the Sun divide the orbit into the four seasons. Seasons on the perihelion side last ~~84.5~~about 85 days and seasons on the aphelion side ~~last 98.0~~98 days, and this nearly two-week contrast swaps sides between the two phases. Day counts ~~are exact~~hold for $e = 0.058$ with a 365-day year, while the drawn eccentricity is exaggerated for clarity.




FIG. 2. Cyclone detection and tracking procedure. (a,c) One 6-hourly sea level pressure field of phase p000 (contours, 8-hPa interval) with all detected cyclone centers poleward of 30°N (dots) and their area-equivalent radii $R = \sqrt{A/\pi}$ (circles), 15 January of one model year. (b,e) The deepest system of that field, with the outermost closed isobar p_e (dashed), the cyclone center (cross), the central and edge pressures and depth annotated, and the binary mask cells (shading) defined by $\text{SLP} < p_e$ inside the $\sqrt{A}$ search window; contour interval 4 hPa. (c,f) All cyclone tracks of the same month (lines) with genesis points (dots). (d) Data processing workflow.




FIG. 3. Annual-mean storm days across the precession cycle. (a)–(l) Annual storm-day anomaly of each phase relative to the cross-phase mean (shading), with the phase-mean climatology outlined at 5 and 8 d month⁻¹ (contours). (m)–(o) Basin-mean annual storm days against ω with interannual standard errors; the across-phase range as a percentage of the phase mean is annotated in each panel.




FIG. 4. Seasonal concentration of the precessional storm response, from the Lagrangian tracks. (a)–(c) Basin-mean track density ($\text{cell}^{-1} \text{ month}^{-1}$) by calendar month for the North Atlantic, the North Pacific, and the Southern Ocean, one line per phase colored by ω , with local summer shaded. (d)–(f) **Anomaly of track density ($\text{p090} - \text{p270}$, percent of Track-density difference between the phase-mean end members p090 and p270 ($\text{cell}^{-1} \text{ month}^{-1}$))** for the annual mean and the 4-four seasons, with the local-summer bar highlighted; **open-symbols give the corresponding ranges of system and genesis densities**. (g)–(i) Basin-mean track-density anomaly relative to the phase mean as a function of calendar month and precession phase (unsmoothed), with the perihelion (red) and aphelion (blue) calendar months of each phase overlaid.




FIG. 5. Precessional response of the maximum Eady growth rate (925–700 hPa). (a) p090 minus p270 difference for JJA and (b) for DJF (day^{-1}). (c) Seasonal response amplitude in each basin, the RMS of the p090 minus p270 difference over the storm box as a percentage of the seasonal climatology, with stars marking local summer. (d) Projection shares of the vertical-shear and static-stability factors in the local-summer difference pattern.




FIG. 6. Difference in storm days (d month^{-1}) between p090 and p270 for (a) the annual mean, (b) JJA, and (c) DJF. Stippling marks grid points where the difference is significant at the 95% level based on a two-sided Welch t test.




FIG. 7. Storm-track latitude across the precession cycle, the storm-day-weighted mean latitude over each basin sector shown as the poleward anomaly from each season's phase mean, for (a) the North Atlantic, (b) the North Pacific, and (c) the Southern Ocean. Lines give the annual mean and the four seasons, with the local summer of each basin drawn heavy with markers. Vertical dotted lines mark the phase with perihelion in the local summer of each basin.




FIG. 8. Phase locking of the storm response to the perihelion calendar. For each calendar month and basin, the phase of the first harmonic across the 12 precession phases of (a) basin-mean storm days, shown as the ω of the activity minimum, and (b) the storm-day-weighted track latitude, shown as the ω of the poleward maximum. Marker size scales with the harmonic amplitude. The red line gives the ω that places perihelion in that calendar month, and dashed lines repeat it delayed by each basin's lag over the amplitude-strong months (annotated in months).




FIG. 9. Storm-attributed precipitation across the precession cycle. (a) North Pacific JJA percent anomalies from the phase mean of basin-mean storm days, per-storm precipitation, and total precipitation; the dotted line marks the phase with perihelion in NH summer. (b) Across-phase range of the storm-attributed share of precipitation, accumulated precipitation inside storm footprints divided by total accumulated precipitation (percentage points), for the annual mean and local summer in each basin.




FIG. 10. Seasonal storm response of the real-orbit experiments, tracked with the same pipeline as the main ensemble and referenced to the preindustrial control. (a)–(c) Anomalies of basin-mean track density for the mid-Holocene (gold) and the Last Interglacial (red), as percentages of the preindustrial mean, for the annual mean and the four seasons, with error bars giving twice the standard error of the 30-year difference and the local summer shaded. Track density is normalized per T63 grid cell, and the annotated preindustrial values give the absolute scale in the units of Fig. 4. (d)–(f) The corresponding change in the track-point mean latitude, positive poleward.




FIG. 11. Storm-day anomalies of the real-orbit experiments relative to the preindustrial control, from the same storm-day product as the main ensemble. (a)–(d) Mid-Holocene minus preindustrial for DJF, MAM, JJA, and SON. (e)–(h) Last Interglacial minus preindustrial. Contours outline the preindustrial climatology at 5 and 8 d month^{-1} . The isolated warm-season maxima over subtropical continental interiors reflect stationary surface heat lows rather than traveling cyclones.

512 *Acknowledgments.* The simulations were performed at the German Climate Computing Center
513 (DKRZ). [TODO funding and grant numbers before submission.]

514 *Data availability statement.* The AWI-ESM simulations of the idealized precession ensemble
515 are described in [?Shi et al. \(2025\)](#). The cyclone-track database, the gridded storm statistics, and
516 the analysis scripts underlying all figures will be archived and assigned a DOI upon acceptance.
517 [TODO repository DOI.]

References

- Bartlein, P. J., and S. L. Shafer, 2019: Paleo calendar-effect adjustments in time-slice and transient climate-model simulations (paleocaladjust v1. 0): Impact and strategies for data analysis. *Geoscientific Model Development*, **12** (9), 3889–3913.
- Berger, A., 1977: Long-term variations of the earth’s orbital elements. *Celestial Mechanics*, **15** (1), 53–74.
- Berger, A., 1978: Long-term variations of daily insolation and quaternary climatic changes. *Journal of Atmospheric Sciences*, **35** (12), 2362–2367.
- Bosmans, J., S. Drijfhout, E. Tuenter, F. Hilgen, and L. Lourens, 2015: Response of the north african summer monsoon to precession and obliquity forcings in the ec-earth gcm. *Climate dynamics*, **44** (1), 279–297.
- Brayshaw, D. J., B. Hoskins, and E. Black, 2010: Some physical drivers of changes in the winter storm tracks over the north atlantic and mediterranean during the holocene. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, **368** (1931), 5185–5223.
- Buiron, D., and Coauthors, 2011: Taldice-1 age scale of the talos dome deep ice core, east antarctica. *Climate of the Past*, **7** (1), 1–16.
- Camargo, S. J., K. A. Emanuel, and A. H. Sobel, 2007: Use of a genesis potential index to diagnose enso effects on tropical cyclone genesis. *Journal of Climate*, **20** (19), 4819–4834.
- Chang, E. K., S. Lee, and K. L. Swanson, 2002: Storm track dynamics. *Journal of climate*, **15** (16), 2163–2183.
- Chang, E. K., C.-G. Ma, C. Zheng, and A. M. Yau, 2016: Observed and projected decrease in northern hemisphere extratropical cyclone activity in summer and its impacts on maximum temperature. *Geophysical Research Letters*, **43** (5), 2200–2208.
- Chen, J., X. Liu, and V. A. Kravchinsky, 2014: Response of the high-resolution chinese loess grain size record to the 50° n integrated winter insolation during the last 500,000 years. *Geophysical Research Letters*, **41** (17), 6244–6251.

545 Coumou, D., J. Lehmann, and J. Beckmann, 2015: The weakening summer circulation in the
546 northern hemisphere mid-latitudes. *Science*, **348** (6232), 324–327.

547 Crawford, A. D., E. A. Schreiber, N. Sommer, M. C. Serreze, J. C. Stroeve, and D. G. Barber,
548 2021: Sensitivity of northern hemisphere cyclone detection and tracking results to fine spatial
549 and temporal resolution using era5. *Monthly Weather Review*, **149** (8), 2581–2598.

550 Curtis, H., 2014: Orbital position as a function of time. *Orbital Mechanics for Engineering*
551 *Students, 3rd edition, Elsevier, Amsterdam, the Netherlands*, 145–186.

552 Danby, J., and T. Burkardt, 1983: The solution of kepler’s equation, i. *Celestial Mechanics*, **31** (2),
553 95–107.

554 Danilov, S., D. Sidorenko, Q. Wang, and T. Jung, 2017: The finite-volume sea ice–ocean model
555 (fesom2). *Geoscientific Model Development*, **10** (2), 765–789.

556 Eady, E. T., 1949: Long waves and cyclone waves. *Tellus*, **1** (3), 33–52.

557 Flückiger, J., E. Monnin, B. Stauffer, J. Schwander, T. F. Stocker, J. Chappellaz, D. Raynaud, and
558 J.-M. Barnola, 2002: High-resolution holocene n2o ice core record and its relationship with ch4
559 and co2. *Global Biogeochemical Cycles*, **16** (1), 10–1.

560 Gertler, C. G., and P. A. O’Gorman, 2019: Changing available energy for extratropical cyclones and
561 associated convection in northern hemisphere summer. *Proceedings of the National Academy of*
562 *Sciences*, **116** (10), 4105–4110.

563 Hess, O., and R. Chemke, 2025: Forced changes in the north atlantic storm track over the last
564 millennium. *Geophysical Research Letters*, **52** (15), e2025GL115 894.

565 Hoskins, B. J., and K. I. Hodges, 2002: New perspectives on the northern hemisphere winter storm
566 tracks. *Journal of the Atmospheric Sciences*, **59** (6), 1041–1061.

567 Hoskins, B. J., and K. I. Hodges, 2005: A new perspective on southern hemisphere storm tracks.
568 *Journal of climate*, **18** (20), 4108–4129.

569 Hoskins, B. J., and P. J. Valdes, 1990: On the existence of storm-tracks. *Journal of Atmospheric*
570 *Sciences*, **47** (15), 1854–1864.

- Huybers, P., 2006: Early pleistocene glacial cycles and the integrated summer insolation forcing. *Science*, **313** (5786), 508–511.
- Joussaume, S., and P. Braconnot, 1997: Sensitivity of paleoclimate simulation results to season definitions. *Journal of Geophysical Research: Atmospheres*, **102** (D2), 1943–1956.
- Kaspar, F., T. Spanghel, and U. Cubasch, 2007: Northern hemisphere winter storm tracks of the eemian interglacial and the last glacial inception. *Climate of the Past*, **3** (2), 181–192.
- Kepler, J., 1953: *Epitome astronomiae copernicanae*. Jo. Plancus.
- Kutzbach, J. E., and P. J. Guetter, 1986: The influence of changing orbital parameters and surface boundary conditions on climate simulations for the past 18 000 years. *Journal of atmospheric sciences*, **43** (16), 1726–1759.
- Laepple, T., M. Werner, and G. Lohmann, 2011: Synchronicity of antarctic temperatures and local solar insolation on orbital timescales. *Nature*, **471** (7336), 91–94.
- Lamy, F., R. Kilian, H. W. Arz, J.-P. Francois, J. Kaiser, M. Prange, and T. Steinke, 2010: Holocene changes in the position and intensity of the southern westerly wind belt. *Nature Geoscience*, **3** (10), 695–699.
- Lamy, F., and Coauthors, 2019: Precession modulation of the south pacific westerly wind belt over the past million years. *Proceedings of the National Academy of Sciences*, **116** (47), 23 455–23 460.
- Laskar, J., P. Robutel, F. Joutel, M. Gastineau, A. C. Correia, and B. Levrard, 2004: A long-term numerical solution for the insolation quantities of the earth. *Astronomy & Astrophysics*, **428** (1), 261–285.
- Lehmann, J., D. Coumou, K. Frieler, A. V. Eliseev, and A. Levermann, 2014: Future changes in extratropical storm tracks and baroclinicity under climate change. *Environmental Research Letters*, **9** (8), 084 002.
- Li, Y., Y. Song, Q. Yin, L. Han, and Y. Wang, 2019: Orbital and millennial northern mid-latitude westerlies over the last glacial period. *Climate Dynamics*, **53** (5), 3315–3324.

597 Lindzen, R., and B. Farrell, 1980: A simple approximate result for the maximum growth rate of
598 baroclinic instabilities. *Journal of Atmospheric Sciences*, **37** (7), 1648–1654.

599 Merlis, T. M., T. Schneider, S. Bordoni, and I. Eisenman, 2013: Hadley circulation response to
600 orbital precession. part i: Aquaplanets. *Journal of Climate*, **26** (3), 740–753.

601 Monnin, E., and Coauthors, 2004: Evidence for substantial accumulation rate variability in antarctica
602 during the holocene, through synchronization of co₂ in the taylor dome, dome c and dml ice
603 cores. *Earth and Planetary Science Letters*, **224** (1-2), 45–54.

604 Otto-Bliesner, B., and Coauthors, 2017: The pmip4 contribution to cmip6–part 2: Two interglacials,
605 scientific objective and experimental design for holocene and last interglacial simulations.
606

607 Pinto, J. G., and P. Ludwig, 2020: Extratropical cyclones over the north atlantic and western europe
608 during the last glacial maximum and implications for proxy interpretation. *Climate of the Past*,
609 **16** (2), 611–626.

610 Priestley, M. D., and J. L. Catto, 2022: Future changes in the extratropical storm tracks and cyclone
611 intensity, wind speed, and structure. *Weather and Climate Dynamics*, **3**, 337–360.

612 Raavi, P. H., J.-E. Chu, A. Timmermann, S.-S. Lee, and K. J. Walsh, 2023: Moisture control
613 of tropical cyclones in high-resolution simulations of paleoclimate and future climate. *Nature
614 Communications*, **14** (1), 6426.

615 Raible, C. C., M. Messmer, F. Lehner, T. F. Stocker, and R. Blender, 2018: Extratropical cyclone
616 statistics during the last millennium and the 21st century. *Climate of the Past*, **14** (10), 1499–1514.

617 Raible, C. C., J. G. Pinto, P. Ludwig, and M. Messmer, 2021: A review of past changes in
618 extratropical cyclones in the northern hemisphere and what can be learned for the future. *Wiley
619 Interdisciplinary Reviews: Climate Change*, **12** (1), e680.

620 Reick, C. H., and Coauthors, 2021: Jsbach 3-the land component of the mpi earth system model:
621 documentation of version 3.2.

Schilt, A., M. Baumgartner, T. Blunier, J. Schwander, R. Spahni, H. Fischer, and T. F. Stocker, 2010: Glacial–interglacial and millennial-scale variations in the atmospheric nitrous oxide concentration during the last 800,000 years. *Quaternary Science Reviews*, **29** (1-2), 182–192.

Schneider, R., J. Schmitt, P. Köhler, F. Joos, and H. Fischer, 2013: A reconstruction of atmospheric carbon dioxide and its stable carbon isotopic composition from the penultimate glacial maximum to the last glacial inception. *Climate of the Past*, **9** (6), 2507–2523.

Shi, X., M. Werner, H. Yang, Q. Gao, J. Liu, and G. Lohmann, 2025: Precession controls on climate and water isotope signals in northern africa. *Paleoceanography and Paleoclimatology*, **40** (1), e2024PA004 999.

Shi, X., and Coauthors, 2021: Calendar effects on surface air temperature and precipitation based on model-ensemble equilibrium and transient simulations from pmip4 and pacmedy. *Climate of the Past Discussions*, **2021**, 1–29.

Sidorenko, D., and Coauthors, 2019: Evaluation of fesom2. 0 coupled to echam6. 3: preindustrial and highresmpip simulations. *Journal of Advances in Modeling Earth Systems*, **11** (11), 3794–3815.

Simmonds, I., and K. Keay, 2000: Mean southern hemisphere extratropical cyclone behavior in the 40-year ncep–ncar reanalysis. *Journal of Climate*, **13** (5), 873–885.

Stevens, B., and Coauthors, 2013: Atmospheric component of the mpi-m earth system model: Echam6. *Journal of Advances in Modeling Earth Systems*, **5** (2), 146–172.

Varma, V., and Coauthors, 2012: Holocene evolution of the southern hemisphere westerly winds in transient simulations with global climate models. *Climate of the Past*, **8** (2), 391–402.

Wang, Y., and Coauthors, 2008: Millennial-and orbital-scale changes in the east asian monsoon over the past 224,000 years. *nature*, **451** (7182), 1090–1093.

Welch, B. L., 1947: The generalization of ‘student’s’ problem when several different population variances are involved. *Biometrika*, **34** (1-2), 28–35.

Zhong, Y., and Coauthors, 2024: Orbital controls on north pacific dust flux during the late quaternary. *Geophysical Research Letters*, **51** (4), e2023GL106 631.