

**Cool summers are stormy summers: precession reorganizes extratropical
cyclone activity through the quiet season**

Xiaoxu Shi^a

^a *Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research, Bremerhaven,
Germany*

Corresponding author: Xiaoxu Shi, shixiaoxu@sml-zhuhai.cn

7 ABSTRACT: Extratropical cyclones peak in winter, while orbital precession reshapes the seasonal
8 insolation cycle mainly in summer. Thus storm tracks might be expected to ignore the precession
9 cycle. This is examined with twelve equilibrium simulations using a coupled climate model AWI-
10 ESM2, spanning a full precession cycle at 30° longitude increments under fixed eccentricity and
11 obliquity. Extratropical cyclones are explicitly tracked based on 6-hourly atmospheric pressure
12 fields for both hemispheres. While the annual-mean cyclone response to precession is small,
13 substantial seasonal response emerges. In the North Atlantic, the North Pacific, and the Southern
14 Ocean the response focuses on local summer. We also find summers spent near perihelion feature
15 suppressed storminess, whereas summers near aphelion experience intensified cyclone activity,
16 yielding antiphase mid-latitude storm variability between the two hemispheres over the precession
17 cycle. This signal is carried by cyclone number and position, and it follows the pronounced
18 summer response of the Eady growth rate driven primarily by vertical wind shear in all three
19 basins. Precession therefore shapes extratropical storm tracks mostly through their quiet season
20 (i.e., summer). Summer-sensitive proxy archives should record a strong and hemispherically
21 antiphased storminess signal at the precession period that annually integrating archives would
22 miss. Moreover, monthly storm responses are phase-locked to the calendar timing of perihelion,
23 lagged by 1-2 months due to surface thermal inertia. This fingerprint also appears when we apply
24 the same cyclone tracking algorithm to pre-industrial (PI, 0 ka, perihelion falling in winter), Last
25 Interglacial (LIG, 127 ka, perihelion falling in summer) and mid-Holocene (MH, 6 ka, perihelion
26 falling in autumn), with weaker amplitudes matching their weaker eccentricity.

27 SIGNIFICANCE STATEMENT: Long orbital cycles drive global climate variability, and the
28 strongest of these, precession, reshapes sunlight mainly in summer, while mid-latitude storms
29 happen mainly in winter. This creates an apparent disconnect between precessional forcing and
30 storm track variability. Climate model simulations covering a full precession cycle show that
31 the orbit nevertheless controls storminess by acting through summer. When perihelion falls in a
32 hemisphere's summer, that summer has fewer mid-latitude storms and the storm belt sits farther
33 poleward, and the two hemispheres evolve in antiphase through the precession cycle. Simulations
34 of the Last Interglacial, when perihelion falls in northern summer, reproduce this pattern. This
35 tells researchers where to look for orbital storm signals in geological records, namely in archives
36 sensitive to summer, not to the yearly average.

37 1. Introduction

38 Extratropical cyclones govern ~~midlatitude~~mid-latitude weather systems. They trans-
39 port the majority of poleward heat and moisture outside the tropics, supply a substan-
40 tial fraction of ~~midlatitude~~mid-latitude precipitation, and aggregate into the canonical
41 storm-track pathways of the North Atlantic, North Pacific, and Southern Ocean ~~(????)~~
42 (Hoskins and Valdes 1990; Chang et al. 2002; Hoskins and Hodges 2002, 2005). These cyclones
43 develop via baroclinic instability driven by the meridional temperature gradient ~~(?)~~(Eady 1949), a
44 gradient that reaches its maximum ~~steepness~~ during winter. Cyclone activities consequently peak
45 in the cold season, with consistent wintertime maxima (e.g., frequency) evident in both the North-
46 ern ~~Hemispheres (?)~~Hemisphere (Hoskins and Hodges 2002) and the Southern Hemisphere ~~(?)~~
47 ~~Midlatitude~~(Simmonds and Keay 2000). Mid-latitude cyclogenesis is therefore most ~~favorable~~
48 favorable during winter.

49 Precession, by contrast, exerts its primary forcing on warm seasons. Slow variations in the
50 longitude of perihelion represent the leading orbital control on seasonal insolation cycles. It
51 redistributes incoming solar radiation, and the amplitude of ~~insolation~~the anomalies scales with a
52 season's background solar ~~irradiance (?)~~Midlatitude radiation (Berger 1978; Laskar et al. 2004)
53 Mid-latitude summertime insolation fluctuates by over 100 W m^{-2} across a full precession cycle,
54 whereas midwinter insolation experiences only minor ~~orbital modulation~~orbital-driven changes.
55 This stark contrast explains why ~~orbital climate frameworks single out summertime solar forcing~~

56 summer solar forcing serves as the most impactful component of climate changes ~~(?)~~(Huybers 2006)
57 . Since extratropical cyclone activity is inherently winter-favored, this invites a simple expectation
58 that precession should exert negligible control over ~~midlatitude~~mid-latitude storm tracks. But this
59 expectation has never been put to a clean test.

60 Previous studies have looked closely at precessional signals, and they find seasonal shifts in
61 climate. This is well established in monsoon research, from the classic orbital experiments
62 of ~~?~~Kutzbach and Guetter (1986) to single-forcing designs that isolate precession explicitly ~~(?)~~
63 (Bosmans et al. 2015; Merlis et al. 2013), and cave records show the northern and southern mon-
64 soons swinging in antiphase as perihelion migrates through the year ~~(?)~~(Wang et al. 2008). The
65 extratropics have received far less attention. ~~?~~Kaspar et al. (2007) simulated a stronger and
66 poleward-shifted North Atlantic winter storm track for the Eemian from two orbital time slices. ~~?~~
67 Brayshaw et al. (2010) linked Holocene orbital shifts to North Atlantic and Mediterranean winter
68 storm tracks via changes in baroclinicity. And ~~?~~Varma et al. (2012) found the Southern Hemi-
69 sphere westerlies ~~migrating-shift~~ under Holocene orbital forcing. Explicit cyclone tracking has
70 entered paleoclimate work mainly at the Last Glacial Maximum, where ice sheets dominate the
71 forcing ~~(?)~~(Pinto and Ludwig 2020; Raible et al. 2021), and in transient simulations of the last
72 millennium ~~(?)~~(Hess and Chemke 2025). For tropical cyclones the precessional response has re-
73 cently been ~~isolated~~examined in high-resolution paleoclimate simulations, where it is governed
74 by atmospheric moisture ~~(?)~~(Raavi et al. 2023). Here our study provides the extratropical counter-
75 part, where cyclone growth is baroclinic. To our knowledge, no previous study has applied explicit
76 cyclone tracking across a full precession cycle to isolate the orbital response of the extratropical
77 storm tracks, and none has asked which season carries the response.

78 ~~This missing piece of research matters beyond just storm dynamics.~~ Many paleoclimate archives
79 that record past storm activity, such as wind-blown dust and sediment, or rainfall linked to westerly
80 winds, only capture signals from specific seasons instead of annual averages. If a proxy record
81 mainly reflects one single season, it can either hide or falsely create an orbital signal, depending
82 on which time of year carries the strongest ~~storm response~~~~(?)~~response (Laepple et al. 2011). Fully
83 understanding which season holds the clearest precessional storm signal is therefore essential for
84 properly interpreting these geological records. Moreover, different metrics for cyclone behavior,
85 e.g., total storm counts, how much rain each storm produces, and storm intensity, react differently

86 to orbital forcing. ~~Researchers~~We must evaluate these quantities separately before drawing
87 conclusions from any single geological proxy.

88 To fill this gap, we run 12 equilibrium climate simulations covering a full precession cycle. Every
89 extratropical cyclone in both hemispheres is tracked individually using 6-hour sea level pressure.
90 Our experimental design isolates precession at amplified eccentricity with all other boundary
91 conditions fixed, so that the seasonal storm response to precession forcing alone can be examined.
92 The results are then tested in ~~simulationso~~simulations of real orbital states, ~~such as~~namely PI,
93 MH, and LIG. Section 2 describes our model and simulation, and the method we use for storm
94 tracking and analysis. Section 3 presents the results. We discuss and conclude in section 4 and 5
95 respectively.

96 2. Methodology

97 *a. Model and experiments*

98 The simulations are performed with the coupled Earth system model AWI-ESM2 ~~(?)~~
99 (Sidorenko et al. 2019). The atmosphere component is ECHAM6 at T63 resolution with 47
100 vertical levels ~~(?)~~(Stevens et al. 2013), which includes the land surface module JSBACH ~~(?)~~
101 (Reick et al. 2021). The ocean and sea ice component is FESOM2, a finite-volume model for-
102 mulated on an unstructured mesh ~~(?)~~(Danilov et al. 2017), with a spatially varying resolution of
103 from 25 km over high latitudes and coastal regions, to 150 km over open oceans.

104 We analyze 12 equilibrium simulations~~are analyzed~~, taken from the idealized precession ensem-
105 ble introduced by ~~?~~Shi et al. (2025). The longitude of perihelion ω is prescribed at values from 0°
106 to 330° in steps of 30° (labeled as p000, p030..., and p330), while eccentricity is fixed at 0.058 and
107 obliquity at 24.5° . The eccentricity value sits near the upper bound reached during the Quaternary
108 and amplifies the precessional ~~modulation of~~effects on insolation. All remaining boundary condi-
109 tions follow the pre-industrial setup, and each phase is branched from a long pre-industrial control.
110 We analyze 30 years of 6-hourly output from the equilibrated part of each simulation~~form the basis~~
111 ~~of the cyclone analysis~~. The calendar month of perihelion for each phase is computed from orbital
112 geometry ~~(?), and the same formulary~~(Berger 1978), which provides the insolation fields shown
113 in Fig. 1c-d.

To test our results under realistic orbital configurations, we additionally conducted three equilibrium simulations including a PI (0 ka) control run, a MH (6 ka) experiment, and a LIG (127 ka) experiment. Boundary conditions follow the protocols of PMIP4 (Otto-Bliesner et al. 2017). Orbital parameters are calculated following Berger (1977), and the greenhouse gas concentrations are taken from multi-archive reconstructions from the Greenland and the Antarctic ice cores (Flückiger et al. 2002; Monnin et al. 2004; Schneider et al. 2013; Schilt et al. 2010; Buiron et al. 2011). Each simulation is integrated for 1,000 model years with the trend of simulated global mean surface temperature in the final 100 model years not exceeding ± 0.05 K/century. We preserved 6-hourly sea level pressure from each simulation. At 127 ka, perihelion occurs close to the June solstice with an eccentricity of 0.039378; at 6 ka, perihelion falls within September, while the pre-industrial control features perihelion in early January.

b. Converting fixed-day calendar into angular calendar

Precession moves perihelion through the seasonal cycle, and Kepler's second law (Kepler 1953) then changes the lengths of the seasons (Fig. 1f-g). This is the calendar effect long known in paleoclimate modeling (Joussaume and Braconnot 1997; Bartlein and Shafer 2019; Shi et al. 2021). Monthly and seasonal averages at different orbital configurations are therefore only comparable when the averaging windows follow the same position of the Earth along its orbit.

Following Shi et al. (2021), we compute the length of a month (season) by calculating how much time the Earth needs to move from the respective starting point to the endpoint. We first define the mean anomaly M as the angle between the perihelion and Earth's position based on the assumption that the orbit describes a perfect circle with the sun at the center by:

$$M = \frac{2\pi}{T} \cdot t_p \quad (1)$$

Here, t_p denotes the time elapsed since Earth passes the perihelion and T is the Earth's revolution period, namely 1 year or 365 days. Taking into account the orbit's eccentricity ϵ , we define the eccentric anomaly E via:

$$E - \epsilon \cdot \sin(E) = M \quad (2)$$

Equation 2 can be solved with the application of Newton's method (Danby and Burkardt 1983), and E follows from the true anomaly θ , the angle at the Sun between perihelion and the actual position of the Earth (see Eq. 3.13b of Curtis 2014):

$$E = 2 \cdot \arctan\left(\sqrt{\frac{1-\epsilon}{1+\epsilon}} \cdot \tan\left(\frac{\theta}{2}\right)\right) \quad (3)$$

The above equations relate t_p to θ by:

$$t_p(\theta) = \frac{MT}{2\pi} = \frac{(E - \epsilon \cdot \sin(E))T}{2\pi} \quad (4)$$

This allows us to define seasons with respect to Earth's position on the orbit rather than relying on a fixed number of days as used in our modern calendar. The insolation panels of Fig. 1 are computed on this angular calendar. Re-binning the 6-hourly cyclone statistics onto the same calendar changes the local-summer response amplitudes by at most 1.5 percentage points and the phase-locking lags by less than half a month, while the autumn response strengthens slightly. All storm diagnostics are therefore reported on the model's own fixed calendar.

c. Cyclone detection

Extratropical cyclones are detected and tracked with the algorithm of Crawford et al. (2021), and Fig. 2 summarizes the procedure ~~with one 6-hourly field of the ensemble~~. 6-hourly sea level pressure is first interpolated onto 100-km equal-area grids centered on each pole, and the two hemispheres are processed separately. A grid cell qualifies as a cyclone center when its pressure p_c lies below that of all 8 neighbors and the surrounding field keeps a sufficient inward gradient. The gradient test compares p_c with the mean pressure $\bar{p}(d)$ on a one-cell ring at distance d from the candidate,

$$\frac{\bar{p}(d) - p_c}{d} \geq \gamma \quad (5)$$

where $d = 1000$ km and $\gamma = 750 \text{ Pa} (1000 \text{ km})^{-1}$, and candidates over terrain above 1500 m are discarded. Centers closer than 1000 km are treated as one multicenter system. The cyclone area A

163 is obtained by growing closed isobars around each center at a 200-Pa interval until a contour fails
 164 to close, and the last closed isobar defines the edge pressure p_e (Fig. 2b). Each cyclone carries the
 165 depth and the area-equivalent radius

$$D = p_e - p_c \quad (6)$$

$$R = \sqrt{A/\pi} \quad (7)$$

166 Tracking links the centers of consecutive 6-hourly fields (Fig. 2c). For every active cyclone a
 167 first-guess position is projected from its previous displacement, damped to three quarters,

$$\mathbf{x}_q = \mathbf{x}_t + 0.75 (\mathbf{x}_t - \mathbf{x}_{t-1}) \quad (8)$$

168 and a center of the next field qualifies as the continuation when it lies within $v_{\max}\Delta t$ of both the
 169 current position $\mathbf{x}_t$ and the projection $\mathbf{x}_q$, with $v_{\max} = 150 \text{ km h}^{-1}$ and $\Delta t = 6 \text{ h}$, a searching radius
 170 of 900 km per step. Among the qualifying candidates the one nearest to the projection continues
 171 the track. Centers left unmatched at the new time start tracks as genesis events, and tracks left
 172 unmatched end as lysis. Only systems poleward of 30° latitude enter the analysis.

173 Several track-based diagnostics are derived. Track density counts every 6-hourly cyclone position,
 174 system density counts each cyclone once at its mean position, and genesis density counts the first
 175 point of each track. ~~Tracks that merely cross the boundaries of the monthly processing windows~~
 176 ~~would appear as spurious genesis or lysis events, and such truncated events are removed.~~ For every
 177 cyclone the lifespan, the maximum depth relative to the outermost closed isobar, the minimum
 178 central pressure, the mean cyclone area, and the propagation speed are recorded. We further define
 179 a ~~storm-day~~ “storm day” field, representing the number of days per month during which a grid point
 180 lies inside a detected cyclone. It therefore measures the storm-influence time of each grid point,
 181 and it is affected by most cyclone metrics, such as the number of cyclones, their area, lifespan, and
 182 propagation speed, ~~though not by their intensity. Frequency statements in this paper therefore rest~~
 183 ~~on track density.~~ ~

184 *d. Baroclinicity diagnostics*

185 The maximum Eady growth rate serves as the measure of baroclinicity ~~(???)~~
 186 (Eady 1949; Lindzen and Farrell 1980; Hoskins and Valdes 1990). It is computed from ~~monthly~~
 187 wind, temperature, and geopotential height in the ~~925–700~~925–700 hPa layer as

$$\sigma = 0.31 |f| \frac{|\partial U / \partial z|}{N}, \quad (9)$$

188 where f is the Coriolis parameter, $|\partial U / \partial z|$ the vertical wind shear, and N the ~~Brunt–Väisälä~~
 189 Brunt–Väisälä frequency. The response to precession is quantified between the end members p090
 190 and p270, which place perihelion at the June and December solstices. ~~Because the difference~~
 191 ~~fields are dipolar, plain box averages cancel, and the response amplitude is instead defined as the~~
 192 ~~root-mean-square of the difference $\Delta\sigma = \sigma_{90} - \sigma_{270}$ over a basin, normalized by the climatological~~
 193 ~~growth rate of the season.~~

194 To separate the roles of vertical shear and stratification we follow the substitution approach of
 195 ~~?Camargo et al. (2007)~~. Writing $\sigma = 0.31 |f| S / N$ with $S = |\partial U / \partial z|$, the difference is split by
 196 changing one factor at a time while holding the other at its p270 value,

$$\Delta\sigma_S = 0.31 |f| \frac{S_{90} - S_{270}}{N_{270}}, \quad \Delta\sigma_N = 0.31 |f| S_{270} \frac{1}{N_{90}} - \frac{1}{N_{270}}, \quad (10)$$

197

$$\Delta\sigma_N = 0.31 |f| S_{270} \left(\frac{1}{N_{90}} - \frac{1}{N_{270}} \right), \quad (11)$$

198 so that $\Delta\sigma = \Delta\sigma_S + \Delta\sigma_N + \varepsilon$, with a small cross term ε . ~~The share of each factor is the projection~~
 199 ~~of its field onto the full difference pattern over the basin sector,~~

$$c_S = \frac{\langle \Delta\sigma_S \Delta\sigma \rangle}{\langle \Delta\sigma \Delta\sigma \rangle}, \quad c_N = \frac{\langle \Delta\sigma_N \Delta\sigma \rangle}{\langle \Delta\sigma \Delta\sigma \rangle},$$

200 where $\langle - \rangle$ denotes the area-weighted average over the storm-track box defined in the next sub-section.

202 *e. Basins and statistics*

203 Three storm-track basins are analyzed, the North Atlantic (30–75°N, 80°W–20°E), the North
204 Pacific (30–75°N, 120°E–120°W), and the Southern Ocean (40–75°S, all longitudes). Basin means
205 are area weighted. ~~The size of the precessional response is reported as the range across the twelve~~
206 ~~phases, the maximum minus the minimum of the 30-year mean, expressed as a percentage of the~~
207 ~~phase mean.~~ Uncertainty estimates use the interannual spread of the 30 individual years in each
208 phase, and the significance of the end-member difference maps is assessed with a two-sided Welch
209 t test ~~(?)~~ (Welch 1947) at each grid point.

210 **3. Results**

211 *a. Summer sensitivity*

212 The 12 precession-phase annual ~~midlatitude~~ mid-latitude storm climatologies exhibit only minor
213 differences, and phase anomalies relative to the multi-phase ensemble mean manifest as weak,
214 slowly evolving dipoles aligned along major storm track pathways (Fig. 3). The basin-averaged
215 time series quantify the muted amplitude of this annual-mean signal, with across-phase ranges
216 of 10%, 17% and 4% in the North Atlantic, the North Pacific and the Southern Ocean, varying
217 smoothly with ω . We also observe an interhemispheric antiphase.

218 There is also a pronounced seasonal cycle. Extratropical cyclone activities peak in DJF in North-
219 ern Hemisphere basins and JJA across the Southern Ocean, with this seasonal timing unchanged
220 across all orbital setups (Fig. 4a–c; ~~Table ??~~). Cold-season cyclones also carry greater intensity,
221 with winter peak cyclone depths exceeding summertime values by 70-100 % within the Northern
222 Hemisphere storm basins (not shown). Overall, winter dominates ~~midlatitude~~ mid-latitude cyclone
223 activity uniformly across hemispheres and all orbital phases.

224 Precession exerts negligible control over annual-mean storm conditions but profoundly reshapes
225 seasonal cyclone variability, and its strongest modulation targets the low-activity warm season.
226 Though each precession phase retains the same wintertime peak in basin-averaged storm track
227 density, cross-phase deviations emerge almost entirely within each basin’s local summer (Fig. 4a–c).
228 ~~Seasonal range statistics~~ Anomalies between end-members (p090 and p270) quantify this stark
229 seasonal contrast ~~in orbital sensitivity~~ (Fig. 4d–f). For the North Pacific, ~~the across-phase range~~
230 ~~of track density accounts for just 7~~ track density drops by 0.10 cell⁻¹ month⁻¹ in boreal summer

(JJA), about 44% of the phase mean ~~on an annual basis, yet rises to 44 % in boreal summer (JJA),~~
 against an annual-mean decrease of only 7%. The North Atlantic follows ~~an identical the same~~
 pattern, with ~~annual variability of 7 % versus a summertime range of 25 %~~ a summertime decrease
near 23% versus 5% annually. The Southern Ocean mirrors this behavior ~~but shifts the sensitive~~
~~season to in~~ its local summer (DJF), ~~with an annual range of only 4 % compared to a 22 % summer~~
~~signal where track density rises by 19%, against a 2% annual increase.~~ All three basins identify
 local summer as the most orbitally sensitive season. ~~The muted annual signal is not a diluted~~
~~version of this summer response but the residual of a partial cancellation. Summer~~ Moreover,
summer and winter vary in antiphase across the precession cycle, with perihelion suppressing the
 summer storms while slightly enhancing the winter ones, ~~so the two largely offset once the year is~~
~~averaged. The transition seasons follow the summer rather than opposing it, weakly in the same~~
~~sense, which is the autumn shoulder that reappears in the phase locking below.~~ Therefore the
annual signal is muted.

This summertime storm response is locked to orbital geometry. When basin-averaged track-
 density anomalies are plotted against both calendar month and precession angle ω , the band of peak
 anomalies traces a diagonal ridge aligned with the calendar timing of perihelion across the full phase
 space (Fig. 4 ~~g-i~~), with a delay of 1-2 months. Summers spent near perihelion exhibit suppressed
 storm activity, whereas aphelion summers feature intensified storminess. ~~A single precessional~~
~~forcing thus drives opposing midlatitude storm responses between the two hemispheres across the~~
~~orbital cycle, as~~ It should be noted that northern summer perihelion aligns with southern summer
 aphelion ~~under orbital precession, and vice versa, reversing the insolation control on warm-season~~
~~storminess between,~~ precessional forcing thus drives antiphased mid-latitude storm signals in the
two hemispheres.

b. Mechanisms

The pronounced seasonal dependence of orbital storm signals originates ~~directly~~ from insolation
 forcing. Averaged across the Northern Hemisphere storm-track band (40–70°N), precessional
 insolation variability peaks at 105 ~~W~~ W m⁻² during boreal summer (JJA) and falls to merely
~~14 W~~ 22 W m⁻² in boreal winter (DJF). The Southern Hemisphere storm band (40–70°S) exhibits
 mirror-symmetric behavior, with a maximum seasonal forcing amplitude of ~~119 W~~ 105 W m⁻² in

austral ~~winter~~ summer (DJF) versus just ~~47 W~~ 22 W m^{-2} in austral ~~summer~~ winter (JJA) (Fig. 1e). The underlying mechanism lies in the multiplicative nature of precessional insolation modulation. Shifting perihelion across the annual cycle alters the Sun–Earth distance for each month, which rescales monthly incoming solar flux by a factor proportional to eccentricity e (approximately $4e$, equaling 23% for $e = 0.058$, Fig. 1a-b). Midwinter at ~~midlatitudes~~ mid-latitudes features minimal climatological insolation, leaving little baseline flux to be modulated by precessional shifts. Precession forcing therefore exerts its strongest modulation on local summer in each hemisphere, and barely perturbs deep winter conditions ~~–(Fig. 1a-e).~~

Orbital insolation anomalies are translated into extratropical storm activity via shifts in ~~midlatitude~~ mid-latitude baroclinicity. Differences in maximum Eady growth rate between the extreme precession phases $\omega = 90^\circ$ and $\omega = 270^\circ$ reach their peak magnitude within each basin’s local summer (Fig. 5). Normalized against seasonal climatological means, summertime Eady growth rate responses hit 31 % in the North Atlantic, a pronounced 92 % in the North Pacific, and 21 % in the Southern Ocean. A decomposition analysis further reveals that vertical wind shear dominates the precessional Eady growth rate anomalies, while static stability contributes negligible signal strength. This is consistent across all storm tracks. ~~Perihelion summers warm the summer hemisphere, with the strongest 850-hPa warming over the subpolar continents and a weaker response over the ice-covered Arctic (Fig. ??). The meridional temperature gradient therefore flattens along the equatorward flank of the storm track and steepens along its poleward flank, the thermal counterpart of the growth-rate dipole. The weakened shear suppresses the baroclinic growth that sustains warm-season cyclones, and the steepened poleward flank supports the poleward shift. The storm anomalies appear somewhat poleward of the growth-rate anomalies, consistent with the downstream development of baroclinic waves (?).~~

b. Meridional shift

Spatial difference maps between the two orbital end-members (p090 minus p270) yields weak responses for the annual means (Fig. 6a). By contrast, boreal summer (JJA) exhibits a prominent dipole pattern across Northern Hemisphere basins, with fewer storm days through the ~~midlatitude~~ mid-latitude core and more along the Arctic flank at perihelion summer, the signature of a poleward shift joined to an overall ~~summer~~ weakening. In DJF the signal migrates to the Southern Ocean and

forms a circumpolar banded structure of opposite sign, marking enhanced ~~midlatitude~~ mid-latitude storm activity and an equatorward shift of the southern storm belt.

The latitudinal displacement of storm tracks driving these dipole anomalies is quantified in Fig. 7. For the North Pacific boreal summer, core storm latitude shifts from 53°N when perihelion falls in northern winter to 61°N when it falls in northern summer, yielding an 8° total poleward migration that peaks precisely at $\omega = 90^{\circ}$. The North Atlantic summertime storm track undergoes a 4.4° poleward shift following identical orbital behavior, while the Southern Ocean mirrors this response with a 3.8° poleward displacement when perihelion coincides with austral summer (near p270). A perihelion summer therefore weakens storm activity and shifts the storm population poleward, consistent with the poleward shift of baroclinic activity evident in the Eady growth rate dipole (Fig. 5a).

By contrast, the annual-mean track latitude varies by no more than 1.3° in any basin. Winter storm-track latitudes only fluctuate by $1.5\text{--}2.8^{\circ}$ across the full precession cycle, and their latitudinal extrema do not align with solstitial orbital phases. This further confirms our earlier conclusion that ~~midlatitude~~ mid-latitude winter storms are orbitally insensitive. Spring and autumn display intermediate latitudinal variability with a coherent phase dependence. For example, autumn poleward displacement maximizes near $\omega = 150^{\circ}$, the orbital configuration that places perihelion in late summer. Latitudinal poleward shifts therefore track the calendar timing of perihelion identically to Fig. 4g–i, with a uniform lag of 1–2 months, consistent with surface thermal inertia. Figure 8 quantifies this basin-dependent lag. All basin-scale storm extrema cluster along basin-specific dashed lines offset from the perihelion reference. The response lags the perihelion calendar by distinct intervals across regions, near 1 month in the North Pacific, 1–2 months in the North Atlantic, and slightly over 2 months in the Southern Ocean.

c. Cyclone precipitation

The hydrological signature of precessional forcing evolves in antiphase with storm dynamical responses (Fig. 9). ~~Detected extratropical cyclones contribute roughly one-third of total midlatitude precipitation across our simulations, consistent with storm precipitation partitioning derived from observational reanalysis datasets (?).~~ During boreal summer over the North Pacific, mean precipitation per individual storm rises by as much as 15 % at the orbital phase where total storm

frequency reaches its minimum, yielding nearly perfectly opposing variability between the two metrics. Perihelion summers feature warmer, more humid ~~midlatitude~~ mid-latitude atmospheres, which amplifies precipitation intensity within the remaining cyclones; ~~nevertheless~~. Nevertheless, the sharp reduction in storm number dominates the total signal, such that basin-wide summertime precipitation drops by 17 % between the two extreme orbital phases. The fractional share of precipitation originating from cyclones also declines markedly, with a full-cycle range of 9.5 % in North Pacific summer compared to just 3 % for the annual mean. Precessional forcing thus exerts two counteracting controls on extratropical hydroclimate simultaneously. It reduces total cyclone counts while boosting precipitation intensity per storm.

d. Realistic paleosimulations (MH and LIG)

Our results based on the 12 idealized simulations can be validated against the realistic orbital-state simulations (PI, MH, and LIG) introduced in Section 2. In LIG, perihelion occurs near June solstice, and the tracked cyclone responses fully reproduce our core theoretical pattern. Relative to PI, summer storm track density decreases by 8% over the North Atlantic and 11% over the North Pacific; winter cyclone changes remain within approximately 3%. Meanwhile, austral summer storm track density across the Southern Ocean rises by 8%. Northern Hemisphere summer storm tracks shift poleward by 1–2 degrees, while the Southern Hemisphere summer storm belt shifts equatorward (Fig. 10). The summer difference composites show identical weakening and poleward displacement of Northern Hemisphere storm tracks as seen in our amplified idealized end-member simulations (Fig. 11). In MH, perihelion happens in September, leaving seasonal forcing dominated by warm September conditions with cool March conditions. Consistent with this seasonal signal, MH storm variability features fewer, poleward-shifted cyclones in autumn and the opposite trend in spring. The magnitude of storm responses in these paleosimulations is much weaker than that of our amplified end-member contrasts (p090 vs. p270, Fig. 6), consistent with their smaller eccentricity values of the real orbits. Notably, these experiments combine realistic precession with their own obliquity and greenhouse gases, so they test our theoretical prediction under multi-factor natural orbital forcing rather than isolating precessional effects.

4. Discussion

The seasonal preference seen in storm responses originates purely from orbital geometry; independent of model physics. Precession redistributes insolation in proportion to the insolation a season already has, meaning winter hemispheres have very little background insolation for orbital changes to modify (Berger 1978). Fig. 1 clearly illustrates this stark seasonal contrast in forcing strength. Winter insolation varies by only roughly 2022 W m^{-2} across the precession cycle, while summer values exceed 125 reach 105 W m^{-2} within mid-latitude storm-track latitudes. This is model independent and will hold in any climate models as it simply reflects the orbital geometry behind the well-known summer-dominated orbital forcing framework (Huybers 2006). Existing monsoon studies have long confirmed that summer bears the strongest precessional signal (Kutzbach and Guetter 1986; Bosmans et al. 2015). But the new insight of our work lies elsewhere. This summer-focused orbital forcing acts on a winter-dominated weather system, reshaping storm activity exclusively during its normally quiet warm season. Note that the term "quiet" here refers to the weaker overall intensity of the summer storm track, not its population. Both hemisphere still hosts abundant storm systems in local summer, and they are merely weaker, shallower and move more slowly compared to winter cyclones.

The mechanism that converts summer insolation into summer storms is the thermal wind. When perihelion falls in local summer, the corresponding hemisphere warms most strongly over the subpolar continents. The temperature gradient flattens on the equatorward side of the storm belt and steepens on its poleward side, so the vertical wind shear that drives baroclinic growth weakens where most storms grow and the growth zone moves poleward (Fig. ??). baroclinic instability. Our factor decomposition analysis confirms that the shear term carries essentially the whole Eady growth rate response in all three ocean basins, while static stability plays a negligible role. We see similar physical mechanism operating in the modern Northern Hemisphere, where greenhouse warming takes the place of perihelion-driven heating. Stronger summer warming at high latitudes has weakened summertime large-scale circulation, eddy kinetic energy, and the number of strong summer cyclones (Coumou et al. 2015; Chang et al. 2016; Gertler and O’Gorman 2019). Future climate projections show the same shear-based decline of Northern Hemisphere summer cyclone activity and baroclinic energy as the globe warms (Lehmann et al. 2014; Priestley and Catto 2022). But

375 a difference should be noted here, as greenhouse warming is a transient forcing that both hemi-
376 spheres ~~feel~~change in phase, whereas our simulations are equilibrated and precession swings both
377 hemispheres in antiphase. Even so, both forcings follow one unified rule that is the core theme of
378 this study: ~~Cooler~~: cooler summers host more storm activity.

379 Compared to summer, storm systems in winter are much more stable. Across all three ocean
380 basins, winter storm activity changes by less than 10% over a full precession cycle, even though
381 winter Eady growth rates shift by 16–18% between the two orbital end-members, ~~and weak~~.
382 Weak winter insolation forcing cannot fully explain this stability. The winter storm track appears
383 saturated in the sense that ample baroclinicity is available in every phase, so that moderate changes
384 in the growth rate do not translate into changes in occurrence. ~~A transient~~Transient simulation
385 of the last millennium likewise found ~~eyelone counts~~winter cyclone nearly insensitive to external
386 forcing ~~(??)~~, (Raible et al. 2018; Hess and Chemke 2025), and those forcings acted to affect storm
387 track in summer (Hess and Chemke 2025). Our findings expand this conclusion to the far larger
388 swings in solar radiation driven by precession, and this insensitivity only holds for wintertime
389 conditions.

390 Geological records sensitive to summertime conditions within ~~midlatitude~~mid-latitude storm
391 belts should preserve prominent precession-scale storm signals, whereas proxies that average cli-
392 mate signals across winter or the full year will show only minor orbital imprint. Just as observed
393 for temperature reconstructions, the seasonal bias of a given archive can either generate arti-
394 ficial orbital storm signals or fully obscure true precessional variations ~~(?)~~(Laepple et al. 2011)
395 . Furthermore, summer-sensitive storm records from both hemispheres will fluctuate oppositely
396 over precessional cycles, forming an extratropical equivalent of the well-documented monsoon
397 seesaw ~~(??)~~(Wang et al. 2008). Current published sediment records already support the feasibil-
398 ity of testing this framework. Distinct precession-period signals have been detected in Northern
399 Hemisphere ~~midlatitude westerly proxies~~(?)mid-latitude westerly proxies (Li et al. 2019), as well
400 as North Pacific dust records. Precessional variation appears specifically in moisture-sensitive
401 dust fractions ~~, while the annually averaged dust flux responds primarily to obliquity forcing~~(?)
402 (Zhong et al. 2024). Meanwhile, dust intensity archives that capture winter-only signals display
403 very weak precessional power ~~(?)~~(Chen et al. 2014). Regarding the Southern Hemisphere, sub-
404 tropical margins of the southern westerly belt exhibit clear precession cycles ~~(?)~~(Lamy et al. 2019)

405 , consistent with our results. ~~And the presence or absence of precession signals at individual sites~~
406 ~~depends on their latitudinal position within the wind belt (?)~~. Records of the southern westerlies
407 also show orbital-band migrations of the belt itself ~~(?)~~ (Lamy et al. 2010; Varma et al. 2012), and
408 glacial-state cyclone modeling has shown how strongly reconstructed storminess depends on which
409 cyclone property a proxy senses ~~(?)~~ (Pinto and Ludwig 2020).

410 ~~Although the three storm basins follow the same physical mechanism, they exhibit different~~
411 ~~regional behaviors, which can be distinguished through separate diagnostic metrics. In local~~
412 ~~summer, storm frequency and track position vary substantially by 16–68% across the precession~~
413 ~~cycle, while storm propagation speed remains largely unchanged. By contrast, storm intensity~~
414 ~~exhibits clear basin-dependent differences. The Southern Ocean shows a 22% swing in storm~~
415 ~~frequency but only a 4% change in storm intensity. The North Pacific features a systematic~~
416 ~~reduction in overall storm activity during perihelion summers. This is consistent with previous~~
417 ~~findings from anthropogenic warming scenarios that storm frequency and intensity respond to~~
418 ~~external forcing through independent physical mechanism (?)~~ (Ulbrich et al. 2009; Catto et al. 2019)
419 ~~. Hydrological responses further amplify this contrast. Individual storms produce stronger~~
420 ~~precipitation in summers with reduced cyclone numbers, indicating that hydrological adjustments~~
421 ~~exceed dynamical changes under orbital forcing, a pattern analogous to that identified under~~
422 ~~greenhouse warming (?)~~. Given that extratropical precipitation is predominantly generated by
423 ~~cyclones and frontal systems (?)~~ (Catto and Pfahl 2013), this competing effect substantially alters
424 ~~precipitation partitioning. So a precipitation proxy tend to record combined signal of reduced~~
425 ~~storm occurrence and intensified single-storm rainfall during perihelion summers.~~

426 Our study is based on idealized simulations designed to isolate pure precessional forcing. The
427 experimental setup holds eccentricity, obliquity, ice sheets, and greenhouse gas concentrations
428 fixed and does not incorporate transient climate evolution or orbital cross-interactions. As such,
429 our results represent dynamical responses to idealized orbital conditions rather than reconstructions
430 of specific geological epochs. Future work could incorporate time-varying boundary conditions
431 and full orbital combinations to further generalize the storm seasonal response across broader
432 paleoclimate backgrounds.

433 5. Conclusions

434 In our study, we carry out a set of equilibrium climate simulations covering a full precession cycle,
435 with explicit tracking of all extratropical cyclones across both hemispheres to isolate precessional
436 forcing. We find that precession regulates extratropical cyclone primarily through local summer,
437 the season with naturally low storm activity, while winter storm tracks has only minor change
438 across all precession phases. Storm activity calms down when perihelion falls in local summer
439 and intensifies when aphelion coincides with local summer. And, as perihelion summer in one
440 hemisphere matches aphelion summer in the other, mid-latitude storminess of the two hemispheres
441 varies in antiphase during a precession cycle. This orbital signal propagates through shear-driven
442 baroclinicity and expresses itself in cyclones numbers and position, both of which are phase locked
443 to the calendar month of perihelion, with a delay of 1-2 months. Our result also bring clear
444 implications for paleoclimate proxy interpretation. Orbital storm signals should be extracted from
445 ~~midlatitude~~mid-latitude archives sensitive to summer climate, and the recorded signals will display
446 opposite variations between the two hemispheres.

580 Three-basin comparison of the precessional storm response. Ranges are max minus min across
 581 the twelve phases as a percentage of the phase mean. Storm and sensitive seasons refer to
 582 local winter and local summer. The storm season is diagnosed from cyclogenesis counts, the
 583 sensitive season from the seasonal storm-day range. The EGR response is the RMS p090–p270
 584 difference over the basin sector relative to its climatology, and the shear share is the projection of
 585 the vertical-shear factor onto the response pattern. Basin StormseasonSensitiveseasonStorm-day
 586 rangeann/summer (%)Summer max/min ω (°)EGR responsesummer (%)Shearshare (%)North
 587 Atlantic DJF JJA 10 / 32 240 / 90 31 107 North Pacific DJF JJA 17 / 68 270 / 90 92 106
 588 Southern Ocean JJA DJF 4 / 16 60 / 210 21 109




FIG. 1. ~~Precessional-insolation forcing from orbital geometry with eccentricity 0.058 and obliquity 24.5°.~~

~~(aa-b) Monthly Range of monthly mean insolation anomaly relative to across the phase mean, averaged over 12 precession phases for the storm-track band 40-70° 40-70°N and 40-70°S bands, computed on the classical fixed-day calendar (red) and on the angular calendar (blue). The dashed line marks the distance factor $4e/(1-e^2)^2$ multiplied by the monthly mean insolation. (c-d) Insolation anomaly as a function of calendar month and precession phase ω on the angular calendar, with the perihelion (red) and aphelion (blue) calendar months of each phase overlaid. (be) As Insolation range in (a), but for 40-70°S. (e) Seasonal forcing amplitude in the two bands, for each season on the seasonal mean of the monthly angular calendar, computed as maximum minus minimum insolation across the twelve 12 phases. Units: $W m^{-2}$. (f),(g) The orbital geometry behind the calendar drift, shown for two precession end-members (i.e., p090 and p270). The equinox and solstice lines through the Sun divide the orbit into the four seasons. Seasons on the perihelion side last about 85 days and seasons on the aphelion side 98 days, and this nearly two-week contrast swaps sides between the two phases. Day counts hold~~




FIG. 2. Cyclone detection and tracking procedure. (a,c) One 6-hourly sea level pressure field of phase p000 (contours, 8-hPa interval) with all detected cyclone centers poleward of 30°N (dots) and their area-equivalent radii $R = \sqrt{A/\pi}$ (circles), 15 January of one model year. (b,e) The deepest system of that field, with the outermost closed isobar p_e (dashed), the cyclone center (cross), the central and edge pressures and depth annotated, and the binary mask cells (shading) defined by $\text{SLP} < p_e$ inside the $\sqrt{A}$ search window; contour interval 4 hPa. (c,f) All cyclone tracks of the same month (lines) with genesis points (dots). (d) Data processing workflow.




FIG. 3. Annual-mean storm days across the precession cycle. (a)–(l) Annual storm-day anomaly of each phase relative to the twelve-phase-cross-phase mean (shading), with the phase-mean climatology outlined at 5 and 8 d month⁻¹ (contours). (m)–(o) Basin-mean annual storm days against ω with interannual standard errors; the across-phase range as a percentage of the phase mean is annotated in each panel.




FIG. 4. Seasonal concentration of the precessional storm response, from the Lagrangian tracks. (a)–(c) Basin-mean track density ($\text{cell}^{-1} \text{ month}^{-1}$) by calendar month for the North Atlantic, the North Pacific, and the Southern Ocean, one line per phase colored by ω , with local summer shaded. (d)–(f) Across-phase range of track density (maximum minus minimum, percent of Track-density difference between the phase-mean end members p090 and p270 ($\text{cell}^{-1} \text{ month}^{-1}$)) for the annual mean and the four seasons, with the local-summer bar highlighted; open-symbols give the corresponding ranges of system and genesis densities. (g)–(i) Basin-mean track-density anomaly relative to the phase mean as a function of calendar month and precession phase (unsmoothed), with the perihelion (red) and aphelion (blue) calendar months of each phase overlaid.




FIG. 5. Precessional response of the maximum Eady growth rate (925–700 hPa). (a) p090 minus p270 difference for JJA and (b) for DJF (day^{-1}). (c) Seasonal response amplitude in each basin, the RMS of the p090 minus p270 difference over the storm box as a percentage of the seasonal climatology, with stars marking local summer. (d) Projection shares of the vertical-shear and static-stability factors in the local-summer difference pattern.




FIG. 6. Difference in storm days (d month^{-1}) between p090 and p270 for (a) the annual mean, (b) JJA, and (c) DJF, on a common color scale. Stippling marks grid points where the difference is significant at the 95% level based on a two-sided Welch t test over the 30 simulated years.




FIG. 7. Storm-track latitude across the precession cycle, the storm-day-weighted mean latitude over each basin sector shown as the poleward anomaly from each season's phase mean, for (a) the North Atlantic, (b) the North Pacific, and (c) the Southern Ocean. Lines give the annual mean and the four seasons, with the local summer of each basin drawn heavy with markers. Vertical dotted lines mark the phase with perihelion in the local summer of each basin.




FIG. 8. Phase locking of the storm response to the perihelion calendar. For each calendar month and basin, the phase of the first harmonic across the 12 precession phases of (a) basin-mean storm days, shown as the ω of the activity minimum, and (b) the storm-day-weighted track latitude, shown as the ω of the poleward maximum. Marker size scales with the harmonic amplitude. The red line gives the ω that places perihelion in that calendar month, and dashed lines repeat it delayed by each basin's median lag over the amplitude-strong months (annotated in months).




FIG. 9. Storm-attributed precipitation across the precession cycle. (a) North Pacific JJA percent anomalies from the phase mean of basin-mean storm days, per-storm precipitation, and total precipitation; the dotted line marks the phase with perihelion in NH summer. (b) Across-phase range of the storm-attributed share of precipitation, accumulated precipitation inside storm footprints divided by total accumulated precipitation (percentage points), for the annual mean and local summer in each basin.





FIG. 10. Seasonal storm response of the real-orbit experiments, tracked with the same pipeline as the main ensemble and referenced to the preindustrial control. (a)–(c) Anomalies of basin-mean track density for the mid-Holocene (gold) and the Last Interglacial (red), as percentages of the preindustrial mean, for the annual mean and the four seasons, with error bars giving twice the standard error of the 30-year difference and the local summer shaded. Track density is normalized per T63 grid cell, and the annotated preindustrial values give the absolute scale in the units of Fig. 4. (d)–(f) The corresponding change in the track-point mean latitude, positive poleward.



575 FIG. 11. Storm-day anomalies of the real-orbit experiments relative to the preindustrial control, from the
 576 same storm-day product as the main ensemble. (a)–(d) Mid-Holocene minus preindustrial for DJF, MAM, JJA,
 577 and SON. (e)–(h) Last Interglacial minus preindustrial. Contours outline the preindustrial climatology at 5 and
 578 8 d month^{-1} . The isolated warm-season maxima over subtropical continental interiors reflect stationary surface
 579 heat lows rather than traveling cyclones.

589 *Acknowledgments.* The simulations were performed at the German Climate Computing Center
590 (DKRZ). [TODO funding and grant numbers before submission.]

591 *Data availability statement.* The AWI-ESM simulations of the idealized precession ensemble
592 are described in [?Shi et al. \(2025\)](#). The cyclone-track database, the gridded storm statistics, and
593 the analysis scripts underlying all figures will be archived and assigned a DOI upon acceptance.
594 [TODO repository DOI.]

595 ~~Thermal footprint of the orbital end members. (a) p090 minus p270 temperature at 850 hPa for~~
596 ~~JJA and (b) for DJF (K). (c) Change in the zonal-mean meridional temperature gradient for JJA~~
597 ~~(red) and DJF (blue), with the 40–70° storm bands shaded. In the Northern Hemisphere negative~~
598 ~~values mark a steepening of the poleward temperature decline, and in the Southern Hemisphere~~
599 ~~positive values do.~~

References

- Bartlein, P. J., and S. L. Shafer, 2019: Paleo calendar-effect adjustments in time-slice and transient climate-model simulations (paleocaladjust v1. 0): Impact and strategies for data analysis. *Geoscientific Model Development*, **12** (9), 3889–3913.
- Berger, A., 1977: Long-term variations of the earth's orbital elements. *Celestial Mechanics*, **15** (1), 53–74.
- Berger, A., 1978: Long-term variations of daily insolation and quaternary climatic changes. *Journal of Atmospheric Sciences*, **35** (12), 2362–2367.
- Bosmans, J., S. Drijfhout, E. Tuenter, F. Hilgen, and L. Lourens, 2015: Response of the north african summer monsoon to precession and obliquity forcings in the ec-earth gcm. *Climate dynamics*, **44** (1), 279–297.
- Brayshaw, D. J., B. Hoskins, and E. Black, 2010: Some physical drivers of changes in the winter storm tracks over the north atlantic and mediterranean during the holocene. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, **368** (1931), 5185–5223.
- Buiron, D., and Coauthors, 2011: Taldice-1 age scale of the talos dome deep ice core, east antarctica. *Climate of the Past*, **7** (1), 1–16.
- Camargo, S. J., K. A. Emanuel, and A. H. Sobel, 2007: Use of a genesis potential index to diagnose enso effects on tropical cyclone genesis. *Journal of Climate*, **20** (19), 4819–4834.
- Catto, J. L., and S. Pfahl, 2013: The importance of fronts for extreme precipitation. *Journal of Geophysical Research: Atmospheres*, **118** (19), 10 791–10 801, <https://doi.org/10.1002/jgrd.50852>.
- Catto, J. L., and Coauthors, 2019: The future of midlatitude cyclones. *Current Climate Change Reports*, **5**, 407–420, <https://doi.org/10.1007/s40641-019-00149-4>.
- Chang, E. K., S. Lee, and K. L. Swanson, 2002: Storm track dynamics. *Journal of climate*, **15** (16), 2163–2183.

- Chang, E. K., C.-G. Ma, C. Zheng, and A. M. Yau, 2016: Observed and projected decrease in northern hemisphere extratropical cyclone activity in summer and its impacts on maximum temperature. *Geophysical Research Letters*, **43** (5), 2200–2208.
- Chen, J., X. Liu, and V. A. Kravchinsky, 2014: Response of the high-resolution chinese loess grain size record to the 50° n integrated winter insolation during the last 500,000 years. *Geophysical Research Letters*, **41** (17), 6244–6251.
- Coumou, D., J. Lehmann, and J. Beckmann, 2015: The weakening summer circulation in the northern hemisphere mid-latitudes. *Science*, **348** (6232), 324–327.
- Crawford, A. D., E. A. Schreiber, N. Sommer, M. C. Serreze, J. C. Stroeve, and D. G. Barber, 2021: Sensitivity of northern hemisphere cyclone detection and tracking results to fine spatial and temporal resolution using era5. *Monthly Weather Review*, **149** (8), 2581–2598.
- Curtis, H., 2014: Orbital position as a function of time. *Orbital Mechanics for Engineering Students, 3rd edition, Elsevier, Amsterdam, the Netherlands*, 145–186.
- Danby, J., and T. Burkardt, 1983: The solution of kepler’s equation, i. *Celestial Mechanics*, **31** (2), 95–107.
- Danilov, S., D. Sidorenko, Q. Wang, and T. Jung, 2017: The finite-volume sea ice–ocean model (fesom2). *Geoscientific Model Development*, **10** (2), 765–789.
- Eady, E. T., 1949: Long waves and cyclone waves. *Tellus*, **1** (3), 33–52.
- Flückiger, J., E. Monnin, B. Stauffer, J. Schwander, T. F. Stocker, J. Chappellaz, D. Raynaud, and J.-M. Barnola, 2002: High-resolution holocene n2o ice core record and its relationship with ch4 and co2. *Global Biogeochemical Cycles*, **16** (1), 10–1.
- Gertler, C. G., and P. A. O’Gorman, 2019: Changing available energy for extratropical cyclones and associated convection in northern hemisphere summer. *Proceedings of the National Academy of Sciences*, **116** (10), 4105–4110.
- Hess, O., and R. Chemke, 2025: Forced changes in the north atlantic storm track over the last millennium. *Geophysical Research Letters*, **52** (15), e2025GL115 894.

- Hoskins, B. J., and K. I. Hodges, 2002: New perspectives on the northern hemisphere winter storm tracks. *Journal of the Atmospheric Sciences*, **59** (6), 1041–1061.
- Hoskins, B. J., and K. I. Hodges, 2005: A new perspective on southern hemisphere storm tracks. *Journal of climate*, **18** (20), 4108–4129.
- Hoskins, B. J., and P. J. Valdes, 1990: On the existence of storm-tracks. *Journal of Atmospheric Sciences*, **47** (15), 1854–1864.
- Huybers, P., 2006: Early pleistocene glacial cycles and the integrated summer insolation forcing. *Science*, **313** (5786), 508–511.
- Joussaume, S., and P. Braconnot, 1997: Sensitivity of paleoclimate simulation results to season definitions. *Journal of Geophysical Research: Atmospheres*, **102** (D2), 1943–1956.
- Kaspar, F., T. Spanghel, and U. Cubasch, 2007: Northern hemisphere winter storm tracks of the eemian interglacial and the last glacial inception. *Climate of the Past*, **3** (2), 181–192.
- Kepler, J., 1953: *Epitome astronomiae copernicanae*. Jo. Plancus.
- Kutzbach, J. E., and P. J. Guetter, 1986: The influence of changing orbital parameters and surface boundary conditions on climate simulations for the past 18 000 years. *Journal of atmospheric sciences*, **43** (16), 1726–1759.
- Laepple, T., M. Werner, and G. Lohmann, 2011: Synchronicity of antarctic temperatures and local solar insolation on orbital timescales. *Nature*, **471** (7336), 91–94.
- Lamy, F., R. Kilian, H. W. Arz, J.-P. Francois, J. Kaiser, M. Prange, and T. Steinke, 2010: Holocene changes in the position and intensity of the southern westerly wind belt. *Nature Geoscience*, **3** (10), 695–699.
- Lamy, F., and Coauthors, 2019: Precession modulation of the south pacific westerly wind belt over the past million years. *Proceedings of the National Academy of Sciences*, **116** (47), 23 455–23 460.
- Laskar, J., P. Robutel, F. Joutel, M. Gastineau, A. C. Correia, and B. Levrard, 2004: A long-term numerical solution for the insolation quantities of the earth. *Astronomy & Astrophysics*, **428** (1), 261–285.

- 679 Lehmann, J., D. Coumou, K. Frieler, A. V. Eliseev, and A. Levermann, 2014: Future changes
680 in extratropical storm tracks and baroclinicity under climate change. *Environmental Research*
681 *Letters*, **9** (8), 084 002.
- 682 Li, Y., Y. Song, Q. Yin, L. Han, and Y. Wang, 2019: Orbital and millennial northern mid-latitude
683 westerlies over the last glacial period. *Climate Dynamics*, **53** (5), 3315–3324.
- 684 Lindzen, R., and B. Farrell, 1980: A simple approximate result for the maximum growth rate of
685 baroclinic instabilities. *Journal of Atmospheric Sciences*, **37** (7), 1648–1654.
- 686 Merlis, T. M., T. Schneider, S. Bordoni, and I. Eisenman, 2013: Hadley circulation response to
687 orbital precession. part i: Aquaplanets. *Journal of Climate*, **26** (3), 740–753.
- 688 Monnin, E., and Coauthors, 2004: Evidence for substantial accumulation rate variability in antarctic
689 during the holocene, through synchronization of co₂ in the taylor dome, dome c and dml ice
690 cores. *Earth and Planetary Science Letters*, **224** (1-2), 45–54.
- 691 Otto-Bliesner, B., and Coauthors, 2017: The pmip4 contribution to cmip6—part 2: Two interglacials,
692 scientific objective and experimental design for holocene and last interglacial simulations.
693
- 694 Pinto, J. G., and P. Ludwig, 2020: Extratropical cyclones over the north atlantic and western europe
695 during the last glacial maximum and implications for proxy interpretation. *Climate of the Past*,
696 **16** (2), 611–626.
- 697 Priestley, M. D., and J. L. Catto, 2022: Future changes in the extratropical storm tracks and cyclone
698 intensity, wind speed, and structure. *Weather and Climate Dynamics*, **3**, 337–360.
- 699 Raavi, P. H., J.-E. Chu, A. Timmermann, S.-S. Lee, and K. J. Walsh, 2023: Moisture control
700 of tropical cyclones in high-resolution simulations of paleoclimate and future climate. *Nature*
701 *Communications*, **14** (1), 6426.
- 702 Raible, C. C., M. Messmer, F. Lehner, T. F. Stocker, and R. Blender, 2018: Extratropical cyclone
703 statistics during the last millennium and the 21st century. *Climate of the Past*, **14** (10), 1499–1514.

704 Raible, C. C., J. G. Pinto, P. Ludwig, and M. Messmer, 2021: A review of past changes in
 705 extratropical cyclones in the northern hemisphere and what can be learned for the future. *Wiley*
 706 *Interdisciplinary Reviews: Climate Change*, **12** (1), e680.

707 Reick, C. H., and Coauthors, 2021: Jsbach 3-the land component of the mpi earth system model:
 708 documentation of version 3.2.

709 Schilt, A., M. Baumgartner, T. Blunier, J. Schwander, R. Spahni, H. Fischer, and T. F. Stocker,
 710 2010: Glacial–interglacial and millennial-scale variations in the atmospheric nitrous oxide
 711 concentration during the last 800,000 years. *Quaternary Science Reviews*, **29** (1-2), 182–192.

712 Schneider, R., J. Schmitt, P. Köhler, F. Joos, and H. Fischer, 2013: A reconstruction of atmospheric
 713 carbon dioxide and its stable carbon isotopic composition from the penultimate glacial maximum
 714 to the last glacial inception. *Climate of the Past*, **9** (6), 2507–2523.

715 Shi, X., M. Werner, H. Yang, Q. Gao, J. Liu, and G. Lohmann, 2025: Precession controls on
 716 climate and water isotope signals in northern africa. *Paleoceanography and Paleoclimatology*,
 717 **40** (1), e2024PA004 999.

718 Shi, X., and Coauthors, 2021: Calendar effects on surface air temperature and precipitation based
 719 on model-ensemble equilibrium and transient simulations from pmip4 and pacmedy. *Climate of*
 720 *the Past Discussions*, **2021**, 1–29.

721 Sidorenko, D., and Coauthors, 2019: Evaluation of fesom2. 0 coupled to echam6. 3: preindustrial
 722 and highresmip simulations. *Journal of Advances in Modeling Earth Systems*, **11** (11), 3794–
 723 3815.

724 Simmonds, I., and K. Keay, 2000: Mean southern hemisphere extratropical cyclone behavior in
 725 the 40-year ncep–ncar reanalysis. *Journal of Climate*, **13** (5), 873–885.

726 Stevens, B., and Coauthors, 2013: Atmospheric component of the mpi-m earth system model:
 727 Echam6. *Journal of Advances in Modeling Earth Systems*, **5** (2), 146–172.

728 Ulbrich, U., G. C. Leckebusch, and J. G. Pinto, 2009: Extra-tropical cyclones in the present and fu-
 729 ture climate: a review. *Theoretical and Applied Climatology*, **96** (1–2), 117–131, [https://doi.org/](https://doi.org/10.1007/s00704-008-0083-8)
 730 [10.1007/s00704-008-0083-8](https://doi.org/10.1007/s00704-008-0083-8).

- 731 Varma, V., and Coauthors, 2012: Holocene evolution of the southern hemisphere westerly winds
732 in transient simulations with global climate models. *Climate of the Past*, **8** (2), 391–402.
- 733 Wang, Y., and Coauthors, 2008: Millennial-and orbital-scale changes in the east asian monsoon
734 over the past 224,000 years. *nature*, **451** (7182), 1090–1093.
- 735 Welch, B. L., 1947: The generalization of ‘student’s’ problem when several different population
736 variances are involved. *Biometrika*, **34** (1-2), 28–35.
- 737 Zhong, Y., and Coauthors, 2024: Orbital controls on north pacific dust flux during the late
738 quaternary. *Geophysical Research Letters*, **51** (4), e2023GL106 631.