
GTEx Cerebellum: comparison before our FDR cutoff
All comparable pairs; no GTEx significance or effect−direction filter
Baseline (0 hr)
1 hr
6 hr
inclusion
skipping
49 pairs; 24 genes
Pearson r = +0.44
Expected signs: 34/49
BH q < 0.05: 8/49
84 pairs; 41 genes
Pearson r = +0.19
Expected signs: 47/84
BH q < 0.05: 14/84
85 pairs; 42 genes
Pearson r = +0.26
Expected signs: 53/85
BH q < 0.05: 13/85
31 pairs; 27 genes
Pearson r = −0.09
Expected signs: 10/31
BH q < 0.05: 8/31
60 pairs; 48 genes
Pearson r = −0.04
Expected signs: 35/60
BH q < 0.05: 16/60
56 pairs; 44 genes
Pearson r = +0.00
Expected signs: 31/56
BH q < 0.05: 12/56
TRAPPC5 T>G
HNRNPK G>C
HNRNPK C>T
ATP5MJ T>C
RPS5 T>C
GTEx beta (normalized intron usage)
isoLASER dPSI: ALT minus REF part inclusion (fraction)
Our permutation FDR
BH q < 0.05
BH q >= 0.05
In previous GTEx significant−pair file
Other tested GTEx pair
Our screen: cis, |dPSI| >=0.10; NO permutation p or BH q cutoff. Exact variant, gene and annotated−event matching.
Inclusion junction: same signs expected. Skipping junction: opposite signs expected. Repeated genes/events are correlated.