

Galois Theory

MATH 440

Steven Xia

April 12, 2023

Galois Theory studies symmetries among roots of polynomials.

— Professor (Spring 2023)

Contents

1 Field Extensions	1
-------------------------------------	----------

1 Field Extensions

Theorem 1.1. *If $F \rightarrow K$ and $K \rightarrow L$ are finite, then $[L : F] = [L : K][K : L]$.*

Proof. Let $\{\alpha_1, \dots, \alpha_n\}$ and $\{\beta_1, \dots, \beta_m\}$ be bases for $F \rightarrow K$ and $K \rightarrow L$ respectively. We claim $S = \{\alpha_i \beta_j\}$ is an F -basis for L . It is immediate that S spans L , so we show that S is linearly independent.

Suppose some linear combination of S is zero, then factoring out by the α_i implies the coefficients of each group of α_i must be zero, but they are all linear combinations of the β_j , hence all coefficients must be zero. \square

Theorem 1.2. *A field extension is finite if and only if it is algebraic and finitely generated.*

Proof. Suppose $F \rightarrow K$ is a field extension. It is trivial to show that

- (i) if $F \rightarrow K$ is not algebraic, then it is not finite, and
- (ii) if $F \rightarrow K$ is not finitely generated, then it is not finite.

Suppose $F \rightarrow K$ is algebraic and finitely generated, and let $\{\alpha_1, \dots, \alpha_n\}$ be a basis for $F \rightarrow K$. Break the extension down by $F \rightarrow F(\alpha_1) \rightarrow F(\alpha_1, \alpha_2) \rightarrow \dots \rightarrow F(\alpha_1, \dots, \alpha_n) = K$, and see that each of these intermediate extensions are finite. Theorem 1.1 asserts $F \rightarrow K$ is finite. \square

Corollary 1.2.1. *Every composition of algebraic field extensions is algebraic.*

Proof. Suppose $F \rightarrow K$ and $K \rightarrow L$ are algebraic. Let $\alpha \in L$ with $m_{\alpha, K} = x^n + c_{n-1}x^{n-1} + \dots + c_0$, and construct $K' = F(c_0, \dots, c_{n-1})$, which is algebraic and finitely generated, hence finite. But $K' \rightarrow K'(\alpha)$ is also finite, so $F \rightarrow K'(\alpha)$ is finite, therefore α is algebraic over F . \square

Theorem 1.3 (Kronecker). *If F is a field and $f \in F[x]$ is non-constant, then there exists a finite $F \rightarrow K$ such that f has a root in K .*

Proof. Without loss of generality, we may assume f is irreducible. Define $K = F[x]/\langle f \rangle$, which is a field because $\langle f \rangle$ is maximal. See that $x + \langle f \rangle \in K$ is a root of f . \square

Theorem 1.4. *A field F is algebraically closed if and only if every algebraic $F \rightarrow K$ has $[K : F] = 1$.*

Proof. The “only if” is trivial, so suppose every $F \rightarrow K$ has $[K : F] = 1$. Let $f \in F[x]$, and Theorem 1.3 asserts there exists a finite $F \rightarrow K$ in which f has a root. But $[K : F] = 1$, so this root is in fact in F . \square

Theorem 1.5. *Every field has an algebraic closure.*

Proof. Suppose F is a field, and define S to be the set of monic and irreducible polynomials in $F[x]$. Also construct $R = F[y_f \mid f \in S]$ and $I = \langle f(y_f) \mid f \in S \rangle$.

We claim $1 \notin I$. Towards a contradiction, suppose $1 \in I$, so we can write $1 = \sum_i a_i f_i(y_{f_i})$ for $a_i \in R$ and $f_i \in S$. Repeating Theorem 1.3 for each f_i generates a field extension in which there exist α_i such that $f_i(\alpha_i) = 0$ for all i , but now we have $1 = \sum_i a_i f_i(\alpha_i) = 0$, a contradiction.

Now we know I is a proper ideal, so it is contained in some maximal ideal M . Define $F \rightarrow K = R/M$, and see that $y_i + \langle M \rangle \in K$ is a root of f_i , so we conclude that K is an algebraic closure of F . \square

Theorem 1.6 (Isomorphism Extension). *Let F and K be fields with isomorphism $\phi : F \rightarrow K$. If $F \rightarrow E$ is algebraic, then there exists an isomorphism ψ between E and a subfield of \bar{K} satisfying $\psi|_F = \phi$.*

Proof. Let S be the set of (E', σ) where E' is a field satisfying $F \subseteq E' \subseteq E$ and σ an isomorphism from E' to a subfield of \bar{K} satisfying $\sigma|_F = \phi$. Define a partial order on S by $(E_1, \sigma_1) \leq (E_2, \sigma_2)$ if and only if $E_1 \subseteq E_2$ and $\sigma_2|_{E_1} = \sigma_1$. We wish to apply Zorn's Lemma to S , so we note

- (i) that $(F, \phi) \in S$ implies S is non-empty, and
- (ii) that every chain $(E_1, \sigma_1) \leq (E_2, \sigma_2) \leq \dots$ in S is bounded above by (E', σ) , where $E' = \bigcup_i E_i$ and σ is defined by simply using whichever σ_i is available, since they are all compatible.

Therefore, there exists a maximal element $(M, \tau) \in S$, and we want to show $M = E$. Towards a contradiction, suppose $M \subsetneq E$ and choose $\alpha \in E \setminus M$ with minimal polynomial $m_{\alpha, M} = x^n + c_{n-1}x^{n-1} + \cdots + c_0 \in M[x]$. Define $L = \tau(M)$ and $f(x) = x^n + \tau(c_{n-1})x^{n-1} + \cdots + \tau(c_0) \in L[x]$, and see that

$$M(\alpha) \rightarrow M[x]/m_{\alpha, M} \rightarrow L[x]/f(x) \rightarrow L(\beta)$$

is an isomorphism for $\beta \in \bar{L} = \bar{K}$ a root of $f(x)$. Moreover, this extends τ , a contradiction. \square

Theorem 1.7. *If F is a field, then all algebraic closures of F are pairwise isomorphic.*

Theorem 1.8 (Symmetric Polynomials). *Every symmetric polynomial can be written uniquely as a polynomial in the elementary symmetric polynomials.*

Theorem 1.9 (Algebra). *The set of complex numbers is algebraically closed.*