Non-Local Magic Resources for Fermionic Gaussian States

Daniele Iannotti,[1,2,3,*] Beatrice Magni,[3,†] Riccardo Cioli,[4,‡] Alioscia Hamma,[1,5,2,§] and Xhek Turkeshi[3,¶]

[1] *Scuola Superiore Meridionale, Largo S. Marcellino 10, 80138 Napoli, Italy*
[2] *INFN Sezione di Napoli, via Cintia, 80126 Napoli, Italy*
[3] *Institut für Theoretische Physik, Universität zu Köln, Zülpicher Strasse 77, 50937 Köln, Germany*
[4] *Dipartimento di Fisica e Astronomia, Università di Bologna and INFN, Sezione di Bologna, via Irnerio 46, I-40126 Bologna, Italy*
[5] *Dipartimento di Fisica 'Ettore Pancini', Università degli Studi di Napoli Federico II, Via Cintia, 80126 Napoli, Italy*

Non-local magic quantifies the nonstabilizerness that local unitaries cannot remove, but its computation requires an intractable optimization. For fermionic Gaussian states and contiguous bipartitions, we prove a closed formula for the Rényi-2 non-local stabilizer entropy over local Gaussian unitaries, set by the reduced Majorana covariance matrix. We derive a Page-like curve for random states, logarithmic scaling at the XY critical point, and a quasiparticle picture after quenches. Our result enables measuring fermionic Gaussian-restricted non-local magic via fermionic shadow tomography.

Introduction.—Entanglement and magic, or nonstabilizerness, are two fundamental notions of quantum complexity that capture complementary aspects of what makes a many-body system genuinely quantum [1–3]. States with low entanglement admit efficient classical simulation via tensor networks [4], whereas states with low magic are accessible to stabilizer-based techniques such as tableau simulation [5, 6] or Pauli propagation [7, 8]. While entanglement is a well-characterized concept in many-body physics [9, 10], the role of magic has only recently come into focus, driven by the development of stabilizer entropies [11, 12], scalable probes of nonstabilizerness whose computational cost is comparable to that of entanglement entropy [13–15] and which have an important operational meaning [16–18]. This advance has stimulated a broad research program, from equilibrium phenomena [19–23] to quantum-circuit and Hamiltonian dynamics [24–32].

A central open question concerns the interplay between entanglement and magic [33–35]. The recently introduced *non-local magic* [36–39] isolates the irreducible nonstabilizerness that cannot be removed by local basis changes, thereby separating genuinely correlated magic from purely local contributions. However, non-local magic is defined through a minimization over all local unitaries acting on the full Hilbert space, rendering it intractable beyond a few qubits. Although estimation schemes exist for matrix-product states [36], where entanglement remains bounded, which classes of quantum matter admit an analytic characterization of non-local magic, especially in regimes with extensive entanglement, has remained an open question.

In this Letter, we answer this question for fermionic Gaussian states and unitaries. We introduce the fermionic Gaussian-restricted non-local (GNL) magic, and prove (Theorem 1) that at Rényi index $\alpha = 2$ it admits a closed-form expression in terms of the eigenvalues of the reduced Majorana covariance matrix of a contiguous subsystem of size ℓ, computable with poly(ℓ) resources. The key insight is that the minimization over local *Gaussian* unitaries is solved exactly by the BCS canonical form [40]. This is in stark contrast with the total magic, which requires Majorana Monte Carlo [41] or perfect sampling [42], both converging in general with an exponential number of samples. We leverage this result to obtain an exact Page-like curve for typical fermionic Gaussian states, corroborated by SYK_2 eigenstates [41, 43–46]; logarithmic scaling of non-local magic at the XY critical point via the Fisher–Hartwig approach; a quasiparticle picture [47] for magic spreading after quantum quenches; and diffusive growth in random Gaussian circuits. Because our result relies solely on two-point correlation functions, it provides a scalable route for the experimental estimation of Gaussian-restricted non-local magic in large-scale quantum processors via fermionic shadow tomography [48].

Setup.—We consider N qubits on a chain $\Lambda = \{1, \dots, N\}$, with Hilbert space dimension $d = 2^N$ [49–52]. The $2N$ Majorana operators are obtained via the Jordan–Wigner (JW) transformation, $\gamma_{2j-1} = \big(\prod_{m<j} Z_m\big) X_j$ and $\gamma_{2j} = \big(\prod_{m<j} Z_m\big) Y_j$, and satisfy $\{\gamma_j, \gamma_k\} = 2\delta_{jk}$. Fermionic Gaussian unitaries (FGU), or matchgates, act on the Majorana operators as rotations, $U_O \gamma_j U_O^\dagger = \sum_m O_{mj}\gamma_m$ with $O \in \mathrm{O}(2N)$; up to the reflection implemented by $X_1 = \gamma_1$, they are generated by quadratic Hamiltonians

$$H = \frac{i}{4}\sum_{m,n=1}^{2N} h_{m,n}\gamma_m\gamma_n\,, \qquad h = -h^{\mathsf{T}}\,. \tag{1}$$

A pure fermionic Gaussian state (FGS) $|\Psi_O\rangle = U_O|\mathbf{0}\rangle$, with $|\mathbf{0}\rangle$ the fermionic vacuum, is fully characterized by its covariance matrix $\Gamma_{mn} = i\langle\Psi_O|[\gamma_m, \gamma_n]|\Psi_O\rangle/2$, given by $\Gamma = O\Gamma_0 O^{\mathsf{T}}$ with $\Gamma_0 = \bigoplus_{m=1}^{N}\left(\begin{smallmatrix} 0 & -1 \\ 1 & 0 \end{smallmatrix}\right)$.

The stabilizer entropy of order α of a pure state is $M_\alpha = (1-\alpha)^{-1}\log_2\zeta_\alpha$ [11], where $\zeta_\alpha = d^{-1}\sum_{P\in\mathcal{P}_N}\langle\Psi|P|\Psi\rangle^{2\alpha}$ is the stabilizer purity and $\mathcal{P}_N$ the set of N-qubit Pauli strings; it is a magic monotone

for integer $\alpha \geq 2$ [12, 53]. Throughout, we consider the bipartition $\Lambda = A \sqcup B$ of the chain into the contiguous blocks $A = \{1, \ldots, \ell\}$ and $B = \{\ell + 1, \ldots, N\}$. The non-local magic is [36]

$$M_\alpha^{\mathrm{NL}} := \min_{U_A, U_B} M_\alpha \left[(U_A \otimes U_B) |\Psi\rangle \right] , \tag{2}$$

where the minimization is over $\mathrm{U}(2^\ell) \otimes \mathrm{U}(2^{N-\ell})$. This is the magic that cannot be erased by local unitaries; for instance, maximally entangled states have vanishing non-local magic, being locally equivalent to stabilizer states. It is distinct from long-range magic, which cannot be removed by finite-depth local circuits [54]. The minimization in Eq. (2) is superexponentially costly; for $\alpha = 2$, a proxy can be obtained from the Schmidt spectrum [36]. For fermionic Gaussian states, we introduce the GNL magic

$$M_\alpha^{\mathrm{GNL}} := \min_{U_{O_A}, U_{O_B}} M_\alpha \left[(U_{O_A} \otimes U_{O_B}) |\Psi\rangle \right] , \tag{3}$$

where the minimization is restricted to the Gaussian unitaries of A and B induced by $O_A \in \mathrm{O}(2\ell)$ and $O_B \in \mathrm{O}(2(N-\ell))$, hence $M_\alpha^{\mathrm{NL}} \leq M_\alpha^{\mathrm{GNL}}$. Since A and B are contiguous, $U_{O_A} \otimes U_{O_B}$ is itself a Gaussian unitary of the whole chain, so that Eq. (3) amounts to a maximization of ζ_α over the local orthogonal orbit $\Gamma \mapsto (O_A \oplus O_B)\Gamma(O_A \oplus O_B)^{\mathsf{T}}$ (End Matter).

Main result.—Let $\pm\lambda_i$, with $\lambda_i \in [0, 1]$ and $i = 1, \ldots, \ell$, be the eigenvalues of the $2\ell \times 2\ell$ reduced covariance matrix $i\Gamma_A$ of the Majorana operators of A.

Theorem 1. *For any pure fermionic Gaussian state $|\Psi_O\rangle$ on N modes and the contiguous blocks $A = \{1, \ldots, \ell\}$ and $B = \{\ell + 1, \ldots, N\}$,*

$$\begin{aligned} M_2^{\mathrm{GNL}}(|\Psi_O\rangle) &= \sum_{i=1}^{\ell} \mathfrak{m}_2(\lambda_i^2) , \\ \mathfrak{m}_2(x) &= -\log_2\left(1 - x + x^2\right) . \end{aligned} \tag{4}$$

Equivalently, the maximum of the stabilizer purity over the local Gaussian orbit is $\max_{O_A, O_B} \zeta_2 = \prod_{i=1}^{\ell} (1 - \lambda_i^2 + \lambda_i^4)$, attained in the canonical modewise axes of Ref. [40].

Local FGU bring any pure FGS into a canonical modewise form of independent two-mode BCS-like pairs, $\cos\theta_i |00\rangle + \sin\theta_i |11\rangle$, and unpaired modes [40], a normal form that underlies several descriptions of fermionic Gaussian entanglement [55–57]. We prove that this representative is the *global* maximizer of ζ_2 on the local Gaussian orbit, from which Eq. (4) follows by a direct computation of the two-qubit stabilizer entropy [36, 37]. The proof is outlined in the End Matter and given in full in Sec. S1 of the Supplemental Material (SM) [58]. It combines a Wick-theorem representation of ζ_2 as the collision probability of a determinantal point process with two ingredients of independent interest: a sharp bound on the collision probability of a Slater determinant in terms of its transpose overlap, and a determinant inequality for positive matrices of rank at most two.

For integer $\alpha > 2$, we conjecture that the same structure survives,

$$\begin{aligned} M_\alpha^{\mathrm{GNL}} &= \sum_{i=1}^{\ell} \mathfrak{m}_\alpha(\lambda_i^2) , \\ \mathfrak{m}_\alpha(x) &= \frac{1}{1-\alpha} \log_2 \frac{(1-x)^\alpha + 1 + x^\alpha}{2} , \end{aligned} \tag{5}$$

which reduces to Eq. (4) at $\alpha = 2$. This conjecture is supported by extensive variational optimization (SM, Sec. S2) and reproduces the same phenomenology as the $\alpha = 2$ case (SM, Sec. S4). Since the proof relies on the quadratic (collision) form of ζ_2, establishing the conjecture for $\alpha > 2$ requires a different strategy; below we focus on $\alpha = 2$.

Several remarks are in order. First, for $\ell \leq 2$ the Gaussian minimization coincides with the full one, $M_2^{\mathrm{NL}} = M_2^{\mathrm{GNL}}$, and Eq. (4) reproduces the expression of Ref. [36]; for larger subsystems, non-Gaussian local unitaries can further decrease the magic, so that Theorem 1 provides an upper bound, $M_2^{\mathrm{NL}} \leq M_2^{\mathrm{GNL}}$. Second, Eq. (4) requires only $\mathrm{poly}(\ell)$ operations, whereas even the total stabilizer entropy of FGS requires in the worst case an exponential number of samples [41, 42]. Third, entanglement is necessary but not sufficient for non-local magic: $\mathfrak{m}_2(\lambda^2)$ vanishes both for unentangled ($\lambda = 1$) and for maximally entangled ($\lambda = 0$) modes. For instance, the rainbow state $\bigotimes_{j=1}^{N/2} (|0_j 0_{N+1-j}\rangle + |1_j 1_{N+1-j}\rangle)/\sqrt{2}$ is a FGS, the ground state of a free-fermion hopping Hamiltonian with long-range pairing, with maximal entanglement across $A = \{1, \ldots, N/2\}$, i.e., $\lambda_j = 0$, yet it is a stabilizer state with $M_2^{\mathrm{NL}} = M_2^{\mathrm{GNL}} = 0$. Local non-Clifford rotations only add local magic; genuine non-local magic requires modes at intermediate entanglement, $0 < \lambda_j < 1$, e.g., generated by a joint non-Clifford Gaussian rotation of each pair [59]. Finally, since Eq. (4) depends only on Γ_A, a polynomial number of two-point functions, GNL magic is accessible via fermionic shadow tomography [48]: random matchgate circuits followed by computational-basis measurements reconstruct Γ_A with accuracy ϵ_Γ and failure probability δ with $O(\ell^2 \log(\ell/\delta)/\epsilon_\Gamma^2)$ shots; the propagation of this error to M_2^{GNL} is discussed in the SM, Sec. S3. This opens a scalable route beyond the recent two-qubit demonstration of non-local magic [60].

Typical states.—We first consider states $|\Psi_O\rangle$ with Haar-random $O \in \mathrm{O}(2N)$, whose statistics is invariant under mode permutations, and $A = \{1, \ldots, \ell\}$, $\ell \leq N/2$. The variables $x_m = \lambda_m^2$ form a Jacobi ensemble [61–63]. Since M_2^{GNL} is a sum of single-mode terms, its quenched average $\overline{M}_2^{\mathrm{GNL}} \equiv \mathbb{E}_O[M_2^{\mathrm{GNL}}]$, performed outside the logarithm and typically harder to evaluate than its annealed counterpart, follows exactly




FIG. 1. **GNL magic of typical fermionic Gaussian states.** (a) Page curve for Haar-random FGS: $\overline{M}_2^{\rm GNL}/N$ vs $r = \ell/N$, compared with the $N \to \infty$ prediction, Eq. (14). (b) Average GNL magic of SYK_2 mid-spectrum eigenstates, in quantitative agreement with Eq. (14).

from the one-point density of the ensemble, both at finite N and in the thermodynamic limit $\ell, N \to \infty$ at fixed $r = \ell/N$ (End Matter). The GNL magic density $\overline{\mathfrak{m}}_2^{\rm GNL} := \overline{M}_2^{\rm GNL}/N$, Eq. (14), is symmetric under $r \to 1-r$, reaches $\overline{\mathfrak{m}}_2^{\rm GNL}(1/2) = 2 - \log_2(2+\sqrt{3})$ at half partition, and grows as $\overline{\mathfrak{m}}_2^{\rm GNL} \simeq r^2/\ln 2$ for $r \to 0$. This *GNL magic Page curve* [Fig. 1(a)] has a quadratic onset for every α (SM, Sec. S4), with a transparent origin: when $\ell = O(1)$ and $N \to \infty$, the modes of A are almost maximally entangled and carry no non-local magic. This is in sharp contrast with the entanglement Page curve, which grows linearly for small r [63–65].

To test universality beyond Haar-random states, we consider mid-spectrum eigenstates of the SYK_2 model, Eq. (1) with $h_{m,n} \sim \mathcal{N}(0, 1/N)$, obtained by flipping a random subset of $N/4$ to $N/2$ single-particle modes in the ground state (SM, Sec. S5). Averaging over 10^3 disorder realizations and 200 eigenstates each, the data converge to Eq. (14) as N increases [Fig. 1(b)].

Ground states.—We now consider the XY chain with periodic boundary conditions,

$$H = -\sum_{j=1}^{N}\left[\frac{1+\eta}{2}X_jX_{j+1} + \frac{1-\eta}{2}Y_jY_{j+1} + \mu Z_j\right], \quad (6)$$

which the JW transformation maps onto a quadratic Majorana chain of the form of Eq. (1). We study it as a *fermionic* model: its ground state is a FGS and the natural notion of locality is that of the fermionic modes, so that $M_2^{\rm GNL}$ is the non-local magic of the free-fermion problem, and we take the contiguous block $A = \{1, \ldots, \ell\}$. For $N \to \infty$ at fixed ℓ, Γ_A is a block-Toeplitz matrix with a 2×2 symbol $g(k)$ fixed by the Bogoliubov angle θ_k (End Matter). Hence, $M_2^{\rm GNL}$ is computed at cost $O(\ell^3)$, and its asymptotics for $\ell \gg 1$ is controlled by the strong Szegő theorem for smooth symbols and by the Fisher–Hartwig formula for singular ones [66, 67]. Since $|\det g(k)| = 1$, volume-law scaling is excluded; in the gapped phases, where θ_k is analytic, $M_2^{\rm GNL}$ saturates with exponentially small corrections [68], while logarithmic growth can only arise from singularities of the symbol, as at the Ising critical point $\mu = 1$, $\eta = 1$.

Off criticality, the single-particle entanglement energies of a semi-infinite cut are known from corner transfer matrices [69, 70]: they are equally spaced, $\varepsilon_n^{\rm ent} = (2n+1)\varepsilon$ for $\mu > 1$ and $\varepsilon_n^{\rm ent} = 2n\varepsilon$ for $\mu < 1$, with $n = 0, 1, \ldots$ and ε given in the End Matter. Since $\lambda_n = \tanh(\varepsilon_n^{\rm ent}/2)$ [71], and the block has two boundaries,

$$M_2^{\rm GNL}(\ell) \simeq 2\sum_{n=0}^{\infty} \mathfrak{m}_2\left[\tanh^2\left(\varepsilon_n^{\rm ent}/2\right)\right], \quad (7)$$

up to corrections exponentially small in ℓ/ξ, with ξ the correlation length. As $\mathfrak{m}_2(x) \simeq (1-x)/\ln 2$ for $x \to 1$, the series converges rapidly. In the ordered phase of the Ising chain ($\eta = 1$, $\mu < 1$), the GNL magic is strongly suppressed: the $n = 0$ mode is maximally entangled ($\lambda_0 = 0$), while all other modes are exponentially close to unentangled ($\lambda_n \to 1$), so that every mode contributes an exponentially small amount, vanishing exactly in the GHZ limit $\mu \to 0$, where the ground state approaches the GHZ state $(|+\rangle^{\otimes N} + |-\rangle^{\otimes N})/\sqrt{2}$, with $X|\pm\rangle = \pm|\pm\rangle$. This is a striking example of a phase with area-law entanglement but negligible non-local magic: the entanglement entropy, in contrast, receives a finite contribution from the maximally entangled $n = 0$ mode.

At $\mu = 1$, the gap closes, $\varepsilon \to 0$, the series in Eq. (7) diverges, and θ_k develops a jump at $k = 0$. The Fisher–Hartwig asymptotic density of the eigenvalues $\lambda \in [0, 1)$ for a block with two boundaries, $\rho(\lambda) \simeq 2\ln\ell\,[\pi^2(1-\lambda^2)]^{-1}$ [67], yields $M_2^{\rm GNL}(\ell) \simeq \beta_{\rm GNL}\ln\ell$, with

$$\beta_{\rm GNL} = \frac{2}{\pi^2}\int_0^1 \frac{\mathfrak{m}_2(\lambda^2)}{1-\lambda^2}\,d\lambda = \frac{\pi^2 - \ln^2\left(7-4\sqrt{3}\right)}{\pi^2\ln 16}, \quad (8)$$

the non-local magic analog of $S \simeq (c/3)\ln\ell$ [72], with the central charge c replaced by $\beta_{\rm GNL}$.

Figure 2 benchmarks these predictions. In panel (a), Eq. (4) agrees with a brute-force minimization over the full group $\mathrm{U}(2^\ell) \otimes \mathrm{U}(2^{N-\ell})$ for $N = 8$, $\ell = 4$. For a general model, however, we do not expect such agreement, and only $M_2^{\rm NL} \le M_2^{\rm GNL}$ is expected to hold (SM, Sec. S2); nevertheless, throughout the XY phase diagram and for the sizes accessible to exact minimization, we find that the agreement persists [73]. In panel (b), exact numerics for $N = 10^6$ agree with Eq. (7). In panel (c), a fit for $\ell > 50$ at $\mu = 1$ gives $\beta_{\rm fit} \approx 0.1072$, in excellent agreement with Eq. (8). Panel (d) shows that over the full XY phase diagram the GNL magic is concentrated near the critical lines.

These results reflect an area law: in unique gapped ground states of one-dimensional local free-fermion Hamiltonians, $M_2^{\rm NL} \le M_2^{\rm GNL} \le 2S(A) \le 2C_{\rm ent}|\partial A|$, with $C_{\rm ent}$ the constant of Hastings' area law [74] (End Matter).




FIG. 2. **Equilibrium GNL magic in the Ising and XY chains.** (a) Ising ground state with $N = 8$, $\ell = 4$: Eq. (4) (crosses) agrees with the brute-force minimization over all local unitaries $U_A \otimes U_B$ (red markers) to numerical precision; the black line is the total stabilizer entropy M_2. (b) $M_2^{\rm GNL}$ vs transverse field μ in the thermodynamic limit for $\ell = 16, 32, 64, 128$; the black curve is the exact $\ell \to \infty$ result, Eq. (7). The GNL magic peaks at the critical point $\mu_c = 1$. (c) $M_2^{\rm GNL}(\ell)$ at criticality ($\mu = 1$, red) grows logarithmically: the dashed line is the fit $\beta_{\rm fit} \ln \ell + b$, with $\beta_{\rm fit} \approx 0.1072$ and $b \approx 0.27$, in agreement with Eq. (8). In the gapped phases ($\mu = 0.5$, 4.0), $M_2^{\rm GNL}$ saturates. (d) $M_2^{\rm GNL}/\ell$ over the XY phase diagram (μ, η), from the exact corner-transfer-matrix formula: the GNL magic concentrates near the critical lines.




FIG. 3. **Dynamics of GNL magic.** (a) Quench at $\eta = 1$ from the fully polarized state ($\mu_0 \to \infty$) to the critical point ($\mu = 1$): exact numerics (circles) agree with the quasiparticle prediction, Eq. (10) (solid lines), for $\ell = 20, 40, 60, 100, 160$. Dashed lines mark the stationary value, Eq. (9). (b) Same for the quench $\mu_0 = 2 \to \mu = 1$, showing weaker GNL magic production due to the smaller Bogoliubov mismatch. (c) Stationary density $M_2^{\rm GNL}(\infty)/\ell$ vs the post-quench field μ ($\mu_0 \to \infty$): Eq. (9) (solid) and GGE numerics at $\ell = 160$ (crosses). (d) Random Gaussian circuits: $\overline{M}_2^{\rm GNL}/N$ vs $\sqrt{t}/N$ for $N = 20, 40, 80$, showing diffusive growth and saturation to the Haar value $\overline{\mathfrak{m}}_2^{\rm GNL}(1/2) = 2 - \log_2(2 + \sqrt{3})$ (dashed line).

Quench dynamics.—We now turn to quantum quenches in the Ising chain ($\eta = 1$): the system is prepared in the ground state of $H(\mu_0)$ and evolved with $H(\mu)$, $\mu \neq \mu_0$. The state remains Gaussian, and $\Gamma_A(t)$ is block-Toeplitz with a time-dependent symbol fixed by the post-quench dispersion ε_k and by the mismatch $\Delta\theta_k = \theta_k - \theta_k^{(0)}$ between post- and pre-quench Bogoliubov angles (End Matter) [75]. At long times, dephasing leads to a generalized Gibbs ensemble (GGE) [75, 76] with stationary eigenvalues $\lambda_k^\infty = |\cos \Delta\theta_k|$, and the Szegő theorem gives

$$M_2^{\rm GNL}(\infty) = \frac{\ell}{\pi} \int_0^\pi dk\, \mathfrak{m}_2\big(\sin^2 \Delta\theta_k\big) \ , \qquad (9)$$

i.e., the quench generically produces volume-law GNL magic, symmetric under $\mu \leftrightarrow \mu_0$. Equation (9) closely parallels the entanglement entropy of the same quench [75], with $\mathfrak{f}(x) = -x \log_2 x - (1-x)\log_2(1-x)$ the binary entropy, $S = (\ell/\pi) \int_0^\pi dk\, \mathfrak{f}\big(\sin^2(\Delta\theta_k/2)\big)$: in terms of λ_k^∞, the entanglement weight $\mathfrak{f}[(1+\lambda_k^\infty)/2]$ is replaced by the magic weight $\mathfrak{m}_2[(\lambda_k^\infty)^2] = \mathfrak{m}_2(\sin^2 \Delta\theta_k)$, where we used $\mathfrak{m}_2(x) = \mathfrak{m}_2(1-x)$. This parallel extends to all times. Writing $M_2^{\rm GNL}$ as a contour integral of $\frac{d}{d\lambda} \ln \det(\lambda - i\Gamma_A)$ and evaluating the moments $\mathrm{Tr}(i\Gamma_A)^{2n}$ via the multidimensional stationary-phase approach of Ref. [77] (End Matter), we derive the quasiparticle formula

$$M_2^{\rm GNL}(\ell, t) \simeq \int_0^\pi \frac{dk}{\pi}\, \min\big(2|v_k|\, t,\ \ell\big)\, \mathfrak{m}_2\big(\sin^2 \Delta\theta_k\big), \quad (10)$$

valid for $\ell, t \gg 1$ at fixed t/ℓ, with $v_k = d\varepsilon_k/dk$ the group velocity. Equation (10) predicts linear growth up to $t^* = \ell/(2v_{\max})$, followed by saturation to Eq. (9), and has a transparent interpretation [75, 77]: the quench creates pairs of entangled quasiparticles that propagate ballistically with velocities $\pm v_k$, and a pair contributes to

the GNL magic of A only when one partner is inside A and the other outside. The magic carried by each pair, $\mathfrak{m}_2(\sin^2\Delta\theta_k)$, vanishes both when $\Delta\theta_k = 0$ (no entanglement production) and when $\Delta\theta_k = \pi/2$ (maximally entangled but stabilizer-like pairs), and is maximal in between. Figures 3(a)–(c) show excellent agreement between Eqs. (9) and (10) and exact numerics, obtained by diagonalizing $\Gamma_A(t)$, for the quenches $\mu_0 = \infty \to 1$ and $\mu_0 = 2 \to 1$ [panels (a), (b)] and for $\mu_0 \to \infty$ at varying μ [panel (c)].

Finally, in brickwork circuits of random nearest-neighbor Gaussian gates [78], where operator spreading is diffusive [79, 80], the half-chain GNL magic grows as $\sqrt{t}$ and its density saturates to the Page value $\overline{\mathfrak{m}}_2^{\rm GNL}(1/2)$ on the timescale $t \sim N^2$ [Fig. 3(d); SM, Sec. S5].

Conclusions.—We have proved an exact closed form for the fermionic Gaussian-restricted non-local magic of Gaussian states at $\alpha = 2$ for contiguous bipartitions, and conjectured its extension to $\alpha > 2$. By reducing the minimization over local Gaussian unitaries to a single diagonalization of the reduced covariance matrix, it establishes fermionic Gaussian states as a broad class of highly entangled many-body systems where Gaussian-restricted non-local magic is exactly tractable, and provides an upper bound on the unrestricted one. We used it to derive an exact Page-like curve, logarithmic scaling at criticality, a quasiparticle picture after quenches and diffusive growth in random circuits. This tractability opens several directions. In higher dimensions, entanglement acquires logarithmic corrections tied to the Fermi surface [81], and it is natural to ask how the scaling rate $\beta_{\rm GNL}$ emerges there. Whether $\beta_{\rm GNL}$ captures universal features of the underlying conformal field theory, as the central charge does for entanglement [72], requires extending our analysis to interacting integrable models, where one can also ask whether the quasiparticle picture of entanglement spreading [82–84] carries over to non-local magic. Our formula applies directly to disordered free fermions, from Anderson localization [85, 86] to strongly disordered chains with Griffiths regions [87–89], and to topological phases such as the Kitaev chain and model [90, 91], whose boundary modes may carry non-local magic reflecting the topological invariant, with implications for coherent errors in fermionic codes [92, 93]. With a U(1) symmetry, the block structure of Γ suggests a symmetry-resolved decomposition of $M_2^{\rm GNL}$ [94–99] and possible quantum Mpemba effects [23, 100, 101]. Finally, in monitored [102–104] and non-Hermitian [105–110] free-fermion dynamics our formula applies along each quantum trajectory, raising the question of whether GNL magic undergoes its own measurement-induced transition.

Note added.—While preparing this manuscript, we became aware of a closely related independent work by M. Collura, B. Beri, and E. Tirrito [111]. When overlapping, our results are consistent.

This work is dedicated to Ada. We are particularly indebted to P. Sierant for spotting a mistake in a prior version of this manuscript, and we thank the anonymous referees for their careful reading and, in particular, for pointing out the role of the contiguity of the bipartition in Theorem 1, the misprint in the Jacobi weight, and a simpler proof of a lemma in the SM. We thank L. Leone, A. Russotto, P. Calabrese, L. Piroli, and J. Odavić for comments and collaborations on related topics. D.I. and B.M. acknowledge the Les Houches Summer School 2025 on Exact Solvability and Quantum Information, which facilitated work on this subject. D.I. acknowledges support from COST Action THEORY-CHALLENGES CA22113 under Short-Term Scientific Mission (STSM) grant E-COST-GRANT-CA22113-6901e967 and Research Grants in Germany, 2026 (57812126) from the German Academic Exchange Service (DAAD). B.M. and X.T. acknowledge support from DFG Emmy Noether Programme proposal "Digital Quantum Matter Out-of-Equilibrium" No. 560726973, DFG under Germany's Excellence Strategy – Cluster of Excellence Matter and Light for Quantum Computing (ML4Q) EXC 2004/2 – 390534769, and DFG Collaborative Research Center (CRC) 183 Project No. 277101999 - project B01. A.H. acknowledges support from PNRR MUR project PE0000023-NQSTI and the PNRR MUR project CN00000013-ICSC. Some support from AI tools was used in the writing and presentation of this manuscript.

Data availability.—The data generated in this study will be made publicly available on Zenodo upon publication.

* d.iannotti@ssmeridionale.it; These two authors contributed equally
† bmagni@uni-koeln.de; These two authors contributed equally
‡ riccardo.cioli3@unibo.it
§ alioscia.hamma@unina.it
¶ xturkesh@uni-koeln.de

[1] R. Horodecki, P. Horodecki, M. Horodecki, and K. Horodecki, Quantum entanglement, Rev. Mod. Phys. **81**, 865 (2009).
[2] E. Chitambar and G. Gour, Quantum resource theories, Rev. Mod. Phys. **91**, 025001 (2019).
[3] V. Veitch, S. A. H. Mousavian, D. Gottesman, and J. Emerson, The resource theory of stabilizer quantum computation, New J. Phys. **16**, 013009 (2014).
[4] U. Schollwöck, The density-matrix renormalization group in the age of matrix product states, Ann. Phys. **326**, 96–192 (2011).
[5] D. Gottesman, The heisenberg representation of quantum computers (1998), arXiv:quant-ph/9807006 [quant-ph].
[6] S. Bravyi, D. Browne, P. Calpin, E. Campbell, D. Gosset, and M. Howard, Simulation of quantum circuits

by low-rank stabilizer decompositions, Quantum **3**, 181 (2019).
[7] N. Dowling, P. Kos, and X. Turkeshi, Magic resources of the heisenberg picture, Phys. Rev. Lett. **135**, 050401 (2025).
[8] M. S. Rudolph, T. Jones, Y. Teng, A. Angrisani, and Z. Holmes, Pauli propagation: A computational framework for simulating quantum systems (2025), arXiv:2505.21606 [quant-ph].
[9] N. Laflorencie, Quantum entanglement in condensed matter systems, Physics Reports **646**, 1–59 (2016).
[10] L. Amico, R. Fazio, A. Osterloh, and V. Vedral, Entanglement in many-body systems, Rev. Mod. Phys. **80**, 517 (2008).
[11] L. Leone, S. F. E. Oliviero, and A. Hamma, Stabilizer Rényi entropy, Phys. Rev. Lett. **128**, 050402 (2022).
[12] L. Leone and L. Bittel, Stabilizer entropies are monotones for magic-state resource theory, Phys. Rev. A **110**, L040403 (2024).
[13] X. Huang, H.-Z. Li, C. H. Lee, and J.-X. Zhong, A fast and exact approach for stabilizer rényi entropy via the xor-fwht algorithm (2026), arXiv:2512.24685 [quant-ph].
[14] Z. Xiao and S. Ryu, Exponentially accelerated sampling of pauli strings for nonstabilizerness (2026), arXiv:2601.00761 [quant-ph].
[15] P. Sierant, J. Vallès-Muns, and A. Garcia-Saez, Computing quantum magic of state vectors (2026), arXiv:2601.07824 [quant-ph].
[16] L. Bittel and L. Leone, Operational interpretation of the stabilizer entropy, Quantum **10**, 2069 (2026).
[17] D. Iannotti, L. Campos Venuti, and A. Hamma, Van hove singularities in stabilizer entropy densities, Journal of Physics A: Mathematical and Theoretical **59**, 075301 (2026).
[18] S. Cusumano, L. C. Venuti, S. Cepollaro, G. Esposito, D. Iannotti, B. Jasser, J. Odavić, M. Viscardi, and A. Hamma, Non-stabilizerness and violations of chsh inequalities (2025), arXiv:2504.03351 [quant-ph].
[19] T. Haug and L. Piroli, Quantifying nonstabilizerness of matrix product states, Phys. Rev. B **107**, 035148 (2023).
[20] P. S. Tarabunga, E. Tirrito, T. Chanda, and M. Dalmonte, Many-body magic via pauli-markov chains—from criticality to gauge theories, PRX Quantum **4**, 040317 (2023).
[21] M. Hoshino, M. Oshikawa, and Y. Ashida, Stabilizer rényi entropy and conformal field theory, Phys. Rev. X **16**, 011037 (2026).
[22] M. Hoshino and Y. Ashida, Stabilizer rényi entropy encodes fusion rules of topological defects and boundaries (2025), arXiv:2507.10656 [quant-ph].
[23] S. Aditya, A. Summer, P. Sierant, and X. Turkeshi, Mpemba effects in quantum complexity (2025), arXiv:2509.22176 [quant-ph].
[24] X. Turkeshi, E. Tirrito, and P. Sierant, Magic spreading in random quantum circuits, Nat. Commun. **16**, 2575 (2025).
[25] J. Odavić, M. Viscardi, and A. Hamma, Stabilizer entropy in nonintegrable quantum evolutions, Phys. Rev. B **112**, 104301 (2025).
[26] E. Tirrito, X. Turkeshi, and P. Sierant, Anticoncentration and nonstabilizerness spreading under ergodic quantum dynamics, Phys. Rev. Lett. **135**, 220401 (2025).
[27] P. R. N. Falcão, P. Sierant, J. Zakrzewski, and E. Tirrito, Nonstabilizerness dynamics in many-body localized systems, Phys. Rev. Lett. **135**, 240404 (2025).
[28] P. Sierant, M. Schirò, M. Lewenstein, and X. Turkeshi, Measurement-induced phase transitions in $(d+1)$-dimensional stabilizer circuits, Phys. Rev. B **106**, 214316 (2022).
[29] B. Magni, A. Christopoulos, A. De Luca, and X. Turkeshi, Anticoncentration in clifford circuits and beyond: From random tensor networks to pseudomagic states, Phys. Rev. X **15**, 031071 (2025).
[30] H. Lóio, G. Lami, L. Leone, M. McGinley, X. Turkeshi, and J. D. Nardis, Quantum state designs via magic teleportation (2026), arXiv:2510.13950 [quant-ph].
[31] S. Aditya, X. Turkeshi, and P. Sierant, Growth and spreading of quantum resources under random circuit dynamics (2025), arXiv:2512.14827 [quant-ph].
[32] M. Viscardi, M. Dalmonte, A. Hamma, and E. Tirrito, Interplay of entanglement structures and stabilizer entropy in spin models, SciPost Physics Core **9**, 10.21468/scipostphyscore.9.1.012 (2026).
[33] E. Tirrito, P. S. Tarabunga, G. Lami, T. Chanda, L. Leone, S. F. E. Oliviero, M. Dalmonte, M. Collura, and A. Hamma, Quantifying nonstabilizerness through entanglement spectrum flatness, Phys. Rev. A **109**, L040401 (2024).
[34] D. Szombathy, A. Valli, C. Moca, L. Farkas, and G. Zaránd, Asymptotically independent fluctuations of stabilizer rényi entropy and entanglement in random unitary circuits, Phys. Rev. Res. **7**, 043072 (2025).
[35] D. Iannotti, G. Esposito, L. Campos Venuti, and A. Hamma, Entanglement and Stabilizer entropies of random bipartite pure quantum states, Quantum **9**, 1797 (2025).
[36] C. Cao, G. Cheng, A. Hamma, L. Leone, W. Munizzi, and S. F. Oliviero, Gravitational backreaction is magical, PRX Quantum **6**, 040375 (2025).
[37] D. Qian and J. Wang, Quantum nonlocal nonstabilizerness, Phys. Rev. A **111**, 052443 (2025).
[38] W. Munizzi and H. J. Schnitzer, Magic and non-clifford gates in topological quantum field theory (2026), arXiv:2604.14271 [hep-th].
[39] C. Cao, Non-trivial area operators require non-local magic, Journal of High Energy Physics **2024**, 105 (2024).
[40] A. Botero and B. Reznik, BCS-like modewise entanglement of fermion Gaussian states, Phys. Lett. A **331**, 39 (2004).
[41] S. Bera and M. Schirò, Non-stabilizerness of Sachdev-Ye-Kitaev model, SciPost Phys. **19**, 159 (2025).
[42] M. Collura, J. D. Nardis, V. Alba, and G. Lami, The non-stabilizerness of fermionic gaussian states (2026), arXiv:2412.05367 [quant-ph].
[43] G. C. Santra, A. Windey, S. Bandyopadhyay, A. Legramandi, and P. Hauke, Complexity transitions in chaotic quantum systems: Nonstabilizerness, entanglement, and fractal dimension in syk and random matrix models (2025), arXiv:2505.09707 [quant-ph].
[44] B. Jasser, J. Odavić, and A. Hamma, Stabilizer entropy and entanglement complexity in the sachdev-ye-kitaev model, Phys. Rev. B **112**, 174204 (2025).
[45] V. Malvimat, M. Sarkis, Y. Suk, and J. Yoon, Multipartite non-local magic and syk model (2026), arXiv:2601.03076 [hep-th].

[46] A. Russomanno, G. Passarelli, D. Rossini, and P. Lucignano, Nonstabilizerness in the unitary and monitored quantum dynamics of xxz-staggered and sachdev-ye-kitaev models, Phys. Rev. B **112**, 064312 (2025).

[47] P. Calabrese and J. Cardy, Entanglement entropy and conformal field theory, J. Phys. A Math. Theor. **42**, 504005 (2009).

[48] K. Wan, W. J. Huggins, J. Lee, and R. Babbush, Matchgate shadows for fermionic quantum simulation, Communications in Mathematical Physics **404**, 629–700 (2023).

[49] G. B. Mbeng, A. Russomanno, and G. E. Santoro, The quantum Ising chain for beginners, SciPost Phys. Lect. Notes , 82 (2024).

[50] J. Surace and L. Tagliacozzo, Fermionic Gaussian states: an introduction to numerical approaches, SciPost Phys. Lect. Notes , 54 (2022).

[51] P. Sierant, P. Stornati, and X. Turkeshi, Fermionic magic resources of quantum many-body systems, PRX Quantum **7**, 010302 (2026).

[52] P. Sierant, X. Turkeshi, and P. S. Tarabunga, Theory of the matchgate commutant (2026), arXiv:2603.12392 [quant-ph].

[53] T. Haug and L. Piroli, Stabilizer entropies and nonstabilizerness monotones, Quantum **7**, 1092 (2023).

[54] D. A. Korbany, M. J. Gullans, and L. Piroli, Long-range nonstabilizerness and phases of matter, Physical Review Letters **135**, 160404 (2025), arXiv:2502.19504 [quant-ph].

[55] C. Spee, K. Schwaiger, G. Giedke, and B. Kraus, Mode entanglement of Gaussian fermionic states, Phys. Rev. A **97**, 042325 (2018).

[56] R. Morral-Yepes, M. Langer, A. Gammon-Smith, B. Kraus, and F. Pollmann, Disentangling strategies and entanglement transitions in unitary circuit games with matchgates (2025), arXiv:2507.05055 [quant-ph].

[57] M. Langer, R. Morral-Yepes, A. Gammon-Smith, F. Pollmann, and B. Kraus, Matchgate circuit representation of fermionic gaussian states: optimal preparation, approximation, and classical simulation (2026), arXiv:2603.05675 [quant-ph].

[58] See Supplemental Material at [URL will be inserted by publisher] for the full proof of Theorem 1, the variational verification of Eqs. (4) and (5), the error analysis for the experimental estimation, the generalization to Rényi index $\alpha > 2$, and further details on the numerical simulations, which includes Refs. [112–117].

[59] J. Huang, R. Lv, X. Qian, and M. Qin, Dismagicker: Unitary gate for non-stabilizerness reduction (2026), arXiv:2604.04046 [quant-ph].

[60] H. G. Ahmad, G. Esposito, V. Stasino, J. Odavic, C. Cosenza, A. Sarno, P. Mastrovito, M. Viscardi, S. Cusumano, F. Tafuri, D. Massarotti, and A. Hamma, Experimental demonstration of non-local magic in a superconducting quantum processor (2025), arXiv:2511.15576 [quant-ph].

[61] M. L. Mehta, *Random Matrices*, 3rd ed. (Academic Press, 2004).

[62] G. Livan, M. Novaes, and P. Vivo, *Introduction to Random Matrices* (Springer, 2018).

[63] E. Bianchi, L. Hackl, and M. Kieburg, Page curve for fermionic gaussian states, Phys. Rev. B **103**, L241118 (2021).

[64] I. Ulčakar and L. Vidmar, Tight-binding billiards, Phys. Rev. E **106**, 034118 (2022).

[65] P. Łydżba, M. Rigol, and L. Vidmar, Eigenstate entanglement entropy in random quadratic hamiltonians, Phys. Rev. Lett. **125**, 180604 (2020).

[66] B.-Q. Jin and V. E. Korepin, Quantum spin chain, toeplitz determinants and the fisher—hartwig conjecture, Journal of Statistical Physics **116**, 79–95 (2004).

[67] A. R. Its, B.-Q. Jin, and V. E. Korepin, Entanglement in thexyspin chain, J. Phys. A Math. Gen. **38**, 2975–2990 (2005).

[68] B. Simon, *Szego's Theorem and Its Descendants: Spectral Theory for L2 Perturbations of Orthogonal Polynomials* (Princeton university press, 2010).

[69] I. Peschel and V. Eisler, Reduced density matrices and entanglement entropy in free lattice models, J. Phys. A: Math. Theor. **42**, 504003 (2009).

[70] V. Eisler and I. Peschel, Analytical results for the entanglement Hamiltonian of a free-fermion chain, J. Phys. A: Math. Theor. **50**, 284003 (2017).

[71] I. Peschel, On the entanglement entropy for an XY spin chain, J. Stat. Mech. Theor. Exp. **2004**, P12005 (2004).

[72] P. Calabrese and J. Cardy, Entanglement entropy and quantum field theory, J. Stat. Mech. Theor. Exp. , P06002 (2004).

[73] Code and data for the paper.

[74] M. B. Hastings, An area law for one-dimensional quantum systems, Journal of Statistical Mechanics: Theory and Experiment **2007**, P08024–P08024 (2007).

[75] P. Calabrese and J. Cardy, Evolution of entanglement entropy in one-dimensional systems, J. Stat. Mech. Theor. Exp. , P04010 (2005).

[76] L. Vidmar, L. Hackl, E. Bianchi, and M. Rigol, Entanglement entropy of eigenstates of quadratic fermionic hamiltonians, Phys. Rev. Lett. **119**, 020601 (2017).

[77] M. Fagotti and P. Calabrese, Evolution of entanglement entropy following a quantum quench: Analytic results for the xy chain in a transverse magnetic field, Phys. Rev. A **78**, 010306 (2008).

[78] E. Tirrito, L. Lumia, A. Paviglianiti, G. Lami, A. Silva, X. Turkeshi, and M. Collura, Magic phase transitions in monitored gaussian fermions (2025), arXiv:2507.07179 [quant-ph].

[79] B. C. Dias, M. Haque, P. Ribeiro, and P. McClarty, Diffusive operator spreading for random unitary free fermion circuits (2021), arXiv:2102.09846 [cond-mat.str-el].

[80] T. Swann, D. Bernard, and A. Nahum, Spacetime picture for entanglement generation in noisy fermion chains, Phys. Rev. B **112**, 064301 (2025).

[81] B. Swingle, Entanglement entropy and the fermi surface, Phys. Rev. Lett. **105**, 050502 (2010).

[82] V. Alba and P. Calabrese, Entanglement and thermodynamics after a quantum quench in integrable systems, Proceedings of the National Academy of Sciences **114**, 7947–7951 (2017).

[83] V. Alba and P. Calabrese, Quantum information scrambling after a quantum quench, Phys. Rev. B **100**, 115150 (2019).

[84] J. Kudler-Flam, Y. Kusuki, and S. Ryu, The quasiparticle picture and its breakdown after local quenches: mutual information, negativity, and reflected entropy, J. High Energ. Phys. **2021** (3), 146.

[85] F. Evers and A. D. Mirlin, Anderson transitions, Rev. Mod. Phys. **80**, 1355 (2008).

[86] P. Sierant, D. Delande, and J. Zakrzewski, Thouless time analysis of anderson and many-body localization transitions, Phys. Rev. Lett. **124**, 186601 (2020).

[87] F. Igloi and C. Monthus, Strong disorder rg approach of random systems, Physics Reports **412**, 277–431 (2005).

[88] C. Monthus, Strong disorder real-space renormalization for the many-body-localized phase of random majorana models, J. Phys. A Math. Theor. **51**, 115304 (2018).

[89] P. Ruggiero and X. Turkeshi, Quantum information spreading in random spin chains, Phys. Rev. B **106**, 134205 (2022).

[90] A. Y. Kitaev, Unpaired Majorana fermions in quantum wires, Physics-Uspekhi **44**, 131 (2001).

[91] A. Kitaev, Anyons in an exactly solved model and beyond, Ann. Phys. **321**, 2–111 (2006).

[92] F. Venn, J. Behrends, and B. Béri, Coherent-error threshold for surface codes from majorana delocalization, Phys. Rev. Lett. **131**, 060603 (2023).

[93] J. Behrends and B. Béri, Statistical mechanical mapping and maximum-likelihood thresholds for the surface code under generic single-qubit coherent errors, PRX Quantum **6**, 040305 (2025).

[94] M. Goldstein and E. Sela, Symmetry-resolved entanglement in many-body systems, Phys. Rev. Lett. **120**, 200602 (2018).

[95] E. Bianchi, L. Hackl, M. Kieburg, M. Rigol, and L. Vidmar, Volume-law entanglement entropy of typical pure quantum states, PRX Quantum **3**, 10.1103/prxquantum.3.030201 (2022).

[96] S. Aditya, E. Tirrito, P. Sierant, and X. Turkeshi, Coherence dynamics in quantum many-body systems with conservation laws (2026), arXiv:2604.23192 [quant-ph].

[97] D. Iannotti, A. Russotto, B. Jasser, J. Odavić, and A. Hamma, Non-stabilizerness and u(1) symmetry in chaotic many-body quantum systems (2026), arXiv:2603.28870 [quant-ph].

[98] S. Cepollaro, G. Cuffaro, M. B. Weiss, S. Cusumano, A. Hamma, and S. Lloyd, Stabilizer entropy of subspaces (2025), arXiv:2512.23013 [quant-ph].

[99] G. Esposito, S. Cepollaro, L. Cappiello, and A. Hamma, Magic of discrete lattice gauge theories, International Journal of Geometric Methods in Modern Physics **22**, 10.1142/s0219887825500033 (2024).

[100] F. Ares, S. Murciano, and P. Calabrese, Entanglement asymmetry as a probe of symmetry breaking, Nat. Commun. **14**, 2036 (2023).

[101] F. Ares, P. Calabrese, and S. Murciano, The quantum mpemba effects, Nat. Rev. Phys. **7**, 451–460 (2025).

[102] M. P. A. Fisher, V. Khemani, A. Nahum, and S. Vijay, Random quantum circuits, Annu. Rev. Condens. Matter Phys. **14**, 335 (2023).

[103] H.-Z. Li, J.-X. Zhong, and X.-J. Yu, Measurement-induced entanglement phase transition in free fermion systems, Journal of Physics: Condensed Matter **37**, 273002 (2025).

[104] H. Oshima, K. Mochizuki, R. Hamazaki, and Y. Fuji, Topology and spectrum in measurement-induced phase transitions, Phys. Rev. Lett. **134**, 240401 (2025).

[105] Y. Ashida, Z. Gong, and M. Ueda, Non-hermitian physics, Advances in Physics **69**, 249–435 (2020).

[106] K. Kawabata and S. Ryu, Non-hermitian disordered systems (2026), arXiv:2603.20393 [cond-mat.mes-hall].

[107] X. Turkeshi and M. Schiró, Entanglement and correlation spreading in non-hermitian spin chains, Phys. Rev. B **107**, L020403 (2023).

[108] Y. L. Gal, X. Turkeshi, and M. Schirò, Volume-to-area law entanglement transition in a non-Hermitian free fermionic chain, SciPost Phys. **14**, 138 (2023).

[109] E. Granet, C. Zhang, and H. Dreyer, Volume-law to area-law entanglement transition in a nonunitary periodic gaussian circuit, Phys. Rev. Lett. **130**, 230401 (2023).

[110] B. Lapierre, L.-H. Mo, and S. Ryu, Entanglement transitions in structured and random nonunitary gaussian circuits (2025), arXiv:2507.03768 [quant-ph].

[111] M. Collura, B. Béri, and E. Tirrito, Nonlocal nonstabilizerness in free fermion models (2026), arXiv:2604.27055 [quant-ph].

[112] R. Bapat, Mixed discriminants of positive semidefinite matrices, Linear Algebra and its Applications **126**, 107–124 (1989).

[113] R. A. Horn and C. R. Johnson, *Matrix Analysis*, 2nd ed. (Cambridge University Press, Cambridge, 2012).

[114] J. B. Hough, M. Krishnapur, Y. Peres, and B. Virág, Determinantal processes and independence, Probab. Surveys **3**, 206 (2006).

[115] M. A. Woodbury, *Inverting modified matrices* (Princeton University, Princeton, NJ, 1950) p. 4, statistical Research Group, Memo. Rep. no. 42,.

[116] R. Bhatia, *Matrix Analysis* (Springer, 1997).

[117] S. F. E. Oliviero, L. Leone, and A. Hamma, Magic-state resource theory for the ground state of the transverse-field ising model, Phys. Rev. A **106**, 042426 (2022).

END MATTER

Local Gaussian orbit.—For $A = \{1, \dots, \ell\}$ and $B = \{\ell + 1, \dots, N\}$, the JW strings of the Majorana operators of A do not leave A, and those of B differ from the ones defined within B only by the parity $P_A = \prod_{m \in A} Z_m$ of A. Since $U_{O_A} P_A U_{O_A}^\dagger = \det(O_A) P_A$, the qubit unitary $U_{O_A} \otimes U_{O_B}$ acts on the Majorana operators of the chain as $O_A \oplus \det(O_A)\, O_B$, i.e., it coincides up to a phase with a Gaussian unitary of the chain in $\mathrm{O}(2\ell) \oplus \mathrm{O}(2(N-\ell))$. As $(O_A, O_B) \mapsto (O_A, \det(O_A)\, O_B)$ is a bijection, Eq. (3) amounts to maximizing ζ_α over the orbit $\Gamma \mapsto (O_A \oplus O_B)\Gamma(O_A \oplus O_B)^\mathsf{T}$. This is where contiguity enters: for non-contiguous bipartitions, the qubits of A must be brought together by a permutation that does not preserve the Gaussian structure, and the identification fails.

Strategy of the proof.—The idea is to work in the fermionic language: since the optimization runs over local Gaussian unitaries, the natural candidate for the optimum is the BCS canonical form, and the proof shows that it is the global maximizer of ζ_2 on the orbit. On this orbit, ζ_2 is the collision probability of a determinantal distribution which depends on two independent rotations and on *all* minors of the matrix parametrizing the state, so that it is not evidently maximized at the canonical point. The proof (SM, Sec. S1) is a chain of steps, each of which either removes one of these compli-

cations exactly, or bounds the functional by a simpler one that is tight at the canonical point. (1) Wick's theorem turns ζ_2 into the collision probability $d\sum_S p_S^2$ of the projection determinantal point process with kernel $K = (\mathbb{I} + i\Gamma)/2$ [42]. (2) For any isometry $Q \in \mathbb{C}^{n\times k}$, the collision probability of the associated Slater determinant obeys $\mathcal{C}(Q) \le 2^{-k}\det(\mathbb{I} + \Sigma^\dagger\Sigma)$, with $\Sigma = Q^{\mathsf{T}}Q$ the transpose overlap; this sharp bound is proved by induction on n. (3) In the BCS parametrization, the dependence on the local rotations is packaged into $Z = O_A Z_0 O_B^{\mathsf{T}}$, with $Z_0 = \bigoplus_j (\mathbb{I}_2 + i\lambda_j J)/\sqrt{1-\lambda_j^2}$ and $J = \left(\begin{smallmatrix} 0 & 1 \\ -1 & 0 \end{smallmatrix}\right)$, and ζ_2 becomes, up to an orbit-invariant constant, the sum $\mathsf{F}(Z) = \sum_{|I|=|I'|} |\det Z_{I,I'}|^4$ of the fourth powers of the moduli of all minors of Z; a Cauchy–Schwarz argument reduces the two rotations to one, $O_A = O_B$. (4) The bound of step (2) gives $\mathsf{F}(Z) \le \mathsf{R}(H)$, a weighted sum over the principal minors of $H = Z^2$ only, with equality at Z_0. (5) A principal-minor identity and a sign average combined with a Cayley transform rewrite $\mathsf{R}(H)$ exactly as an orbit-invariant prefactor times a sum of squared principal minors of a single Hermitian matrix. (6) Interpreting this sum as the probability of an event for another determinantal process, a change of basis within pairs of coordinates reduces it, up to an orbit-invariant prefactor, to $\mathsf{L}(U) = \sum_S |\det U_S|^2$ for a unitary U. (7) A determinant inequality for positive matrices of rank at most two, $[z_1 \cdots z_n]\det(\sum_i z_i B_i) \le \prod_i \|B_i\|_{\mathrm{HS}}$, with $[z_1 \cdots z_n]$ the coefficient of $z_1 \cdots z_n$, proved with fermionic creation operators, bounds $\mathsf{L}(U)$ through the diagonal of U, and hence by the orbit invariant $\det(\mathbb{I} + T^\dagger T)$, $T = (U + U^{\mathsf{T}})/2$. (8) The canonical form saturates every inequality of the chain, which proves Theorem 1; a continuity argument covers $\lambda_j = 1$, and padding with ancillas removes the assumption $N = 2\ell$.

Page curve.—For Haar-random $O \in \mathrm{O}(2N)$ and $|A| = \ell \le N/2$, the joint distribution of $x_m = \lambda_m^2$ is [63]

$$\mathbb{P}(x_1, \dots, x_\ell) = \frac{1}{Z_\ell} \prod_{1\le i<j\le \ell} (x_i - x_j)^2 \prod_{m=1}^{\ell} \frac{(1-x_m)^{N-2\ell}}{\sqrt{x_m}} , \tag{11}$$

with Z_ℓ a normalization. The one-point density $\varrho^{(\ell)}(x) = \mathbb{E}_{\mathbb{P}}[\sum_m \delta(x - x_m)]$, normalized to $\int_0^1 \varrho^{(\ell)} = \ell$, is given by the orthogonal-polynomial kernel [51, 52, 61]

$$\varrho^{(\ell)}(x) = w(x) \sum_{n=0}^{\ell-1} p_n(x)^2 , \qquad w(x) = \frac{(1-x)^{N-2\ell}}{\sqrt{x}} , \tag{12}$$

where $p_n(x) = \mathrm{P}_n^{(-1/2, N-2\ell)}(1-2x)/\sqrt{h_n}$ are orthonormal with respect to w on $[0,1]$, $\mathrm{P}_n^{(a,b)}$ are Jacobi polynomials, and

$$h_n = \frac{\Gamma(n+\frac12)\Gamma(n+N-2\ell+1)}{(2n+N-2\ell+\frac12)\Gamma(n+1)\Gamma(n+N-2\ell+\frac12)} . \tag{13}$$

Equation (12) gives the exact quenched average $\overline{M}_2^{\mathrm{GNL}} = \int_0^1 \mathfrak{m}_2(x)\varrho^{(\ell)}(x)dx$ at finite N, which we benchmarked against direct sampling of Haar-random $O \in \mathrm{O}(2N)$, including the data of Fig. 1(a), finding agreement within statistical errors. For $\ell, N \to \infty$ at fixed $r = \ell/N \le 1/2$, $\varrho^{(\ell)}/N$ converges to the free Jacobi law $\varrho_r(x) = \sqrt{(c-x)x}/[2\pi x(1-x)]$ on $[0,c]$, with $c = 4r(1-r)$ and total mass r; $O(1)$ shifts of the exponents of $w(x)$ only affect $O(1/N)$ corrections to $\overline{M}_2^{\mathrm{GNL}}/N$. Setting $x = c\sin^2\theta$,

$$\overline{\mathfrak{m}}_2^{\mathrm{GNL}}(r) \simeq \frac{c}{\pi}\int_0^{\pi/2} \frac{\cos^2\theta}{1 - c\sin^2\theta}\, \mathfrak{m}_2(c\sin^2\theta)\, d\theta , \tag{14}$$

which gives $2 - \log_2(2+\sqrt{3})$ at $r = 1/2$ and, with $c \simeq 4r$, $r^2/\ln 2$ for $r \to 0$.

XY chain.—After the JW mapping, Eq. (6) takes the form of Eq. (1) with $h_{2m-1,2m} = 2\mu$, $h_{2m,2m+1} = 1+\eta$, $h_{2m+2,2m-1} = 1-\eta$, the remaining entries being fixed by antisymmetry [49]. We take N divisible by 4, so that the ground state lies in the Neveu–Schwarz sector, $c_{N+1} = -c_1$ (i.e., $\gamma_{2N+1} = -\gamma_1$, $\gamma_{2N+2} = -\gamma_2$), with momenta $k = \pi(2n+1)/N$, $n = 0, \dots, N/2-1$. A Fourier transform followed by a Bogoliubov rotation at each k diagonalizes ih, whose eigenvalues are $\pm 2\varepsilon_k$, with Bogoliubov angle and dispersion

$$\begin{aligned} e^{i\theta_k} &= (\mu - \cos k + i\eta \sin k)/\varepsilon_k , \\ \varepsilon_k &= \sqrt{(\mu - \cos k)^2 + \eta^2 \sin^2 k} . \end{aligned} \tag{15}$$

For $N \to \infty$, the reduced covariance matrix of $A = \{1, \dots, \ell\}$ is block Toeplitz,

$$\begin{aligned} (\Gamma_A)_{mn} &= \int_{-\pi}^{\pi} \frac{dk}{2\pi}\, e^{ik(m-n)}\, g(k) , \\ g(k) &= \begin{pmatrix} 0 & e^{i\theta_k} \\ -e^{-i\theta_k} & 0 \end{pmatrix} . \end{aligned} \tag{16}$$

The reduced state is $\rho_A \propto \exp(-H_{\mathrm{ent}})$, with $H_{\mathrm{ent}} = \frac{i}{2}\sum_n \varepsilon_n^{\mathrm{ent}} \tilde\gamma_{2n-1}\tilde\gamma_{2n}$ for suitable Majorana operators $\tilde\gamma$, so that $\lambda_n = \tanh(\varepsilon_n^{\mathrm{ent}}/2)$ [71]; $\lambda_n = 1$ corresponds to an unentangled mode. For a semi-infinite cut, the corner transfer matrix gives the equally spaced spectrum quoted in the main text [69, 70], with $\varepsilon = \pi K(\kappa')/K(\kappa)$, where K is the complete elliptic integral of the first kind, $\kappa' = \sqrt{1-\kappa^2}$, and

$$\kappa = \begin{cases} \eta/\sqrt{\eta^2+\mu^2-1} & \mu > 1 , \\ \sqrt{\eta^2+\mu^2-1}/\eta & \mu < 1,\ \eta^2+\mu^2 > 1 , \\ \sqrt{(1-\eta^2-\mu^2)/(1-\mu^2)} & \mu < 1,\ \eta^2+\mu^2 < 1 . \end{cases} \tag{17}$$

At $\mu = 1$ ($\eta = 1$), $\kappa \to 1$ and $\varepsilon \to 0$, with correlation length $\xi \simeq |\mu - 1|^{-1}$.

Area law.—With $\mathfrak{f}(x) := -x\log_2 x - (1-x)\log_2(1-x)$, the Gaussian entanglement entropy of the contiguous

block $A = \{1, \dots, \ell\}$ of Theorem 1 is $S(A) = \sum_i \mathfrak{f}[(1+\lambda_i)/2]$. Since $\mathfrak{m}_\alpha(x) = H_\alpha\big(\frac{1}{2}, \frac{x}{2}, \frac{1-x}{2}\big) - 1$, with H_α the Rényi entropy (in bits), which is non-increasing in α, we have $\mathfrak{m}_\alpha(x) \le \mathfrak{m}_1(x) = \frac{1}{2}\mathfrak{f}(x)$ for $\alpha \ge 1$. Together with $\frac{1}{2}\mathfrak{f}(\lambda^2) \le 2\,\mathfrak{f}[(1+\lambda)/2]$ for $\lambda \in [0,1)$, this gives $M_\alpha^{\rm GNL} \le 2S(A)$, the Gaussian analog of the bound of Ref. [36], rigorous at $\alpha = 2$ by Theorem 1. Hastings' theorem [74], $S(A) \le C_{\rm ent}|\partial A|$ for unique gapped ground states of one-dimensional local Hamiltonians, then yields $M_\alpha^{\rm NL} \le M_\alpha^{\rm GNL} \le 2C_{\rm ent}|\partial A|$.

Quench dynamics.—After the quench $\mu_0 \to \mu$ at $\eta = 1$, with time measured in the units of Refs. [75, 77], i.e., for the evolution generated by $H(\mu)/2$, $\Gamma_A(t)$ is block Toeplitz as in Eq. (16) (with $k \in [0, 2\pi)$), with symbol [75]

$$g(k,t) = \begin{pmatrix} -q(k,t) & p(k,t) \\ -p(-k,t) & q(k,t) \end{pmatrix}, \tag{18}$$

where $p(k,t) = e^{-i\theta_k}\,[\cos\Delta\theta_k + i \sin\Delta\theta_k \cos(2\varepsilon_k t)]$, $q(k,t) = i \sin\Delta\theta_k \sin(2\varepsilon_k t)$, ε_k and θ_k refer to the post-quench Hamiltonian, and

$$\cos\Delta\theta_k = \frac{(\mu - \cos k)(\mu_0 - \cos k) + \sin^2 k}{\varepsilon_k\, \varepsilon_k^{(0)}}\,. \tag{19}$$

At $t = 0$, $g(k,0)$ coincides with the ground-state symbol of $H(\mu_0)$ at $-k$, i.e., up to the reflection $m \to \ell+1-m$ of the block, which leaves the spectrum of $i\Gamma_A$ unchanged, while for $t \to \infty$ the oscillating terms dephase, leaving $p(k,\infty) = e^{-i\theta_k}\cos\Delta\theta_k$ and $q = 0$, i.e., the GGE leading to Eq. (9). At finite times, we write

$$M_2^{\rm GNL} = \frac{1}{4\pi i}\oint_{\mathcal{C}} d\lambda\; \mathfrak{m}_2(\lambda^2)\; \frac{d}{d\lambda} \ln\det(\lambda \mathbb{I}_{2\ell} - i\Gamma_A)\,, \tag{20}$$

where $\mathcal{C}$ encircles all eigenvalues counterclockwise and the factor $1/2$ compensates the double counting of the $\pm\lambda_n$ pairs. Expanding in λ^{-1} expresses $M_2^{\rm GNL}$ through the moments $\mathrm{Tr}(i\Gamma_A)^{2n} = \mathrm{Tr}\, W_+^{2n} + \mathrm{Tr}\, W_-^{2n}$, where $W_\pm = \mathcal{H} \pm i\mathcal{T}$ are Hermitian, with $\mathcal{T}_{jl} = \int \frac{dk}{2\pi} e^{-ik(j-l)} q(k,t)$ Toeplitz and $\mathcal{H}_{jl} = \int \frac{dk}{2\pi} e^{-ik(j+l-\ell-1)} p(k,t)$ Hankel. Following Ref. [77], a multidimensional stationary-phase analysis of the resulting momentum integrals gives, for $\ell \gg 1$,

$$\mathrm{Tr}(i\Gamma_A)^{2n} \simeq \frac{2\ell}{\pi}\int_0^\pi dk \Big[1 - \big(1 - \cos^{2n}\Delta\theta_k\big) \times \min\Big(1, \frac{2|v_k|t}{\ell}\Big)\Big], \tag{21}$$

which interpolates ballistically between the pure-state value 2ℓ and the GGE moments. Inserting Eq. (21) into Eq. (20) yields Eq. (10).