

# Assignment 1 of MATP6610/4820

(Due on Jan-31-2020 in class)

## Problem 1

Show that the sequence  $x^{(k)} = (\frac{3}{4})^{2^k} - 2$  is Q-quadratically convergent to  $-2$ .

---

## Problem 2

Consider the sequence  $\{x^{(k)}\}_{k=1}^{\infty}$  defined by

$$x^{(k)} = \begin{cases} (\frac{1}{2})^{2^k}, & \text{if } k \text{ is odd} \\ \frac{1}{k}x^{(k-1)}, & \text{if } k \text{ is even} \end{cases}$$

Is this sequence Q-superlinearly convergent? Q-quadratically convergent? R-quadratically convergent? Justify your answer.

---

### Problem 3

Find all stationary points of  $f(x_1, x_2) = 8x_1 + 12x_2 + x_1^2 - 2x_2^2$  and verify if they are local minimizers, or local maximizers, or saddle points.

---

## Problem 4

Consider the function

$$f(x_1, x_2) = \frac{1}{2} \left( 2x_1 + \frac{1}{2}x_2^2 \right)^2.$$

1. Give the gradient of  $f$  at the point  $(1, 0)$ ;
2. Show that  $\mathbf{p} = (-1, 2)$  is a descent direction, i.e.,  $\langle \mathbf{p}, \nabla f(1, 0) \rangle < 0$ , where  $\langle \cdot, \cdot \rangle$  denotes vector inner product.
3. Let  $c_1 = 0.1$ . Does  $\alpha = 1$  satisfy the Armijo's condition?