



*National Aeronautics and Space  
Administration Goddard Earth Science  
Data Information and Services Center (GES  
DISC)*

# README Document for NASA GLDAS Version 2 Data Products

---

**Last Revised February 27, 2020**

Goddard Earth Sciences Data and Information Services Center (GES DISC)  
<http://disc.gsfc.nasa.gov>  
NASA Goddard Space Flight Center, Code 610.2  
Greenbelt, MD 20771 USA

# README for NASA GLDAS Version 2 Data

Prepared By:

**Hualan Rui**\_\_\_\_\_

GES DISC  
GSFC Code 610.2

**Hiroko Beaudoin**\_\_\_\_\_

Hydrological Sciences Laboratory (HSL)  
GSFC Code 617

August 6, 2019\_\_\_\_\_

Date

Reviewed By:

**Carlee Loeser**\_\_\_\_\_

GES DISC  
GSFC Code 610.2

February 27, 2020\_\_\_\_\_

Date

**Goddard Space Flight  
Center Greenbelt,  
Maryland**

## Revision History

---

Revision	Change	Author
05/19/2010	Initial version based on information from Hiroko Beaudoin.	Hualan Rui
06/17/2010	Review and revise	Hiroko Beaudoin
11/21/2011	Update GES DISC Helpdesk email address	Hualan Rui
10/16/2012	Add information for GLDAS-2.0	Hiroko Beaudoin
11/01/2012	Add information for GLDAS-2.0	Hualan Rui
11/15/2013	Add information for GLDAS-2.0 0.25° product	Hiroko Beaudoin
11/18/2014	Update Table 3 for additional fields	Hualan Rui
05/12/2015	Add Table 2: DOIs for GLDAS-2 products	Hualan Rui
06/16/2015	Add information for GLDAS-2.0 products	Hualan Rui
07/03/2015	Review and revise	Hiroko Beaudoin
07/22/2016	Add information for GLDAS-2.1 Noah 1.0 ° products	Hiroko Beaudoin
08/01/2016	Add GLDAS-1 and GLDAS-2.1 differences	Hiroko Beaudoin
11/11/2016	Add information for GLDAS-2.1 Noah 0.25 °	Hualan Rui
02/08/2017	Convert to comply with the newer README	Hualan Rui
07/18/2017	Update URLs to comply with GES DISC new Web	Hualan Rui
11/16/2017	Add more information regarding accumulation	Carlee Loeser
12/20/2017	Add information for Catchment model data	Hiroko Beaudoin
02/02/2018	Add data interpretation for TWS	Hiroko Beaudoin
02/06/2018	Review and revise	Bailing Li
03/29/2018	Add DOI for Catchment model data product	Hualan Rui
04/05/2019	Update Data Interpolation for TWS and GWS	Hualan Rui
08/06/2019	Update References and add snow density note	Carlee Loeser
09/18/2019	Update for the reprocessed data	Hiroko Beaudoin
11/29/2019	Review and revise	Carlee Loeser
02/27/2020	Add new GLDAS-2.1 and GLDAS-2.2 products	Carlee Loeser

# Table of Contents

---

1.0 Introduction.....	6
1.1 Basic Characteristics of GLDAS-2 .....	6
1.2 Specifications of GLDAS-2.....	7
1.2.1 Land Surface Models.....	7
1.2.2 GLDAS-2.0.....	7
1.2.3 GLDAS-2.1 .....	7
1.2.4 GLDAS-2.2.....	8
1.3. Digital Object Identifier (DOI) and Citations.....	8
1.3 Contact Information .....	11
1.4 What's New?.....	12
1.4.1 What is new about the reprocessed GLDAS-2.0 and GLDAS-2.1?.....	12
1.4.2 What are the newest datasets for GLDAS-2?.....	12
1.4.3 What are the differences between GLDAS-1 and GLDAS-2.1?.....	12
1.4.4 What is GLDAS-2.2? .....	13
2.0 Data Organization .....	14
2.1 File Naming Convention .....	14
2.2 File Format and Structure.....	15
3.0 Data Contents.....	15
3.1 GLDAS-2.0 Data.....	15
3.2 GLDAS-2.1 Data.....	17
3.3 GLDAS-2.2 Data.....	21
3.4 Data Interpretation.....	22
4.0 Options for Reading the Data .....	23
4.1 Utilities.....	23
4.2 Panoply .....	23
4.3 GrADS.....	24
5.0 Data Services .....	24
5.1 HTTPS .....	24
5.2 Earthdata Search.....	24
5.3 GES DISC Subsetter/Regridder.....	25
5.4 OPeNDAP.....	25

## README for NASA GLDAS Version 2 Data

5.5 GrADS Data Server (GDS).....	25
5.6 Giovanni.....	25
6.0 More Information .....	26
7.0 Acknowledgements .....	26
References.....	26
Acronyms.....	28
Appendix A .....	29

# 1.0 Introduction

---

This document provides the basic information for using NASA GLDAS Version 2 products.

The goal of the NASA Global Land Data Assimilation System (GLDAS) is to generate optimal fields of land surface states and fluxes, by ingesting satellite- and ground-based observational data products, using advanced land surface modeling and data assimilation techniques (Rodell et al., 2004). GLDAS drives multiple, offline (not coupled to the atmosphere) land surface models, integrates a huge quantity of observation-based data, and executes globally at high resolutions (2.5° to 1 km), enabled by the Land Information System (LIS) (Kumar et al., 2006). Currently, GLDAS drives four land surface models (LSMs): Noah, Catchment (CLSM), the Community Land Model (CLM), and the Variable Infiltration Capacity (VIC). More information is available at the [Land Data Assimilation Systems \(LDAS\)](#) and [Land Information System \(LIS\)](#) websites.

This document specifically describes the reprocessed data products of Version 2 of the Global Land Data Assimilation System (hereafter, GLDAS-2).

## 1.1 Basic Characteristics of GLDAS-2

NASA GLDAS-2 has three components: GLDAS-2.0, GLDAS-2.1, and GLDAS-2.2. GLDAS-2.0 is forced entirely with the Princeton meteorological forcing input data and provides a temporally consistent time series from 1948 through 2014. GLDAS-2.1 is forced with a combination of model and observation data from 2000 to present. GLDAS-2.2 product suites use data assimilation (DA), whereas the GLDAS-2.0 and GLDAS-2.1 products are “open-loop” (i.e., no data assimilation). Choice of forcing data, as well as DA observation source, variable, and scheme vary for different GLDAS-2.2 products. Currently, the GLDAS-2.2 products include data assimilation from the Gravity Recovery and Climate Experiment (GRACE) from 2003 to present.

The temporal resolutions for the GLDAS-2 products are 3-hourly and daily. The monthly products are generated through temporal averaging of the 3-hourly products. Table 1 lists some basic characteristics of the GLDAS-2 data. Please check up on the newest hydrology-related alert messages at [GES DISC Alerts](#).

Table 1. Basic characteristics of the NASA GLDAS-2 data.

<b>Contents</b>	<b>Outputs from Land Surface Models</b>
Format	NetCDF
Latitude Extent	-60° to 90°
Longitude Extent	-180° to 180°
Spatial Resolution	1.0°, 0.25°
Temporal Resolution	3-hourly, daily, monthly
Temporal Coverage	GLDAS-2.0: 03Z January 1, 1948 – 21Z December 31, 2014

	GLDAS-2.1: 03Z January 1, 2000 – Present
	GLDAS-2.2: February 1, 2003 – Present
Dimensions	360 (lon) x 150 (lat) for the 1.0° x 1.0° data
	1440 (lon) x 600 (lat) for the 0.25° x 0.25° data
Origins (1 <sup>st</sup> grid center)	(179.5 W, 59.5 S) for the 1.0° x 1.0° data
	(179.875 W, 59.875 S) for the 0.25° x 0.25° data
Land Surface Models	Noah-3.6, CLSM-F2.5, VIC-4.1.2

## 1.2 Specifications of GLDAS-2

### 1.2.1 Land Surface Models

The Noah model uses the Modified IGBP MODIS 20-category vegetation classification and the soil texture based on the Hybrid STATSGO/FAO datasets. The Catchment model uses the Mosaic land cover classification, together with soils, topographic, and other model-specific parameters that were derived in a manner consistent with that of the NASA/GMAO's GEOS-5 climate modeling system. The VIC model uses the UMD land cover classification, and the parameters are derived from the 0.5-degree Global VIC dataset (Nijssen et al., 2014).

### 1.2.2 GLDAS-2.0

The GLDAS-2.0 model simulations were initialized on simulation date January 1, 1948, using soil moisture and other state fields from the LSM climatology for that day of the year. The simulations were forced by the global meteorological forcing dataset from Princeton University (Sheffield et al., 2006). Each simulation uses the common GLDAS datasets for land water mask (MOD44W: Carroll et al., 2009) and elevation (GTOPO30), along with the model default land cover and soils datasets. The MODIS-based land surface parameters are used in the current GLDAS-2.0 and GLDAS-2.1 products, while the AVHRR-based parameters were used in GLDAS-1 and previous GLDAS-2 products (prior to October 2012). The land mask was modified to accommodate the river routing scheme included in the simulations in the November 2019-December 2020 reprocessing update.

### 1.2.3 GLDAS-2.1

The GLDAS-2.1 model simulation started on January 1, 2000 using the conditions from the GLDAS-2.0 simulation. This simulation was forced with National Oceanic and Atmospheric Administration (NOAA)/Global Data Assimilation System (GDAS) atmospheric analysis fields (Derber et al., 1991), the disaggregated Global Precipitation Climatology Project (GPCP) V1.3 Daily Analysis precipitation fields (Adler et al., 2003; Huffman et al., 2001), and the Air Force Weather Agency's AGRicultural METeorological modeling system (AGRMET) radiation fields. The simulation was only used with GDAS and GPCP from January 2000 to February 2001, followed by the addition of AGRMET from March 1, 2001 onwards.

## README for NASA GLDAS Version 2 Data

GLDAS-2.1 data products are available in two production streams: one forced with combined forcing data including the GPCP version 1.3 (main production stream), and one without this forcing data (the early production stream). Since the GPCP version 1.3 data have a 3-4 month latency, the GLDAS-2.1 data products are first created without it, and are designated as Early Products (EPs), with about 1.5 month latency. Once the GPCP version 1.3 data become available, the GLDAS-2.1 data products are processed in the main production stream and are removed from the Early Products archive.

### 1.2.4 GLDAS-2.2

The GLDAS-2.2 Daily Catchment model simulation started on February 1, 2003 using the conditions from the GLDAS-2.0 Daily Catchment model simulation. This simulation was forced with the meteorological analysis fields from the operational European Centre for Medium-Range Weather Forecasts (ECMWF) Integrated Forecasting System (<https://www.ecmwf.int/en/publications/ifs-documentation>). The total terrestrial water anomaly observation from Gravity Recovery and Climate Experiment (GRACE) was assimilated (Li et al., 2019). The GRACE RL06 and GRACE Follow-On data were provided by the Center for Space Research at the University of Texas (Save et al., 2012; Save et al., 2016). GRACE data have a latency of 2-6 months; thus, the simulation extends through the present without DA in recent months and the GLDAS-2.2 data are reprocessed and replaced as the GRACE data become available. The Daily Catchment model simulations use the UMD land cover scheme from AVHRR land cover map. Due to the data agreement with the ECMWF that prohibits dissemination of the IFS product, this GLDAS-2.2 Daily product does not include the meteorological fields.

GLDAS-2.2 data products are available in two production streams: one with GRACE data assimilation outputs (main production stream), and one without GRACE data (the early production stream). Since the GRACE data have a 2-6 month latency, the GLDAS-2.2 data products are first created without it, and are designated as Early Products (EPs), with about 1.5 month latency. Once the GRACE data become available, the GLDAS-2.2 data products are processed in the main production stream and are removed from the Early Products archive.

## 1.3. Digital Object Identifier (DOI) and Citations

Users of GLDAS data products should cite the data in their research papers with the Digital Object Identifiers (DOIs). A DOI is a unique alphanumeric string used to identify a digital object and provide a permanent link online. DOIs are often used in online publications in citations. Table 2 provides the DOIs for each GLDAS-2 data product.



# README for NASA GLDAS Version 2 Data

Table 2. DOIs for NASA GLDAS-2 Data Products

	<b>Data Product Name</b>	<b>DOI</b>
<b>GLDAS-2.0</b>	GLDAS Catchment Land Surface Model L4 daily 0.25 x 0.25 degree V2.0 (GLDAS_CLSM025_D_2.0)	<a href="https://doi.org/10.5067/LYHA9088MFWQ">10.5067/LYHA9088MFWQ</a>
	GLDAS Noah Land Surface Model L4 3 hourly 0.25 x 0.25 degree V2.0 (GLDAS_NOAH025_3H_2.0)	<a href="https://doi.org/10.5067/3420HQM9AK6Q">10.5067/3420HQM9AK6Q</a>
	GLDAS Noah Land Surface Model L4 monthly 0.25 x 0.25 degree V2.0 (GLDAS_NOAH025_M_2.0)	<a href="https://doi.org/10.5067/9SQ1B3ZXP2C5">10.5067/9SQ1B3ZXP2C5</a>
	GLDAS Noah Land Surface Model L4 3 hourly 1.0 x 1.0 degree V2.0 (GLDAS_NOAH10_3H_2.0)	<a href="https://doi.org/10.5067/L0JGCNVBNRAX">10.5067/L0JGCNVBNRAX</a>
	GLDAS Noah Land Surface Model L4 monthly 1.0 x 1.0 degree V2.0 (GLDAS_NOAH10_M_2.0)	<a href="https://doi.org/10.5067/QN80TO7ZHFJZ">10.5067/QN80TO7ZHFJZ</a>
<b>GLDAS-2.1</b>	GLDAS Catchment Land Surface Model L4 3 hourly 1.0 x 1.0 degree V2.1 (GLDAS_CLSM10_3H_2.1)	<a href="https://doi.org/10.5067/VCO8OCV72XOQ">10.5067/VCO8OCV72XOQ</a>
	GLDAS Catchment Land Surface Model L4 3 hourly 1.0 x 1.0 degree Early Product V2.1 (GLDAS_CLSM10_3H_EP_2.1)	<a href="https://doi.org/10.5067/W024WFHJXZ0E">10.5067/W024WFHJXZ0E</a>
	GLDAS Catchment Land Surface Model L4 monthly 1.0 x 1.0 degree V2.1 (GLDAS_CLSM10_M_2.1)	<a href="https://doi.org/10.5067/FOUXNLXFAZNY">10.5067/FOUXNLXFAZNY</a>
	GLDAS Catchment Land Surface Model L4 monthly 1.0 x 1.0 degree Early Product V2.1 (GLDAS_CLSM10_M_EP_2.1)	<a href="https://doi.org/10.5067/1LQF1ORIE8OW">10.5067/1LQF1ORIE8OW</a>
	GLDAS Noah Land Surface Model L4 3 hourly 0.25 x 0.25 degree V2.1 (GLDAS_NOAH025_3H_2.1)	<a href="https://doi.org/10.5067/E7TYRXPJKWOQ">10.5067/E7TYRXPJKWOQ</a>
	GLDAS Noah Land Surface Model L4 3 hourly 0.25 x 0.25 degree Early Product V2.1 (GLDAS_NOAH025_3H_EP_2.1)	<a href="https://doi.org/10.5067/G90R32A924YM">10.5067/G90R32A924YM</a>
	GLDAS Noah Land Surface Model L4 monthly 0.25 x 0.25 degree V2.1 (GLDAS_NOAH025_M_2.1)	<a href="https://doi.org/10.5067/SXAVCZFAQLNO">10.5067/SXAVCZFAQLNO</a>
	GLDAS Noah Land Surface Model L4 monthly 0.25 x 0.25 degree Early Product V2.1 (GLDAS_NOAH025_M_EP_2.1)	<a href="https://doi.org/10.5067/5OVHMF2IAV3">10.5067/5OVHMF2IAV3</a>

# README for NASA GLDAS Version 2 Data

	GLDAS Noah Land Surface Model L4 3 hourly 1.0 x 1.0 degree V2.1 (GLDAS_NOAH10_3H_2.1)	<a href="https://doi.org/10.5067/IIG8FHR17DA9">10.5067/IIG8FHR17DA9</a>
	GLDAS Noah Land Surface Model L4 3 hourly 1.0 x 1.0 degree Early Product V2.1 (GLDAS_NOAH10_3H_EP_2.1)	<a href="https://doi.org/10.5067/7FK9SEEE6VP3">10.5067/7FK9SEEE6VP3</a>
	GLDAS Noah Land Surface Model L4 monthly 1.0 x 1.0 degree V2.1 (GLDAS_NOAH10_M_2.1)	<a href="https://doi.org/10.5067/LWTYSMP3VM5Z">10.5067/LWTYSMP3VM5Z</a>
	GLDAS Noah Land Surface Model L4 monthly 1.0 x 1.0 degree Early Product V2.1 (GLDAS_NOAH10_M_EP_2.1)	<a href="https://doi.org/10.5067/MCM8JKVDO3W3">10.5067/MCM8JKVDO3W3</a>
	GLDAS VIC Land Surface Model L4 3 hourly 1.0 x 1.0 degree V2.1 (GLDAS_VIC10_3H_2.1)	<a href="https://doi.org/10.5067/ZOG6BCSE26HV">10.5067/ZOG6BCSE26HV</a>
	GLDAS VIC Land Surface Model L4 3 hourly 1.0 x 1.0 degree Early Product V2.1 (GLDAS_VIC10_3H_EP_2.1)	<a href="https://doi.org/10.5067/KMPD4R2A549N">10.5067/KMPD4R2A549N</a>
	GLDAS VIC Land Surface Model L4 monthly 1.0 x 1.0 degree V2.1 (GLDAS_VIC10_M_2.1)	<a href="https://doi.org/10.5067/VWTH7S6218SG">10.5067/VWTH7S6218SG</a>
	GLDAS VIC Land Surface Model L4 monthly 1.0 x 1.0 degree Early Product V2.1 (GLDAS_VIC10_M_EP_2.1)	<a href="https://doi.org/10.5067/472GKYTU73QR">10.5067/472GKYTU73QR</a>
<b>GLDAS-2.2</b>	GLDAS Catchment Land Surface Model L4 daily 0.25 x 0.25 degree GRACE-DA1 V2.2 (GLDAS_CLSM025_DA1_D_2.2)	<a href="https://doi.org/10.5067/TXBMLX370XX8">10.5067/TXBMLX370XX8</a>
	GLDAS Catchment Land Surface Model L4 daily 0.25 x 0.25 degree GRACE-DA1 Early Product V2.2 (GLDAS_CLSM025_DA1_D_EP_2.2)	<a href="https://doi.org/10.5067/IIU5JWU2AGRP">10.5067/IIU5JWU2AGRP</a>

Each of the DOIs in Table 2 is linked to its corresponding dataset landing page. On the page, the tab labeled “Data Citation” provides the recommended citation for that product. If you use a GLDAS data product(s) in your research or applications, please include the corresponding reference(s) in your publication(s).

For example, the following is the citation for GLDAS\_NOAH025\_3H\_2.1:  
Beaudoin, H. and M. Rodell, NASA/GSFC/HSL (2016), GLDAS Noah Land Surface Model L4 3 hourly 0.25 x 0.25 degree V2.1, Greenbelt, Maryland, USA, Goddard Earth

## README for NASA GLDAS Version 2 Data

Sciences Data and Information Services Center (GES DISC), Accessed: [*Data Access Date*], [10.5067/E7TYRXPJKWOQ](https://disc.gsfc.nasa.gov/datasets/10.5067/E7TYRXPJKWOQ)

Please also cite the primary reference for GLDAS-2:

Rodell, M., P.R. Houser, U. Jambor, J. Gottschalck, K. Mitchell, C.-J. Meng, K. Arsenault, A. Cosgrove, J. Radakovich, M. Bosilovich, J. K. Entin, J. P. Walker, D. Lohmann, and D. Toll, 2004. The Global Land Data Assimilation System, *Bull. Amer. Meteor. Soc.*, 85(3): 381-394, 10.1175/BAMS-85-3-381.

### 1.3 Contact Information

For information about or assistance in using any GES DISC data, please contact the GES DISC Help Desk at:

Goddard Earth Sciences Data and Information Services Center, Code 610.2  
NASA Goddard Space Flight Center  
Greenbelt, MD 20771  
Email: [gsfc-help-disc@lists.nasa.gov](mailto:gsfc-help-disc@lists.nasa.gov)  
Phone: 301-614-5224  
Fax: 301-614-5268

For general science questions and comments, please contact:

Matthew Rodell, Ph.D.  
Hydrological Sciences Laboratory, Code 617  
NASA Goddard Space Flight Center  
Greenbelt, MD 20771  
Email: [Matthew.Rodell@nasa.gov](mailto:Matthew.Rodell@nasa.gov)  
Phone: 301-286-9143

-or-

Hiroko Kato Beaudoin, M.S.  
Earth System Science Interdisciplinary Center  
University of Maryland, College Park  
Hydrological Sciences Laboratory, Code 617  
NASA Goddard Space Flight Center  
Greenbelt, MD 20771  
Email: [Hiroko.Kato-1@nasa.gov](mailto:Hiroko.Kato-1@nasa.gov)

-or-

Bailing Li, Ph.D.  
Earth System Science Interdisciplinary Center  
University of Maryland, College Park  
Hydrological Sciences Laboratory, Code 617  
NASA Goddard Space Flight Center  
Greenbelt, MD 20771  
Email: [bailing.li-1@nasa.gov](mailto:bailing.li-1@nasa.gov)

## 1.4 What's New?

### 1.4.1 What is new about the reprocessed GLDAS-2.0 and GLDAS-2.1?

In November and December 2019, the GLDAS-2.0 Noah products were reprocessed with updated Princeton Forcing V2.2 Data, an upgraded version of Noah model (V3.6), and upgraded Land Information System (LIS) software. The reprocessed GLDAS-2 data are archived in NetCDF-4 format. Additional model outputs are included and described for each model (see Tables 3.1, 3.2, and 3.3). The land surface characteristics (i.e., land cover, soil texture) over some grid cells were modified due to the routing. Details of the changes and the new land surface parameter datasets are available at <https://ldas.gsfc.nasa.gov/gldas/>. The GLDAS-2.0 data were extended through December 2014 during this time.

In January and February 2020, the GLDAS-2.1 Noah products were also reprocessed with these same upgrades. During this time, an additional production stream was added for GLDAS-2.1. Since the GPCP version 1.3 data have a 3-4 month latency, the GLDAS-2.1 data products are first created without it, and are now designated as Early Products (EPs), with about 1.5 month latency. Once the GPCP version 1.3 data become available, the GLDAS-2.1 data products are processed in the main production stream and are removed from the Early Products archive.

### 1.4.2 What are the newest datasets for GLDAS-2?

In March 2018, Daily Catchment LSM outputs at 0.25-degree resolution were added to the GLDAS-2.0 suite (Li et al., 2019).

In February 2020, GLDAS-2.1 VIC and CLSM LSMs simulation outputs were publicly released. GLDAS-2.1 extends from 2000 to present with about 1.5-month latency and is updated monthly.

In February 2020, the GLDAS-2.2 daily data products with GRACE data assimilation (known as DA1) were publicly released. GLDAS-2.2 extends from February 1, 2003 to present.

In March 2020, the GLDAS-1 forward stream will end and all products will be decommissioned in June 2020. GLDAS-2.1 will serve as the replacement for GLDAS-1.

### 1.4.3 What are the differences between GLDAS-1 and GLDAS-2.1?

The main objective of GLDAS-2.1 is to provide up-to-date global land surface model outputs, using observation-based forcing, while preserving consistency of the long-term climatology (i.e., GLDAS-2.0) to the extent possible. Two major issues were found in the GLDAS-1 forcing fields. First, the AGRMET shortwave downward radiation flux displayed sharp, unnatural gradient lines in the Northern Hemisphere during certain

years. Second, there was a dramatic change in precipitation in certain locations starting in 2009. Furthermore, comparisons of GLDAS-1 radiation and precipitation fields revealed that GLDAS-1 had high bias relative to the well-validated Surface Radiation Budget (SRB) dataset (Stackhouse et al., 2011), and GLDAS-1 precipitation (i.e., CMAP) had low bias relative to the Global Precipitation Climatology Project (GPCP) dataset. Similar biases were observed compared to GLDAS-2.0 (i.e., Princeton forcing), whose radiation fields were bias corrected to the SRB dataset and precipitation fields were disaggregated using the GPCP and Tropical Rainfall Measuring Mission (TRMM) datasets.

GLDAS-2.1 addressed these issues as follows. The AGRMET radiation flux fields were bias-corrected using the period of overlap between AGRMET and SRB (2002-2007) to compute monthly, gridded scale factors that are applied for the overlapping period of AGRMET data. Similarly, GDAS radiation fields were bias-corrected to SRB for the period of 2000-2001/02. Because AGRMET displayed high bias compared to SRB, the fluxes for 2008 onwards are adjusted by applying another set of gridded scale factors that are computed from the annual mean climatology of 2002-2007, thus avoiding a discontinuity in the GLDAS-2.1 data. The bias-corrected AGRMET forcing data should be consistent with the climatology of SRB; however, due to the short overlapping period, the scaling approach is unable to correct an apparent shift (of unknown origin) in the AGRMET climatology after 2011. For the precipitation fields, we used the GPCP 1-degree Daily dataset (Huffman et al., 2001) and an updated disaggregation routine (making use of GDAS precipitation fields) to prepare 3-hourly GPCP fields. GLDAS-2.1 products obtained before January and February 2020 used the GPCP 1DD v1.2 data for January 2000 through October 2015, and the GDAS was used for November 2015 onwards, in order to run GLDAS-2.1 up to present. In the reprocessed GLDAS-2.1 data, a new version 1.3 of the GPCP daily data is used, in which extends through present with 3-4 months latency. GDAS precipitation data fill the gap between the last available GPCP data to present; therefore, GLDAS-2.1 will be reprocessed and replaced for the recent months as the GPCP data updates each month.

### 1.4.4 What is GLDAS-2.2?

GLDAS-2.2 explores the data assimilation capabilities in the LIS (Kumar et al., 2016; Kumar et al., 2019). The Catchment land surface model provides the model design that is suitable for assimilating the GRACE TWS (Terrestrial water storage) anomaly observation as shown in past studies. The GLDAS-2.2 Daily Catchment product is the outcome of the study by Li et al., 2019.

## 2.0 Data Organization

---

GLDAS-2.0 consists of 3-hourly, daily, and monthly products at 0.25° x 0.25° and 1.0° x 1.0° spatial resolutions. GLDAS-2.1 consists of 3-hourly and monthly products at 0.25° x 0.25° and 1.0° x 1.0° spatial resolutions. GLDAS-2.2 consists of a daily product at a 0.25° x 0.25° spatial resolution.

### 2.1 File Naming Convention

NASA GLDAS-2.0 and GLDAS-2.1 data files are named in accordance with the following convention:

GLDAS\_<Model><SpatialResolution>\_<TemporalResolution>.A<Date><ProductVersion>.nc4

Attribute	Description
<Model>	"CLSM" for the Catchment Model
	"NOAH" for the Noah Model
	"VIC" for the Variable Infiltration Capacity Model
<SpatialResolution>	"025" for 0.25 degree
	"010" for 1.0 degree
<TemporalResolution>	"3H" for 3-hourly
	"D" for daily
	"M" for monthly
<Date>*	<YYYYMMDD>.<HHHH> for 3-hourly
	<YYYYMMDD> for daily
	<YYYYMM> for monthly
<ProductVersion>	"020" for GLDAS-2.0
	"021" for GLDAS-2.1
	"022" for GLDAS-2.2

\*Date represented as 4-digit year, 2-digit month, 2-digit day of month, 4-digit GMT hour and minute of day.

NASA GLDAS-2.1 Early Product data files are named in accordance with the following convention:

GLDAS\_<Model><SpatialResolution>\_<TemporalResolution>\_EP.A<Date><ProductVersion>.nc4

...where all attributes are the same as above, but with the added EP designation.

NASA GLDAS-2.2 data files are named in accordance with the following convention:

GLDAS\_<Model><SpatialResolution>\_<DataAssimilationReference>\_<TemporalResolution>\_A<Date>.<ProductVersion>.nc4

...where all attributes are the same as above, but with the added DataAssimilationReference designation.

## README for NASA GLDAS Version 2 Data

### Examples:

The file name for the monthly 1.0 degree GLDAS-2.0 Noah data for January 1948 is:  
GLDAS\_NOAH10\_M\_A194801.020.nc4

The file name for the 3-hourly 0.25 degree GLDAS-2.1 Noah data at 03:00Z on January 1, 2000 is:

GLDAS\_NOAH025\_3H.A20000101.0300.021.nc4

The file name for the daily 0.25 degree GLDAS-2.0 Catchment data on January 1, 1948 is:

GLDAS\_CLSM025\_D.A19480101.020.nc4

The file name for the daily 0.25 degree GLDAS-2.2 Catchment with GRACE-DA1 on February 1, 2003 is:

GLDAS\_CLSM025\_DA1\_D.A20030201.022.nc4

## 2.2 File Format and Structure

All GLDAS-2 data files are in NetCDF format, which is a set of software libraries and self-describing, machine-independent data formats that support the creation, access, and sharing of array-oriented scientific data. More information can be found here:

<https://www.unidata.ucar.edu/software/netcdf/docs/>.

## 3.0 Data Contents

---

### 3.1 GLDAS-2.0 Data

GLDAS-2.0 includes five data products:

- GLDAS Catchment Land Surface Model L4 daily 0.25 x 0.25 degree V2.0 (GLDAS\_CLSM025\_D\_2.0)
- GLDAS Noah Land Surface Model L4 3 hourly 0.25 x 0.25 degree V2.0 (GLDAS\_NOAH025\_3H\_2.0)
- GLDAS Noah Land Surface Model L4 monthly 0.25 x 0.25 degree V2.0 (GLDAS\_NOAH025\_M\_2.0)
- GLDAS Noah Land Surface Model L4 3 hourly 1.0 x 1.0 degree V2.0 (GLDAS\_NOAH10\_3H\_2.0)
- GLDAS Noah Land Surface Model L4 monthly 1.0 x 1.0 degree V2.0 (GLDAS\_NOAH10\_M\_2.0)

The daily product from the GLDAS-2.0 Catchment model contains 33 parameters; the 3-hourly and monthly data products from the GLDAS-2.0 Noah model contain 36 parameters each. The parameters included in the GLDAS-2.0 data are listed in Table 3.1.

# README for NASA GLDAS Version 2 Data

Table 3.1 Parameters in the GLDAS-2.0 data.

CLSM	Noah	Short Name	Long Name	Units
X		ACond_tavg	Aerodynamic conductance	m s-1
	X	Albedo_inst	Albedo	%
X*	X	AvgSurfT_inst	Average surface skin temperature	K
X*	X	CanopInt_inst	Plant canopy surface water	kg m-2
X	X	ECanop_tavg	Canopy water evaporation	W m-2
X	X	ESoil_tavg	Direct evaporation from bare soil	W m-2
X	X	Evap_tavg	Evapotranspiration	kg m-2 s-1
X		EvapSnow_tavg	Snow evaporation	kg m-2 s-1
X		GWS_tavg	Ground water storage	mm
X	X	LWdown_f_tavg	Downward longwave radiation	W m-2
X	X	Lwnet_tavg	Net longwave radiation flux	W m-2
	X	PotEvap_tavg	Potential evaporation rate	W m-2
X*	X	Psurf_f_inst	Surface air pressure	Pa
X*	X	Qair_f_inst	Specific humidity	kg kg-1
X	X	Qg_tavg	Ground heat flux	W m-2
X	X	Qh_tavg	Sensible heat net flux	W m-2
X	X	Qle_tavg	Latent heat net flux	W m-2
X*	X	Qs_acc	Storm surface runoff	kg m-2 per 3-hour
X*	X	Qsb_acc	Baseflow-groundwater runoff	kg m-2 per 3-hour
X*	X	Qsm_acc	Snow melt	kg m-2 per 3-hour
X	X	Rainf_f_tavg	Total precipitation rate	kg m-2 s-1
X	X	Rainf_tavg	Rain precipitation rate	kg m-2 s-1
	X	RootMoist_inst	Root zone soil moisture	kg m-2
X*	X	SnowDepth_inst	Snow depth	m
X	X	Snowf_tavg	Snow precipitation rate	kg m-2 s-1
X		SnowT_tavg	Snow surface temperature	K
	X	SoilMoi0_10cm_inst	Soil moisture content (0-10 cm underground)	kg m-2



## README for NASA GLDAS Version 2 Data

	X	SoilMoi10_40cm_inst	Soil moisture content (10-40 cm underground)	kg m-2
	X	SoilMoi40_100cm_inst	Soil moisture content (40-100 cm underground)	kg m-2
	X	SoilMoi100_200cm_inst	Soil moisture content (100-200 cm underground)	kg m-2
X		SoilMoist_P_tavg	Profile soil moisture	kg m-2
X		SoilMoist_RZ_tavg	Root zone soil moisture	kg m-2
X		SoilMoist_S_tavg	Surface soil moisture	kg m-2
	X	SoilTMP0_10cm_inst	Soil temperature (0-10 cm underground)	K
	X	SoilTMP10_40cm_inst	Soil temperature (10-40 cm underground)	K
	X	SoilTMP40_100cm_inst	Soil temperature (40-100 cm underground)	K
	X	SoilTMP100_200cm_inst	Soil temperature (100-200 cm underground)	K
X	X	SWdown_f_tavg	Downward shortwave radiation flux	W m-2
X*	X	SWE_inst	Snow depth water equivalent	kg m-2
X	X	Swnet_tavg	Net shortwave radiation flux	W m-2
X*	X	Tair_f_inst	Air temperature	K
X	X	Tveg_tavg	Transpiration	W m-2
X		TWS_tavg	Terrestrial water storage	mm
	X	Wind_f_inst	Wind speed	m s-1

The short names with extension “\_tavg” are 24-hr or past 3-hr averaged variables.

The short names with extension “\_acc” are past 3-hr accumulated variables.

The short names with extension “\_inst” are instantaneous variables.

The short names with “\_f” are forcing variables.

\*All variables from GLDAS-2.0 Catchment model are 24-hr averaged variables (\_tavg).

## 3.2 GLDAS-2.1 Data

GLDAS-2.1 includes 16 data products:

- GLDAS Catchment Land Surface Model L4 3 hourly 1.0 x 1.0 degree V2.1 (GLDAS\_CLSM10\_3H\_2.1)
- GLDAS Catchment Land Surface Model L4 3 hourly 1.0 x 1.0 degree Early Product V2.1 (GLDAS\_CLSM10\_3H\_EP\_2.1)
- GLDAS Catchment Land Surface Model L4 monthly 1.0 x 1.0 degree V2.1 (GLDAS\_CLSM10\_M\_2.1)
- GLDAS Catchment Land Surface Model L4 monthly 1.0 x 1.0 degree Early Product V2.1 (GLDAS\_CLSM10\_M\_EP\_2.1)

## README for NASA GLDAS Version 2 Data

- GLDAS Noah Land Surface Model L4 3 hourly 0.25 x 0.25 degree V2.1 (GLDAS\_NOAH025\_3H\_2.1)
- GLDAS Noah Land Surface Model L4 3 hourly 0.25 x 0.25 degree Early Product V2.1 (GLDAS\_NOAH025\_3H\_EP\_2.1)
- GLDAS Noah Land Surface Model L4 monthly 0.25 x 0.25 degree V2.1 (GLDAS\_NOAH025\_M\_2.1)
- GLDAS Noah Land Surface Model L4 monthly 0.25 x 0.25 degree Early Product V2.1 (GLDAS\_NOAH025\_M\_EP\_2.1)
- GLDAS Noah Land Surface Model L4 3 hourly 1.0 x 1.0 degree V2.1 (GLDAS\_NOAH10\_3H\_2.1)
- GLDAS Noah Land Surface Model L4 3 hourly 1.0 x 1.0 degree Early Product V2.1 (GLDAS\_NOAH10\_3H\_EP\_2.1)
- GLDAS Noah Land Surface Model L4 monthly 1.0 x 1.0 degree V2.1 (GLDAS\_NOAH10\_M\_2.1)
- GLDAS Noah Land Surface Model L4 monthly 1.0 x 1.0 degree Early Product V2.1 (GLDAS\_NOAH10\_M\_EP\_2.1)
- GLDAS VIC Land Surface Model L4 3 hourly 1.0 x 1.0 degree V2.1 (GLDAS\_VIC10\_3H\_2.1)
- GLDAS VIC Land Surface Model L4 3 hourly 1.0 x 1.0 degree Early Product V2.1 (GLDAS\_VIC10\_3H\_EP\_2.1)
- GLDAS VIC Land Surface Model L4 monthly 1.0 x 1.0 degree V2.1 (GLDAS\_VIC10\_M\_2.1)
- GLDAS VIC Land Surface Model L4 monthly 1.0 x 1.0 degree Early Product V2.1 (GLDAS\_VIC10\_M\_EP\_2.1)

The GLDAS-2.1 Catchment model data products contain 38 parameters; the GLDAS-2.1 Noah model data products contain 36 parameters; the GLDAS-2.1 VIC model data products contain 34 parameters. The parameters included in the GLDAS-2.1 data are listed in Table 3.2.

Table 3.2 Parameters in the GLDAS-2.1 data.

CLSM	Noah	VIC	Short Name	Long Name	Units
X		X	ACond_tavg	Aerodynamic conductance	m s <sup>-1</sup>
X	X	X	Albedo_inst	Albedo	%
X	X	X	AvgSurfT_inst	Average surface skin temperature	K
X	X	X	CanopInt_inst	Plant canopy surface water	kg m <sup>-2</sup>
X	X	X	ECanop_tavg	Canopy water evaporation	W m <sup>-2</sup>

# README for NASA GLDAS Version 2 Data

X	X	X	ESoil_tavg	Direct evaporation from bare soil	W m-2
X	X	X	Evap_tavg	Evapotranspiration	kg m-2 s-1
X	X	X	LWdown_f_tavg	Downward longwave radiation	W m-2
X	X	X	Lwnet_tavg	Net longwave radiation flux	W m-2
	X		PotEvap_tavg	Potential evaporation rate	W m-2
X	X	X	Psurf_f_inst	Surface air pressure	Pa
X	X	X	Qair_f_inst	Specific humidity	kg kg-1
X	X	X	Qg_tavg	Ground heat flux	W m-2
X	X	X	Qh_tavg	Sensible heat net flux	W m-2
X	X	X	Qle_tavg	Latent heat net flux	W m-2
X	X	X	Qs_acc	Storm surface runoff	kg m-2 per 3-hour
X	X	X	Qsb_acc	Baseflow-groundwater runoff	kg m-2 per 3-hour
X	X	X	Qsm_acc	Snow melt	kg m-2 per 3-hour
X	X	X	Rainf_f_tavg	Total precipitation rate	kg m-2 s-1
X	X	X	Rainf_tavg	Rain precipitation rate	kg m-2 s-1
	X	X	RootMoist_inst	Root zone soil moisture	kg m-2
X	X	X	SnowDepth_inst	Snow depth	m
X	X	X	Snowf_tavg	Snow precipitation rate	kg m-2 s-1
X			SnowT_tavg	Snow surface temperature	K
	X		SoilMoi0_10cm_inst	Soil moisture content (0-10 cm underground)	kg m-2
		X	SoilMoi0_30cm_inst	Soil moisture content of surface layer	kg m-2
	X		SoilMoi10_40cm_inst	Soil moisture content (10-40 cm underground)	kg m-2
	X		SoilMoi40_100cm_inst	Soil moisture content (40-100 cm underground)	kg m-2
	X		SoilMoi100_200cm_inst	Soil moisture content (100-200 cm underground)	kg m-2
		X	SoilMoi_depth2_inst	Soil moisture content of second layer	kg m-2
		X	SoilMoi_depth3_inst	Soil moisture content of bottom layer	kg m-2

# README for NASA GLDAS Version 2 Data

X			SoilMoist_P_inst	Profile soil moisture	kg m-2
X			SoilMoist_RZ_inst	Root zone soil moisture	kg m-2
X			SoilMoist_S_inst	Surface soil moisture	kg m-2
X	X		SoilTMP0_10cm_inst	Soil temperature (0-10 cm underground)	K
		X	SoilTMP0_30cm_inst	Soil temperature of surface layer	K
X			SoilTMP10_29cm_inst	Soil temperature (10-29 cm underground)	K
	X		SoilTMP10_40cm_inst	Soil temperature (10-40 cm underground)	K
X			SoilTMP29_68cm_inst	Soil temperature (29-68 cm underground)	K
	X		SoilTMP40_100cm_inst	Soil temperature (40-100 cm underground)	K
X			SoilTMP68_144cm_inst	Soil temperature (68-144 cm underground)	K
	X		SoilTMP100_200cm_inst	Soil temperature (100-200 cm underground)	K
X			SoilTMP144_295cm_inst	Soil temperature (144-295 cm underground)	K
X			SoilTMP295_1295cm_inst	Soil temperature (295-1295 cm underground)	K
		X	SoilTMP_depth2_inst	Soil temperature of second layer	K
		X	SoilTMP_depth3_inst	Soil temperature of bottom layer	K
X	X	X	SWdown_f_tavg	Downward shortwave radiation flux	W m-2
X	X	X	SWE_inst	Snow depth water equivalent	kg m-2
X	X	X	Swnet_tavg	Net shortwave radiation flux	W m-2
X	X	X	Tair_f_inst	Air temperature	K
X	X	X	Tveg_tavg	Transpiration	W m-2
X			TWS_tavg	Terrestrial water storage	mm
X	X	X	Wind_f_inst	Wind speed	m s-1

The short names with extension “\_tavg” are 24-hr or past 3-hr averaged variables.

The short names with extension “\_acc” are past 3-hr accumulated variables.

The short names with extension “\_inst” are instantaneous variables.

The short names with “\_f” are forcing variables.

### 3.3 GLDAS-2.2 Data

GLDAS-2.2 includes two data products:

- GLDAS Catchment Land Surface Model L4 daily 0.25 x 0.25 degree GRACE-DA1 V2.2 (GLDAS\_CLSM025\_DA1\_D\_2.2)
- GLDAS Catchment Land Surface Model L4 daily 0.25 x 0.25 degree GRACE-DA1 Early Product V2.2 (GLDAS\_CLSM025\_DA1\_D\_EP\_2.2)

The daily data assimilation product from GLDAS-2.2 Catchment model contains 24 parameters, with the forcing variables, including Rainf\_tavg and Snowf\_tavg, excluded. The parameters included in the GLDAS-2.2 data are listed in Table 3.3.

Table 3.3 Parameters in the GLDAS-2.2 data.

Short Name	Long Name	Units
ACond_tavg	Aerodynamic conductance	m s-1
AvgSurfT_tavg	Average surface skin temperature	K
CanopInt_tavg	Plant canopy surface water	kg m-2
ECanop_tavg	Canopy water evaporation	kg m-2 s-1
ESoil_tavg	Direct evaporation from bare soil	kg m-2 s-1
Evap_tavg	Evapotranspiration	kg m-2 s-1
EvapSnow_tavg	Snow evaporation	kg m-2 s-1
GWS_tavg	Ground water storage	mm
Lwnet_tavg	Net longwave radiation flux	W m-2
Qg_tavg	Ground heat flux	W m-2
Qh_tavg	Sensible heat net flux	W m-2
Qle_tavg	Latent heat net flux	W m-2
Qs_tavg	Storm surface runoff	kg m-2 s-1
Qsb_tavg	Baseflow-groundwater runoff	kg m-2 s-1
Qsm_tavg	Snow melt	kg m-2 s-1
SnowDepth_tavg	Snow depth	m
SnowT_tavg	Snow surface temperature	K
SoilMoist_P_tavg	Profile soil moisture	kg m-2
SoilMoist_RZ_tavg	Root zone soil moisture	kg m-2
SoilMoist_S_tavg	Surface soil moisture	kg m-2
SWE_tavg	Snow depth water equivalent	kg m-2
Swnet_tavg	Net shortwave radiation flux	W m-2

TVeg_tavg	Transpiration	kg m <sup>-2</sup> s <sup>-1</sup>
TWS_tavg	Terrestrial water storage	mm

All variables are 24-hr averaged variables (\_tavg).

### 3.4 Data Interpretation

1. Due to unreliable Greenland forcing data and the lack of a glacier/ice sheet model, snow water equivalent accumulates indefinitely in Greenland and a few other Arctic points. Therefore, it is highly recommended that Greenland and other points with abnormally large snow water equivalent values be masked out when performing global analyses.
2. Total precipitation rate is the sum of rain and snow precipitation rates. The forcing variable "Rainf\_f\_tavg" is the total precipitation rate, whereas "Rainf\_tavg" and "Snowf\_tavg" are the liquid precipitation rate and frozen precipitation rate, respectively.
3. Total runoff is the sum of subsurface runoff "Qsb\_tavg" and surface runoff "Qs\_tavg".
4. The number of vertical levels for soil moisture (SoilMoi) and soil temperature (SoilTMP) is model specific. Please follow the list below for the correct depths of the soil layers.
  - Noah (4 layers): 0-0.1, 0.1-0.4, 0.4-1.0, 1.0-2.0 m
  - VIC (3 layers): 0.0-0.3 m surface, variable depth for the second and bottom layers. The map of depths are available to download from: <https://ldas.gsfc.nasa.gov/gldas/specifications>.
  - CLSM does not have explicit vertical levels for soil moisture. Instead, soil moisture is represented in Surface (0-2 cm), Root Zone (0-100 cm), and Profile (varies grid-by-grid) reservoirs. They are inclusive: Profile includes Surface and Root Zone, and Root Zone includes Surface.
  - CLSM has six layers for soil temperature: 0-0.1, 0.1-0.29, 0.29-0.68, 0.68-1.44, 1.44-2.95, and 2.95-12.95 m.
5. CLSM does simulate shallow groundwater, so Terrestrial Water Storage (TWS) in CLSM is the sum of soil water, snow water equivalent, canopy water, and groundwater. Ground Water Storage (GWS) in CLSM is already included in TWS. Ground Water Storage (GWS) in CLSM was computed using formula:  

$$GWS = TWS - \text{RootZoneSoilMoisture} - \text{SnowWaterEquivalent} - \text{CanopyInterception}.$$
 Noah TWS in CLSM is the sum of soil moisture in all layers, accumulated snow, and plant canopy surface water.
6. Use temporal averaging, not accumulation, to upscale the data to different temporal resolutions. For example, rainfall and snowfall are provided as rates, i.e., kg/m<sup>2</sup>/s. The correct method of upscaling is averaging, which does not change the units.
7. Monthly average files contain straight averages of 3-hourly data, so each monthly average has units PER 3 HOURS. For example, total evapotranspiration (Evap\_tavg) for April 1979 is the average 3-hour mean rate of evapotranspiration over all 3-hour intervals in April 1979. It is NOT the accumulated evapotranspiration in April 1979. To compute the latter, use this formula:

$$\text{Evap\_tavg (April)}\{\text{kg/m}^2\} = \text{Evap\_tavg (April)}\{\text{kg/m}^2/\text{sec}\} * 10800\{\text{sec}/3\text{hr}\} * 8\{3\text{hr}/\text{day}\} * 30\{\text{days}\}$$

For accumulated variables such as Qs\_acc, the monthly mean surface runoff is the average 3-hour accumulation over all 3-hour intervals in April 1979. To compute monthly accumulation, use this formula:

$$\text{Qs\_acc (April)}\{\text{kg/m}^2\} = \text{Qs\_acc (April)}\{\text{kg/m}^2/3\text{hr}\} * 8\{3\text{hr}/\text{day}\} * 30\{\text{days}\}$$

This would be irrelevant, and the above formula should not be used, if the field of interest is an instantaneous state.

8. Heights of forcing fields depend on the datasets used to drive the simulation. Presently, all the GLDAS datasets use the 2-meter temperature and specific humidity and the 10-meter wind for the entire time span.
9. The mean fields in monthly data (e.g., evapotranspiration, see Table 3.1) contain straight average of 3-hour accumulation from 0300z on the 1st to 0000z on the first day of the next month. The instantaneous fields are averaged over 0000z on the 1st day of the month to 2100z on the last day of the month.
10. Snow density computed using the snow water equivalent and snow depth included in the current GLDAS-2.0 Daily CLSM product is not valid, because snow depth had not accounted for the grid fraction of snow cover at the time the simulation was done. This problem doesn't apply to the 3-hourly and monthly GLDAS-2.1 and GLDAS-2.2 products.

## 4.0 Options for Reading the Data

---

### 4.1 Utilities

NASA GLDAS-2 data are archived in self-describing and machine-independent NetCDF format. This Unidata page provides a list of software for manipulating or displaying NetCDF data: <https://www.unidata.ucar.edu/software/netcdf/software.html>.

### 4.2 Panoply

Panoply (<https://www.giss.nasa.gov/tools/panoply/>) is a cross-platform application that plots geo-referenced and other arrays from NetCDF, HDF, GRIB, and other data formats. The [HowTo](#) page from the GES DISC provides a recipe for [How to View Remote Data in OPeNDAP with Panoply](#).

## 4.3 GrADS

The Grid Analysis and Display System (GrADS) is an interactive desktop tool for easy access, manipulation, and visualization of earth science data. GrADS supports several data formats, such as binary, NetCDF, HDF, and GRIB. The documentation and software for GrADS can be found at <http://cola.gmu.edu/grads/grads.php>.

Each individual GLDAS-2 NetCDF can be opened by the GrADS utility [sdfopen](#) directly without a data descriptor file (i.e., a ctl file). After calling sdfopen, GrADS commands, such as “q file”, “d [VariableName]”, etc. can be used to query file information, and read and display the data. An example showing how to use sdfopen to read a GLDAS-2 NetCDF file and query for its dimensions and variables is in Appendix A.

## 5.0 Data Services

---

As of August 1, 2016, access to GES DISC data requires all users to be registered with the NASA Earthdata Login. Data continue to be free of charge and accessible via HTTPS. As of October 3, 2016, access to data via FTP is no longer available. Detailed instructions on how to register and receive authorization to access all GES DISC data are provided at <https://disc.gsfc.nasa.gov/data-access>.

GES DISC users who deploy scripting methods to list and download data in bulk are advised to review the instructions from the link above that provide examples of GNU wget commands for listing and downloading data via HTTPS.

If you need assistance or wish to report a problem, please contact us:

Goddard Earth Sciences Data and Information Services Center, Code 610.2  
NASA Goddard Space Flight Center  
Greenbelt, MD 20771  
Email: [gsfc-help-disc@lists.nasa.gov](mailto:gsfc-help-disc@lists.nasa.gov)  
Phone: 301-614-5224  
Fax: 301-614-5268

## 5.1 HTTPS

Access the online archive via HTTPS for direct download entire files:  
<https://hydro1.gesdisc.eosdis.nasa.gov/data/GLDAS/>.

## 5.2 Earthdata Search

Use the Earthdata Search interface to find and retrieve datasets from the GES DISC and other NASA data centers: <https://search.earthdata.nasa.gov/search?q=GLDAS>.



## 5.3 GES DISC Subsetter/Regridder

Access the GES DISC Regridder and Subsetter tool through the [GLDAS search results page](#) or any GLDAS dataset landing page by selecting the Subset/Get Data link. This tool allows for spatial, temporal, and variable subsetting, as well as re-gridding the data to various other grids through several interpolation methods.

## 5.4 OPeNDAP

Access the data via the OPeNDAP protocol for parameter and spatial subsetting, with several output formats: <https://hydro1.gesdisc.eosdis.nasa.gov/opensap/GLDAS/>.

## 5.5 GrADS Data Server (GDS)

The GrADS Data Server (GDS) is another form of OPeNDAP that provides subsetting and some analysis services across the internet:  
<https://hydro1.gesdisc.eosdis.nasa.gov/dods/>.

## 5.6 Giovanni

The GES-DISC Interactive Online Visualization And aNalysis Infrastructure (Giovanni) is a web-based tool that allows users to interactively visualize and analyze data:  
<https://giovanni.gsfc.nasa.gov/giovanni/#dataKeyword=GLDAS>

The sample image below is generated by NASA Giovanni.




Figure 1. Soil moisture content (0-10 cm underground) map for January 2000, from GLDAS-2.1 Noah Land Surface Model L4 monthly 0.25 x 0.25 degree data.

## 6.0 More Information

---

Land Data Assimilation System (LDAS) Project: <https://ldas.gsfc.nasa.gov/>

## 7.0 Acknowledgements

---

The GLDAS data are produced by the NASA GSFC Hydrological Sciences Laboratory (HSL). Please refer to Rodell et al. (2004) for more information about the GLDAS project.

## References

---

- Rodell, M., P. R. Houser, U. Jambor, J. Gottschalck, K. Mitchell, C.-J. Meng, K. Arsenault, A. Cosgrove, J. Radakovich, M. Bosilovich, J. K. Entin, J. P. Walker, D. Lohmann, and D. Toll, 2004. The Global Land Data Assimilation System, *Bull. Amer. Meteor. Soc.*, 85(3): 381-394.
- Adler, R.F., G.J. Huffman, A. Chang, R. Ferraro, P. Xie, J. Janowiak, B. Rudolf, U. Schneider, S. Curtis, D. Bolvin, A. Gruber, J. Susskind, P. Arkin, E. Nelkin, 2003: The Version 2 Global Precipitation Climatology Project (GPCP) Monthly Precipitation Analysis (1979-Present). *J. Hydrometeor.*, 4,1147-1167.
- Carroll, M.L., Townshend, J.R., DiMiceli, C.M., Noojipady, P, Sohlberg, R.A. (2009). A new global raster water mask at 250 m resolution. *International Journal of Digital Earth*, 2(4), 291-308.
- Derber, J. C., D. F. Parrish, and S. J. Lord, 1991: The new global operational analysis system at the National Meteorological Center. *Weather Forecasting*, 6, 538-547.
- Friedl, M.A., Sulla-Menashe, D., Tan, B., Schneider, A., Ramankutty, N., Sibley, A., & Huang, X. (2010). MODIS Collection 5 global land cover: Algorithm refinements and characterization of new datasets. *Remote Sensing of Environment*, 114,168-182.
- Huffman, G. J., R. F. Adler, M. Morrissey, D. T. Bolvin, S. Curtis, R. Joyce, B. McGavock, J. Susskind, 2001: Global Precipitation at One-Degree Daily Resolution from Multi-Satellite Observations. *J. Hydrometeor.*, 2(1), 36-50.

## README for NASA GLDAS Version 2 Data

Kumar, S.V., M. Jasinski, D.M. Mocko, M. Rodell, J. Borak, B. Li, H.K. Beaudoin, and C.D. Peters-Lidard, 2019: [NCA-LDAS Land Analysis: Development and Performance of a Multisensor, Multivariate Land Data Assimilation System for the National Climate Assessment](https://doi.org/10.1175/JHM-D-17-0125.1). *J. Hydrometeorol.*, **20**, 1571–1593, <https://doi.org/10.1175/JHM-D-17-0125.1>

Kumar, S. V., Zaitchik, B. F., Peters-Lidard, C., Rodell, M., Reichle, R., Li, B., Jasinski, M., Mocko, D., Getirana, A., de Lannoy, G., Cosh, M. H., Hain, C. R., Anderson, M., Arsenault, K. R., Xia, Y., & Ek, M. (2016). Assimilation of gridded GRACE terrestrial water storage estimates in the North American Land Data Assimilation System, 2016. *Journal of Hydrometeorology*, **17**( 7), 1951– 1972.

Li, B., Rodell, M., Kumar, S., Beaudoin, H. K., Getirana, A., Zaitchik, B. F., et al. (2019). Global GRACE data assimilation for groundwater and drought monitoring: Advances and challenges. *Water Resources Research*, **55**, 7564– 7586. <https://doi.org/10.1029/2018WR024618>

Li, B., Rodell, M., Sheffield, J., Wood, E., and Sutanudjaja, E. (2019). Long-term, non- anthropogenic groundwater storage changes simulated by three global-scale hydrological models, *Scientific Reports*, **9**, 10746.

Nijssen, B., S. Shukla, C. Lin, H. Gao, T. Zhou, Ishottama, J. Sheffield, E. F. Wood, and D. P. Lettenmaier, 2014: A prototype Global Drought Information System based on multiple land surface models. *Journal of Hydrometeorology*, doi:10.1175/jhm-d-13-090.1.

Reynolds C.A., T. J. Jackson, and W.J. Rawls. (1999). Estimating Available Water Content by Linking the FAO Soil Map of the World with Global Soil Profile Databases and Pedo-transfer Functions. *Proceedings of the AGU 1999 Spring Conference*, Boston, MA. May 31-June 4, 1999.

Save, H., Bettadpur, S., & Tapley, B. D. (2016). High resolution CSR GRACE RL05 mascons. *Journal of Geophysical Research: Solid Earth*, **121**, 7547– 7569. <https://doi.org/10.1002/2016JB013007>

Save, H., Bettadpur, S., & Tapley, B. D. (2012). Reducing errors in the GRACE gravity solutions using regularization. *Journal of Geodesy*, **86**( 9), 695– 711. <https://doi.org/10.1007/s00190-012-0548-5>

Sheffield, J., G. Goteti, and E. F. Wood, 2006: Development of a 50-yr high-resolution global dataset of meteorological forcings for land surface modeling, *J. Climate*, **19** (13), 3088-3111

## Acronyms

---

The following acronyms and abbreviations are used in this document.

AGRMET	AGRIcultural METeorological Modeling System
CAPE	Convective Available Potential Energy
CMAP	CPC Merged Analysis of Precipitation
CMORPH	CPCF precipitation MORPHing technique
CPC	NCEP's Climate Prediction Center
CPPA	Climate Prediction Program for the Americas
EMC	NCEP's Environmental Modeling Center
GDAS	Global Data Assimilation System
GDS	GrADS Data Server
GES DISC	Goddard Earth Sciences Data and Information Services Center
Giovanni	GES-DISC Interactive On-line Visualization and Analysis Infrastructure
GLDAS	Global Land Data Assimilation System
GRACE	Gravity Recovery and Climate Experiment
GrADS	Grid Analysis and Display System
GPCP	Global Precipitation Climatology Project
GRIB	GRIdded Binary
HDF	Hierarchical Data Format
HDISC	Hydrology Data and Information Services Center
LDAS	Land Data Assimilation System
LIS	Land Information System
LSM	Land Surface Model
NARR	North American Regional Reanalysis
NASA	National Aeronautics and Space Administration
NCEP	National Centers for Environmental Prediction
NetCDF	Network Common Data Form
NIDIS	National Integrated Drought Information System
NLDAS	North American Land Data Assimilation System
NOAA	National Oceanic and Atmospheric Administration
OHD	NOAA's Office of Hydrologic Development
PRISM	Parameter-elevation Regressions on Independent Slopes Model
SAC	Sacramento model
SRB	Surface Radiation Budget
SVAT	Soil Vegetation Atmosphere Transfer model
VIC	Variable Infiltration Capacity macroscale model

## Appendix A

---

Below is an example showing how to use sdfopen to read a GLDAS-2 NetCDF file and query for its dimensions and variables.

```
ga -> xdfopen GLDAS_NOAH10_M.2.0.xdf
Scanning Descriptor File: GLDAS_NOAH10_M.2.0.xdf
SDF file
/ftp/data/s4pa/GLDAS_GLDA20_NOAH10_M.2.0/%y4/GLDAS_NOAH10_M.A%y4%m2.
020.nc4
is open as file 1
LON set to 0 360
LAT set to -59.5 89.5
LEV set to 0 0
Time Values set: 1948:1:1:0 1948:1:1:0
E set to 1
1 ga -> q file
File 1: GLDAS2.0 LIS land surface model output monthly mean
Descriptor: GLDAS_NOAH10_M.2.0.XDF
Binary: GLDAS_NOAH10_M.2.0/%y4/GLDAS_NOAH10_M.A%y4%m2.020.nc4
Type = Gridded
Xsize = 360 Ysize = 150 Zsize = 1 Tsize = 780 Esize = 1
Number of Variables = 36
    swnet_tavg 0 t,y,x Net short wave radiation flux
    lwnet_tavg 0 t,y,x Net long-wave radiation flux
    qle_tavg 0 t,y,x Latent heat net flux
    qh_tavg 0 t,y,x Heat flux
    snowf_tavg 0 t,y,x Snow precipitation rate
    rainf_tavg 0 t,y,x Rain precipitation rate
    evap_tavg 0 t,y,x Evapotranspiration
    qs_acc 0 t,y,x Storm surface runoff
    qsb_acc 0 t,y,x Baseflow-groundwater runoff
    qsm_acc 0 t,y,x Snow melt
    avgsurf_inst 0 t,y,x Average Surface Skin temperature
    albedo_inst 0 t,y,x Albedo
    swe_inst 0 t,y,x Snow depth water equivalent
    snowdepth_inst 0 t,y,x Snow depth
    soilmoi0_10cm_i 0 t,y,x Soil moisture
    soilmoi10_40cm_ 0 t,y,x Soil moisture
    soilmoi100_200c 0 t,y,x Soil moisture
    soiltmp0_10cm_i 0 t,y,x Soil temperature
    soiltmp10_40cm_ 0 t,y,x Soil temperature
    soiltmp40_100cm 0 t,y,x Soil temperature
    soiltmp100_200c 0 t,y,x Soil temperature
    potevap_tavg 0 t,y,x Potential evaporation rate
```

## README for NASA GLDAS Version 2 Data

```
ecanop_tavg 0 t,y,x Canopy water evaporation
tveg_tavg 0 t,y,x Transpiration
esoil_tavg 0 t,y,x Direct Evaporation from Bare Soil
rootmoist_inst 0 t,y,x Root zone soil moisture
canopint_inst 0 t,y,x Plant canopy surface water
wind_f_inst 0 t,y,x Wind speed
rainf_f_tavg 0 t,y,x Total precipitation rate
tair_f_inst 0 t,y,x Temperature
qair_f_inst 0 t,y,x Specific humidity
psurf_f_inst 0 t,y,x Pressure
swdown_f_tavg 0 t,y,x Downward short-wave radiation flux
lwdown_f_tavg 0 t,y,x Downward long-wave radiation flux
ga ->
```

With a GrADS descriptor file, by using GrADS command `xdfopen`, multiple GLDAS-2 NetCDF files can be opened. Therefore, time aggregation-related visualization and data analysis can be done by GrADS. Below is a GrADS sample descriptor file for 3-hourly 1.0 x 1.0 degree Noah data product, GLDAS\_NOAH10\_3H.2.0.

```
GLDAS_NOAH10_M.2.0.xdf, a sample data descriptor file
DSET ./GLDAS_NOAH10_M.2.0/%y4/GLDAS_NOAH10_M.A%y4%m2.020.nc4
OPTIONS template
TDEF time 780 LINEAR jan1948 1mo
***variable name may not appear completely (max 15 characters)
```

An example for using `xdfopen` to open GLDAS\_NOAH10\_3H.2.0.xdf:

```
ga -> xdfopen GLDAS_NOAH10M.2.0.xdf
Scanning Descriptor File: GLDAS_NOAH10_M.2.0.xdf
SDF file
/ftp/data/s4pa/GLDAS/GLDAS_NOAH10_M.2.0/%y4/GLDAS_NOAH10_M.A%y4%m2.0
20.nc4
is open as file 1
LON set to 0 360
LAT set to -59.5 89.5
LEV set to 0 0
Time values set: 1948:1:1:0 1948:1:1:0
E set to 1 1
ga -> q file
File 1: GLDAS2.0 LIS land surface model output monthly mean
Descriptor: GLDAS_NOAH10_M.2.0.XDF
Binary: GLDAS_NOAH10_M.2.0/%y4/GLDAS_NOAH10_M.A%y4%m2.020.nc4
Type = Gridded
Xsize = 360 Ysize = 150 Zsize = 1 Tsize = 780 Esize = 1
Number of Variables = 36
```

## README for NASA GLDAS Version 2 Data

swnet\_tavg 0 t,y,x Net short wave radiation flux  
lwnet\_tavg 0 t,y,x Net long-wave radiation flux  
qle\_tavg 0 t,y,x Latent heat net flux  
qh\_tavg 0 t,y,x Heat flux  
snowf\_tavg 0 t,y,x Snow precipitation rate  
rainf\_tavg 0 t,y,x Rain precipitation rate  
evap\_tavg 0 t,y,x Evapotranspiration  
qs\_acc 0 t,y,x Storm surface runoff  
qsb\_acc 0 t,y,x Baseflow-groundwater runoff  
qsm\_acc 0 t,y,x Snow melt  
avgsurft\_inst 0 t,y,x Average Surface Skin temperature  
albedo\_inst 0 t,y,x Albedo  
swe\_inst 0 t,y,x Snow depth water equivalent  
snowdepth\_inst 0 t,y,x Snow depth  
soilmoi0\_10cm\_i 0 t,y,x Soil moisture  
soilmoi10\_40cm\_ 0 t,y,x Soil moisture  
soilmoi100\_200c 0 t,y,x Soil moisture  
soiltmp0\_10cm\_i 0 t,y,x Soil temperature  
soiltmp10\_40cm\_ 0 t,y,x Soil temperature  
soiltmp40\_100cm 0 t,y,x Soil temperature  
soiltmp100\_200c 0 t,y,x Soil temperature  
potevap\_tavg 0 t,y,x Potential evaporation rate  
ecanop\_tavg 0 t,y,x Canopy water evaporation  
tveg\_tavg 0 t,y,x Transpiration  
esoil\_tavg 0 t,y,x Direct Evaporation from Bare Soil  
rootmoist\_inst 0 t,y,x Root zone soil moisture  
canopint\_inst 0 t,y,x Plant canopy surface water  
wind\_f\_inst 0 t,y,x Wind speed  
rainf\_f\_tavg 0 t,y,x Total precipitation rate  
tair\_f\_inst 0 t,y,x Temperature  
qair\_f\_inst 0 t,y,x Specific humidity  
psurf\_f\_inst 0 t,y,x Pressure  
swdown\_f\_tavg 0 t,y,x Downward short-wave radiation flux  
lwdown\_f\_tavg 0 t,y,x Downward long-wave radiation flux

ga ->