

Forward KL Can Be Better: Normalizing Flow Boltzmann Sampling with Log-Ratio Variation

Qi Feng¹, Rongjie Lai², Di Qi², and Xuda Ye²

¹Department of Mathematics, Florida State University, Tallahassee, FL 32306

²Department of Mathematics, Purdue University, West Lafayette, IN 47907

July 10, 2026

Abstract

Forward KL training of a normalizing flow Boltzmann generator is mass-covering and usually trains well. Its weakness lies in the annealing chain that supplies its target samples, which starts at the flow’s own pushforward and mixes only among the modes the flow already covers. The objective can then miss the remaining modes while still reporting a high effective sample size, a fake ESS where the importance weights look benign but target mass is absent. The chain’s samples are also biased, and the forward KL controls only the mean of the log-ratio, so this bias passes into the trained flow. We address these failures with the log-ratio variation, a family of X functionals that penalize pairwise variation of the flow’s own log-ratio. Three weights give complementary roles: accuracy under the target, mode discovery against a wide-coverage measure built by quench and temper, and suppression of leakage outside the target. The balanced loss $KL + X_\mu + X_{(\bar{\mu} + \bar{\nu})/2}$ combines them at the cost of one extra sampling batch. A Fisher–Rao analysis shows that the forward KL still converges under the X_μ term and that this term can lower the accuracy floor left by biased samples. The model problems confirm the intended robustness at small batch and show a second regime: on a difficult multimodal annealing path, the balanced loss retains a clear aggregate advantage over forward KL through the largest tested batch, although that advantage need not hold at every stage.

1 Introduction

Drawing samples from a target distribution $\mu \propto \exp(-U)$ on $\mathbb{R}^d$, known only up to its normalization constant, is a recurring problem in molecular simulation, Bayesian inference, and statistical physics. When μ is high-dimensional and multimodal, its mass splits among basins of U separated by barriers. A local Markov chain Monte Carlo (MCMC) sampler that proposes small moves is then trapped: the time to cross a barrier of free-energy height ΔF grows like $e^{\Delta F}$, so within any practical budget the chain explores the basin it began in and leaves the rest of the mass unseen (Metropolis et al., 1953; Hastings, 1970). The estimates such a chain returns are biased toward that basin. The bias is silent, since the chain reports nothing about the mass it never visited.

The classical responses soften the barriers rather than remove them. Parallel tempering (PT) and replica exchange run replicas at a ladder of temperatures and swap configurations between neighbors, carrying a basin crossed hot down to the temperature of interest (Swendsen and Wang, 1986; Geyer, 1991; Hukushima and Nemoto, 1996). Annealed importance sampling (AIS) bridges a tractable proposal to μ along intermediate distributions while accumulating importance weights (Neal, 2001). Sequential Monte Carlo (SMC) carries a weighted particle population along the same kind of path (Doucet et al., 2001; Del Moral et al., 2006). Nested sampling recasts the normalizing integral as a one-dimensional quadrature over a rising likelihood threshold (Skilling, 2006). A line meeting the learned-sampler program partway replaces the fixed hot reference by a map, sharpening the swap schedule, tuning the endpoint, or training a flow to carry the hot reference toward the cold target (Syed et al., 2022; Surjanovic et al., 2022; Zhang et al., 2025). This line is the closest to ours; we differ in that our mode discovery is built off the ladder, rather than carried by the

tempering chain itself. These methods are the workhorses of multimodal sampling and remain the references against which a learned sampler is judged. They share one limit: they move mass only between basins a chain can already traverse within budget, so a basin no replica reaches contributes nothing, and raising the temperature enough to bridge every barrier flattens the structure one is trying to resolve (Woodard et al., 2009; Chehab et al., 2025). We do not discard this machinery; we use it. Our training samples come from an AIS ladder. The wide-coverage measure that drives mode discovery is built by a quench and temper construction that borrows the temperature-annealing idea while avoiding the reliance on a chain crossing every barrier; we supply the modes the chain cannot reach from a separate source.

Against this classical backdrop, Boltzmann generators (BG) replace the chain by a learned transport map (Noé et al., 2019). A normalizing flow is an invertible parametric map $G : \mathbb{R}^d \rightarrow \mathbb{R}^d$ (Rezende and Mohamed, 2015; Papamakarios et al., 2017; Kingma and Dhariwal, 2018). It is trained so that the pushforward $\nu := (G^{-1})_{\#}\mu_0$ of an easily samplable reference $\mu_0 \propto \exp(-U_0)$, typically a Gaussian, approximates μ . By convention G maps target to source, $G_{\#}\mu \approx \mu_0$, so G^{-1} draws a batch of independent samples by pushing μ_0 through it in one forward pass. Invertibility gives ν in closed form by the change of variables, so each sample carries its own importance weight μ/ν against the unnormalized target. Reweighting then turns the samples into Monte Carlo estimates with controlled finite-sample error. The effective sample size (ESS) of those weights is a standard diagnostic of how well the flow matches μ on the support it has reached (Liu, 2001; Doucet et al., 2001; Elvira et al., 2022).

The choice of training objective then divides the field. The original BG (Noé et al., 2019) and its variational-inference predecessor (Rezende and Mohamed, 2015) minimize the reverse Kullback–Leibler (KL) divergence

$$\text{(reverse)} \quad \text{KL}(\nu\|\mu) = \int_{\mathbb{R}^d} \nu(x) \log \frac{\nu(x)}{\mu(x)} dx, \quad (1)$$

which needs only samples from ν and energy evaluations, and is therefore cheap. It is also mode-seeking (Wainwright and Jordan, 2008): on multimodal μ the flow tends to collapse onto a subset of the modes and cannot recover from a mode it never visits, since a missed mode contributes no gradient (Wu et al., 2020; Felardos et al., 2023). Several lines soften this without removing it, interleaving MCMC kernels with the deterministic layers, folding the symmetries of μ into the architecture, lowering the gradient variance near the optimum, or annealing from a hot landscape toward μ (Wu et al., 2020; Köhler et al., 2020; Vaitl et al., 2022; Schopmans and Friederich, 2025). None restores a dropped mode, because the reverse KL places no sample where ν has no mass. Our $X_{\bar{\nu}}$ term below plays the leakage-suppression role one wants from a reverse KL-like penalty, restraining the mass ν places where μ has none. It reads this from a stop-gradient copy of ν , so the flow’s own sampling is never differentiated through, the source of the reverse KL’s instability on sharp targets.

The opposing objective minimizes the forward KL,

$$\text{(forward)} \quad \text{KL}(\mu\|\nu) = \int_{\mathbb{R}^d} \mu(x) \log \frac{\mu(x)}{\nu(x)} dx, \quad (2)$$

which is mass-covering, diverging wherever ν misses mass that μ carries (Wainwright and Jordan, 2008), but needs samples from μ , the very object one is trying to produce. Existing work breaks this circularity by choosing where the target samples come from, and each choice fixes what the method can see: maximum likelihood on configurations from a reference thermodynamic state (Wirnsberger et al., 2020), a flow trained on a running chain and used as a global proposal (Gabrié et al., 2022), fitting to a molecular dynamics (MD) trajectory (Klein et al., 2023), amortizing coverage across many molecules through a shared architecture (Klein and Noé, 2024), or supplying the intractable gradient from a persistent chain or an ℓ_2 fit to coupled targets (Naesseth et al., 2020; Rehman et al., 2026). In every case the forward KL is silent about mass its sample source never reaches: a mode absent from the reference data, the running chain, or the trajectory set leaves no trace in the gradient, as a missed mode is invisible to a chain. These remedies change the training signal or amortize coverage across systems. By contrast we keep the bare forward KL and confront the circularity directly, adding the log-ratio variation below, which reuses the same samples on a single target. Our AIS ladder supplies the target samples. Because that ladder inherits the classical limit, the regularizer draws on an independent sample source built to cover modes the chain misses.

The line closest to ours couples an annealing ladder to flow training so the flow improves the sampler that trains it. The flow annealed importance sampling bootstrap (FAB) (Midgley et al.) replaces the forward KL

by the α -divergence at $\alpha = 2$, whose integrand is proportional to μ^2/ν and whose minimizer also minimizes the variance of the importance weights. It draws training points from AIS chains targeting this minimum-variance surrogate and recycles them through a prioritized replay buffer. Related work weaves a learned flow into each stage of an SMC annealing path, reweights down a temperature ladder, adds an inference-time SMC correction at molecular scale, or curbs the mass teleportation these geometric paths allow (Arbel et al., 2021; Matthews et al., 2022; Schopmans and Friederich, 2025; Tan et al., 2025; von Klitzing et al., 2026). We take this AIS-coupled forward KL line as our point of departure and depart from it on three counts, set side by side in Remark 3. First, we keep the bare forward KL and add the log-ratio variation rather than change the divergence, so the minimizer is unchanged and the regularizer reuses the same samples and the same autograd graph. Second, our AIS ladder targets μ itself and is score-free in the flow: every chain moves under the target score $-\nabla U$, the flow enters only through evaluations of the log-ratio, and no replay buffer is kept. By contrast, the surrogate μ^2/ν has a score that carries the positional gradient of the pushforward density at every chain step. Third, mode discovery is delegated to a quench and temper measure $\hat{\mu}$ rather than asked of a chain that would still have to cross the barriers of μ . The trade-offs are real. The order-2 Rényi divergence behind μ^2/ν carries a stronger tail-coverage incentive than the forward KL, and with a sufficient annealing budget FAB needs no separate wide-coverage construction. Where an inference-time SMC pass buys scale, we keep the correction inside the tractable per-sample importance weight, so the score-free ladder and the absent replay buffer leave a small memory footprint and let the method run on a single consumer GPU. The off-policy log-dispersion regularizer penalizes a scalar, variance-type spread of $\log(\mu/\nu)$ (Schopmans et al., 2026). The log-ratio variation below instead penalizes the localized pairwise variation of $\log(\mu/\nu)$ under a chosen weight. This weight lets one family take three complementary roles from a single pairwise integral: mode discovery through $\hat{\mu}$, leakage suppression through $\bar{\nu}$, and shape control through μ .

A separate tradition drops the bijection and trains a stochastic process directly: a controlled stochastic differential equation, a denoising diffusion, or a flow-matching velocity carries the reference to μ along a continuous-time path, learning the drift rather than an invertible map (Zhang and Chen, 2022; Vargas et al., 2023; Berner et al., 2024; Nüsken and Richter, 2021; Richter and Berner, 2024). Energy-only variants such as iterated denoising energy matching (iDEM) alternate a sampling loop with a score- or velocity-fitting loop and a replay buffer (Akhound-Sadegh et al., 2024; Woo and Ahn, 2024). The non-equilibrium transport sampler augments AIS with a learned drift and so sits between the AIS-coupled and diffusion lines while still offering no tractable terminal density (Albergo and Vanden-Eijnden, 2025), and successors scale the idea to larger many-body systems (Havens et al., 2025; Akhound-Sadegh et al., 2025). These samplers are expressive and train from the energy alone. The cost is decisive for our purpose: once the bijection is gone, the terminal density ν is no longer available in closed form, so the importance weight μ/ν , the ESS read from it, and the reweighting that corrects residual bias are out of reach without a further per-sample ordinary-differential-equation solve (He et al., 2025). We commit to invertible flows for that reason: the invertibility of G keeps the log-ratio $\log(\mu/\nu)$ exact and inexpensive per sample, and with it the finite-sample importance-sampling pipeline on which both the forward KL and the log-ratio variation rest. The two lines are complementary rather than competing: one buys scale through dynamics at the cost of a tractable density, the other keeps the density and the reweighting at the cost of invertibility. Table 1 places these training lines side by side.

	objective	AIS required	importance weights	mode source	replay buffer	score-free flow
reverse KL	$\text{KL}(\nu\ \mu)$	—	✓	none	—	✓
forward KL	$\text{KL}(\mu\ \nu)$	✓	✓	AIS chain	—	✓
FAB	$\alpha=2$ divergence, μ^2/ν	✓	✓	AIS chain	✓	—
iDEM	energy matching	—	—	sampling loop	✓	—
ours	$\text{KL} + \text{X}_\mu + \text{X}_{(\hat{\mu} + \bar{\nu})/2}$	✓	✓	$\hat{\mu}$, no chain	—	✓

Table 1: Where the balanced $\text{KL} + \text{X}_\mu + \text{X}_{(\hat{\mu} + \bar{\nu})/2}$ flow sits among Boltzmann generator training lines. Entries are qualitative; ✓marks a property the line has by construction, — its absence.

What is new. The forward KL inherits the limits of its target surrogate: an annealing procedure initialized at ν may mix only among modes ν already covers. This leaves two core weaknesses, together with a third mechanism the same regularizer addresses. The first is silent missed modes, the fake ESS pitfall. If ν misses a mode of μ , the sampler initialized at ν may remain in the reached basins; its observed importance weights can be nearly uniform and the ESS close to one while target mass is absent (Noé et al., 2019; Midgley et al.). The second is accuracy under an imperfect target surrogate. Finite batches add noise, and the score-free annealing procedure of Section 3.2 is biased by construction. Because the forward KL constrains only the mean of $\log(\mu/\nu)$ under that surrogate and leaves its spread free, the approximation error passes directly into the trained flow. The third mechanism is leakage: where μ vanishes, the forward KL integrand $\mu \log(\mu/\nu)$ vanishes too, so the gradient carries no signal about mass that ν places outside the support of μ . Better calibration on the modes already reached does not cure a missed mode.

Our remedy is to regularize the forward KL with the *log-ratio variation*, a family of X functionals X_ω named for the pairwise variation of the log-ratio $z = \log(\mu/\nu)$ they penalize. For a non-negative weight ω on $\mathbb{R}^d$, not necessarily normalized, let

$$X_\omega(\mu\|\nu) = \iint_{\mathbb{R}^d \times \mathbb{R}^d} \omega(x) \omega(y) \left| \log \frac{\mu(x)}{\nu(x)} - \log \frac{\mu(y)}{\nu(y)} \right| dx dy, \quad (3)$$

against which both arguments are integrated. The idea is direct: matching ν to μ is the same as flattening z , and the pairwise difference cancels the unknown normalizer of μ . Where the forward KL constrains $\nu \approx \mu$ only in a global, Jensen-averaged sense, X_ω penalizes the variation of z wherever ω places mass and localizes the constraint to the support of ω . It vanishes whenever z is constant on that support. In particular, adding this non-negative term to the exact forward KL preserves $\nu = \mu$ as a global minimizer. For $\omega = \mu$ its integrand is built from the same log-ratio field as the forward KL, reusing the same samples and the same autograd graph.

Three choices of ω , with complementary roles, jointly address these failures. With $\omega = \mu$, the functional X_μ controls the spread of z under the target where the forward KL controls only its mean, tightening the density match once ν has reached the modes and trimming the bias inherited from imperfect annealed samples. With $\omega = \hat{\mu}$, a coarse, over-spread approximation of μ , the functional $X_{\hat{\mu}}$ drives mode discovery: by the insensitivity above, the only role of $\hat{\mu}$ is to place mass on modes and tails the current ν has not yet covered, and the gradient then pulls the flow toward them. We build $\hat{\mu}$ once, before training, with a quench and temper construction that bridges basins, collapses onto mode centers, and rejuvenates with a short Langevin chain. It errs toward broad coverage, and inherits a blind spot on heavy outer tails. With $\omega = \bar{\nu}$, a stop-gradient copy of the current pushforward, the functional $X_{\bar{\nu}}$ plays the reverse KL role without its instability, penalizing the mass ν places outside the support of μ and so suppressing the leakage the other two terms cannot see. Our preferred loss combines all three with the forward KL,

$$\mathcal{L}[\nu] = \text{KL}(\mu\|\nu) + \lambda X_\mu(\mu\|\nu) + X_{\alpha\hat{\mu} + \beta\bar{\nu}}(\mu\|\nu), \quad (4)$$

with $\lambda \geq 0$ and either $\alpha = \beta = 0$ or $\alpha, \beta \geq 0$ with $\alpha + \beta = 1$. The third term is X_ω at the mixture weight $\omega = \alpha\hat{\mu} + \beta\bar{\nu}$. Putting the mixture inside a single pairwise integral does more than add the two self-terms: expanding the product measure also generates a cross term that anchors the in-support and out-of-support log-ratios to each other. Written through the change of variables, every term reads off one scalar field along the flow,

$$z(y) = U_0(G(y)) - U(y) - \log |\det J_G(y)|, \quad (5)$$

up to an additive constant that cancels in each pairwise difference, so the loss in terms of the flow map G is

$$\mathcal{L}[G] = \underbrace{\mathbb{E}_{y \sim \mu} [z(y)]}_{\text{KL}(\mu\|\nu)} + \lambda \underbrace{\mathbb{E}_{y, y' \sim \mu} [|z(y) - z(y')|]}_{X_\mu(\mu\|\nu)} + \underbrace{\mathbb{E}_{y, y' \sim \alpha\hat{\mu} + \beta\bar{\nu}} [|z(y) - z(y')|]}_{X_{\alpha\hat{\mu} + \beta\bar{\nu}}(\mu\|\nu)}. \quad (6)$$

The X_μ term reuses the target batch and log-ratio evaluations of the forward KL, while the mixture term draws and evaluates one further batch. Adding either term preserves $\nu = \mu$ as a global minimizer of the exact forward KL.

Contributions. This paper makes the following contributions. We introduce the log-ratio variation X_ω , a family of X functionals that augment the forward KL by penalizing the pairwise variation of the flow’s own log-ratio $z = \log(\mu/\nu)$. We study three instances in complementary roles, X_μ , $X_{\hat{\mu}}$, and $X_{\bar{\nu}}$, together with the balanced loss $\text{KL} + X_\mu + X_{(\hat{\mu} + \bar{\nu})/2}$, which folds accuracy, mode discovery, and leakage suppression into one objective. The construction keeps the AIS target score free of the flow and requires no replay buffer; the balanced mixture adds one sampling batch beyond a plain forward KL step. We give a Fisher–Rao analysis showing that the forward KL decays along the $\text{KL} + \lambda X_\mu$ flow under no structural assumption on μ and, under a bound on λ , that the X_μ term lowers the accuracy floor left by biased target samples. We pair the ESS with a coverage diagnostic (Naeem et al., 2020), so density calibration on the reached support is not confused with mode coverage. Finally, we build an adaptive-temperature Boltzmann generator that grows its temperature ladder from per-stage sampling diagnostics. The numerical study isolates the three roles of the X functionals and separates two empirical regimes. Small batches and high dimension expose the intended accuracy robustness; on the multimodal clock ladder, the balanced loss also reduces the aggregate error-propagation factor at every tested batch size, including the largest, while the last stages at the smallest batch delimit the claim.

Organization. Section 2 states the main Fisher–Rao gradient-flow results for the regularized loss, with the full derivations, convergence proof, and biased-target accuracy analysis deferred to Appendix A. Section 3 turns the loss into a training algorithm for the flow map G , detailing the annealing procedure that approximates μ , the quench and temper construction of the wide-coverage measure $\hat{\mu}$, and the ESS and coverage diagnostics. Section 4 builds the adaptive-temperature Boltzmann generator on top of this loss. Section 5 reports the numerical study on model problems, and Section 7 concludes.

2 Fisher–Rao gradient flow analysis

Section 1 introduced the forward KL with log-ratio variation. Before turning it into a training algorithm in Section 3, we summarize its gradient-flow properties on the space of densities; Appendix A gives the full derivations and proofs. This continuous-time population idealization exposes the mechanism of the loss but does not imply a convergence rate for the discrete optimizer: we view ν as a probability density evolving along the gradient flow and ask how fast $\text{KL}(\mu\|\nu_t)$ decays to zero.

For a tractable analysis we keep only the forward KL and the shape term X_μ , the log-ratio variation at weight $\omega = \mu$. The mode-discovery and leakage terms $X_{\hat{\mu}}$ and $X_{\bar{\nu}}$ are left out: their wide-coverage and stop-gradient weights are training devices that do not enter this density-space idealization. The loss is then

$$\mathcal{L}[\nu] = \text{KL}(\mu\|\nu) + \lambda X_\mu(\mu\|\nu), \quad (7)$$

with $\lambda > 0$ fixed. Write $w = \mu/\nu$ for the importance weight and $z = \log w$ for the log-ratio. The statements below apply along any positive solution of the Fisher–Rao flow; for $\lambda \leq \frac{1}{2}$, the explicit birth term in Appendix A is non-negative and preserves positivity.

The choice of Riemannian metric matters. In the Wasserstein-2 geometry the forward KL is not displacement convex, so the Otto–Villani / Bakry–Émery machinery that yields exponential convergence for the reverse KL does not apply. We therefore work in the Fisher–Rao (FR) geometry, in which the forward KL admits an unconditional exponential decay and X_μ adds a nonlocal birth–death modulation: the update at a point is driven by how its importance weight ranks against the population of weights under μ . Writing the ν -mean as $\langle \cdot \rangle_\nu := \int(\cdot)\nu dx$, the FR gradient of a functional $\mathcal{F}$ and the flow it generates are

$$\text{grad}^{\text{FR}}\mathcal{F}(\nu) = \nu \left(\frac{\delta\mathcal{F}}{\delta\nu} - \left\langle \frac{\delta\mathcal{F}}{\delta\nu} \right\rangle_\nu \right), \quad \partial_t \nu_t = -\text{grad}^{\text{FR}}\mathcal{L}(\nu_t), \quad (8)$$

and the mean-subtraction makes $\int \text{grad}^{\text{FR}}\mathcal{F} = 0$, so the flow conserves probability mass.

Differentiating $\text{KL}(\mu\|\nu_t)$ along this flow, and reducing the X_μ term by a rank-correlation identity, gives the dissipation

$$\frac{d}{dt}\text{KL}(\mu\|\nu_t) = -\chi^2(\mu\|\nu_t) - \lambda \mathbb{E}_{x,y \sim \mu} \left[\left| \frac{\mu(x)}{\nu_t(x)} - \frac{\mu(y)}{\nu_t(y)} \right| \right], \quad (9)$$

with χ^2 the chi-squared divergence. The regularizer therefore contributes an additional non-positive term to the instantaneous KL dissipation. Since $\chi^2(\mu\|\nu_t) \geq \text{KL}(\mu\|\nu_t)$, Grönwall’s lemma yields the exponential rate

$$\text{KL}(\mu\|\nu_t) \leq e^{-t} \text{KL}(\mu\|\nu_0), \quad t \geq 0, \quad (10)$$

under no structural assumption on μ — no log-concavity, spectral gap, or growth condition. The rate does not claim forward KL is intrinsically faster. It is a population statement on the space of densities, agnostic to the cost of producing the target samples that AIS must supply. The contribution of X_μ to (9) is structural — it amplifies mass where ν_t undercovers μ — rather than a faster transient.

The convergence above assumes the outer expectation is taken under the exact target. In practice AIS supplies only an approximation $\tilde{\mu}$, and the realized loss replaces μ by $\tilde{\mu}$ in its outer integrands. At $\lambda = 0$ a stationary point is $\nu^* = \tilde{\mu}$: pure forward KL inherits the AIS bias verbatim, leaving the residual error $\text{KL}(\mu\|\tilde{\mu})$ as a hard floor. The shape term lowers this floor.

Theorem 1 *Assume μ is absolutely continuous with respect to $\tilde{\mu}$ and $\lambda \in [0, \frac{1}{2})$. Then every stationary density ν^* of the loss $\tilde{\mathcal{L}} = \text{KL}(\mu\|\cdot) + \lambda X_\mu(\mu\|\cdot)$ taken under $\tilde{\mu}$ satisfies*

$$\text{KL}(\mu\|\nu^*) \leq \text{KL}(\mu\|\tilde{\mu}) - \lambda \iint_{\mathbb{R}^d \times \mathbb{R}^d} \tilde{\mu}(x) \tilde{\mu}(y) \left| \frac{\mu(x)}{\nu^*(x)} - \frac{\mu(y)}{\nu^*(y)} \right| dx dy. \quad (11)$$

The rank-based feedback inflates ν^* where it undercovers μ and deflates it elsewhere, pulling the stationary density off $\tilde{\mu}$ back toward μ by an amount set by the spread of the importance weights μ/ν^* across the support of $\tilde{\mu}$. Thus X_μ is not merely a stabilizer for the forward KL flow: within the stated range of λ , it lowers the asymptotic accuracy floor left by biased target samples, a correction especially relevant at small batches and early stages. Our experiments use $\lambda = 1$ and show robust performance there, but the theorem itself guarantees the bound only for $\lambda < \frac{1}{2}$. The full FR-gradient derivation, the rank-correlation lemma, the convergence proof, and the accuracy proof are in Appendix A. This analysis lives on the space of densities and assumes access to μ ; the next section supplies approximations of μ and the wide-coverage measure $\hat{\mu}$ that make the loss trainable.

3 Flow training with log-ratio variations

This section turns the abstract loss of the introduction into a training algorithm for the flow map G . Let U_0 and U be the source and target potentials on $\mathbb{R}^d$, so that $\mu_0 \propto \exp(-U_0)$ is the samplable reference and $\mu \propto \exp(-U)$ is the target. We parameterize $G : \mathbb{R}^d \rightarrow \mathbb{R}^d$ so that $G_\# \mu \approx \mu_0$ — mapping target to source, so no inversion is required during training — and write $\nu := (G^{-1})_\# \mu_0$ for the induced approximation to μ . We first assemble the training loss and identify its sampling ingredients (Section 3.1), then supply them in turn: a score-free annealing approximation of μ (Section 3.2), the quench and temper construction of the wide-coverage measure $\hat{\mu}$ (Section 3.3), and the ESS and coverage diagnostics used to evaluate the trained flow (Section 3.4).

3.1 The log-ratio-variation training loss

By the change-of-variables formula the pullback density of μ_0 under G is $\nu(y) = \mu_0(G(y)) |\det J_G(y)|$, so the log-ratio between the target and the model along the flow is the single scalar field $z(y)$ of (5). Both ingredients of the loss are functionals of z . The forward KL is its mean under the target, $\text{KL}(\mu\|\nu) = \mathbb{E}_{y \sim \mu}[z(y)] + \text{const}$ (cf. (2)). Each log-ratio variation X_ω (3) is the mean absolute pairwise variation of z under a non-negative weight ω , $X_\omega(\mu\|\nu) = \mathbb{E}_{y, y' \sim \omega} |z(y) - z(y')|$ when ω is normalized (in general it is the double integral (3)). Combining the shape term $\omega = \mu$ with the balanced mixture $\omega = \alpha \hat{\mu} + \beta \bar{\nu}$ that handles mode discovery and leakage suppression, the total training loss in terms of the flow map G is

$$\mathcal{L}[G] = \underbrace{\mathbb{E}_{y \sim \mu}[z(y)]}_{\text{KL}(\mu\|\nu)} + \lambda \underbrace{\mathbb{E}_{y, y' \sim \mu}[|z(y) - z(y')|]}_{X_\mu(\mu\|\nu)} + \underbrace{\mathbb{E}_{y, y' \sim \alpha \hat{\mu} + \beta \bar{\nu}}[|z(y) - z(y')|]}_{X_{\alpha \hat{\mu} + \beta \bar{\nu}}(\mu\|\nu)}. \quad (12)$$

We refer to (12), previewed as (6), as forward KL with log-ratio variation. The balanced method used in the staged experiments is named by its loss, $\text{KL} + X_\mu + X_{(\hat{\mu} + \bar{\nu})/2}$. Note the two pushforward measures in

(12). The model distribution $\nu = (G^{-1})_{\#}\mu_0$, with density $\nu(y) = \mu_0(G(y))|\det J_G(y)|$, is the flow’s live approximation to the target μ and depends on the trainable parameters of G . The barred measure $\bar{\nu}$ is the *same* pushforward frozen at the current parameters, a stop-gradient copy of ν , which enters the mixture term only as a sampling weight and is never differentiated through.

The three sampling measures entering (12) address complementary failure modes of the bare forward KL:

- μ : *shape accuracy and robustness*. The shape term X_μ compares log-importance ratios pairwise across the target batch and lowers the asymptotic accuracy floor under biased target samples (Section 2). Small or noisy batches are its primary design regime; difficult high-dimensional or staged targets can retain an aggregate advantage at larger batches.
- $\hat{\mu}$: *mode discovery*. The wide-coverage measure built by quench and temper (Section 3.3) places sample pairs across modes that the score-free annealing surrogate did not reach within its budget, pulling ν toward mass the bare loss never sees.
- $\bar{\nu}$: *leakage suppression*. The frozen pushforward penalizes the mass the flow places outside the support of μ , preventing the overfit that target samples alone cannot detect: where μ provides no samples, the bare forward KL provides no signal.

Adding these non-negative terms to the exact forward KL preserves $\nu = \mu$ as a global minimizer. Their purpose is to improve numerical robustness without changing the desired population solution.

Remark (Robustness across regimes) *As discussed in Section 2, the exponential rate (10) is a statement about the idealized FR flow. The discrete Adam dynamics carries no corresponding speed guarantee. The loss was designed primarily to stabilize training with small or imperfect target batches. On a difficult staged target, however, the benefit need not be confined to that regime: in the clock experiment, $\text{KL}+X_\mu+X_{(\hat{\mu}+\bar{\nu})/2}$ reduces the aggregate propagation factor by 2.6–3.6 across every tested batch size, including $B = 2000$. This is benchmark-specific evidence, not a universal large-batch guarantee; at $B = 125$, the bare forward KL leads on the final four stages where inter-sector mass transport becomes limiting.*

The forward KL and X_μ share one target batch: evaluate $\mathbf{z} = (z(y_1), \dots, z(y_B))$, take $\text{mean}(\mathbf{z})$ for the forward KL, and $\text{mean}(|\mathbf{z} - \mathbf{z}[\sigma]|)$ for a random permutation σ . Thus X_μ adds only a permutation and pairwise difference to the same autograd graph. The balanced mixture evaluates one additional batch drawn from $(\hat{\mu} + \bar{\nu})/2$ and applies the same randomized pairwise estimator.

The loss draws on three measures, of which only two must be built. The $\bar{\nu}$ batch is the raw pushforward $\bar{y}_i = G^{-1}(x_i)$ of source samples $x_i \sim \mu_0$, taken with the gradient through G detached. This avoids differentiating through the sampled pushforward. The other two require construction: an approximation of the target μ for the forward KL and X_μ terms, supplied by the annealing procedure of Section 3.2, and samples of the wide-coverage measure $\hat{\mu}$ for the mixture term, supplied by quench and temper (Section 3.3). With equal target and mixture batch sizes, $\text{KL}+X_\mu+X_{(\hat{\mu}+\bar{\nu})/2}$ uses about twice the raw per-step sample evaluation of forward KL. Inference uses the same flow architecture for both methods.

We use two fixed settings. The *X_μ -only setting* $\lambda = 1, \alpha = \beta = 0$ adds only X_μ to the forward KL and reuses its target batch and autograd graph. The *balanced setting* $\lambda = 1, \alpha = \beta = \frac{1}{2}$ draws the mixture batch equally from $\hat{\mu}$ and $\bar{\nu}$: the $\hat{\mu}$ half pulls ν toward modes discovered by quench and temper that the annealing procedure did not reach within its budget, while the $\bar{\nu}$ half restrains mass leaked between or outside target modes. We use this balanced setting whenever the source μ_0 may not cover all modes of μ and report it as $\text{KL}+X_\mu+X_{(\hat{\mu}+\bar{\nu})/2}$.

3.2 Approximating μ by score-free annealing

Evaluating the μ -expectations in (12) requires samples from μ , which are not directly available. The natural surrogate draws $y_1, \dots, y_B$ from the pushforward $\nu = (G^{-1})_{\#}\mu_0$ and reweights them by

$$w(y) = \frac{\mu(y)}{\nu(y)} = \exp(z(y)) = \exp\left(U_0(G(y)) - U(y) - \log |\det J_G(y)|\right), \quad y \in \mathbb{R}^d, \quad (13)$$

followed by a resampling step and a short Langevin simulation to diffuse the duplicates. In high dimensions, however, μ and ν can differ greatly, so the weights $w(y_i)$ collapse onto a few outliers and the samples are of low quality. We temper the reweighting along the geometric path of $M + 1$ distributions

$$\pi_m(y) \propto \nu(y)^{1-\frac{m}{M}} \mu(y)^{\frac{m}{M}} = ((G^{-1})_{\#} \mu_0)(y) w(y)^{\frac{m}{M}}, \quad m = 0, 1, \dots, M, \quad (14)$$

so that $\pi_0 = \nu$ and $\pi_M = \mu$. Each step reweights by only $w(y)^{\frac{1}{M}}$, reducing the degeneracy of direct importance sampling.

Algorithm 1: Score-free annealing from $(G^{-1})_{\#} \mu_0$ toward μ

Input: samples $y_1, \dots, y_B$ from $(G^{-1})_{\#} \mu_0$, ladder length M , intermediate distributions π_m (14)

Output: a biased sample approximation $y_1, \dots, y_B$ of μ

```

1 for  $m = 1, \dots, M$  do
    // reweight along the geometric path
2   Compute the incremental weights on  $y_1, \dots, y_B$ ,  $w(y_i)^{\frac{1}{M}} = \pi_m(y_i)/\pi_{m-1}(y_i) = \exp(z(y_i)/M)$ ;
    // resample
3   Resample from the weighted set,  $y'_1, \dots, y'_B \leftarrow \text{resample}(y_{1:B}, (w(y_i)^{\frac{1}{M}})_{i=1}^B)$ ;
    // rejuvenate under the target
4   Diffuse the resampled duplicates by a short Langevin chain on  $\mu$ ,  $y_{1:B} \leftarrow \text{langevin}(y'_{1:B}, U)$ ;
5 end
```

An exact SMC sampler on the path (14) would rejuvenate at step m with a Markov kernel invariant for π_m . Its score contains $\nabla_y \log \nu(y)$ and hence positional derivatives of the flow density, including $\nabla_y \log |\det J_G(y)|$. To keep the training sampler score-free in the flow, Algorithm 1 instead rejuvenates directly under μ , using only $-\nabla U$. This substitution is a deliberate approximation: the target-invariant kernel does not preserve the intermediate π_m , so the output is a biased target surrogate rather than an exact AIS/SMC sample. The construction trades exact intermediate invariance for a cheap sampler that pulls resampled particles toward the final target. Section 2 models precisely this situation through the surrogate measure $\tilde{\mu}$, and the numerical study evaluates the resulting trained flow by independent importance weights and coverage diagnostics.

Remark (Choosing the ladder length M) For simple targets close to the source, $M = 1$ suffices. The choice $M = d$ is typical but not necessary in every test — an empirical rule by which the ladder length scales with the dimension of the deformation the flow must bridge, of which d is the worst case. For instance, annealing $\mu_0 \propto e^{-|x|^2}$ to $\mu \propto e^{-2|x|^2}$ on $\mathbb{R}^d$ deforms all d coordinates at once, and a ladder of $M = d$ steps keeps the weighted empirical distribution at a finite ESS that does not degrade with d . When the deformation concentrates on fewer directions, a much shorter ladder suffices.

Remark (Comparison with FAB) FAB (Midgley et al.) also couples AIS to flow training, but places the flow inside the chains. Its transitions target the mode-seeking surrogate μ^2/ν , whose score

$$\nabla_y \log \frac{\mu^2(y)}{\nu(y)} = -2\nabla U(y) - \nabla_y \log \nu(y)$$

carries the positional gradient of the pushforward density at every MCMC step, the term avoided by diffusing under μ above. Training then feeds on a replay buffer of past AIS samples reweighted against staleness. The scheme of this paper is score-free in the flow: every chain targets μ with score $-\nabla U$, the flow enters only through evaluations of the log-ratio z , and each gradient step runs Algorithm 1 afresh on a new source batch, so no buffer is kept. The one persistent set, the quench and temper pool of Section 3.3, is built once and covered by sample robustness. FAB explores through its AIS chain, which must still cross the barriers of μ ; here mode discovery is delegated to the quench, which uses no such chain. The trade-offs are real: the order-2 Rényi divergence behind μ^2/ν carries a stronger tail-coverage incentive than the forward KL, and with sufficient AIS budget FAB needs no wide-coverage construction.

3.3 Constructing $\hat{\mu}$ by quench and temper

The mixture term needs samples from the wide-coverage measure $\hat{\mu}$. For a single-flow problem these are prepared once before training; in the staged generator of Section 4, one pool is prepared for each stage and then held fixed during that stage’s optimization. At each training step a mini-batch is drawn from the pool and refreshed by a short Langevin chain before entering the X_ω computation of Section 3.1 with μ replaced by $\hat{\mu}$. The set is produced by quench and temper (QT), a map

$$\hat{\mathcal{Y}} = \text{QT}(U)$$

from the target potential U to a wide-coverage set $\hat{\mathcal{Y}} = \{x_1, \dots, x_P\}$ distributed as $\hat{\mu}$, proceeding in three phases: diffusion (melt), optimization (quench), and rejuvenation (temper).

Algorithm 2: Quench and temper (QT): proactive mode discovery

Input: target potential U (with fixed pool size P and melt scale m_e)
Output: wide-coverage set $\hat{\mathcal{Y}} = \{x_1, \dots, x_P\}$ forming $\hat{\mu}$, written $\hat{\mathcal{Y}} = \text{QT}(U)$

- 1 Draw an initial pool $x_1, \dots, x_P \sim \mu_0$;
// diffusion (melt)
- 2 Scatter the samples to a wider region of $\mathbb{R}^d$, $x_i \leftarrow x_i + m_e \zeta_i$ with $\zeta_i \sim \mathcal{N}(0, I_d)$ i.i.d. and melt scale $m_e > 0$;
// optimization (quench)
- 3 Drive each sample toward a mode center of μ by L-BFGS on U , updating $x_1, \dots, x_P$;
// rejuvenation (temper)
- 4 Spread the samples around each mode, $x_{1:P} \leftarrow \text{langevin}(x_{1:P}, U)$;

The output samples concentrate near mode centers of μ , while the initial Gaussian scatter broadens the set of basins the quench can reach. The $X_{\hat{\mu}}$ term therefore supplies training pairs on modes where the current pushforward ν places too little mass, including modes the score-free annealing procedure did not reach within its allotted budget.

A large melt scale m_e can bias the QT pool toward modes far from the bulk of μ_0 . This is intentional: $\hat{\mu}$ is a discovery measure rather than a calibrated approximation of target mode weights. Adding $X_{\hat{\mu}}$ to the exact forward KL still preserves $\nu = \mu$ as a global minimizer, while the empirical mode weights in the finite QT pool can affect the optimization path.

Under typical conditions the wide-coverage set $\hat{\mathcal{Y}}$ lies within the support of μ , and the temper step samples it directly. On extreme targets — heavy tails or stiff potentials — the L-BFGS quench can instead drive particles outside the support, into high-energy regions whose weights then destabilize training (the Langevin temper itself, like the AIS rejuvenation, stays well behaved). A remedy, a regularization we call δ -reweighted QT, tempers the rejuvenation: run the temper step on the flattened potential $(1 - \delta)U$,

$$x_{1:P} \leftarrow \text{langevin}(x_{1:P}, (1 - \delta)U), \quad \delta \in (0, 1),$$

whose stationary law $\propto e^{-(1-\delta)U}$ is broader, and attach to each sample the self-normalized importance weight $w_i = e^{-\delta U(x_i)} / \sum_j e^{-\delta U(x_j)}$. If the temper chain has equilibrated to the flattened law, this reweighting recovers expectations under e^{-U} ; at finite chain length it is an importance-weighted approximation. In either case the weights suppress leaked, high-energy samples in the $X_{\hat{\mu}}$ estimate without a hard cutoff. We write this δ -reweighted construction as

$$\hat{\mathcal{Y}} = \text{QT}_\delta(U),$$

recovering plain $\text{QT}(U) = \text{QT}_0(U)$ at $\delta = 0$.

3.4 Importance sampling at evaluation: ESS and coverage

At evaluation time the trained flow G supplies the model distribution $\nu = (G^{-1})_{\#}\mu_0$ as an approximation to μ , together with the tractable density $\nu(y) = \mu_0(G(y))|\det J_G(y)|$ that any normalizing flow makes available. The importance weight $w(y) = \mu(y)/\nu(y) = \exp(z(y))$ of (13) can therefore be evaluated on any pushforward sample without fitting an additional density model. Drawing $\mathbf{y} = (y_1, \dots, y_N)$ with $y_i = G^{-1}(x_i)$ and $x_i \sim \mu_0$ i.i.d., the weights $w(y_i)$ serve two uses. First, they diagnose how well ν approximates μ , through the effective sample size defined below. Second, they correct residual bias by reweighting or resampling the pushforward, followed when desired by a short Langevin refresh under μ . The better the flow, the more uniform the weights and the less correction the resampling performs.

We assess this pipeline through two metrics: the effective sample size, which measures how uniform the importance weights are on the support reached by the flow, and the coverage, which checks whether every mode of μ has been reached at all.

The first metric is the effective sample size, normalized to $[0, 1]$ by dividing the classical ESS by the sample count N . Let $\mathbf{y} = (y_1, \dots, y_N) \in \mathbb{R}^{N \times d}$ be i.i.d. samples from ν , with importance weights $w(y_i)$ given by (13). The ESS is defined as

$$\text{ESS}(\mathbf{y}) = \frac{(\sum_{i=1}^N w(y_i))^2}{N \sum_{i=1}^N w(y_i)^2} \in [0, 1], \quad (15)$$

where the upper bound 1 follows from the Cauchy–Schwarz inequality and is attained when all weights are equal. The factor N in the denominator distinguishes our normalized ESS from the classical effective sample size $(\sum_i w(y_i))^2 / \sum_i w(y_i)^2 \in [1, N]$ standard in the literature (Elvira et al., 2022): that quantity counts an effective number of samples and scales with the batch, whereas dividing by N leaves the per-sample efficiency, a fraction in $[0, 1]$ comparable across batch sizes. When G is perfectly trained the weights are constant and the ESS reaches 1. As noted in the introduction, however, this conclusion is one-way: a high ESS indicates that the sampled weights are uniform *on the support reached by ν* , but modes of μ that the pushforward never reaches contribute nothing to the sum and so cannot lower the ESS, which gives rise to the *fake ESS* pitfall.

To detect such missing modes, we pair the ESS with the coverage metric of Naeem et al. (2020). Let $\mathbf{x} = (x_1, \dots, x_P) \in \mathbb{R}^{P \times d}$ be the samples of $\hat{\mu}$ produced by the quench and temper algorithm (Algorithm 2). For each $1 \leq i \leq P$, the κ -nearest-neighbor radius of x_i (with $\kappa = 5$) is

$$\text{NND}_{\kappa}(x_i; \mathbf{x}) = \inf \left\{ r \geq 0 : \#\{1 \leq j \leq P : 0 < |x_i - x_j| < r\} \geq \kappa \right\}, \quad (16)$$

i.e., the distance from x_i to its κ -th nearest neighbor in $\mathbf{x}$. The coverage of $\mathbf{x}$ by the pushforward samples $\mathbf{y}$ is then

$$\text{Coverage}_{\kappa}(\mathbf{y}; \mathbf{x}) = \frac{1}{P} \sum_{i=1}^P \mathbf{1}\left\{ \exists 1 \leq j \leq N : |y_j - x_i| \leq \text{NND}_{\kappa}(x_i; \mathbf{x}) \right\}, \quad (17)$$

i.e., the fraction of points in $\mathbf{x}$ whose κ -nearest-neighbor ball contains at least one pushforward sample. A coverage close to 1 indicates that the finite QT reference set is well covered by the flow, while a coverage well below 1 exposes missing regions that the ESS alone cannot see. Because $\hat{\mu}$ is the same wide-coverage measure that $X_{\hat{\mu}}$ trains toward, this diagnostic checks only the modes found by QT; it cannot reveal a mode absent from the reference pool.

4 Adaptive-temperature Boltzmann generator

Section 3 introduced forward KL with log-ratio variation (12). For a difficult high-dimensional or strongly multimodal target, training one flow map directly from the source $\mu_0 \propto \exp(-U_0)$ to the target $\mu \propto \exp(-U)$ can be impractical. We instead decompose the transport into a sequence of easier stages along a temperature ladder. Each flow learns one stage, a controlled change in the potential, and the ladder is built adaptively from sampling diagnostics.

4.1 Construction of temperature ladder

To bridge the simple source $\mu_0 \propto \exp(-U_0)$ to the target $\mu \propto \exp(-U)$, we seek an increasing sequence

$$0 = t_0 < t_1 < \dots < t_K = 1$$

defining a sequence of intermediate distributions

$$\mu_k \propto \exp(-U_{t_k}), \quad U_{t_k}(x) = (1 - t_k) U_0(x) + t_k U_1(x), \quad k = 0, 1, \dots, K,$$

with $U_1 := U$ an alias for the target potential, so that μ_0 is the source and μ_K the target, together with K flow maps $\{G_k\}_{k=1}^K$ trained so that $(G_k^{-1})_{\#} \mu_{k-1} \approx \mu_k$. The number of stages K is flexible. The design choice is each increment $\Delta t_k = t_k - t_{k-1}$. Too large an increment degrades the training of stage k ; too small an increment wastes computation. We want the largest step each flow can still learn well.

Training the flow G_k requires a target surrogate for μ_k , supplied during optimization by Section 3.2. Separately, classical SMC on a linear potential bridge proposes the outer increment Δt_k . Its smallest per-step ESS is a heuristic measure of how difficult that candidate bridge is for the current particle population, so we shrink the increment until the estimate clears a fixed floor.

Algorithm 3: Adaptive selection of the temperature step Δt_k

Input: potentials U_0, U , previous level t_{k-1} , selection pool $\mathcal{P}$ of P samples from μ_{k-1} , floor $\tau = 0.7$, shrink factor $\gamma = 0.7$, initial guess t_{init}

Output: next level t_k , giving the step $\Delta t_k = t_k - t_{k-1}$

```

1  $U_{t_{k-1}} \leftarrow (1 - t_{k-1})U_0 + t_{k-1}U_1$ ;
   // start from the supplied initial guess
2  $t_k \leftarrow t_{\text{init}}$ ;
3 while true do
   // run classical SMC across the candidate step
4   Run  $M$ -step classical SMC on the pool  $\mathcal{P}$ , using the linear potential bridge from  $U_{t_{k-1}}$  to  $U_{t_k}$ 
     and a Langevin kernel invariant for each intermediate distribution ( $M$  as in Remark 2);
   // score the step by its worst per-step ESS
5   Record  $\text{ESS}_{\min}$ , the smallest per-step ESS over the  $M$  steps;
6   if  $\text{ESS}_{\min} \geq \tau$  then
7     | return  $t_k$ ;
8   else
9     | // shrink the step toward  $t_{k-1}$ 
9     |  $t_k \leftarrow t_{k-1} + \gamma(t_k - t_{k-1})$ ;
10  end
11 end
```

These SMC runs involve only the potentials U_0 and U , never the flow map. They use no QT samples, so their per-step ESS can overlook a mode the SMC population never reaches. We therefore use this ESS only to propose an increment; it is a heuristic measure of bridge difficulty, not a coverage certificate. The trained-flow validation in step (v) supplies a second acceptance gate, while QT supplies the independent discovery pool. The floor $\tau = 0.7$ and shrink factor $\gamma = 0.7$ are empirical. Starting from t_{init} , the procedure returns the first candidate along the shrink sequence whose estimated ESS clears the floor. As t_k approaches t_{k-1} , the bridge becomes trivial and its population ESS approaches 1; in finite samples we also impose the outer loop's retry and stage limits.

4.2 The full training procedure

The two ingredients assemble into a single loop (Algorithm 4) that grows the ladder one stage at a time, from U_0 to U , carrying a validation set $\mathcal{Y}_k \approx \mu_k$ across stages. Stage k runs five steps. Steps (iii)–(v) are wrapped in an acceptance loop that shrinks the increment and retrains whenever the trained flow’s validation ESS falls below a floor τ_v . The outer loop stops once t_k reaches 1, so the number of stages K is selected automatically.

- (i) *Source samples.* Stage k consumes source samples in three ways, governed by two size parameters: the *pool size* P and the *batch size* B . First, a selection pool of P samples supplies the SMC particles for the adaptive step selection of step (ii); it is drawn from the validation set $\mathcal{Y}_{k-1} \approx \mu_{k-1}$ carried over from the previous stage and refreshed by a short Langevin rejuvenation on $U_{t_{k-1}}$. Second, the quench and temper of step (iii) melts an initial pool of P samples from μ_0 (Algorithm 2). Third, the training batches of step (iv) are drawn at run time, B samples per gradient step, from the same $\mathcal{Y}_{k-1}$; no separate training set is built.
- (ii) *Adaptive step.* The next level t_k is fixed by the adaptive selection scheme (Algorithm 3). Starting from an initial guess t_{init} and shrinking, it takes the largest step below the guess whose classical SMC ESS from μ_{k-1} to μ_k , run on the selection pool of step (i), stays above the floor τ . The first stage uses the safe start $t_{\text{init}} = t_{\text{safe}}$. The leading increment faces the largest deformation, and an over-aggressive start is exposed only after a full training, so the first stage begins from a level it can handle. Later stages extrapolate the accepted increments, $t_{\text{init}} = \min(1, t_{k-1} + \Gamma(t_{k-1} - t_{k-2}))$ with enlarge factor Γ ; the step sizes grow, since only the first few stages are difficult. This sets the stage target $\mu_k \propto \exp(-U_{t_k})$ with $U_{t_k} = (1 - t_k)U_0 + t_k U_1$.
- (iii) *Wide-coverage set.* A wide-coverage set $\hat{\mathcal{Y}}_k$ of P samples, distributed as $\hat{\mu}_k$, is drawn by quench and temper (Algorithm 2) on the stage target U_{t_k} , spreading samples across its modes. It supplies the wide-coverage half of the mixture term in the training loss.
- (iv) *Training loss.* The flow G_k is trained by $\text{KL} + \text{X}_{\mu_k} + \text{X}_{(\hat{\mu}_k + \bar{\nu}_k)/2}$, the balanced loss of Section 3.1 specialized to the stage $\mu_{k-1} \rightarrow \mu_k$. With model $\nu_k = (G_k^{-1})_{\#} \mu_{k-1}$ and its stop-gradient copy $\bar{\nu}_k$, the stage log-ratio is

$$z_k(y) = U_{t_{k-1}}(G_k(y)) - U_{t_k}(y) - \log |\det J_{G_k}(y)|,$$

and the loss reads

$$\mathcal{L}[G_k] = \underbrace{\mathbb{E}_{y \sim \mu_k} [z_k(y)]}_{\text{KL}(\mu_k \| \nu_k)} + \lambda \underbrace{\mathbb{E}_{y, y' \sim \mu_k} [z_k(y) - z_k(y')]}_{\text{X}_{\mu_k}(\mu_k \| \nu_k)} + \underbrace{\mathbb{E}_{y, y' \sim (\hat{\mu}_k + \bar{\nu}_k)/2} [z_k(y) - z_k(y')]}_{\text{X}_{(\hat{\mu}_k + \bar{\nu}_k)/2}(\mu_k \| \nu_k)}, \quad (18)$$

its μ_k -expectations estimated from the biased score-free annealing surrogate: the level- $(k-1)$ batch $\mathcal{X}$ is pushed to $\bar{\mathcal{X}}_\nu = G_k^{-1}(\mathcal{X})$, distributed as ν_k , and annealed toward μ_k by Algorithm 1 with $G = G_k$. The wide-coverage measure $\hat{\mu}_k$ is realized by the set $\hat{\mathcal{Y}}_k$ of step (iii).

- (v) *Validation set update.* Once G_k is trained, the validation set advances from $\mathcal{Y}_{k-1}$ to $\mathcal{Y}_k \approx \mu_k$ by importance sampling. Each $y \in \mathcal{Y}_{k-1}$ is pushed through the trained inverse to $\tilde{y} = G_k^{-1}(y)$, a sample of $\nu_k = (G_k^{-1})_{\#} \mu_{k-1}$, and carries the importance weight

$$w_k(\tilde{y}) = \frac{\mu_k(\tilde{y})}{\nu_k(\tilde{y})} = \exp(z_k(\tilde{y})).$$

Its effective sample size $\text{ESS}(w_k)$ is the final, flow-based check on the increment. The adaptive selection of step (ii) sizes t_k from a classical SMC ESS that uses no QT samples and may report a fake ESS, whereas $\text{ESS}(w_k)$ measures the trained flow directly. If $\text{ESS}(w_k) < \tau_v$ (we use $\tau_v = 0.3$), the increment was too aggressive: we discard G_k , shrink to $t_k \leftarrow t_{k-1} + \gamma(t_k - t_{k-1})$, and repeat steps (iii)–(v) at the reduced step. Otherwise the validation set advances to $\mathcal{Y}_k = \text{langevin}(\text{resample}(\tilde{\mathcal{Y}}, w_k), U_{t_k}) \approx \mu_k$: resample $\tilde{\mathcal{Y}}$ by its weights w_k , then rejuvenate with a short Langevin chain on U_{t_k} .

Algorithm 4: Adaptive-temperature Boltzmann generator training

Input: potentials U_0, U , pool size P , batch size B , validation ESS floor $\tau_v = 0.3$, shrink factor $\gamma = 0.7$,
enlarge factor $\Gamma = 2$, safe start $t_{\text{safe}} = 0.2$

Output: ladder $0 = t_0 < \dots < t_K = 1$ and flow maps $\{G_k\}_{k=1}^K$ with $(G_k^{-1})_{\#} \mu_{k-1} \approx \mu_k$

// validation set, $\mathcal{Y}_k \approx \mu_k$

```
1  $\mathcal{Y}_0 \leftarrow$  samples from  $\mu_0$ ,  $t_0 \leftarrow 0$ ,  $k \leftarrow 0$ ;  
2 while  $t_k < 1$  do  
3    $k \leftarrow k + 1$ ;  
4   Form the stage source potential,  $U_{t_{k-1}} \leftarrow (1 - t_{k-1})U_0 + t_{k-1}U_1$ ;  
   // (i) source samples (selection pool)  
5   Draw a pool  $\mathcal{P}$  of  $P$  samples from  $\mathcal{Y}_{k-1}$  and rejuvenate,  $\mathcal{P} \leftarrow \text{langevin}(\mathcal{P}, U_{t_{k-1}})$ ;  
   // (ii) adaptive temperature selection  
6   if  $k = 1$  then  
7     Set  $t_{\text{init}} = t_{\text{safe}}$ ;  
8   else  
9     Set  $t_{\text{init}} = \min(1, t_{k-1} + \Gamma(t_{k-1} - t_{k-2}))$ ;  
10  end  
11  Choose  $t_k$  by adaptive selection (Algorithm 3) on the pool  $\mathcal{P}$  with initial guess  $t_{\text{init}}$ ;  
12  Form the target potential  $U_{t_k} \leftarrow (1 - t_k)U_0 + t_kU_1$ ;  
13  repeat  
14    Initialize  $G_k$  to the identity map;  
    // (iii) wide-coverage set  
15    Draw the wide-coverage set of  $P$  samples by quench and temper,  $\hat{\mathcal{Y}}_k \leftarrow \text{QT}(U_{t_k})$  (Algorithm 2);  
    // (iv) balanced log-ratio-variation loss  
16    for each gradient step do  
17      Draw a training batch of  $B$  samples  $\mathcal{X}$  from the validation set  $\mathcal{Y}_{k-1}$ ;  
18      Form the detached model batch,  $\tilde{\mathcal{X}}_\nu \leftarrow G_k^{-1}(\mathcal{X})$ , distributed as  $\nu_k = (G_k^{-1})_{\#} \mu_{k-1}$ ;  
19      Form the biased target surrogate  $\mathcal{X}_\mu$  by score-free annealing (Algorithm 1, from  $\nu_k$  toward  $\mu_k$ );  
20      Draw a wide-coverage batch  $\hat{\mathcal{X}}_\mu \subset \hat{\mathcal{Y}}_k$  and rejuvenate,  $\hat{\mathcal{X}}_\mu \leftarrow \text{langevin}(\hat{\mathcal{X}}_\mu, U_{t_k})$ ;  
21      Update  $G_k$  by a gradient step on  $\text{KL} + \text{X}_{\mu_k} + \text{X}_{(\hat{\mu}_k + \bar{\nu}_k)/2}$  in (18), from  $\mathcal{X}_\mu$  and the balanced  
      mixture of  $\hat{\mathcal{X}}_\mu$  and  $\tilde{\mathcal{X}}_\nu$ ;  
22    end  
    // (v) validation set update  
23    Push the validation set  $\tilde{\mathcal{Y}} \leftarrow G_k^{-1}(\mathcal{Y}_{k-1})$ ;  
24    Compute the weights  $w_k \leftarrow \exp(U_{t_{k-1}}(\mathcal{Y}_{k-1}) - U_{t_k}(\tilde{\mathcal{Y}}) - \log |\det J_{G_k}(\tilde{\mathcal{Y}})|)$ ;  
25    if  $\text{ESS}(w_k) < \tau_v$  then  
26      // abort the step and shrink toward  $t_{k-1}$   
       $t_k \leftarrow t_{k-1} + \gamma(t_k - t_{k-1})$ , then re-form  $U_{t_k} \leftarrow (1 - t_k)U_0 + t_kU_1$ ;  
27    else  
28      Resample  $\mathcal{Y}_k \leftarrow \text{resample}(\tilde{\mathcal{Y}}, w_k)$ ;  
29      Employ Langevin rejuvenation to obtain  $\mathcal{Y}_k \leftarrow \text{langevin}(\mathcal{Y}_k, U_{t_k})$ ;  
30    end  
31  until  $\text{ESS}(w_k) \geq \tau_v$ ;  
32 end  
33  $K \leftarrow k$ ;  
34 return  $\{t_k\}_{k=0}^K$  and  $\{G_k\}_{k=1}^K$ ;
```

The trained ladder is a Boltzmann generator for μ . Starting from $x \sim \mu_0$, each stage applies its map G_k^{-1} , then reweights and resamples by the importance weights of Section 3.4, as in step (v), carrying the samples from source to target one bridge at a time.

Setting every stage flow G_k to the identity reduces the adaptive-temperature generator to annealed SMC, a Monte Carlo sampler that requires no training and serves as the benchmark against the flow-based methods. With no flow to train, steps (iii) and (iv) (the wide-coverage set and balanced training loss) fall away, leaving the source pool, adaptive temperature selection, and reweight-resample-rejuvenate update. The log-Jacobian then vanishes, so the stage weight reduces to the classical tempering weight $w_k = \exp(U_{t_{k-1}}(\mathcal{Y}_{k-1}) - U_{t_k}(\mathcal{Y}_{k-1}))$. Run on the same adaptive ladder and acceptance standard, this baseline isolates the value added by the trained stage maps.

5 Numerical experiments on model problems

We test the three roles of the log-ratio-variation terms before applying the full objective to an annealed lattice model. The 2D targets isolate accuracy, mode discovery, and leakage control; the product multi-well measures how the accuracy gain scales with dimension; and the tilted ϕ^4 theory exposes a collective-barrier failure that ESS alone cannot detect. The p -state clock model then tests $\text{KL} + \text{X}_\mu + \text{X}_{(\hat{\mu} + \bar{\nu})/2}$ along a shared annealing ladder. Throughout, ESS is paired with a diagnostic that detects missing modes: coverage on the multimodal benchmarks, the minority-phase weight on ϕ^4 , and sector occupancy on the clock model.

5.1 2D benchmark distributions

We consider four 2D targets: Two-Moon, Three-Well, Himmelblau, and Sparse. Two-Moon has two distorted crescent components and isolates finite-batch accuracy because both components lie within reach of the source. The remaining targets isolate mode collapse: Three-Well contains a low-probability basin, Himmelblau has four separated wells, and Sparse places two tight modes far into the source tails.

For every target, we initialize the same neural spline flow at the identity and compare four objectives:

$$\text{KL}, \quad \text{KL} + \text{X}_\mu, \quad \text{KL} + \text{X}_\mu + \text{X}_{\hat{\mu}}, \quad \text{KL} + \text{X}_\mu + \text{X}_{(\hat{\mu} + \bar{\nu})/2}.$$

The flow has 32 spline bins, 6 transforms, and conditioner widths (128, 128). Each training step constructs a target surrogate by a single-stage AIS update through the current flow followed by 50 MALA steps of size 2×10^{-3} . Quench and temper uses melt scale 2, an Armijo L-BFGS quench with initial step 0.5 and 200 iterations, and a separate reference pool for coverage. The final ESS is computed from the flow importance weights on the full fixed source set. Coverage, with $\kappa = 5$ against the quench-and-temper reference pool, measures the fraction of target mass represented by the pushforward. Hence a collapsed flow can have ESS near one but cannot have full coverage.

Two-Moon. The source is $\mathcal{N}(0, I_2)$ and both crescents lie within its reach, so all four losses attain coverage 1. With batch size 200, 500 Adam steps, and learning rate 2×10^{-3} , the bare forward KL reaches ESS 0.8021. Adding X_μ raises it to 0.9157 without additional flow evaluations, while the equal-weight mixture reaches 0.9446 (Figure 1). This benchmark isolates the accuracy role of X_μ : the pairwise term constrains the spread of the log-ratio left uncontrolled by the mean-only forward KL objective.

Three-Well. The source is $\mathcal{N}(0, 0.25^2 I_2)$, narrower than the well separation and initially covering no basin. Training uses batch size 1000, 500 steps, and learning rate 10^{-3} . Forward KL and $\text{KL} + \text{X}_\mu$ fill the two favored wells and report ESS 0.9830 and 0.9933, respectively, while their coverage remains 0.7525. Quench and temper discovers the suppressed third well. The $\hat{\mu}$ and equal-weight mixtures raise coverage to 0.9763 and 0.9660 while retaining ESS 0.9771 and 0.9858 (Figure 2).

Himmelblau. The source is $\mathcal{N}(0, I_2)$, whereas the four wells lie at radius approximately 3–4. Training uses batch size 1000, 1000 steps, and learning rate 10^{-3} . Forward KL and $\text{KL} + \text{X}_\mu$ capture only two wells, giving coverage 0.5140 despite ESS 0.9722 and 0.9818. Both discovery mixtures attain coverage 1. However, the




Figure 1: Two-Moon target. Columns, from left to right, are forward KL, $\text{KL}+X_\mu$, $\text{KL}+X_\mu+X_{\hat{\mu}}$, and $\text{KL}+X_\mu+X_{(\hat{\mu}+\bar{\nu})/2}$. Each panel overlays the learned pushforward on the target energy and reports its final ESS and coverage. Gray points show the source.




Figure 2: Three-Well target. Methods and annotations are as in Figure 1.

$\hat{\mu}$ -only flow leaks thin trails of mass between wells and has ESS 0.9516, whereas adding the $\bar{\nu}$ half suppresses this leakage and raises the ESS to 0.9757 (Figure 3).

Sparse. The target is a four-component Gaussian mixture with component standard deviation 0.08: two modes lie near the source center and two lie in opposite corners, more than five source standard deviations away. With source standard deviation 0.6, batch size 1000, 1000 steps, and learning rate 5×10^{-3} , forward KL and $\text{KL}+X_\mu$ cover only the central pair. Their coverage is 0.5038, while their ESS remains 0.9590 and 0.9572. The $\hat{\mu}$ -only mixture restores coverage but pays a large leakage cost, with ESS 0.8112. The equal-weight mixture also attains coverage 1 and recovers most of the calibration, reaching ESS 0.9417.

target	KL	$\text{KL}+X_\mu$	$\text{KL}+X_\mu+X_{\hat{\mu}}$	$\text{KL}+X_\mu+X_{(\hat{\mu}+\bar{\nu})/2}$
Two-Moon	0.802 (1.000)	0.916 (1.000)	0.887 (1.000)	0.945 (1.000)
Three-Well	0.983 (0.753)	0.993 (0.753)	0.977 (0.976)	0.986 (0.966)
Himmelblau	0.972 (0.514)	0.982 (0.514)	0.952 (1.000)	0.976 (1.000)
Sparse	0.959 (0.504)	0.957 (0.504)	0.811 (1.000)	0.942 (1.000)

Table 2: Final ESS, with $\kappa = 5$ coverage in parentheses, on the four 2D targets. Bold marks the equal-weight mixture, which is the strongest overall setting: it attains essentially full coverage and lies within 0.02 of the best ESS on every target.

The four targets separate the objective’s three mechanisms. The X_μ term improves finite-batch accuracy when discovery is not limiting. The $X_{\hat{\mu}}$ term reveals modes absent from the live AIS surrogate. The $\bar{\nu}$ half then penalizes the low-probability bridges introduced by aggressive discovery. The equal-weight mixture




Figure 3: Himmelblau target. Methods and annotations are as in Figure 1.




Figure 4: Sparse target. Methods and annotations are as in Figure 1.

combines all three effects and is the only setting that gives essentially full coverage with near-best calibration throughout Table 2.

5.2 High-dimensional product multi-well

We next isolate accuracy scaling in dimension $d = 2^k$, $k = 1, \dots, 8$, using

$$U(\mathbf{x}) = \frac{1}{2} \|\mathbf{x}\|^2 + 12 \sum_{i=1}^k \exp(-x_i^2). \quad (19)$$

The first k coordinates are symmetric double wells with minima at $\pm\sqrt{\ln 24}$ and barrier approximately 9.9, while the remaining $d - k$ coordinates are standard Gaussian. The target therefore has exactly $2^k = d$ equal-weight modes. Multimodality occupies only a k -dimensional subspace, but the log-weight error accumulates across all d coordinates.

All four objectives use the same identity-initialized neural spline flow on $[-4, 4]^d$, with 16 bins, 6 transforms, and conditioner widths (256, 256). At every dimension, training uses batch size 250, 2000 Adam steps at learning rate 10^{-3} , gradient clipping at 10^2 , and a source pool of size $10^4 2^k$. The live target surrogate uses a single AIS hop followed by 50 MALA steps of size 2×10^{-3} . Quench and temper uses a pool of size $100 2^k$, melt scale 2, and an Armijo L-BFGS quench with initial step 0.5 and 200 iterations. Final ESS is evaluated on 20000 fresh source samples. Strict sign-pattern coverage counts a mode only if its empirical share exceeds half of the uniform share.

Every loss finds every mode, so ESS is meaningful here. The forward KL degrades from 0.974 at $d = 2$ to 0.420 at $d = 256$ as coordinatewise fitting errors accumulate in the log weights. Adding X_μ improves every dimension and reaches 0.562 at $d = 256$. The equal-weight mixture is best in every column, retaining ESS 0.941 at $d = 32$ and 0.596 at $d = 256$, a factor 1.42 above the bare forward KL.

loss \ d	2	4	8	16	32	64	128	256
forward KL	0.974	0.942	0.906	0.878	0.796	0.704	0.615	0.420
KL+ X_μ	0.997	0.983	0.958	0.957	0.909	0.850	0.744	0.562
KL+ X_μ + $X_{\tilde{\mu}}$	0.996	0.983	0.959	0.952	0.923	0.875	0.744	0.585
KL+ X_μ + $X_{(\tilde{\mu}+\tilde{\nu})/2}$	0.998	0.985	0.963	0.960	0.941	0.887	0.791	0.596

Table 3: Final ESS on the product multi-well target (19). Every loss finds all 2^k modes under the strict coverage threshold; bold marks the best ESS in each dimension.




Figure 5: Per-step training ESS at $d = 256$. Faint curves show raw histories and bold curves their moving averages. Forward KL rises quickly but plateaus near 0.44; the objectives containing log-ratio variations continue improving, and the quench-and-temper mixtures finish in the highest band.

Figure 5 separates transient speed from final accuracy. Forward KL climbs fastest during roughly the first 500 steps but saturates early. The X_μ objective initially trails it and then overtakes it, while the two quench-and-temper mixtures combine the fast initial climb with the highest final ESS. Thus the gain in Table 3 is not obtained by sacrificing the early training regime.

5.3 Lattice ϕ^4 field theory: fake ESS at a collective barrier

The tilted scalar ϕ^4 theory provides a sharp fake-ESS test. On a periodic $L \times L$ lattice, with $d = L^2$ and magnetization $m(\phi) = d^{-1} \sum_j \phi_j$, the action is

$$U(\phi) = -0.8 \sum_{\langle j,l \rangle} \phi_j \phi_l + \sum_{j=1}^d \left[\phi_j^2 + \frac{1}{2} (\phi_j^2 - 1)^2 \right] + \begin{cases} 0.0257 \sum_{j=1}^d \phi_j, & L = 6, \\ 0.0144 \sum_{j=1}^d \phi_j, & L = 8. \end{cases}$$

In the broken phase the field orders near two vacua $\pm v \mathbf{1}$, separated by a collective domain-wall barrier. The positive linear term gives the two phases unequal probability, so the minority weight $p_+ = \mathbb{P}(m > 0)$ is not fixed by symmetry and must be learned.

Parallel tempering gives the references $p_+ = 0.125$ and 0.128 , consistent with an independent symmetry-confined MALA reweighting. Each loss is run at seeds 0, 1, 2 using an identity-initialized neural spline flow on $[-3, 3]^d$, with 16 bins, 6 transforms, and conditioner widths (256, 256). The source is $\mathcal{N}(0, 0.25I_d)$, centered on the barrier. Training uses a fixed set of 10^5 source samples, batch size 500, 2000 Adam steps at learning rate 10^{-3} , and gradient clipping at 10^3 . The live surrogate uses one AIS hop and 50 MALA steps of size 2×10^{-3} . Quench and temper uses 2000 particles, melt scale 2, and an Armijo L-BFGS quench with initial step 0.1 and 200 iterations.

	$L = 6$ ($d = 36$)		$L = 8$ ($d = 64$)	
	ESS	p_+	ESS	p_+
forward KL	0.78/0.82/0.82	1/1/0	0.67/0.68/0.67	1/0/1
KL+ X_μ	0.93/0.93/0.94	1/1/0	0.81/0.82/0.82	1/0/0
KL+ X_μ + $X_{\hat{\mu}}$	0.88/0.87/0.87	0.127/0.128/0.126	0.62/0.61/0.63	0.127/0.126/0.124
KL+ X_μ + $X_{(\hat{\mu}+\bar{\nu})/2}$	0.90/0.89/0.88	0.128/0.129/0.128	0.67/0.68/0.71	0.130/0.129/0.130
PT reference	—	0.125	—	0.128

Table 4: Final ESS and reweighted minority-phase weight p_+ for three seeds on the tilted ϕ^4 lattice. Integer values 0 and 1 indicate collapse onto one vacuum. Bold marks the highest ESS among the objectives that recover both phases.

Forward KL and KL+ X_μ collapse at every size and seed. Because the AIS chain starts from the flow pushforward and cannot cross the collective barrier, the second vacuum remains absent and reweighting gives $p_+ = 0$ or 1. Nevertheless, KL+ X_μ has the highest ESS at both sizes: sharpening the captured vacuum improves calibration on the reached support while making the global failure no less severe.




Figure 6: Magnetization density on the $L = 8$ lattice at seed 0. Each panel shows the pushforward density (gray), the reweighted density (color), and the parallel-tempering reference (dotted); the title reports ESS and the reweighted p_+ .

Both quench-and-temper objectives populate both vacua at every seed. Their recovered p_+ lies between 0.124 and 0.130, within 0.004 of the corresponding reference. This recovery has an unavoidable ESS cost: a diffeomorphism maps the connected Gaussian source to a connected pushforward, so covering both vacua forces a low-weight bridge across the suppressed inter-phase region. At $L = 8$, the equal-weight mixture recovers part of this cost, with ESS 0.67–0.71 rather than 0.61–0.63 for the $\hat{\mu}$ -only objective. The $\bar{\nu}$ component penalizes precisely the leaked bridge mass while preserving discovery.

5.4 Boltzmann generator on p -state clock model

The preceding benchmarks use a single flow. We now test the full annealed construction on the $p = 6$ clock model on a periodic 8×8 lattice,

$$U(\boldsymbol{\theta}) = - \sum_{\langle jl \rangle} \cos(\theta_j - \theta_l) - \frac{1}{2} \sum_{j=1}^d \cos(6\theta_j), \quad \boldsymbol{\theta} \in [-\pi, \pi)^d, \quad d = 64, \quad (20)$$

where $\langle jl \rangle$ runs over nearest-neighbor pairs. The source is uniform on the torus and the bridge is $U_t = tU$. At low temperature, the target separates into six symmetry-related clock sectors.

Each stage uses an identity-initialized neural circular spline flow with 16 bins, 6 autoregressive transforms, and conditioner widths (256, 256). The validation set contains 4×10^5 particles and the SMC and quench-and-temper pools each contain 10^5 particles. Every stage uses 3000 Adam steps at learning rate 10^{-3} , a

four-stage training AIS surrogate with 100 MALA steps of size 10^{-3} , energy clipping at 10^3 , and gradient clipping at 10^2 . We test batch sizes $B = 2000, 1000, 500, 250, 125$.

The comparison is schedule-matched. Forward KL first constructs an adaptive ladder using SMC preselection and an ESS acceptance gate. The loss $\text{KL} + X_\mu + X_{(\hat{\mu} + \bar{\nu})/2}$ is then trained on the identical accepted values of t , without a second gate. Hence the two methods see the same bridge increments within each batch-size pair. Every configuration is a single run, so individual stage values carry single-run fluctuation.

The principal metric is the validation ESS of each accepted increment. We define

$$\text{propagation factor } F := \prod_k \text{ESS}_k^{-1}$$

estimates the factor by which staged reweighting enlarges Monte Carlo error across the full ladder. Smaller F is better. Table 5 reports every accepted stage and the resulting F .

stage k	1	2	3	4	5	6	7	8	9	10	F
t_k ($B = 2000$)	0.250	0.495	0.663	0.779	0.887	1.000	×	×	×	×	
forward KL	0.752	0.572	0.499	0.506	0.527	0.630	—	—	—	—	27.7
$\text{KL} + X_\mu + X_{(\hat{\mu} + \bar{\nu})/2}$	0.919	0.767	0.643	0.587	0.578	0.664	—	—	—	—	9.8
t_k ($B = 1000$)	0.250	0.495	0.663	0.779	0.887	1.000	×	×	×	×	
forward KL	0.678	0.492	0.419	0.450	0.487	0.585	—	—	—	—	55.7
$\text{KL} + X_\mu + X_{(\hat{\mu} + \bar{\nu})/2}$	0.863	0.659	0.539	0.507	0.522	0.610	—	—	—	—	20.2
t_k ($B = 500$)	0.250	0.421	0.590	0.705	0.818	0.945	1.000	×	×	×	
forward KL	0.580	0.556	0.428	0.464	0.401	0.412	0.784	—	—	—	120.9
$\text{KL} + X_\mu + X_{(\hat{\mu} + \bar{\nu})/2}$	0.778	0.705	0.530	0.535	0.430	0.416	0.784	—	—	—	45.7
t_k ($B = 250$)	0.250	0.421	0.539	0.654	0.734	0.811	0.887	1.000	×	×	
forward KL	0.470	0.464	0.512	0.428	0.522	0.493	0.534	0.476	—	—	319.6
$\text{KL} + X_\mu + X_{(\hat{\mu} + \bar{\nu})/2}$	0.662	0.603	0.625	0.503	0.584	0.540	0.564	0.476	—	—	94.1
t_k ($B = 125$)	0.175	0.295	0.413	0.528	0.607	0.685	0.761	0.835	0.908	1.000	
forward KL	0.455	0.506	0.472	0.434	0.533	0.514	0.514	0.547	0.601	0.515	891.1
$\text{KL} + X_\mu + X_{(\hat{\mu} + \bar{\nu})/2}$	0.656	0.698	0.637	0.553	0.634	0.559	0.509	0.501	0.559	0.496	246.3

Table 5: Validation ESS on every accepted stage of the schedule-matched clock-model comparison between forward KL and $\text{KL} + X_\mu + X_{(\hat{\mu} + \bar{\nu})/2}$. Within each batch-size block, both losses traverse the same ladder. The final column is $F = \prod_k \text{ESS}_k^{-1}$; bold marks the better loss on each stage and the smaller F .

The loss $\text{KL} + X_\mu + X_{(\hat{\mu} + \bar{\nu})/2}$ raises the geometric mean stage ESS by 0.07–0.11 at every batch size and reduces F by a factor between 2.6 and 3.6. The advantage therefore persists at the largest batch $B = 2000$, where F decreases from 27.7 to 9.8. As the batch shrinks, both ladders lengthen, from 6 stages at $B = 2000$ to 10 at $B = 125$, and error propagation grows sharply.

The improvement is concentrated on early and middle stages. There the bridge deformation is large but smooth, so fitting accuracy is the binding constraint. Near $t = 1$, the six sectors separate and the problem shifts from fitting within sectors to transporting their relative mass. Reweighting, common to both losses, performs that redistribution. At $B = 125$, forward KL is slightly better on the final four stages, where quench-and-temper discovers sectors but does not assign their exact weights. This is the main limitation observed in the ladder experiment. By $B \geq 1000$, $\text{KL} + X_\mu + X_{(\hat{\mu} + \bar{\nu})/2}$ again leads on every stage.

All six sectors are found at every batch size. To test whether staged resampling preserves their symmetry, we rebuild fresh samples through the trained $B = 2000$ ladders at sizes $N = 10^4 2^k$, $k = 0, \dots, 8$, using equal total work of 2.56×10^6 particles per row. We define

$$\text{occupancy bias} := \frac{1}{6} \sum_{s=0}^5 \left| p_s - \frac{1}{6} \right|.$$




Figure 7: Angle densities from 2×10^6 fresh samples advanced through the trained $B = 2000$ $\text{KL} + \text{X}_\mu + \text{X}_{(\hat{\mu} + \bar{\nu})/2}$ ladder by map, reweighting, resampling, and MALA. The single-site marginal peaks at the six clock angles. Pair-difference densities at increasing lattice separation retain an alignment peak at zero while transferring mass toward neighboring clock wells and the tails.

Both losses approach the Monte Carlo scaling rate: fitted log-log slopes are -0.474 for forward KL and -0.584 for $\text{KL} + \text{X}_\mu + \text{X}_{(\hat{\mu} + \bar{\nu})/2}$. The $\text{KL} + \text{X}_\mu + \text{X}_{(\hat{\mu} + \bar{\nu})/2}$ ladder has lower occupancy bias at every tested size and reaches 6.4×10^{-4} at $N = 2.56 \times 10^6$. The largest- N point of each curve is a single rebuild and therefore influences the fitted slopes. Together with Table 5, this shows that the per-stage ESS improvement survives composition without hiding a sector imbalance, while the late-stage reversal remains confined to the smallest-batch corner.

6 Boltzmann generators on molecular potential energies

Each model problem of the previous section isolates one failure mode of forward KL training on a smooth target: the 2D benchmarks, the high-dimensional multi-well, the lattice ϕ^4 theory, and the p -state clock model. The application that motivates a Boltzmann generator is a molecule. A molecular target exercises the whole pipeline at once. This section applies the adaptive-temperature, X-regularized generator of the previous section to three molecules of increasing difficulty in full internal coordinates, $d = 36$ to 60 , on a single consumer GPU. It has five parts: the motivation and the two difficulties that make a molecular potential hard (Section 6.1); *sharpening*, the simplification that makes the framework tractable on a singular target (Section 6.2); the molecule-specific engineering it requires (Section 6.3); the numerical results on glycerol, diethanolamine, and alanine dipeptide (Section 6.4); and the limitations of the molecular study (Section 6.5).

6.1 Motivation and background

Concretely, the target is the equilibrium (Boltzmann) distribution $\mu \propto e^{-U}$ of a molecular potential, with U its dimensionless reduced form. Its samples are the conformers that set binding affinities, reaction rates, and free energies in computational chemistry and drug design (Noé et al., 2019). The targets studied so far are smooth by design; a molecular potential is not. Drawing *independent* samples from μ is the central computational problem. MD and MCMC mix slowly across the high torsion barriers that separate conformers. A trained flow that emits independent conformers with a tractable density supports the exact importance-sampling pipeline of the previous sections, and this payoff justifies the training cost.

We take the potential energy to be the Amber molecular-mechanics force field (Cornell et al., 1995), the standard classical model. Its raw force-field energy $U_{\text{FF}}(x)$ in kJ/mol sums bonded terms, harmonic bonds




Figure 8: Clock-sector occupancy bias versus fresh rebuild size. Error bars are ± 2 standard errors, and the dashed line is an $N^{-1/2}$ reference anchored at the smallest $\text{KL} + X_\mu + X_{(\bar{\mu} + \bar{\nu})/2}$ point. Each curve replays one trained ladder, so the cross-loss comparison retains the single-run qualification.

and angles and periodic torsions, and nonbonded Lennard-Jones and Coulomb terms,

$$U_{\text{FF}}(x) = \sum_{\text{bonds}} k_b(r - r_0)^2 + \sum_{\text{angles}} k_\theta(\theta - \theta_0)^2 + \sum_{\text{torsions}} \frac{V_n}{2} [1 + \cos(n\phi - \gamma)] + \sum_{i < j} \left[4\epsilon_{ij} \left(\frac{\sigma_{ij}^{12}}{r_{ij}^{12}} - \frac{\sigma_{ij}^6}{r_{ij}^6} \right) + \frac{q_i q_j}{r_{ij}} \right], \quad (21)$$

the last sum running over nonbonded atom pairs, with charges in units that absorb the Coulomb constant. The Boltzmann distribution at temperature T is $\mu \propto e^{-U}$ in the dimensionless reduced potential

$$U(x) = \frac{1}{k_B T} U_{\text{FF}}(x),$$

with k_B the Boltzmann constant. We evaluate U_{FF} for the *isolated molecule in vacuum*, with no solvent and no nonbonded cutoff. Section 6.5 discusses this and two further simplifications of the molecular study.

We work in *whitened internal coordinates* (Noé et al., 2019). The six rigid-body degrees of freedom, global translation and rotation, leave U unchanged. We drop them and describe the shape by bond lengths, bond angles, and torsions. A z-matrix gives a smooth, invertible map $x \mapsto \xi \in \mathbb{R}^d$ with $d = 3N - 6$ for an N -atom molecule and a tractable volume Jacobian. Bonds and angles are stiff and near-Gaussian about their equilibria, so each is standardized to unit variance; torsions stay periodic on $[-\pi, \pi)$. The source $\mu_0 \propto e^{-U_0}$ is then a product base, standard normal on the bond/angle directions and uniform on the torsions. The target, pulled back through the change of variables, is

$$\mu(\xi) \propto \exp(-U(x(\xi))) |\det(\partial x / \partial \xi)|,$$

so U_0 and U as functions of ξ already absorb the change-of-variables log-Jacobian. This makes the target tractable for a flow. The rigid-body symmetries and the stiff, correlated bond/angle vibrations are factored out, leaving the combinatorially multimodal torsional landscape, the conformers, as the structure the diffeomorphism must transport.

Two features make (21) harder than a smooth model target, and they compound. First, the dimension grows with the molecule ($d = 36$ to 60 here), and the conformer landscape is *combinatorially multimodal*: every rotatable torsion is multi-well (gauche $^\pm$ /trans), so the joint over several torsions has dozens of basins the flow must discover. Second, the Lennard-Jones r^{-12} repulsion makes U *singular*: as any pair distance $r_{ij} \rightarrow 0$ the energy diverges, so the target has a hard, near-singular clash boundary that a smooth diffeomorphism cannot represent without enormous Jacobians. Neither difficulty is decisive alone, since high-dimensional smooth targets and low-dimensional singular ones are both routine. Together they are worse than either:

the singular clash core sits *inside* the high-dimensional multimodal landscape, and a single flow asked to discover modes *and* resolve the singularity at once trains badly at both. This compounding, absent from every model problem of the previous section, is what the rest of this section addresses. On alanine dipeptide ($d = 60$) below it is decisive: the clash core, not the torsion profile, becomes the binding constraint.

6.2 Sharpening: the core simplification

Sharpening keeps the singular part out of the flow, by annealing a second index the flow never sees. Annealing runs from the source U_0 of Section 6.1 to the target, for which we write the alias $U_1 = U$. A *regularization* index $s \in [0, 1]$ defines a soft-core target $U_{1,s}$ by taming the $r \rightarrow 0$ clash of the raw energy U_{FF} (21) before its reduction by $k_B T$. Two soft-core parameters act on U_{FF} : an energy cap e_{cap} (kJ/mol) that log-compresses the energy above its threshold to $e_{\text{cap}} + a \log(1 + (U_{\text{FF}} - e_{\text{cap}})/a)$ with fixed scale $a = 50$ kJ/mol, a C^1 wall of finite height and gradient (a hard clamp would zero the escape force); and a distance floor r_{floor} (nm) that replaces each nonbonded pair distance by $\max(r_{ij}, r_{\text{floor}})$, removing the r_{ij}^{-12} pole. Writing $U_{\text{FF}}|_{e_{\text{cap}}, r_{\text{floor}}}$ for the regularized energy, the reduced soft-core target is

$$U_{1,s} = \frac{1}{k_B T} U_{\text{FF}}|_{e_{\text{cap}}(s), r_{\text{floor}}(s)}, \quad (22)$$

with the cap threshold $e_{\text{cap}}(s)$ and floor $r_{\text{floor}}(s)$ annealed from soft ($s = 0$) to sharp ($s = 1$) on a geometric or arithmetic schedule,

$$e_{\text{cap}}(s) = e_{\min}(e_{\max}/e_{\min})^s \quad \text{or} \quad e_{\min} + (e_{\max} - e_{\min})s, \quad r_{\text{floor}}(s) = r_{\max} + (r_{\min} - r_{\max})s,$$

between a low cap $e_{\min}$ and high cap $e_{\max}$, and a loose floor $r_{\max}$ and the physical floor $r_{\min}$. Both controls act only on the high-energy clash region, inactive on the equilibrium basins where $U_{\text{FF}} \leq e_{\text{cap}}$ and $r_{ij} > r_{\text{floor}}$, so $U_{1,1} = U_1$ there and $U_{0,s} = U_0$ for all s . The bridge potential carries both indices,

$$U_{t,s} = (1 - t)U_0 + tU_{1,s},$$

and the whole run is a single anneal from the soft source $U_{0,0}$ to the sharp target $U_{1,1}$, the regularization reaching the temperature at every accepted stage, $s_k = t_k$. The two indices coincide only at the accepted nodes. Within a stage they differ: t first advances to t_k while s lags at s_{k-1} , then s catches up to s_k , as described next.

The change from Algorithm 4 is one extra, flow-free step per stage. Algorithm 4 advances the single temperature index t on the bridge U_{t_k} ; with sharpening each stage advances the pair (t, s) in two sub-steps,

$$\underbrace{(t_{k-1}, s_{k-1}) \xrightarrow{G_k} (t_k, s_{k-1})}_{\text{flow step: advance } t} \xrightarrow{w_k^{\text{sharp}}} \underbrace{(t_k, s_{k-1}) \rightarrow (t_k, s_k)}_{\text{sharpening step: advance } s}.$$

The *flow step* advances the temperature at the *lagged* regularization, $U_{t_{k-1}, s_{k-1}} \rightarrow U_{t_k, s_{k-1}}$. At fixed s_{k-1} the soft-core bridge $U_{t_k, s_{k-1}}$ is a single-temperature bridge to the target $U_{1, s_{k-1}}$, so the trained map G_k and its validation update are steps (i)–(v) of Algorithm 4 with U_1 replaced by $U_{1, s_{k-1}}$; the diffeomorphism transports only the smooth surrogate. The *Monte Carlo sharpening step* then advances the regularization at fixed temperature, $U_{t_k, s_{k-1}} \rightarrow U_{t_k, s_k}$, by a single importance reweight with weight

$$w_k^{\text{sharp}}(\xi) = \exp(U_{t_k, s_{k-1}}(\xi) - U_{t_k, s_k}(\xi)) = \exp(-t_k[U_{1, s_k}(\xi) - U_{1, s_{k-1}}(\xi)]), \quad (23)$$

followed by resampling and a short Langevin rejuvenation on U_{t_k, s_k} , with no flow. The importance step asks nothing the singularity forbids. It never inverts or differentiates U_1 ; it reweights and rejuvenates samples inside the basins they already occupy. The stiff repulsive wall costs some weight variance rather than the unbounded Jacobian it would impose on a smooth map. This is the asymmetry the split exploits: the flow handles the smooth, high-dimensional part, the Monte Carlo step the singular part, confined to the thin clash boundary.

Each stage thus yields two effective sample sizes: the flow’s validation $\text{ESS}_k^{\text{val}}$ from the flow step, the ESS of the step-(v) validation weights w_k of Algorithm 4, and the sharpening $\text{ESS}_k^{\text{sharp}}$ of w_k^{sharp} (23). Each is

the normalized effective sample size of Section 3.4, a fraction in $(0, 1]$. The Monte Carlo error-propagation factor through the two per-stage reweights,

$$F = \prod_k \frac{1}{\text{ESS}_k^{\text{val}}} \frac{1}{\text{ESS}_k^{\text{sharp}}}, \quad F = 1 \text{ ideal}, \quad (24)$$

charges the cost of both; it is smaller-is-better, and a run with no sharpening step has $\text{ESS}_k^{\text{sharp}} \equiv 1$. On the clash-dominated target a few near-singular weights collapse the raw validation ESS, so before forming it we saturate the log-weights about their median. With m the median of $\{\log w_i\}$ and $\sigma_w = c \text{median}_i |\log w_i - m|$, the *smooth* validation effective sample size is

$$\text{ESS}_k^{\text{val}*} = \text{ESS}(\{\tilde{w}_i\}), \quad \log \tilde{w}_i = m + \sigma_w \tanh((\log w_i - m)/\sigma_w), \quad (25)$$

the normalized ESS of Section 3.4 evaluated on the saturated weights, with $c = 6$ on alanine dipeptide as detailed in Section 6.3. Its error-propagation factor is

$$F^* = \prod_k \frac{1}{\text{ESS}_k^{\text{val}*}} \frac{1}{\text{ESS}_k^{\text{sharp}}}. \quad (26)$$

We report F where the raw validation ESS is well behaved, and F^* , written with a star, where the clash singularity forces the smooth estimator; the two are the same product evaluated with raw or smooth validation ESS. This factor is the headline metric for every result below: the variance amplification a downstream importance-sampling estimator inherits from the full ladder.

6.3 Engineering for the singularity

The singular targets require three further devices, all inert on the smooth ones (glycerol and diethanolamine ran with none of them). (i) *Particle drop*: each validation and resample ESS discards the worst 2% of particles before the ESS is formed, so a few clash-singular outliers, finite but enormous under the soft cap, cannot hijack the importance weights. (ii) *Best ESS gate snapshots*: within a stage the flow is snapshotted at each ESS gate, and the stage keeps the *highest* validation ESS snapshot rather than the last training step, which overfits the soft cap and sheds ESS; a step-0 identity snapshot is always included as a Monte Carlo fallback. (iii) *Distance-floor anneal*: for the worst clash core (alanine dipeptide) r_{floor} is annealed alongside the cap, softening the Lennard-Jones wall early and tightening it to the physical value only at $s = 1$, at the cost of a longer ladder. The wide-coverage pool $\hat{\mu}$ comes from the δ -reweighted quench and temper of Section 3.3 ($\delta = 0.1$), unchanged. The sharpening split remains the core mechanism; these three devices are the supporting engineering the singularity demands.

A metric-level choice accompanies them on the singular target: the per-stage validation ESS that gates and scores each stage is the smooth estimator $\text{ESS}_k^{\text{val}*}$ of (25), with $c = 6$ on alanine dipeptide. Saturating the log-weights about their median bounds the few clash-singular dominators that otherwise collapse the raw weighted ESS; a raw ESS of 0.4 then reads as a smooth ≈ 0.5 . Glycerol and diethanolamine, free of such dominators, keep the raw ESS and report F .

6.4 Numerical results

We treat three molecules of increasing difficulty in full internal coordinates on a single consumer GPU (RTX 5070 Ti): glycerol ($d = 36$), diethanolamine ($d = 48$), and alanine dipeptide ($d = 60$, the canonical Boltzmann generator benchmark). One comparison runs through all three: the X-regularized δ -reweighted loss against an annealed SMC reference, the same adaptive sharpening scheme with the flow set to the identity at every stage, which selects its own longer ladder. Because the factor is a product over stages, the *ratio* of trained-flow to annealed SMC F , not the absolute value, is the cross-molecule comparison. The trained flow cuts F by ~ 30 – $40\times$ on the two clash-mild, combinatorially multimodal targets, but only $\sim 2\times$ on alanine dipeptide, where the singular clash core rather than the multimodality dominates, the core that the Monte Carlo sharpening already resolves. The per-molecule subsections below give the full ablation, ladder, and reference figures for each. The baselines here, the bare forward KL and the identity-flow annealed SMC, are the natural same-cost comparisons for a forward KL-trained generator.

6.4.1 Glycerol ($d = 36$)

We ablate the generator on the full internal-coordinate glycerol target along three axes: the loss (forward KL versus the X-regularized $\text{KL} + \text{X}_\mu + \text{X}_{(\hat{\mu} + \bar{\nu})/2}$); the regularization schedule (*sharpen*: the soft-core cap e_{cap} is geometrically annealed from 100 to 200 kJ/mol along the ladder, with a Monte Carlo sharpening step per stage; *raw*: fixed $e_{\text{cap}} = 200$ kJ/mol, no anneal or sharpening); and the wide-coverage sampler (plain quench and temper versus its δ -reweighted variant, $\delta = 0.1$, defined only for the X loss). The headline metric is the error-propagation factor F (24). The *raw* runs take no sharpening step ($\text{ESS}^{\text{sharp}} \equiv 1$).

loss (raw pushforward)	K	wall (min)	F
annealed SMC	10	2	635.56
forward KL	7	215	40.86
$\text{KL} + \text{X}_\mu + \text{X}_{(\hat{\mu} + \bar{\nu})/2}$	6	162	21.96
$\text{KL} + \text{X}_\mu + \text{X}_{(\hat{\mu} + \bar{\nu})/2}$, δ -rw.	6	158	16.56
loss (sharpening)	K	wall (min)	F
annealed SMC	10	5	463.02
forward KL	4	88	44.00
$\text{KL} + \text{X}_\mu + \text{X}_{(\hat{\mu} + \bar{\nu})/2}$	4	73	22.51
$\text{KL} + \text{X}_\mu + \text{X}_{(\hat{\mu} + \bar{\nu})/2}$, δ -rw.	4	73	14.16

Table 6: Glycerol ($d = 36$) ablation, raw pushforward (top) and sharpening (bottom): ladder length K , training wall-clock, and the error-propagation factor F (24) (smaller better; the sharpening rows include the per-stage Monte Carlo sharpening reweight). δ -rw. = δ -reweighted quench and temper.

The X regularizer is the dominant factor (Table 6), roughly halving F at either schedule. The δ -reweighted quench and temper lowers F further in both schedules. Once the sharpening reweight is charged to F , the *sharpen* schedule no longer beats the fixed-cap *raw* run on F alone; its advantage is feasibility and speed. The lowest- F and fastest run is $\text{KL} + \text{X}_\mu + \text{X}_{(\hat{\mu} + \bar{\nu})/2}$, δ -reweighted, under the sharpen schedule. Turning the trained flow off entirely (the identity map at every stage, i.e. plain adaptive annealed SMC over the same ladder and acceptance standard) gives a far larger F in $K = 10$ stages. The trained flow thus cuts the propagation factor by ~ 30 – $40\times$ and shortens the ladder from $K = 10$ to $K = 4$ – 6 (Figure 9). This is the value the transport map adds over annealing alone.

Glycerol is achiral (each torsion marginal is even, $p(\phi) = p(-\phi)$; Figure 10), and every rotatable torsion is three-fold: gauche $^\pm$ near $\pm 60^\circ$ and trans near 180° . The $\pm\phi$ symmetry pairs gauche $^- \leftrightarrow$ gauche $^+$ and leaves trans self-symmetric. The joint target over the five rotatable torsions is therefore combinatorially multimodal (dozens of conformers), which the wide-coverage pool must cover.




Figure 9: Glycerol per-stage validation ESS as annealing-ladder arcs, raw (top) and sharpen (bottom). In each row forward KL is drawn above the axis (blue), $\text{KL} + X_\mu + X_{(\hat{\mu} + \bar{\nu})/2}$ below (purple), and its δ -reweighted variant further below (red). Each arc is an accepted stage; the apex label is that stage’s validation ESS; a cross marks a bridge the ladder failed to reach.




Figure 10: Glycerol torsion marginals of both generators against a converged annealed SMC reference (grey): forward KL (blue) and $\text{KL} + X_\mu + X_{(\hat{\mu} + \bar{\nu})/2}$, δ -reweighted (red), both sharpened with the cap annealed to $e_{\text{cap}} = 200$ kJ/mol. Both recover the multimodal gauche $^\pm$ /trans structure of the characteristic glycerol torsions.

6.4.2 Diethanolamine ($d = 48$)

We test the generator on the full internal-coordinate diethanolamine target ($\text{HN}(\text{CH}_2\text{CH}_2\text{OH})_2$, $d = 48$), comparing the two best losses under the same sharpening schedule (the soft cap geometrically annealed $e_{\text{cap}} : 100 \rightarrow 200$ kJ/mol, a Monte Carlo sharpening step per stage): forward KL, and the X-regularized $\text{KL} + \text{X}_\mu + \text{X}_{(\hat{\mu} + \bar{\nu})/2}$ with the δ -reweighted quench and temper ($\delta = 0.1$). Forward KL reaches $t = 1$ in $K = 6$ stages, the X-regularized loss in $K = 5$, a single direct $0.65 \rightarrow 1$ step.

loss	K	wall (min)	F
annealed SMC	10	7	1455.44
forward KL	6	896	92.60
$\text{KL} + \text{X}_\mu + \text{X}_{(\hat{\mu} + \bar{\nu})/2}$, δ -rw.	5	782	35.90

Table 7: Diethanolamine ($d = 48$): two trained-flow losses and an annealed SMC reference, all with a per-stage Monte Carlo sharpening reweight (cap e_{cap} geometrically annealed $100 \rightarrow 200$ kJ/mol): ladder length K , training wall-clock, and the error-propagation factor F (24) (smaller better; F includes the sharpening reweight). δ -rw. = δ -reweighted quench and temper.

The X-regularized loss with δ -reweighting reaches a lower F than forward KL (Table 7), a $\sim 2.6\times$ reduction in importance-weight waste at identical settings (glycerol showed a comparable $\sim 3.1\times$ gap). The advantage shows in the late, sharp stages and in the ladder length. Forward KL’s per-stage validation ESS stalls and it needs 6 stages to reach $t = 1$, whereas the $\text{KL} + \text{X}_\mu + \text{X}_{(\hat{\mu} + \bar{\nu})/2}$, δ -reweighted run holds a higher ESS and reaches $t = 1$ in 5 (Figure 11). This target is singular, high-dimensional, and combinatorially multimodal, so each run is a multi-hour computation. Closing it efficiently is the practical contribution of the X-regularized loss.




Figure 11: Diethanolamine per-stage ESS as annealing-ladder arcs (sharpening schedule): forward KL above the axis (blue), $\text{KL} + \text{X}_\mu + \text{X}_{(\hat{\mu} + \bar{\nu})/2}$, δ -reweighted, below (red). Each arc is an accepted stage; the apex label is that stage’s validation ESS, with the sharpening ESS in parentheses.

Diethanolamine has a plane of symmetry (each torsion marginal is even, $p(\phi) = p(-\phi)$). Its rotatable backbone torsions, O-C-C-N and C-C-N-C along the ethanolamine arms plus the hydroxyl rotations, are each multi-well (gauche $^\pm$ /trans), so the joint target is combinatorially multimodal, which the wide-coverage pool must cover.

Against the annealed SMC reference (Figure 12) the two losses are visually indistinguishable: both recover the multimodal backbone torsions, and neither is closer to the reference. The separation is in efficiency, not in the sampled marginals. $\text{KL} + \text{X}_\mu + \text{X}_{(\hat{\mu} + \bar{\nu})/2}$ reaches $t = 1$ in fewer stages, with a higher final-step ESS, a $\sim 2.6\times$ smaller error-propagation factor, and fewer rejected stage attempts in the early ladder. Switching the trained flow off entirely (the identity map at every stage, i.e. plain adaptive annealed SMC over the same ladder) gives a far larger F . The trained X-regularized flow therefore cuts the propagation factor by $\sim 40\times$ and halves the ladder length (5 vs 10). On this singular target the transport map adds even more over annealing alone than on glycerol, a justification for training a flow.

diethanolamine ($d = 48$) — torsion marginals: BG vs annealed SMC reference




Figure 12: Diethanolamine torsion marginals of both generators against a converged annealed SMC reference (grey): forward KL (blue) and $\text{KL} + X_\mu + X_{(\hat{\mu} + \bar{\nu})/2}$, δ -reweighted (red), both sharpened with the cap annealed to $e_{\text{cap}} = 200$ kJ/mol. Both recover the multimodal backbone torsions and track the annealed SMC reference.

6.4.3 Alanine dipeptide ($d = 60$)

We push the generator to alanine dipeptide (Ac-Ala-NHMe), the canonical Boltzmann generator benchmark, in full internal coordinates ($d = 60$). The step up from glycerol and diethanolamine is the target’s character. Those are dominated by smooth rotatable-bond multimodality. Alanine dipeptide folds the backbone tightly enough that the Lennard-Jones clash core, not the torsion profile, sets the hardest part of the anneal. We report the X-regularized loss only ($\text{KL} + X_\mu + X_{(\hat{\mu} + \bar{\nu})/2}$, δ -reweighted, $\delta = 0.1$), with a per-stage Monte Carlo sharpening in which *both* the cap $e_{\text{cap}}: 100 \rightarrow 200$ kJ/mol and the clash floor $r_{\text{floor}}: 0.2 \rightarrow 0.1$ nm are annealed arithmetically, unlike the geometric e_{cap} schedule of glycerol and diethanolamine. Alanine dipeptide is also the only target that anneals r_{floor} , which glycerol and diethanolamine hold fixed at 0.1 nm. The run reaches $t = 1$ in $K = 11$ stages, every stage holding a high validation and sharpening ESS (Table 8, Figure 13). The bare composed flow’s end-to-end ESS against the sharp target, from a single importance reweight of source samples pushed through the fully composed K -stage map and scored against $U_{1,1}$, is only $\sim 5 \times 10^{-6}$, so the per-stage sharpening, not a raw pushforward, is the deliverable.

loss	K	wall (min)	F^*
annealed SMC	12	23	129.70*
$\text{KL} + X_\mu + X_{(\hat{\mu} + \bar{\nu})/2}$, δ -rw.	11	~ 1000	67.95*

Table 8: Alanine dipeptide ($d = 60$), the X-regularized loss with a per-stage Monte Carlo sharpening reweight (the cap e_{cap} and floor r_{floor} annealed arithmetically): ladder length K , wall-clock, and the error-propagation factor F^* (26), starred throughout because every entry uses the smooth validation ESS (smaller better; F^* includes the sharpening reweight). The first row is an annealed SMC reference, the identity map over the same ladder. δ -rw. = δ -reweighted quench and temper.

Compared with the lighter targets, alanine dipeptide’s larger F^* is the price of its singular nonbonded core, not of its multimodality. Three alanine-dipeptide-only factors make this F^* read smaller than a strict like-for-like extension of that trend: the per-stage ESS is the smooth estimator $\text{ESS}_k^{\text{val}*}$ (25) rather than the raw weighted ESS; r_{floor} is annealed alongside the cap; and the ladder is longer ($K = 11$ versus 4 and 5), its smaller per-step jumps keeping each per-stage ESS high. All three raise the per-stage ESS entering F^* , and none applied to the lighter targets.

The instability is localized to a clash bottleneck near $t \approx 0.3$ –0.43. The stages spanning $t \approx 0.29$ –0.43 (soft cap $e_{\text{cap}} \approx 129$ –143 kJ/mol) are where the nonbonded core first bites: the run minimum validation ESS sits here, and these are the only stages that shrink the step or retry heavily. Below $t \approx 0.3$ the torsions dominate and the flow trains in a single attempt. Above $t \approx 0.5$ the cap is sharp enough that the basins separate cleanly and ESS recovers to ≈ 0.8 . The binding constraint for alanine dipeptide is the




Figure 13: Alanine dipeptide per-stage ESS history: each panel is the per-step effective sample size during that stage’s X-regularized δ -reweighted flow training (red), against the identity-map baseline at $t = 0$ (blue dashed). Below each panel is that stage’s accepted validation ESS with the sharpening ESS in parentheses. The dip across $t \approx 0.29$ – 0.43 is the clash bottleneck, with ESS recovering to ≈ 0.8 once the cap is sharp enough to separate the basins.

clash singularity, not the multimodality. This is the qualitative difference from the $d \leq 48$ molecules, and the reason the three devices of Section 6.3 (particle drop, best ESS gate snapshots, r -floor anneal), inert on glycerol and diethanolamine, were necessary here.

The per-stage ESS history (Figure 13) exposes two coupled instabilities that the best ESS gate works around but does not cure. First, the within-stage ESS is *non-monotone and unstable*: it can rise, plateau, then drop abruptly (the collapse late in the $t = 0.65$ stage and the volatility at $t = 1$ are the clearest cases), so the last training step is often far below the best. Second, near the clash bottleneck ($t \approx 0.3$ – 0.5) several flows *degenerate toward the identity map*: the best achievable ESS barely exceeds the $t = 0$ identity baseline, i.e. training fails to improve on doing nothing. Both follow from the *singularity of the potential*: the Lennard-Jones $r \rightarrow 0$ divergence leaves the soft-capped target ill-conditioned where the nonbonded core first bites, so the gradient signal that would pull the flow off identity is weakest there. We report this as a known limitation; remedies (stronger or adaptive regularization annealing, parameterizations tailored to the repulsive wall) are left to future work.

Alanine dipeptide’s backbone ϕ, ψ are multi-basin (Figure 14). A second, alanine-dipeptide-specific feature is chirality: the classical force field is mirror-symmetric (every term in (21) is invariant under reflection), so the L- and D-forms are isoenergetic, and the internal-coordinate flow, carrying no chirality constraint,

alanine dipeptide ($d = 60$) — torsion marginals: BG vs annealed SMC reference




Figure 14: Alanine dipeptide torsion marginals of the BG deliverable (red) against a converged annealed SMC reference (grey). The four most multimodal proper torsions are shown; the densities are $\pm\phi$ -symmetrised, the achiral target having $p(\phi) = p(-\phi)$. The BG recovers the multimodal backbone torsions and tracks the reference; the peptide-bond torsion ω (C-C-N-C) is trans-dominant in the generator, the one torsion whose high rotation barrier also makes the uniform-init annealed SMC reference slow to converge.

emits *both* where a physical trajectory would stay in one. In the Ramachandran (Figure 15) this appears as the $(\phi, \psi) \rightarrow (-\phi, -\psi)$ point symmetry, the D-basins mirroring the L-basins, each populated.

The X-regularized BG closes the full internal-coordinate alanine dipeptide target to $t = 1$ in 11 stages, recovering the multimodal backbone torsions against a converged annealed SMC reference and emitting both chiral forms of the mirror-symmetric target. The lesson from the lighter $d \leq 48$ molecules is inverted: alanine dipeptide’s difficulty is not its combinatorial torsion multimodality but its nonbonded clash core, concentrated in the $t \approx 0.3$ – 0.43 stages, which the particle drop, best ESS gate snapshots, and r -floor anneal together make tractable. Across all three molecules the pattern is consistent: the sharpening split makes the singular target trainable at all, and the X-regularized flow then earns a ~ 30 – $40\times$ reduction in importance-weight waste where the landscape is multimodal but clash-mild, narrowing to $\sim 2\times$ where the singularity dominates.


6.5 Limitations

Four simplifications bound the molecular experiments, each a direction for future work. First, the targets are evaluated in vacuum: U_{FF} is the isolated-molecule Amber potential (21) with no solvent and no nonbonded cutoff. Solvation only reshapes U_{FF} , not the sampler, and the vacuum targets already display the well-separated potential wells the generator must discover and resolve; implicit or explicit solvent and larger peptides are left untreated. Second, the internal-coordinate generator carries no chirality constraint, so for the achiral Amber force field it samples both the L- and D-enantiomers (Section 6.4.3), whereas a physical trajectory stays in one; isolating a single enantiomer by a filter or restraint is left to future work. Third, we do not benchmark against FAB (Midgley et al.): its large per-step batches and replay buffer make a like-for-like comparison under the same compute budget as these forward KL-type losses prohibitive, so the natural same-cost baselines here are the bare forward KL and the identity-flow annealed SMC. Fourth, on the alanine dipeptide clash core the per-stage flow training is unstable and can degenerate toward the identity map (Section 6.4.3); the best ESS gate works around this but does not cure it, and a regularization or parameterization that conditions the repulsive wall remains open.

7 Conclusions

Forward KL training of a normalizing flow Boltzmann generator usually trains well. But it can miss modes and report a fake effective sample size. Noisy or biased annealed samples limit its accuracy. A smaller leakage failure rides alongside, where the target vanishes. We close these gaps by regularizing the forward KL with a term that reuses the flow’s own log-ratio.

alanine dipeptide ($d = 60$) — Ramachandran: BG vs annealed SMC reference



BG: forward $\text{KL} + X_\mu + X_{(\hat{\mu} + \bar{\nu})/2}$ (δ -reweighted)

Figure 15: Alanine dipeptide Ramachandran (ϕ, ψ), shared log-density scale: the BG deliverable (right) against the annealed SMC reference (left). Red dots mark the maximum-probability conformer of each enantiomer (L at $\phi < 0, \psi > 0$; D at $\phi > 0, \psi < 0$). Both panels populate the same backbone basins with the $(\phi, \psi) \rightarrow (-\phi, -\psi)$ point symmetry: the achiral vacuum force field makes the L- and D-enantiomers isoenergetic, and the generator, carrying no chirality constraint, samples both.

- We propose the log-ratio variation, a family of X functionals. They augment the forward KL by penalizing the pairwise variation of $\log(\mu/\nu)$ under a non-negative weight. One forward pass of the flow computes them. The shape term costs little beyond a plain forward KL step; the mixture adds one sampling batch.
- One regularizer takes three roles through its weight. X_μ controls the shape of the log-ratio and stabilizes training. $X_{\hat{\mu}}$ drives mode discovery, drawing the flow toward a wide-coverage measure that quench and temper prepares before training. $X_{\bar{\nu}}$ suppresses leakage.
- The balanced loss $\text{KL} + X_\mu + X_{(\hat{\mu} + \bar{\nu})/2}$ combines the three roles in one objective. It folds mode discovery and leakage suppression into a single pairwise integral. It serves as a default and needs little tuning. A Fisher–Rao analysis shows the forward KL still decays along the X_μ -regularized flow. Under a bound on the regularization strength, the X_μ term can lower the accuracy floor that biased AIS samples leave behind. The forward KL is sensitive to these samples early in training, so the X -regularized losses help most when the training batch is small. The loss applies across smooth benchmarks, a high-dimensional multi-well, and lattice field theories, where it tends to discover modes and sharpen accuracy.
- We build an adaptive-temperature Boltzmann generator on the loss. It grows its ladder from the SMC effective sample size. On molecular potentials, a sharpening reweight pulls each stage toward the physical potential. In our experiments it tends to improve per-stage quality and reduce importance-weight waste over annealing alone. Gains are smaller where a singular clash core, not multimodality, is the difficulty.

Several directions invite further study. First, a comparison with FAB (Midgley et al.), and a possible combination with it, would weigh our quench against its replay buffer; the same comparison extends to other AIS-based flow samplers. Second, the training instability at the singular clash core calls for a remedy

that conditions the repulsive wall, since the present best ESS gate works around the instability rather than curing it. Third, larger and solvated systems are a natural target, where the potential reshapes but the sampler need not change. Finally, a JAX implementation in place of the present PyTorch one would speed up training and inference.

Acknowledgement

The normalizing flows are built on the Python packages `zflows` and `zflows.md`, written by the author Xuda Ye and installable through Anaconda. The source code reproducing all numerical experiments is openly available. Both are listed below:

```
conda install xudaye::zflows xudaye::zflows.md  
https://github.com/xuda-ye-math/X-regularization
```

Numerical tests run on an NVIDIA RTX 5070 Ti GPU with 16 GB of VRAM.

A Fisher–Rao gradient flow with log-ratio variation

Section 2 states the main gradient-flow results used in the paper. This appendix supplies their derivations and proofs on the space of densities. We view ν as a generic probability density evolving along the gradient flow of the loss under an appropriate Riemannian metric on the space of densities (Ambrosio et al., 2008; Villani, 2009) and ask how fast ν_t converges to the target μ in $\text{KL}(\mu\|\nu_t)$.

In this section we restrict to the loss with only the forward KL and the shape regularizer X_μ , the log-ratio variation at weight $\omega = \mu$,

$$\mathcal{L}[\nu] = \text{KL}(\mu\|\nu) + \lambda X_\mu(\mu\|\nu) = \int_{\mathbb{R}^d} \mu(x) \log \frac{\mu(x)}{\nu(x)} dx + \lambda \iint_{\mathbb{R}^d \times \mathbb{R}^d} \mu(x) \mu(y) \left| \log \frac{\mu(x)}{\nu(x)} - \log \frac{\mu(y)}{\nu(y)} \right| dx dy, \quad (27)$$

where $\lambda > 0$ is a fixed constant. The calculations below hold along any positive solution of the Fisher–Rao flow; when $\lambda \leq \frac{1}{2}$, the birth term in (33) is non-negative and hence preserves positivity. Throughout this section we abbreviate the importance weight by

$$w(x) := \frac{\mu(x)}{\nu(x)},$$

so that the log-ratio is $z(x) = \log w(x) = \log \mu(x) - \log \nu(x)$, the field z of (5). The importance weight w is distinct from the localization measure ω in X_ω .

The choice of Riemannian metric matters. In the Wasserstein-2 geometry, induced by optimal transport (Jordan et al., 1998; Otto, 2001), the forward KL fails to be displacement convex along Wasserstein-2 geodesics, and the standard Otto–Villani / Bakry–Émery machinery (Otto and Villani, 2000; Bakry et al., 2014) that yields exponential convergence for the reverse KL does not apply. We therefore work in the Fisher–Rao (FR) geometry (Amari, 2016; Bauer et al., 2016), in which the forward KL admits an unconditional exponential decay rate and the regularizer X_μ contributes a qualitatively new, nonlocal birth–death modulation (Lu et al., 2019): the local update at x is driven by how the importance weight at x ranks against the population of weights under μ .

Denoting the ν -mean by $\langle \cdot \rangle_\nu := \int (\cdot) \nu dx$, the Fisher–Rao gradient of a functional $\mathcal{F}[\nu]$ and the associated dissipation identity along the FR gradient flow $\partial_t \nu_t = -\text{grad}^{\text{FR}} \mathcal{F}(\nu_t)$ are

$$\text{grad}^{\text{FR}} \mathcal{F}(\nu) = \nu \left(\frac{\delta \mathcal{F}}{\delta \nu} - \left\langle \frac{\delta \mathcal{F}}{\delta \nu} \right\rangle_\nu \right), \quad \frac{d}{dt} \mathcal{F}[\nu_t] = - \int_{\mathbb{R}^d} \left(\frac{\delta \mathcal{F}}{\delta \nu} - \left\langle \frac{\delta \mathcal{F}}{\delta \nu} \right\rangle_{\nu_t} \right)^2 \nu_t dx. \quad (28)$$

The mean-subtraction enforces $\int \text{grad}^{\text{FR}} \mathcal{F} = 0$, so the FR flow conserves probability mass.

A.1 Derivation of X_μ -regularized Fisher–Rao gradient flow

The first variation of the forward KL with respect to ν admits the form

$$\frac{\delta \text{KL}(\mu\|\nu)}{\delta \nu}(x) = -\frac{\mu(x)}{\nu(x)} = -w(x), \quad (29)$$

so by (28), using $\langle w \rangle_\nu = \int_{\mathbb{R}^d} \mu dx = 1$, the FR gradient of the forward KL is

$$\text{grad}^{\text{FR}} [\text{KL}(\mu\|\nu)](x) = \nu(x) - \mu(x). \quad (30)$$

For the X_μ functional, the inner log-ratio difference has first variation

$$\frac{\delta (\log w(x) - \log w(y))}{\delta \nu(x)} = \frac{1}{\nu(y)} \delta(x - y) - \frac{1}{\nu(x)}.$$

By the symmetry of X_μ in x and y ,

$$\begin{aligned}\frac{\delta X_\mu(\mu\|\nu)}{\delta\nu}(x) &= 2 \int_{\mathbb{R}^d} \mu(x)\mu(y) \operatorname{sgn}(w(x) - w(y)) \frac{\delta(\log w(x) - \log w(y))}{\delta\nu(x)} dy \\ &= 2 \int_{\mathbb{R}^d} \mu(x)\mu(y) \operatorname{sgn}(w(x) - w(y)) \left(\frac{1}{\nu(y)} \delta(x - y) - \frac{1}{\nu(x)} \right) dy \\ &= -2w(x) \int_{\mathbb{R}^d} \mu(y) \operatorname{sgn}(w(x) - w(y)) dy,\end{aligned}$$

where the contribution of $\delta(x - y)$ vanishes because $\operatorname{sgn}(w(x) - w(y)) = 0$ at $y = x$. Hence

$$\frac{\delta X_\mu(\mu\|\nu)}{\delta\nu}(x) = -2w(x)S(x), \quad S(x) := \mathbb{E}_{y \sim \mu} [\operatorname{sgn}(w(x) - w(y))]. \quad (31)$$

We call S the *rank field* of the importance weight under μ : $S(x) > 0$ indicates that $w(x)$ is comparatively large within the population of weights, and hence that ν pointwise undercovers μ relative to that population; $S(x) < 0$ has the opposite interpretation, and $S \equiv 0$ at $\nu = \mu$. By antisymmetry,

$$\mathbb{E}_\mu[S] = \mathbb{E}_{x, y \sim \mu} [\operatorname{sgn}(w(x) - w(y))] = 0,$$

and the FR gradient of X_μ with respect to ν is

$$\operatorname{grad}^{\text{FR}}[X_\mu(\mu\|\nu)](x) = -2\mu(x)S(x). \quad (32)$$

Since the FR gradient is linear in the functional, the FR gradient of $\mathcal{L}[\nu] = \text{KL}(\mu\|\nu) + \lambda X_\mu(\mu\|\nu)$ is the direct sum of (30) and (32):

$$\operatorname{grad}^{\text{FR}}[\mathcal{L}](x) = \nu(x) - \mu(x) - 2\lambda\mu(x)S(x).$$

The regularized Fisher–Rao gradient flow $\partial_t \nu_t = -\operatorname{grad}^{\text{FR}}[\mathcal{L}]$ therefore reads

$$\partial_t \nu_t(x) = \mu(x) - \nu_t(x) + 2\lambda\mu(x)S_t(x), \quad \text{where } S_t(x) := \mathbb{E}_{y \sim \mu} \left[\operatorname{sgn} \left(\frac{\mu(x)}{\nu_t(x)} - \frac{\mu(y)}{\nu_t(y)} \right) \right]. \quad (33)$$

Equation (33) is a *nonlocal* PDE: its right-hand side at x depends on $S_t(x)$, which in turn requires the entire profile of the importance weight μ/ν_t over the support of μ . Mass is conserved along the flow because

$$\frac{d}{dt} \int_{\mathbb{R}^d} \nu_t(x) dx = 2\lambda \int_{\mathbb{R}^d} \mu(x) S_t(x) dx = 0,$$

and the flow is stationary at $\nu_t = \mu$, where $\mu(x)/\nu_t(x) \equiv 1$ and hence $S_t \equiv 0$. When $\lambda = 0$, (33) reduces to the pure Fisher–Rao gradient flow of the forward KL.

A.2 Convergence analysis of the X_μ -regularized flow

To analyze the convergence rate of the regularized flow (33), we take the forward KL $\text{KL}(\mu\|\nu_t)$ as the error metric and differentiate it along the flow:

$$\frac{d}{dt} \text{KL}(\mu\|\nu_t) = - \int_{\mathbb{R}^d} \frac{\mu^2}{\nu_t} (1 + 2\lambda S_t) dx + \int_{\mathbb{R}^d} \mu dx = -\chi^2(\mu\|\nu_t) - 2\lambda \int_{\mathbb{R}^d} \frac{\mu^2}{\nu_t} S_t dx,$$

where the chi-squared divergence is

$$\chi^2(\mu\|\nu) := \int_{\mathbb{R}^d} (w - 1)^2 \nu dx = \int_{\mathbb{R}^d} w^2 \nu dx - 1.$$

The sign of the second integral is settled by the following rank-correlation identity.

Lemma 1 For any probability density ν on $\mathbb{R}^d$,

$$\int_{\mathbb{R}^d} \frac{\mu(x)^2}{\nu(x)} S(x) dx = \frac{1}{2} \mathbb{E}_{x,y \sim \mu} [|w(x) - w(y)|] \geq 0,$$

with equality if and only if w is μ -almost-everywhere constant.

Proof. Since $\mu^2/\nu = \mu w$, the definition of S gives

$$\int_{\mathbb{R}^d} \frac{\mu^2}{\nu} S dx = \mathbb{E}_\mu [w(x) S(x)] = \mathbb{E}_{x,y \sim \mu} [w(x) \operatorname{sgn}(w(x) - w(y))].$$

Symmetrizing in $x \leftrightarrow y$ gives

$$\int_{\mathbb{R}^d} \frac{\mu^2}{\nu} S dx = \frac{1}{2} \mathbb{E}_{x,y \sim \mu} [(w(x) - w(y)) \operatorname{sgn}(w(x) - w(y))] = \frac{1}{2} \mathbb{E}_{x,y \sim \mu} [|w(x) - w(y)|].$$

□

Applying Lemma 1 with $\nu = \nu_t$ and $S = S_t$, the dissipation rewrites as

$$\frac{d}{dt} \operatorname{KL}(\mu \| \nu_t) = -\chi^2(\mu \| \nu_t) - \lambda \mathbb{E}_{x,y \sim \mu} \left[\left| \frac{\mu(x)}{\nu_t(x)} - \frac{\mu(y)}{\nu_t(y)} \right| \right]. \quad (34)$$

The elementary inequality $w \log w \leq w^2 - w$ applied to $w = \mu/\nu_t$ gives $\chi^2(\mu \| \nu_t) \geq \operatorname{KL}(\mu \| \nu_t)$, hence $\frac{d}{dt} \operatorname{KL}(\mu \| \nu_t) \leq -\operatorname{KL}(\mu \| \nu_t)$, and Grönwall's lemma yields the exponential decay

$$\operatorname{KL}(\mu \| \nu_t) \leq e^{-t} \operatorname{KL}(\mu \| \nu_0), \quad t \geq 0, \quad (35)$$

with no structural assumption on μ : neither log-concavity, nor a spectral gap, nor any growth condition. This contrasts with the Wasserstein-2 geometry, where the forward KL is not displacement convex and an analogous exponential rate is unavailable without further assumptions.

The uniform convergence rate (35) should *not* be read as a blanket statement that minimizing the forward KL is intrinsically faster or more efficient than minimizing the reverse KL. The FR flow is a continuous-time idealization on the space of densities and is agnostic to the sampling cost. In practice, the difficulty of training has been partially transported to the AIS step: evaluating the forward KL and its gradient requires accurate samples of μ , and producing such samples in high dimensions is itself a hard problem. This is the reason that AIS is needed, and that the $X_{\hat{\mu}}$ regularizer must compensate for modes that AIS alone fails to reach. The exponential rate (35) quantifies how fast the flow *can* converge given accurate target samples; it does not eliminate the burden of producing them.

A naive comparison of dissipation rates between KL-only and KL + λX_μ would also be misleading, since scaling the loss by a constant is mere time reparametrization with no qualitative gain. The substantive contribution of X_μ is structural rather than rate-based: while the pure KL FR gradient is local in x , the rank field S_t is global, so the regularized flow (33) couples the update at x to the entire distribution of importance weights under μ . This nonlocal coupling amplifies the birth rate on undercovered regions of μ and suppresses it on overcovered ones, a global comparison that pure forward KL, which knows only whether $\mu(x)/\nu_t(x)$ exceeds or falls below 1 pointwise, cannot produce.

Two further cautions concern the discrete realization of the flow. First, the time-reparametrization argument cuts in both directions once the optimizer enters. Training follows Adam steps, not the FR flow: the per-coordinate second-moment normalization absorbs a global rescaling of the loss, so a larger gradient buys no speedup by itself, while a regularizer that reshapes the relative magnitudes of the gradient components does change the optimizer's effective time scale. Part of any observed difference in training speed between KL and KL + λX_μ may therefore be optimizer scaling rather than information geometry, and we claim no universal transient speedup on this basis. Second, the dissipation (34) is a population statement, while each gradient step sees an estimate from a finite, generally biased target surrogate. Small batches are the primary regime for which the regularizer was designed. On the difficult staged clock target, however, KL + $X_\mu + X_{(\hat{\mu} + \bar{\nu})/2}$ retains an aggregate advantage across $B = 125$ –2000, including a 2.8-fold smaller propagation factor at $B = 2000$. This does not imply uniform dominance: at $B = 125$, the final four stages favor the bare forward KL.

Remark (Monotone decay of X_μ) The regularizer X_μ also decreases monotonically along the regularized flow: pairing the variation $\delta X_\mu / \delta \nu = -2wS$ (31) against the flow (33) gives

$$\frac{d}{dt} X_\mu(\mu \| \nu_t) = -2 \int_{\mathbb{R}^d} \frac{\mu^2}{\nu_t} S_t dx - 4\lambda \int_{\mathbb{R}^d} \frac{\mu^2}{\nu_t} S_t^2 dx \leq 0,$$

with strict inequality unless μ/ν_t is μ -almost-everywhere constant, the first term being non-positive by Lemma 1 and vanishing only in that case. Thus X_μ is itself driven to zero in the long-time limit.

A.3 Asymptotic accuracy under biased AIS samples

The convergence analysis of Section A.2 assumes that the outer expectation in $\text{KL}(\mu \| \nu) + \lambda X_\mu(\mu \| \nu)$ is taken under the exact target μ . In practice, the algorithm only has access to an approximation $\tilde{\mu}$ of μ supplied by AIS, and the realized loss reads

$$\tilde{\mathcal{L}}[\nu] = \int_{\mathbb{R}^d} \tilde{\mu}(x) \log w(x) dx + \lambda \iint_{\mathbb{R}^d \times \mathbb{R}^d} \tilde{\mu}(x) \tilde{\mu}(y) |\log w(x) - \log w(y)| dx dy, \quad (36)$$

where the pointwise ratio $w(x) = \mu(x)/\nu(x)$ is still evaluated against the exact target, since it is computed per sample from the target potential U and the flow's own density. Repeating the derivation of Section A.1 with μ replaced by $\tilde{\mu}$ in the outer integrands gives the Fisher–Rao gradient

$$\text{grad}^{\text{FR}}[\tilde{\mathcal{L}}](x) = \nu(x) - \tilde{\mu}(x)(1 + 2\lambda \tilde{S}(x)), \quad \tilde{S}(x) := \mathbb{E}_{y \sim \tilde{\mu}} [\text{sgn}(w(x) - w(y))].$$

A stationary point ν^* of (36) therefore satisfies the self-consistent equation

$$\nu^*(x) = \tilde{\mu}(x) \left[1 + 2\lambda \int_{\mathbb{R}^d} \tilde{\mu}(y) \text{sgn} \left(\frac{\mu(x)}{\nu^*(x)} - \frac{\mu(y)}{\nu^*(y)} \right) dy \right]. \quad (37)$$

At $\lambda = 0$ the equation collapses to $\nu^* = \tilde{\mu}$: pure forward KL training inherits the bias of the target surrogate $\tilde{\mu}$ verbatim, and the residual error is exactly $\text{KL}(\mu \| \tilde{\mu})$. For $\lambda > 0$, however, the rank-based correction $1 + 2\lambda \tilde{S}^*$, where $\tilde{S}^*$ denotes $\tilde{S}$ evaluated at $\nu = \nu^*$, inflates ν^* above $\tilde{\mu}$ where the importance weight μ/ν^* is high, that is, where ν^* still undercovers μ , and deflates it where the weight is low, providing a global feedback that pulls ν^* from $\tilde{\mu}$ back toward μ . Two facts prepare the statement. For $\lambda < \frac{1}{2}$ the bracket in (37) is strictly positive, because $\tilde{S}^* \in [-1, 1]$. Moreover, the antisymmetry of the sign in x, y implies

$$\int_{\mathbb{R}^d} \tilde{\mu}(x) \tilde{S}^*(x) dx = \mathbb{E}_{x, y \sim \tilde{\mu}} \left[\text{sgn} \left(\frac{\mu(x)}{\nu^*(x)} - \frac{\mu(y)}{\nu^*(y)} \right) \right] = 0,$$

hence every stationary point $\nu^* = \tilde{\mu}(1 + 2\lambda \tilde{S}^*)$ is a probability density.

Theorem 2 Assume that μ is absolutely continuous with respect to $\tilde{\mu}$ and that $\lambda \in [0, \frac{1}{2})$. Then every stationary density ν^* of the loss $\tilde{\mathcal{L}}$ (36) satisfies

$$\text{KL}(\mu \| \nu^*) \leq \text{KL}(\mu \| \tilde{\mu}) - \lambda \iint_{\mathbb{R}^d \times \mathbb{R}^d} \tilde{\mu}(x) \tilde{\mu}(y) \left| \frac{\mu(x)}{\nu^*(x)} - \frac{\mu(y)}{\nu^*(y)} \right| dx dy. \quad (38)$$

Proof. Substituting the stationary identity $\nu^* = \tilde{\mu}(1 + 2\lambda \tilde{S}^*)$ into the difference of forward KLs and using $\mu = \nu^* \cdot (\mu/\nu^*) = \tilde{\mu}(1 + 2\lambda \tilde{S}^*)(\mu/\nu^*)$,

$$\text{KL}(\mu \| \tilde{\mu}) - \text{KL}(\mu \| \nu^*) = \int_{\mathbb{R}^d} \mu(x) \log(1 + 2\lambda \tilde{S}^*(x)) dx = \int_{\mathbb{R}^d} \tilde{\mu}(x) \frac{\mu(x)}{\nu^*(x)} u(x) \log u(x) dx,$$

where $u(x) := 1 + 2\lambda \tilde{S}^*(x) \geq 1 - 2\lambda > 0$, since $\tilde{S}^* \in [-1, 1]$ (a $\tilde{\mu}$ -average of signs) and $\lambda < \frac{1}{2}$. The elementary inequality $u \log u \geq u - 1$ for $u > 0$ then yields

$$\text{KL}(\mu \| \tilde{\mu}) - \text{KL}(\mu \| \nu^*) \geq 2\lambda \int_{\mathbb{R}^d} \tilde{\mu}(x) \frac{\mu(x)}{\nu^*(x)} \tilde{S}^*(x) dx.$$

Applying the rank-correlation argument of Lemma 1 verbatim with μ replaced by $\tilde{\mu}$ in the outer measure,

$$\int_{\mathbb{R}^d} \tilde{\mu}(x) \frac{\mu(x)}{\nu^*(x)} \tilde{S}^*(x) dx = \frac{1}{2} \iint_{\mathbb{R}^d \times \mathbb{R}^d} \tilde{\mu}(x) \tilde{\mu}(y) \left| \frac{\mu(x)}{\nu^*(x)} - \frac{\mu(y)}{\nu^*(y)} \right| dx dy,$$

which gives (38). $\square$

The right-hand side of (38) (Theorem 2) contrasts the two roles of X_μ training. At $\lambda = 0$ the bound is tight at $\nu^* = \tilde{\mu}$ and the residual KL is fixed at $\text{KL}(\mu \parallel \tilde{\mu})$; the surrogate bias is a hard floor. Switching on the regularizer with $\lambda < \frac{1}{2}$ lowers this floor (strictly, unless μ/ν^* is constant on the support of $\tilde{\mu}$) by an amount controlled by the discrepancy of the importance weights μ/ν^* across the support of $\tilde{\mu}$: the larger this discrepancy, the more the rank-based feedback pulls ν^* toward μ . The X_μ regularizer is therefore not merely a numerical stabilizer for the forward KL flow: within the theorem’s assumptions, it lowers the asymptotic accuracy floor of the trained density when the target surrogate is biased.

The threshold $\lambda < \frac{1}{2}$ is a sufficient condition enforcing strict positivity of ν^* uniformly in x . The experiments use $\lambda = 1$ and show robust performance, but they do not extend this accuracy-floor theorem beyond its stated range. Within that range, the result shows that X_μ alone can reduce the bias floor inherited from an inaccurate target surrogate.

References

- Tara Akhound-Sadeh, Jarrod Rector-Brooks, Avishek Joey Bose, Sarthak Mittal, Pablo Lemos, Cheng-Hao Liu, Marcin Sendera, Siamak Ravanbakhsh, Gauthier Gidel, Yoshua Bengio, Nikolay Malkin, and Alexander Tong. Iterated denoising energy matching for sampling from Boltzmann densities. In *International Conference on Machine Learning*, 2024.
- Tara Akhound-Sadeh, Jungyoon Lee, Avishek Joey Bose, Valentin De Bortoli, Arnaud Doucet, Michael Bronstein, Dominique Beaini, Siamak Ravanbakhsh, Kirill Neklyudov, and Alexander Tong. Progressive inference-time annealing of diffusion models for sampling from Boltzmann densities. In *Advances in Neural Information Processing Systems (NeurIPS)*, 2025.
- Michael Samuel Albergo and Eric Vanden-Eijnden. NETS: A non-equilibrium transport sampler. In *International Conference on Machine Learning*, pages 1026–1055. PMLR, 2025. arXiv preprint arXiv:2410.02711.
- Shun-ichi Amari. *Information Geometry and Its Applications*, volume 194 of *Applied Mathematical Sciences*. Springer, 2016.
- Luigi Ambrosio, Nicola Gigli, and Giuseppe Savaré. *Gradient Flows in Metric Spaces and in the Space of Probability Measures*. Lectures in Mathematics ETH Zürich. Birkhäuser, 2 edition, 2008.
- Michael Arbel, Alexander G. D. G. Matthews, and Arnaud Doucet. Annealed flow transport Monte Carlo. In *International Conference on Machine Learning (ICML)*, 2021.
- Dominique Bakry, Ivan Gentil, and Michel Ledoux. *Analysis and Geometry of Markov Diffusion Operators*, volume 348 of *Grundlehren der mathematischen Wissenschaften*. Springer, 2014.
- Martin Bauer, Martins Bruveris, and Peter W. Michor. Uniqueness of the Fisher–Rao metric on the space of smooth densities. *Bulletin of the London Mathematical Society*, 48(3):499–506, 2016.
- Julius Berner, Lorenz Richter, and Karen Ullrich. An optimal control perspective on diffusion-based generative modeling. *Transactions on Machine Learning Research*, 2024.
- Omar Chehab, Anna Korba, Austin Stromme, and Adrien Vacher. Provable convergence and limitations of geometric tempering for Langevin dynamics. In *International Conference on Learning Representations*, volume 2025, pages 5451–5493, 2025.

- Wendy D. Cornell, Piotr Cieplak, Christopher I. Bayly, Ian R. Gould, Kenneth M. Merz, David M. Ferguson, David C. Spellmeyer, Thomas Fox, James W. Caldwell, and Peter A. Kollman. A second generation force field for the simulation of proteins, nucleic acids, and organic molecules. *Journal of the American Chemical Society*, 117(19):5179–5197, 1995.
- Pierre Del Moral, Arnaud Doucet, and Ajay Jasra. Sequential Monte Carlo samplers. *Journal of the Royal Statistical Society: Series B*, 68(3):411–436, 2006.
- Arnaud Doucet, Nando De Freitas, and Neil Gordon. An introduction to sequential Monte Carlo methods. In *Sequential Monte Carlo methods in practice*, pages 3–14. Springer, 2001.
- Víctor Elvira, Luca Martino, and Christian P Robert. Rethinking the effective sample size. *International Statistical Review*, 90(3):525–550, 2022.
- Loris Felardos, Jérôme Hénin, and Guillaume Charpiat. Designing losses for data-free training of normalizing flows on Boltzmann distributions. *arXiv preprint arXiv:2301.05475*, 2023.
- Marylou Gabrié, Grant M Rotskoff, and Eric Vanden-Eijnden. Adaptive Monte Carlo augmented with normalizing flows. *Proceedings of the National Academy of Sciences*, 119(10):e2109420119, 2022.
- Charles J. Geyer. Markov chain Monte Carlo maximum likelihood. In *Computing Science and Statistics: Proc. 23rd Symposium on the Interface*, pages 156–163, 1991.
- W. K. Hastings. Monte Carlo sampling methods using Markov chains and their applications. *Biometrika*, 57(1):97–109, 1970.
- Aaron Havens, Benjamin Kurt Miller, Bing Yan, Carles Domingo-Enrich, Anuroop Sriram, Brandon Wood, Daniel Levine, Bin Hu, Brandon Amos, Brandon Karrer, Xiang Fu, Guan-Hong Liu, and Ricky T. Q. Chen. Adjoint sampling: highly scalable diffusion samplers via adjoint matching. In *International Conference on Machine Learning (ICML)*, 2025.
- Jiajun He, Yuanqi Du, Francisco Vargas, Dinghuai Zhang, Shreyas Padhy, RuiKang OuYang, Carla Gomes, and José Miguel Hernández-Lobato. No trick, no treat: pursuits and challenges towards simulation-free training of neural samplers. In *ICLR 2025 Workshop on Frontiers in Probabilistic Inference*, 2025.
- Koji Hukushima and Koji Nemoto. Exchange Monte Carlo method and application to spin glass simulations. *Journal of the Physical Society of Japan*, 65(6):1604–1608, 1996.
- Richard Jordan, David Kinderlehrer, and Felix Otto. The variational formulation of the Fokker–Planck equation. *SIAM Journal on Mathematical Analysis*, 29(1):1–17, 1998.
- Durk P Kingma and Prafulla Dhariwal. Glow: Generative flow with invertible 1x1 convolutions. *Advances in neural information processing systems*, 31, 2018.
- Leon Klein and Frank Noé. Transferable Boltzmann generators. In *Advances in Neural Information Processing Systems (NeurIPS)*, 2024.
- Leon Klein, Andreas Krämer, and Frank Noé. Equivariant flow matching. In *Advances in Neural Information Processing Systems (NeurIPS)*, 2023.
- Jonas Köhler, Leon Klein, and Frank Noé. Equivariant flows: exact likelihood generative learning for symmetric densities. In *International Conference on Machine Learning (ICML)*, 2020.
- Jun S Liu. *Monte Carlo strategies in scientific computing*, volume 10. Springer, 2001.
- Yulong Lu, Jianfeng Lu, and James Nolen. Accelerating Langevin sampling with birth–death. *arXiv preprint arXiv:1905.09863*, 2019.
- Alex Matthews, Michael Arbel, Danilo J. Rezende, and Arnaud Doucet. Continual repeated annealed flow transport Monte Carlo. In *ICML*, 2022.

- Nicholas Metropolis, Arianna W. Rosenbluth, Marshall N. Rosenbluth, Augusta H. Teller, and Edward Teller. Equation of state calculations by fast computing machines. *The Journal of Chemical Physics*, 21(6):1087–1092, 1953.
- Laurence Illing Midgley, Vincent Stimper, Gregor NC Simm, Bernhard Schölkopf, and José Miguel Hernández-Lobato. Flow annealed importance sampling bootstrap. In *NeurIPS 2022 AI for Science: Progress and Promises*.
- Muhammad Ferjad Naeem, Seong Joon Oh, Youngjung Uh, Yunjey Choi, and Jaejun Yoo. Reliable fidelity and diversity metrics for generative models. In *International conference on machine learning*, pages 7176–7185. PMLR, 2020.
- Christian A. Naesseth, Fredrik Lindsten, and David M. Blei. Markovian score climbing: variational inference with $\text{KL}(p||q)$. In *Advances in Neural Information Processing Systems (NeurIPS)*, 2020.
- Radford M. Neal. Annealed importance sampling. *Statistics and Computing*, 11(2):125–139, 2001.
- Frank Noé, Simon Olsson, Jonas Köhler, and Hao Wu. Boltzmann generators: Sampling equilibrium states of many-body systems with deep learning. *Science*, 365(6457):eaaw1147, 2019.
- Nikolas Nüsken and Lorenz Richter. Solving high-dimensional Hamilton–Jacobi–Bellman PDEs using neural networks: perspectives from controlled diffusions and measures on path space. *Partial Differential Equations and Applications*, 2:48, 2021.
- Felix Otto. The geometry of dissipative evolution equations: the porous medium equation. *Communications in Partial Differential Equations*, 26(1–2):101–174, 2001.
- Felix Otto and Cédric Villani. Generalization of an inequality by Talagrand and links with the logarithmic Sobolev inequality. *Journal of Functional Analysis*, 173(2):361–400, 2000.
- George Papamakarios, Theo Pavlakou, and Iain Murray. Masked autoregressive flow for density estimation. *Advances in neural information processing systems*, 30, 2017.
- Danyal Rehman, Oscar Davis, Jiarui Lu, Jian Tang, Michael Bronstein, Yoshua Bengio, Alexander Tong, and Avishek Joey Bose. Efficient regression-based training of normalizing flows for Boltzmann generators. In *International Conference on Learning Representations (ICLR)*, 2026.
- Danilo Rezende and Shakir Mohamed. Variational inference with normalizing flows. In *International conference on machine learning*, pages 1530–1538. PMLR, 2015.
- Lorenz Richter and Julius Berner. Improved sampling via learned diffusions. In *International Conference on Learning Representations (ICLR)*, 2024.
- Henrik Schopmans and Pascal Friederich. Temperature-annealed Boltzmann generators. In *International Conference on Machine Learning (ICML)*, 2025. arXiv:2501.19077.
- Henrik Schopmans, Annika von Klitzing, and Pascal Friederich. Efficient training of Boltzmann generators using off-policy log-dispersion regularization. *arXiv preprint arXiv:2602.03729*, 2026.
- John Skilling. Nested sampling for general Bayesian computation. *Bayesian Analysis*, 1(4):833–859, 2006.
- Nikola Surjanovic, Saifuddin Syed, Alexandre Bouchard-Côté, and Trevor Campbell. Parallel tempering with a variational reference. In *Advances in Neural Information Processing Systems (NeurIPS)*, 2022.
- Robert H. Swendsen and Jian-Sheng Wang. Replica Monte Carlo simulation of spin-glasses. *Physical Review Letters*, 57(21):2607–2609, 1986.
- Saifuddin Syed, Alexandre Bouchard-Côté, George Deligiannidis, and Arnaud Doucet. Non-reversible parallel tempering: a scalable highly parallel MCMC scheme. *Journal of the Royal Statistical Society Series B*, 84(2):321–350, 2022.

- Charlie B. Tan, Avishek Joey Bose, Chen Lin, Leon Klein, Michael M. Bronstein, and Alexander Tong. Scalable equilibrium sampling with sequential Boltzmann generators. In *International Conference on Machine Learning (ICML)*, 2025.
- Lorenz Vaitl, Kim A. Nicoli, Shinichi Nakajima, and Pan Kessel. Gradients should stay on path: better estimators of the reverse- and forward-KL divergence for normalizing flows. *Machine Learning: Science and Technology*, 2022.
- Francisco Vargas, Will Grathwohl, and Arnaud Doucet. Denoising diffusion samplers. In *International Conference on Learning Representations*, 2023.
- Cédric Villani. *Optimal Transport: Old and New*, volume 338 of *Grundlehren der mathematischen Wissenschaften*. Springer, 2009.
- Annika von Klitzing, Denis Blessing, Henrik Schopmans, Pascal Friederich, and Gerhard Neumann. Learning Boltzmann generators via constrained mass transport. In *International Conference on Learning Representations (ICLR)*, 2026.
- Martin J. Wainwright and Michael I. Jordan. Graphical models, exponential families, and variational inference. *Foundations and Trends in Machine Learning*, 1(1–2):1–305, 2008.
- Peter Wirnsberger, Andrew J. Ballard, George Papamakarios, Stuart Abercrombie, Sébastien Racanière, Alexander Pritzel, Danilo Jimenez Rezende, and Charles Blundell. Targeted free energy estimation via learned mappings. *The Journal of Chemical Physics*, 153(14), 2020.
- Dongyeop Woo and Sungsoo Ahn. Iterated energy-based flow matching for sampling from Boltzmann densities. In *NeurIPS*, 2024. arXiv:2408.16249.
- Dawn B. Woodard, Scott C. Schmidler, and Mark Huber. Conditions for rapid mixing of parallel and simulated tempering on multimodal distributions. *The Annals of Applied Probability*, 19(2):617–640, 2009.
- Hao Wu, Jonas Köhler, and Frank Noé. Stochastic normalizing flows. *Advances in neural information processing systems*, 33:5933–5944, 2020.
- Leo Zhang, Peter Potapchik, Jiajun He, Yuanqi Du, Arnaud Doucet, Francisco Vargas, Hai-Dang Dau, and Saifuddin Syed. Accelerated parallel tempering via neural transports. *arXiv preprint arXiv:2502.10328*, 2025.
- Qinsheng Zhang and Yongxin Chen. Path integral sampler: A stochastic control approach for sampling. In *International Conference on Learning Representations*, 2022.