

Hebbian Learning

Neuroinformatics Tutorial 5

Duc Duy Pham¹

¹Intelligent Systems, Faculty of Engineering,
University of Duisburg-Essen, Germany

Content

- Revision: Mahalanobis Classifier
- Revision: Lecture
- Hebbian Learning

Content

- Revision: Mahalanobis Classifier
- Revision: Lecture
- Hebbian Learning

Revision: Mahalanobis Distance

- What is the main idea of the Mahalanobis distance?

Revision: Mahalanobis Distance

- What is the main idea of the Mahalanobis distance?
 1. Measure Manhattan distance in uncorrelated space
 2. Use Chomsky matrix to transform into uncorrelated space
 3. Use square root of Manhattan distance in uncorrelated space
 4. Measure euclidean distance in linearly uncorrelated space

Revision: Mahalanobis Distance

- What is the main idea of the Mahalanobis distance?
 1. Measure Manhattan distance in uncorrelated space
 2. Use Chomsky matrix to transform into uncorrelated space
 3. Use square root of Manhattan distance in uncorrelated space
 4. Measure euclidean distance in linearly uncorrelated space

A: 2,3

B: 1,2

C: 2,4

D: 4

Revision: Mahalanobis Distance

- What is the main idea of the Mahalanobis distance?
 1. Measure Manhattan distance in uncorrelated space
 2. Use Chomsky matrix to transform into uncorrelated space
 3. Use square root of Manhattan distance in uncorrelated space
 4. Measure euclidean distance in linearly uncorrelated space

A: 2,3



B: 1,2

C: 2,4

D: 4

Inverse Cholesky Transform

$$L := U^T$$



Revision: Mahalanobis Distance

- What statement regarding the Mahalanobis distance is true?

Revision: Mahalanobis Distance

- What statement regarding the Mahalanobis distance is true?
 1. It is the same as the likelihood
 2. It represents a probability distribution
 3. It is not the same as the likelihood
 4. It can be calculated for any data set

A: 1

B: 2

C: 3

D: 4

Revision: Mahalanobis Distance

- What statement regarding the Mahalanobis distance is true?
 1. It is the same as the likelihood
 2. It represents a probability distribution
 3. It is not the same as the likelihood
 4. It can be calculated for any data set

A: 1

B: 2

C: 3

D: 4

Revision: Mahalanobis Distance

- What statement regarding the Mahalanobis distance is true?
 - It is the same as the likelihood
 - It represents a probability distribution
 - It is not the same as the likelihood
 - It can be calculated for any data set

A: 1

B: 2

C: 3

D: 4

$$d(x, \mu) := \sqrt{(x - \mu)^T \Sigma^{-1} (x - \mu)}$$

Revision: Mahalanobis Distance

- What do you need to calculate the Mahalanobis distance in practice?

Revision: Mahalanobis Distance

- What do you need to calculate the Mahalanobis distance in practice?
 1. Chomsky Matrix
 2. Cholesky Matrix
 3. Class mean(s)
 4. Covariance Matrix

Revision: Mahalanobis Distance

- What do you need to calculate the Mahalanobis distance in practice?
 1. Chomsky Matrix
 2. Cholesky Matrix
 3. Class mean(s)
 4. Covariance Matrix

A: all

B: 2,3,4

C: 3,4

D: 4

Revision: Mahalanobis Distance

- What do you need to calculate the Mahalanobis distance in practice?
 - Chomsky Matrix
 - Cholesky Matrix
 - Class mean(s)
 - Covariance Matrix

A: all

B: 2,3,4

C: 3,4

D: 4

$$d(x, \mu) := \sqrt{(x - \mu)^T \Sigma^{-1} (x - \mu)}$$

Revision: Mahalanobis Distance

- How can you use the Mahalanobis distance for classification?

Revision: Mahalanobis Distance

- How can you use the Mahalanobis distance for classification?
 1. Estimate something similar to feature likelihood
 2. Incorporation to McCulloch Pitts Neuron
 3. Choose class with least Mahalanobis distance to class mean
 4. Maximum A-Posteriori Classification

Revision: Mahalanobis Distance

- How can you use the Mahalanobis distance for classification?
 1. Estimate something similar to feature likelihood
 2. Incorporation to McCulloch Pitts Neuron
 3. Choose class with least Mahalanobis distance to class mean
 4. Maximum A-Posteriori Classification

A: 1,3

B: 3

C: 2,3,4

D: 3,4

Revision: Mahalanobis Distance

- How can you use the Mahalanobis distance for classification?
 1. Estimate something similar to feature likelihood
 2. Incorporation to McCulloch Pitts Neuron
 3. Choose class with least Mahalanobis distance to class mean
 4. Maximum A-Posteriori Classification

A: 1,3

B: 3

C: 2,3,4

D: 3,4

Revision: Mahalanobis Distance

- How do you use the Mahalanobis distance for classification (step by step)?

Revision: Mahalanobis Distance

- „Training: “
 - Calculate mean for each class

Revision: Mahalanobis Distance

- „Training: “
 - Calculate mean for each class
 - Calculate covariance matrix for each class

$$\text{Cov}(\mathbf{X}_k) = \frac{1}{N_k} \sum_{i=1}^{N_k} (\mathbf{x}_i - \mu_k)(\mathbf{x}_i - \mu_k)^T$$

Revision: Mahalanobis Distance

- „Training: “
 - Calculate mean for each class
 - Calculate covariance matrix for each class

$$\text{Cov}(\mathbf{X}_k) = \frac{1}{N_k} \sum_{i=1}^{N_k} (\mathbf{x}_i - \mu_k)(\mathbf{x}_i - \mu_k)^T$$

- Invert covariance matrices

Revision: Mahalanobis Distance

- „Training: “
 - Calculate mean for each class
 - Calculate covariance matrix for each class

$$\text{Cov}(\mathbf{X}_k) = \frac{1}{N_k} \sum_{i=1}^{N_k} (\mathbf{x}_i - \mu_k)(\mathbf{x}_i - \mu_k)^T$$

- Invert covariance matrices
- Inference:
 - Calculate Mahalanobis distance to each class mean
 - Choose class with least distance

Mahalanobis Classifier: Jupyter

Content

- Revision: Mahalanobis Classifier
- **Revision: Lecture**
- Hebbian Learning

Revision: Lecture

- When are two sets $\mathcal{P}, \mathcal{N} \subset \mathbb{R}^n$ linearly separable?

Revision: Lecture

- When are two sets $\mathcal{P}, \mathcal{N} \subset \mathbb{R}^n$ linearly separable?
 1. If you can fit a manifold in between both sets
 2. If you can fit a line in between both sets
 3. If you can fit a hypeplane in between both sets
 4. If you one set contains the other

Revision: Lecture

- When are two sets $\mathcal{P}, \mathcal{N} \subset \mathbb{R}^n$ linearly separable?
 1. If you can fit a manifold in between both sets
 2. If you can fit a line in between both sets
 3. If you can fit a hypeplane in between both sets
 4. If you one set contains the other

A: 1,2,3

B: 2,3

C: 3

D: 2,4

Revision: Lecture

- When are two sets $\mathcal{P}, \mathcal{N} \subset \mathbb{R}^n$ linearly separable?
 1. If you can fit a manifold in between both sets
 2. If you can fit a line in between both sets
 3. If you can fit a hypeplane in between both sets
 4. If you one set contains the other

A: 1,2,3


B: 2,3

C: 3

D: 2,4


Revision: Lecture

- When are two sets $\mathcal{P}, \mathcal{N} \subset \mathbb{R}^n$ linearly separable?




Revision: Lecture

- When are two sets $\mathcal{P}, \mathcal{N} \subset \mathbb{R}^n$ linearly separable?



Revision: Lecture

- When are two sets $\mathcal{P}, \mathcal{N} \subset \mathbb{R}^n$ linearly separable?



Revision: Lecture

- Formal definition of (absolute) linear separability of two sets $\mathcal{P}, \mathcal{N} \subset \mathbb{R}^n$ (must know!):

Revision: Lecture

- Formal definition of (absolute) linear separability of two sets $\mathcal{P}, \mathcal{N} \subset \mathbb{R}^n$ (must know!):
- If there exist $(w_1, \dots, w_n)^T \in \mathbb{R}^n, \Theta \in \mathbb{R}$

Revision: Lecture

- Formal definition of (absolute) linear separability of two sets $\mathcal{P}, \mathcal{N} \subset \mathbb{R}^n$ (must know!):

- If there exist $(w_1, \dots, w_n)^T \in \mathbb{R}^n, \Theta \in \mathbb{R}$

such that $\sum_{i=1}^n w_i x_i \geq (>) \Theta \quad \forall (x_1, \dots, x_n)^T \in \mathcal{P}$

Revision: Lecture

- Formal definition of (absolute) linear separability of two sets $\mathcal{P}, \mathcal{N} \subset \mathbb{R}^n$ (must know!):


- If there exist $(w_1, \dots, w_n)^T \in \mathbb{R}^n, \Theta \in \mathbb{R}$

such that $\sum_{i=1}^n w_i x_i \geq (>) \Theta \quad \forall (x_1, \dots, x_n)^T \in \mathcal{P}$

and $\sum_{i=1}^n w_i x_i < \Theta \quad \forall (x_1, \dots, x_n)^T \in \mathcal{N}$

Revision: Lecture


- Interpretation of formulas



Revision: Lecture

- Interpretation of formulas

$$H_{w,\Theta} := \{x \in \mathbb{R}^n \mid w^T x = \Theta\}, w \in \mathbb{R}^n, \Theta \in \mathbb{R}$$




Revision: Lecture

- Interpretation of formulas

$$H_{w,\Theta} := \{x \in \mathbb{R}^n \mid w^T x = \Theta\}, w \in \mathbb{R}^n, \Theta \in \mathbb{R}$$

$$\Rightarrow \Theta = \|w\| \cdot \|x\| \cdot \cos(\varphi)$$




Revision: Lecture

- Interpretation of formulas

$$H_{w,\Theta} := \{x \in \mathbb{R}^n | w^T x = \Theta\}, w \in \mathbb{R}^n, \Theta \in \mathbb{R}$$


$$\Rightarrow \Theta = \|w\| \cdot \|x\| \cdot \cos(\varphi)$$

$$\Rightarrow \|x\| \cdot \cos(\varphi) = \frac{\Theta}{\|w\|}$$



Revision: Lecture

- Interpretation of formulas




$$H_{w, \Theta} := \{x \in \mathbb{R}^n | w^T x = \Theta\}, w \in \mathbb{R}^n, \Theta \in \mathbb{R}$$

$$\Rightarrow \Theta = \|w\| \cdot \|x\| \cdot \cos(\varphi)$$

$$\Rightarrow \|x\| \cdot \cos(\varphi) = \frac{\Theta}{\|w\|}$$

Revision: Lecture

- Interpretation of formulas




$$H_{w, \Theta} := \{x \in \mathbb{R}^n | w^T x = \Theta\}, w \in \mathbb{R}^n, \Theta \in \mathbb{R}$$

$$\Rightarrow \Theta = \|w\| \cdot \|x\| \cdot \cos(\varphi)$$

$$\Rightarrow \|x\| \cdot \cos(\varphi) = \frac{\Theta}{\|w\|}$$


Revision: Lecture

- Interpretation of formulas



Revision: Lecture

- Interpretation of formulas




$$H_{w, \Theta} := \{x \in \mathbb{R}^n \mid w^T x = \Theta\}, w \in \mathbb{R}^n, \Theta \in \mathbb{R}$$

$$\Rightarrow \Theta = \|w\| \cdot \|x\| \cdot \cos(\varphi)$$

$$\Rightarrow \|x\| \cdot \cos(\varphi) = \frac{\Theta}{\|w\|}$$

Revision: Lecture

- Interpretation of formulas




$$H_{w,\Theta} := \{x \in \mathbb{R}^n \mid w^T x = \Theta\}, w \in \mathbb{R}^n, \Theta \in \mathbb{R}$$

$$\Rightarrow \Theta = \|w\| \cdot \|x\| \cdot \cos(\varphi)$$

$$\Rightarrow \|x\| \cdot \cos(\varphi) = \frac{\Theta}{\|w\|}$$

Revision: Lecture

- Interpretation of formulas




$$H_{w,\Theta} := \{x \in \mathbb{R}^n \mid w^T x = \Theta\}, w \in \mathbb{R}^n, \Theta \in \mathbb{R}$$

$$\Rightarrow \Theta = \|w\| \cdot \|x\| \cdot \cos(\varphi)$$

$$\Rightarrow \|x\| \cdot \cos(\varphi) = \frac{\Theta}{\|w\|}$$

Revision: Lecture

- Interpretation of formulas




$$H_{w, \Theta} := \{x \in \mathbb{R}^n | w^T x = \Theta\}, w \in \mathbb{R}^n, \Theta \in \mathbb{R}$$

$$\Rightarrow \Theta = ||w|| \cdot ||x|| \cdot \cos(\varphi)$$

$$\Rightarrow ||x|| \cdot \cos(\varphi) = \frac{\Theta}{||w||}$$

Revision: Lecture

- Interpretation of formulas




$$H_{w,\Theta} := \{x \in \mathbb{R}^n | w^T x = \Theta\}, w \in \mathbb{R}^n, \Theta \in \mathbb{R}$$

$$\Rightarrow \Theta = \|w\| \cdot \|x\| \cdot \cos(\varphi)$$

$$\Rightarrow \|x\| \cdot \cos(\varphi) = \frac{\Theta}{\|w\|}$$

Revision: Lecture

- Interpretation of formulas




$$H_{w, \Theta} := \{x \in \mathbb{R}^n \mid w^T x = \Theta\}, w \in \mathbb{R}^n, \Theta \in \mathbb{R}$$

$$\Rightarrow \Theta = \|w\| \cdot \|x\| \cdot \cos(\varphi)$$

$$\Rightarrow \|x\| \cdot \cos(\varphi) = \frac{\Theta}{\|w\|}$$

Revision: Lecture

- Interpretation of formulas




$$H_{w,\Theta} := \{x \in \mathbb{R}^n | w^T x = \Theta\}, w \in \mathbb{R}^n, \Theta \in \mathbb{R}$$

$$\Rightarrow \Theta = ||w|| \cdot ||x|| \cdot \cos(\varphi)$$

$$\Rightarrow ||x|| \cdot \cos(\varphi) = \frac{\Theta}{||w||}$$

Revision: Lecture

- Interpretation of formulas




$$H_{w, \Theta} := \{x \in \mathbb{R}^n | w^T x = \Theta\}, w \in \mathbb{R}^n, \Theta \in \mathbb{R}$$

$$\Rightarrow \Theta = ||w|| \cdot ||x|| \cdot \cos(\varphi)$$

$$\Rightarrow ||x|| \cdot \cos(\varphi) = \frac{\Theta}{||w||}$$

Revision: Lecture

- Interpretation of formulas



$$H_{w,\Theta} := \{x \in \mathbb{R}^n | w^T x = \Theta\}, w \in \mathbb{R}^n, \Theta \in \mathbb{R}$$


$$\Rightarrow \Theta = ||w|| \cdot ||x|| \cdot \cos(\varphi)$$

$$\Rightarrow ||x|| \cdot \cos(\varphi) = \frac{\Theta}{||w||}$$

$$\Rightarrow w^T x_1 = ||w|| ||x_1|| \cos(\varphi_1) < \Theta$$

Revision: Lecture

- Interpretation of formulas




$$H_{w,\Theta} := \{x \in \mathbb{R}^n | w^T x = \Theta\}, w \in \mathbb{R}^n, \Theta \in \mathbb{R}$$

$$\Rightarrow \Theta = \|w\| \cdot \|x\| \cdot \cos(\varphi)$$

$$\Rightarrow \|x\| \cdot \cos(\varphi) = \frac{\Theta}{\|w\|}$$

Revision: Lecture

- Interpretation of formulas




$$H_{w,\Theta} := \{x \in \mathbb{R}^n | w^T x = \Theta\}, w \in \mathbb{R}^n, \Theta \in \mathbb{R}$$

$$\Rightarrow \Theta = ||w|| \cdot ||x|| \cdot \cos(\varphi)$$

$$\Rightarrow ||x|| \cdot \cos(\varphi) = \frac{\Theta}{||w||}$$

Revision: Lecture

- Interpretation of formulas




$$H_{w, \Theta} := \{x \in \mathbb{R}^n \mid w^T x = \Theta\}, w \in \mathbb{R}^n, \Theta \in \mathbb{R}$$

$$\Rightarrow \Theta = \|w\| \cdot \|x\| \cdot \cos(\varphi)$$

$$\Rightarrow \|x\| \cdot \cos(\varphi) = \frac{\Theta}{\|w\|}$$

Revision: Lecture

- Interpretation of formulas




$$H_{w,\Theta} := \{x \in \mathbb{R}^n | w^T x = \Theta\}, w \in \mathbb{R}^n, \Theta \in \mathbb{R}$$

$$\Rightarrow \Theta = \|w\| \cdot \|x\| \cdot \cos(\varphi)$$

$$\Rightarrow \|x\| \cdot \cos(\varphi) = \frac{\Theta}{\|w\|}$$

Revision: Lecture

- Interpretation of formulas



$$H_{w, \Theta} := \{x \in \mathbb{R}^n | w^T x = \Theta\}, w \in \mathbb{R}^n, \Theta \in \mathbb{R}$$

$$\Rightarrow \Theta = ||w|| \cdot ||x|| \cdot \cos(\varphi)$$

$$\Rightarrow ||x|| \cdot \cos(\varphi) = \frac{\Theta}{||w||}$$

$$\Rightarrow w^T x_2 = ||w|| ||x_2|| \cos(\varphi_2) > \Theta$$

Revision: Lecture

- Formal definition of (absolute) linear separability of two sets $\mathcal{P}, \mathcal{N} \subset \mathbb{R}^n$ (must know!):

- If there exist $(w_1, \dots, w_n)^T \in \mathbb{R}^n, \Theta \in \mathbb{R}$

such that $\sum_{i=1}^n w_i x_i \geq (>) \Theta \quad \forall (x_1, \dots, x_n)^T \in \mathcal{P}$

and $\sum_{i=1}^n w_i x_i < \Theta \quad \forall (x_1, \dots, x_n)^T \in \mathcal{N}$

Revision: Lecture

- What is the formal definition of a Rosenblatt Perceptron?

Revision: Lecture

- What is the formal definition of a Rosenblatt Perceptron?

1. Real input vector

$$(x_1, \dots, x_n)^T \in \mathbb{R}^n$$

Revision: Lecture

- What is the formal definition of a Rosenblatt Perceptron?

1. Real input vector
2. Real weights

$$(x_1, \dots, x_n)^T \in \mathbb{R}^n$$

$$(w_1, \dots, w_n)^T \in \mathbb{R}^n$$

Revision: Lecture

- What is the formal definition of a Rosenblatt Perceptron?

1. Real input vector
2. Real weights
3. Propagation function is linear associator

$$(x_1, \dots, x_n)^T \in \mathbb{R}^n$$

$$(w_1, \dots, w_n)^T \in \mathbb{R}^n$$

$$\sum_{i=1}^n w_i x_i$$

Revision: Lecture

- What is the formal definition of a Rosenblatt Perceptron?

1. Real input vector
2. Real weights
3. Propagation function is linear associator
4. Real Threshold

$$(x_1, \dots, x_n)^T \in \mathbb{R}^n$$

$$(w_1, \dots, w_n)^T \in \mathbb{R}^n$$

$$\sum_{i=1}^n w_i x_i$$

$$\Theta \in \mathbb{R}$$

Revision: Lecture

- What is the formal definition of a Rosenblatt Perceptron?

1. Real input vector
2. Real weights
3. Propagation function is linear associator
4. Real Threshold
5. Activation function is step function

$$(x_1, \dots, x_n)^T \in \mathbb{R}^n$$

$$(w_1, \dots, w_n)^T \in \mathbb{R}^n$$

$$\sum_{i=1}^n w_i x_i$$

$$\Theta \in \mathbb{R}$$

$$y := f_a := \begin{cases} 1 & \text{if } \sum_{i=1}^n w_i x_i \geq \Theta \\ 0 & \text{else} \end{cases}$$

Revision: Lecture

- Reformulation of step function

$$x_0 = 1$$

$$w_0 = -\Theta$$

Revision: Lecture

- Reformulation of step function

$$x_0 = 1$$

$$w_0 = -\Theta$$

$$y := f_a := \begin{cases} 1 & \text{if } \sum_{i=0}^n w_i x_i \geq 0 \\ 0 & \text{else} \end{cases}$$

Revision: Lecture


- Reformulation of step function

$$x_0 = 1$$

$$w_0 = -\Theta$$

$$y := f_a := \begin{cases} 1 & \text{if } \sum_{i=0}^n w_i x_i \geq 0 \\ -1 & \text{else} \end{cases}$$

Scheme of Artificial Neuron



$f_p|f_a$ wird oft weggelassen, wenn aus dem Zusammenhang klar.

Revision: Lecture

- Which statements regarding Rosenblatt Perceptron and McCulloch Pitts Neuron are true?

Revision: Lecture

- Which statements regarding Rosenblatt Perceptron and McCulloch Pitts Neuron are true?
 1. The McCulloch Pitts Neuron can process real valued input
 2. The Rosenblatt Perceptron can process real valued input
 3. Both neuron models have inhibiting and excitatory edges
 4. The Rosenblatt Perceptron can model McCulloch Pitts Neurons

Revision: Lecture

- Which statements regarding Rosenblatt Perceptron and McCulloch Pitts Neuron are true?
 1. The McCulloch Pitts Neuron can process real valued input
 2. The Rosenblatt Perceptron can process real valued input
 3. Both neuron models have inhibiting and excitatory edges
 4. The Rosenblatt Perceptron can model McCulloch Pitts Neurons

A: all

B: 1,2,4

C: 2

D: 2,4

Revision: Lecture

- Which statements regarding Rosenblatt Perceptron and McCulloch Pitts Neuron are true?
 1. The McCulloch Pitts Neuron can process real valued input
 2. The Rosenblatt Perceptron can process real valued input
 3. Both neuron models have inhibiting and excitatory edges
 4. The Rosenblatt Perceptron can model McCulloch Pitts Neurons

A: all


B: 1,2,4

C: 2

D: 2,4


Revision: Lecture

- How can you use a Rosenblatt Perceptron to represent a McCulloch Pitts Neuron?





Revision: Lecture

- How can you use a Rosenblatt Perceptron to represent a McCulloch Pitts Neuron?





Revision: Lecture

- How can you use a Rosenblatt Perceptron to represent a McCulloch Pitts Neuron?





Revision: Lecture

- How can you use a Rosenblatt Perceptron to represent a McCulloch Pitts Neuron?





Revision: Lecture

- How can you use a Rosenblatt Perceptron to represent a McCulloch Pitts Neuron?



Revision: Lecture

- How can you use a Rosenblatt Perceptron to represent a McCulloch Pitts Neuron?



Content

- Revision: Mahalanobis Classifier
- Revision: Lecture
- **Hebbian Learning**

Hebbian Learning

- Postulated by Donald Olding Hebb in 1949
- Foundation of many learning rules


Hebbian Learning

- Postulated by Donald Olding Hebb in 1949
- Foundation of many learning rules
- Idea:
 - If neuron j receives a signal from neuron i and both neurons are strongly activated, then the connection of j and i should be strong!

Hebbian Learning

- Postulated by Donald Olding Hebb in 1949
- Foundation of many learning rules
- Idea:
 - If neuron j receives a signal from neuron i and both neurons are strongly activated, then the connection of j and i should be strong!
 - In artificial neural networks the strength of the connection is usually represented by the edge weight

Calculation of propagated value



$$h_1 = \sum_{i=0}^3 w_{i1} x_i$$

$$h_2 = \sum_{i=0}^3 w_{i2} x_i$$

$$x = \begin{bmatrix} x_0 \\ x_1 \\ x_2 \\ x_3 \end{bmatrix} \quad W = \begin{bmatrix} w_{01} & w_{02} \\ w_{11} & w_{12} \\ w_{21} & w_{22} \\ w_{31} & w_{32} \end{bmatrix}$$

$$\begin{bmatrix} h_1 \\ h_2 \end{bmatrix} = W^T \cdot x$$

Hebbian Learning Rule

- Given:
 - Outputs of previous neurons (can also be input neurons)

Hebbian Learning Rule

- Given:
 - Outputs of previous neurons (can also be input neurons)
 - In case of scheme : input neurons, i.e:

$$x_0, x_1, x_2, \dots, x_n \in \{-1, 1\}$$

Hebbian Learning Rule

- Given:
 - Outputs of previous neurons (can also be input neurons)
 - In case of scheme : input neurons, i.e:
$$x_0, x_1, x_2, \dots x_n \in \{-1, 1\}$$
 - Output of current neurons
 - In case of scheme: $h_1, h_2 \in \{-1, 1\}$

Hebbian Learning Rule

- Given:
 - Outputs of previous neurons (can also be input neurons)
 - In case of scheme : input neurons, i.e:

$$x_0, x_1, x_2, \dots x_n \in \{-1, 1\}$$
 - Output of current neurons
 - In case of scheme: $h_1, h_2 \in \{-1, 1\}$
- Weights in between all neurons

Hebbian Learning Rule

- Given:
 - Outputs of previous neurons (can also be input neurons)
 - In case of scheme : input neurons, i.e:
 $x_0, x_1, x_2, \dots x_n \in \{-1, 1\}$
 - Output of current neurons
 - In case of scheme: $h_1, h_2 \in \{-1, 1\}$
 - Weights in between all neurons
 - Learning rate α

Hebbian Learning Rule

- Given:
 - Outputs of previous neurons (can also be input neurons)
 - In case of scheme : input neurons, i.e:

$$x_0, x_1, x_2, \dots x_n \in \{-1, 1\}$$
 - Output of current neurons
 - In case of scheme: $h_1, h_2 \in \{-1, 1\}$
 - Weights in between all neurons
 - Learning rate α
- Learning Rule:
 - Update weights by comparing similarity of outputs

Hebbian Learning Rule

- Given:
 - Outputs of previous neurons (can also be input neurons)
 - In case of scheme : input neurons, i.e:

$$x_0, x_1, x_2, \dots, x_n \in \{-1, 1\}$$
 - Output of current neurons
 - In case of scheme: $h_1, h_2 \in \{-1, 1\}$
 - Weights in between all neurons
 - Learning rate α
- Learning Rule:
 - Update weights by comparing similarity of outputs
 - $\Delta w_{i,j} := \alpha \cdot x_i \cdot h_j$

Hebbian Learning Rule

- Given:
 - Outputs of previous neurons (can also be input neurons)
 - In case of scheme : input neurons, i.e:

$$x_0, x_1, x_2, \dots x_n \in \{-1, 1\}$$
 - Output of current neurons
 - In case of scheme: $h_1, h_2 \in \{-1, 1\}$
 - Weights in between all neurons
 - Learning rate α
- Learning Rule:
 - Update weights by comparing similarity of outputs
 - $\Delta w_{i,j} := \alpha \cdot x_i \cdot h_j$
 - $w_{i,j} \leftarrow w_{i,j} + \Delta w_{i,j}$

