

Smooth Manifolds and Lie Groups

Directed Reading Program Fall 2023

Estella Xu

Colin Fan

December 1, 2023

Northwestern University Dept. of Mathematics

1. An Introduction to Manifolds
2. Lie Groups
3. A Non-Example: S^2
4. The Hairy Ball Theorem and Parallelizability

Definition (Manifold). Let M be a topological space. M is a manifold if it is

- (1) Hausdorff,
- (2) second-countable, and
- (3) locally Euclidean.

Today, we'll only focus on (3).

Topological Spaces


Being in a topological space allows us to define continuity of functions.

Locally Euclidean

Definition (Locally Euclidean). Given any point on a manifold M of dimension n , we can find an open set around the point and a homeomorphism (a continuous bijective map with a continuous inverse) between that open set and \mathbb{R}^n .

Intuitively...

A manifold is an object that looks like \mathbb{R}^n in a small area around each point.



Smooth Manifolds

We will look at **smooth manifolds**, which will allow us to do calculus on manifolds.

Smooth manifolds “smoothly” look like \mathbb{R}^n locally; instead of finding a continuous map from an open set to \mathbb{R}^n , we find a smooth (infinitely differentiable) one.¹

¹More precisely, given any two charts on M , the transition map is smooth.

Manifold Examples

- \mathbb{R}^n where we take the identity map as our homeomorphism.
- The torus (we'll touch on this later).
- The 2-dimensional sphere, given by

$$S^2 = \{(x, y, z) \in \mathbb{R}^3 : x^2 + y^2 + z^2 = 1\}.$$

It is *embedded* in \mathbb{R}^3 .

Let's provide more details on the sphere.

Example (Sphere)

The sphere is a smooth manifold. Consider the open set $U = \{(x, y, z) : z > 0\}$.

The “Flattening Map”²


$$\phi : U \rightarrow \mathbb{R}^2$$

$$(x, y, z) \mapsto (x, y),$$

with inverse

$$\phi^{-1} : \mathbb{R}^2 \rightarrow U$$

$$(x, y) \mapsto \left(x, y, \sqrt{1 - (x^2 + y^2)} \right).$$




²Another map we can use is stereographic projection.

Nonexample

Consider the x - and y -axes (embedded in \mathbb{R}^2). This is **not** a manifold.

The key reasoning:

- Homeomorphisms on a space are still homeomorphisms on a subspace of the space.
- Continuous maps preserve connectedness (and homeomorphisms preserve the number of connected components).



Definition (Lie Group). G is a Lie group if it is a smooth manifold that is a group such that

- the multiplication map $m : G \times G \rightarrow G$ and
- the inverse map $(\)^{-1} : G \rightarrow G$

are smooth.

Example (\mathbb{R}^n)

Consider \mathbb{R}^n .


- We've seen that \mathbb{R}^n is a smooth manifold.
- \mathbb{R}^n is a group under addition; in particular, addition and negation (its inverse map) are both smooth.

Example (Tori)


As a group, the n -torus is $\mathbb{R}^n / \mathbb{Z}^n$. Consider the group operation

$$(x_1, \dots, x_n) + (y_1, \dots, y_n) = (x_1 + y_1 \bmod 1, \dots, x_n + y_n \bmod 1).$$

The 1-Torus



The 2-Torus



Example (Matrix Groups)

- The general linear group: $GL(n) = \{A \in M(n \times n) : A \text{ is invertible}\}$.
- The rotation group: $SO(3) = \{A \in M(3 \times 3) : A^T A = A A^T = I\}$.
- The special linear group: $SL(2) = \{A \in M(2 \times 2) : \det(A) = 1\}$.

Nonexample (Sphere)

We show that the sphere is not a Lie group.

How can you show something is not a Lie group?

It's tricky...

You'd have to show that no group operation on the space will be smooth.

Instead, let's use a much nicer method.


Vector Fields

We create a **vector field** on a smooth manifold by placing tangent vectors at every point on the manifold.

- If the vectors *continuously vary*, the vector field is continuous.
- If the vectors *smoothly vary*, the vector field is smooth.

Example


$$-y \frac{\partial}{\partial x} + x \frac{\partial}{\partial y}$$



The Hairy Ball Theorem

Theorem (Hairy Ball). Every continuous vector field on a sphere has to vanish somewhere, i.e., there is some point on the sphere whose vector is 0.

The Hairy Ball Theorem



The Tangent Bundle of a Lie Group (Parallelizability)

Theorem. Let G be a Lie group of dimension n . Then,

$$TG \cong G \times \mathbb{R}^n.$$

The Tangent Bundle

The **tangent bundle** of a manifold M (written TM) is, loosely speaking, the collection of all smooth vector fields on M .

Diffeomorphic

Two smooth manifolds are **diffeomorphic** (\cong) if there exists a smooth, bijective map with a smooth inverse between them. We can *identify* the two spaces with each other.

A Contradiction

Suppose the sphere is a Lie group. Then, by the previous theorem,

$$TS^2 \cong S^2 \times \mathbb{R}^2.$$

Then, there would be a vector field consisting of a *nonzero* constant tangent vector at every point on the sphere. $X = \{ (p, (1,1)) : p \in S^2 \}$

However, this is impossible under the Hairy Ball Theorem.

“Parallelizability”

The tangent bundle theorem tells us that Lie groups have a *nonvanishing vector field*. This gives us a way to show if something is *not* a Lie group.

Some Fun Facts

It turns out that...

- S^1 and S^3 are Lie groups.
 - S^1 can be endowed with the operation of angle addition.
 - $S^3 \cong SU(2)$.
- S^7 is parallelizable, i.e.,

$$TS^7 \cong S^7 \times \mathbb{R}^7,$$

but *not* a Lie group (parallelizability is not an “if and only if” relation!).

Conclusion

1. A manifold is locally similar to \mathbb{R}^n .
2. A Lie group is a smooth manifold that is also a topological group.
3. The tangent bundle of a Lie group is trivial.