

Proto-Dilithium as a ZKP

attempt #1: let $R_q = \mathbb{Z}_q[x] / \langle x^n + 1 \rangle$, S_η denote the set of polynomials in R_q whose coefficients are $[-\eta, \eta]$.

KeyGen	Commitment	Response(c)
$\vec{s} \leftarrow \# S_\eta^L$	$\vec{\gamma} \leftarrow \# S_{\gamma_1}^L$	$\vec{z} \leftarrow c \cdot \vec{s} + \vec{\gamma}$
$A \leftarrow \# R_q^{k \times L}$	$\vec{w} \leftarrow A \vec{\gamma}$	return \vec{z}
$\vec{t} \leftarrow A \vec{s}$	return \vec{w}	
$pk \leftarrow (A, \vec{t})$		<u>Verify(\vec{z})</u>
$sk \leftarrow \vec{s}$	<u>Challenge</u>	assert $A \vec{z} == c \cdot \vec{t} + \vec{w}$
return (pk, sk)	$c \leftarrow \text{Ball}(\chi)$	
	return c	

SIS ZKP Attempt #1

for each ZKP we need 3 properties:

Completeness: honest verifier will accept proof from an honest prover

Soundness: dishonest prover cannot make honest verifier accept

(honest verifier) **zero knowledge**: the transcript of the proof does not reveal information about the secret

Attempt #1 is not sound because solving $A \vec{z} = c \cdot \vec{t} + \vec{w}$ is easy where there is no constraint on \vec{z} and $A \in R_q^{k \times L}$ is wide ($L \geq k$).

solution: estimate a bound on \vec{z} then pose a constraint

recall that $\vec{z} \leftarrow c \cdot \vec{s} + \vec{\gamma}$, let $i \in \{1, 2, \dots, L\}$ denote the index of the polynomial and $j \in \{1, 2, \dots, n\}$ denote the index of a coefficient, then we know

$$z_i[j] = (c \cdot s_i)[j] + \gamma_i[j]$$

we already know $-\gamma_1 \leq \gamma_i[j] \leq \gamma_1$, it remains to establish bound on $(c \cdot s_i)[j]$.

$$c = c[0] + c[1]x + \dots + c[n-1]x^{n-1}$$

$$s_i = s_i[0] + s_i[1]x + \dots + s_i[n-1]x^{n-1}$$

$$(c \cdot s_i)[j] = \sum_{\substack{0 \leq a, b < n \\ a+b \equiv j \pmod n}} c[a] s_i[b]$$

* only partly correct since $x^n \equiv -1 \pmod{x^n+1}$
so some coeffs need to mul with -1

for each of j the summation has n terms. Among them exactly $\tilde{\alpha}$ of them are non-zero, and each of $S_i[j] \in [-\eta, \eta]$, so we have that $(C \cdot S_i)[j] \in [-\eta\tilde{\alpha}, \eta\tilde{\alpha}]$, which gives the bound for \tilde{z} .

KeyGen	Commitment	Response(C)
$\vec{s} \xleftarrow{\$} S_n^L$	$\vec{\gamma} \xleftarrow{\$} S_{\delta_1}^L$	$\vec{z} \leftarrow C \cdot \vec{s} + \vec{\gamma}$
$A \xleftarrow{\$} R_q^{k \times L}$	$\vec{w} \leftarrow A \vec{\gamma}$	return \vec{z}
$\vec{t} \leftarrow A \vec{s}$	return \vec{w}	
$pk \leftarrow (A, \vec{t})$		<u>Verify(\vec{z})</u>
$sk \leftarrow \vec{s}$	<u>Challenge</u>	assert $A \vec{z} = C \cdot \vec{t} + \vec{w}$
return (pk, sk)	$c \leftarrow \text{Ball}(\tilde{\alpha})$	and $\ \vec{z}\ _{\infty} \leq \delta_1 + \eta\tilde{\alpha}$
	return c	

SIS ZKP Attempt #2

Attempt #2 is somewhat sound now. This is because A is uniform random, so finding a $\vec{z} \in R_q^L$ such that $A \vec{z} = (\dots)$ under the constraint $\|\vec{z}\|_{\infty} \leq \beta$ is equivalent to solving an instance of the inhomogeneous Module-SIS problem. \Rightarrow still need to tune parameters!

Attempt #2 is not zero-knowledge: certain values of \vec{z} can leak information of the secret key \vec{s} . Consider an extreme example where $z_i[j] = \eta\tilde{\alpha} + \delta_1$, then we know $S_i[j]$ must be $\pm\eta$ where $C[j]$ is ± 1 . Less extreme values give less but still substantial amount of information, "until within a bound, then \vec{z} gives no information".

Recall that $z_i[j] = (C S_i)[j] + \gamma_i[j]$, we will focus on a single coeff since the coeffs of S_i and γ_i are all iid. Without knowing $z_i[j]$, an adversary only knows that $S_i \xleftarrow{\$} R_{[-\eta, \eta]}$ and $\gamma_i[j] \xleftarrow{\$} [-\delta_1, \delta_1]$

if $z_i[j] = \delta_1 + \eta\tilde{\alpha}$, then it must be $(C S_i)[j] = \eta\tilde{\alpha}$ and $\gamma_i[j] = \delta_1$
 if $z_i[j] = \delta_1 + \eta\tilde{\alpha} - 1$, then $(C S_i)[j] = \eta\tilde{\alpha}$ and $\gamma_i[j] = \delta_1 - 1$
 or $(C S_i)[j] = \eta\tilde{\alpha} - 1$ and $\gamma_i[j] = \delta_1$

...
 if $z_i[j] = \delta_1 + \eta\tilde{\alpha} - 2\eta\tilde{\alpha}$ then $(C S_i)[j] = \eta\tilde{\alpha}$ and $\gamma_i[j] = \delta_1 - 2\eta\tilde{\alpha}$

...
 $(C S_i)[j] = -\eta\tilde{\alpha}$ and $\gamma_i[j] = \delta_1$

intuitively speaking, when $z_i[j] > \delta - \eta \tau$, some values for $cs_i[j]$ are impossible because for such value, $y_i[j] = z_i[j] - cs_i[j]$ will be outside the allowed range $[-\delta, \delta]$. On the other hand when $z_i[j] = \delta - \eta \tau$, all values within $[-\eta \tau, \eta \tau]$ are possible for $cs_i[j]$ because all corresponding $y_i[j]$ values fall within the allowed range. The formal notion is expressed as follows:

Lemma if $\|\vec{z}\|_\infty \leq \delta - \eta \tau$, then the distribution of s_i is identical to the distribution of $s_i | C, z_i$

Proof:
$$P[s_i | C, z_i] = \sum_{s'} P[s_i \cap cs_i = s' | C, z_i]$$
$$= \sum_{s'} P[cs_i = s' | C, z_i] \cdot P[s_i | C, z_i, cs_i = s'] \quad \dots (1)$$

observe that:

(a) $P[s_i | C, z_i, cs_i = s'] = P[s_i | C, cs_i = s']$ since z_i does not give extra information when C, s' are already given

(b)
$$P[s' | C, z_i] = \frac{P[s', C, z_i]}{P[C, z_i]} = \frac{P[z_i | s', C] \cdot P[C, s']}{P[C, z_i]}$$
$$= \frac{P[z_i | s', C] \cdot P[s' | C]}{P[z_i | C]} \quad \left. \vphantom{\frac{P[s', C, z_i]}{P[C, z_i]}} \right\} \text{Bayes rule}$$

(c) $P[z_i | s', C] = P[y_i] = \frac{1}{2\delta + 1}$ since $z = s' + y$ and for $z \in [-\delta + \eta \tau, \delta - \eta \tau]$ and $s' \in [-\eta \tau, \eta \tau]$, y is always in the allowed range of values \Rightarrow the main reason that this proof works

(d)
$$P[z_i | C] = \sum_{s'} P[z_i \cap s' | C]$$
$$= \sum_{s'} P[s' | C] \cdot P[z_i | s', C] \quad \left. \vphantom{\sum_{s'} P[s' | C] \cdot P[z_i | s', C]} \right\} \text{because (c)}$$
$$= P[y_i] \cdot \sum_{s'} P[s' | C]$$
$$= P[y_i]$$

Putting (a) ~ (d) together:

$$P[s_i | C, z_i] = \sum_{s'} P[s_i | s', C] \cdot \frac{P[y_i] \cdot P[s' | C]}{P[y_i]}$$
$$= \sum_{s'} P[s_i | s', C] \cdot P[s' | C]$$
$$= \sum_{s'} P[s_i, s' | C]$$
$$= P[s_i | C] \quad \left. \vphantom{\sum_{s'} P[s_i, s' | C]} \right\} s_i \perp C \text{ are independent}$$
$$= P[s_i]$$

□

now we have a complete, sound, and zk Σ protocol:

KeyGen	Commitment	Response(c)
$\vec{s} \leftarrow S_n^L$	$\vec{\gamma} \leftarrow S_{\delta_1}^L$	$\vec{z} \leftarrow c \cdot \vec{s} + \vec{\gamma}$
$A \leftarrow R_q^{k \times L}$	$\vec{w} \leftarrow A \vec{\gamma}$	if $\ \vec{z}\ _\infty > \delta_1 - \eta \tau$ then
$\vec{t} \leftarrow A \vec{s}$	return \vec{w}	return \perp
$pk \leftarrow (A, \vec{t})$		return \vec{z}
$sk \leftarrow \vec{s}$	Challenge	Verify(\vec{z})
return (pk, sk)	$c \leftarrow \text{Ball}(\tau)$	assert $A \vec{z} == c \cdot \vec{t} + \vec{w}$
	return c	and $\ \vec{z}\ _\infty \leq \delta_1 - \eta \tau$

Module-SIS ZKP attempt #3

Attempt #3 captures the idea from "Fiat-Shamir w/ abort" (Ly'09): in the Σ -protocol, if the combination of $\vec{\gamma}$ and c is such that \vec{z} will leak information, then the prover simply refuse to release \vec{z} (in practice prover can make many commits, or verifier can send many challenges).

We can estimate the probability of no abort:

$$\begin{aligned}
 P[\|\vec{z}\|_\infty \leq \delta_1 - \eta \tau] &= \prod_{i,j} P[|z_{i,j}| \leq \delta_1 - \eta \tau] \\
 &= \prod_{i,j} \frac{\delta_1 - \eta \tau}{\delta_1} \\
 &= \left(1 - \frac{\eta \tau}{\delta_1}\right)^{L \cdot n}
 \end{aligned}$$

Proto-Dilithium

parameters: $R_q = \mathbb{Z}_q[x]/\langle x^n + 1 \rangle$, $S_\eta = R_{[-\eta, \eta]}$, $L \geq k$, $\text{Ball}(\alpha)$, δ_1

KeyGen	Sign(sk= \vec{s} , m)	Verify(pk, m, σ)
$\vec{s} \xleftarrow{\$} S_\eta^L$	$\vec{y} \xleftarrow{\$} S_{\delta_1}^L$	$(\vec{w}, \vec{z}) \leftarrow \sigma$
$A \xleftarrow{\$} R_q^{k \times L}$	$\vec{w} \leftarrow A\vec{y}$	if $\ \vec{z}\ _\infty > \delta_1 - \eta\alpha$ then
$\vec{t} \leftarrow A\vec{s}$	$c \leftarrow H(\vec{w}, m)$	return 0
$sk \leftarrow \vec{s}$	$\vec{z} \leftarrow c \cdot \vec{s} + \vec{y}$	$\hat{c} \leftarrow H(\vec{w}, m)$
$pk \leftarrow (A, \vec{t})$	if $\ \vec{z}\ _\infty > \delta_1 - \eta\alpha$ then	if $A\vec{z} = \hat{c} \cdot \vec{t} + \vec{w}$ then
return (pk, sk)	return \perp	return 1
	$\sigma \leftarrow (\vec{w}, \vec{z})$	return 0
	return σ	

proto-Dilithium derived from applying Fiat-Shamir to the ZKP in attempt #3

problem: signing routine might fail, the probability of success:

$$P[\text{Sign} \neq \perp] = \left(1 - \frac{\eta\alpha}{\delta_1}\right)^{\ln}$$

to decrease the probability of abort we want:

- a smaller value for η , but a smaller value for η means the bound for $\|\vec{z}\|_\infty \leq \delta_1 - \eta\alpha$ is relaxed, which makes the underlying Module-SIS problem easier to solve \Rightarrow less security

solution: use LWE, where $A \xleftarrow{\$} R_q^{k \times L}$ but $\vec{t} \leftarrow A\vec{s} + \vec{e}$. Commit in still $\vec{w} \leftarrow A\vec{y}$ and response is still $\vec{z} \leftarrow c \cdot \vec{s} + \vec{y}$.

Verification will need tweaking:

$$A\vec{z} = c \cdot A\vec{s} + A\vec{y} = c \cdot (\vec{t} - \vec{e}) + \vec{w} = c \cdot \vec{t} + \vec{w} - \underline{\underline{c \cdot \vec{e}}}$$

the verifier instead checks whether $\|A\vec{z} - c \cdot \vec{t} - \vec{w}\|_\infty$ is "small enough"