

Kontsevich's Characteristic Classes as Topological Invariants of Configuration Space Bundles

Xujia Chen

September 21, 2023

Abstract

Kontsevich's characteristic classes are invariants of framed smooth fiber bundles with homology sphere fibers. It was shown by Watanabe that they can be used to distinguish smooth S^4 -bundles that are all trivial as topological fiber bundles. In this article we show that this ability of Kontsevich's classes is a manifestation of the following principle: the “real blow-up” construction on a smooth manifold essentially depends on its smooth structure and thus, given a smooth manifold (or smooth fiber bundle) M , the topological invariants of spaces constructed from M by real blow-ups could potentially differentiate smooth structures on M . The main theorem says that Kontsevich's characteristic classes of a smooth framed bundle π are determined by the topology of the 2-point configuration space bundle of π and framing data.

Contents

1	Introduction	2
1.1	Statement of main result	3
1.2	Outline of the proof	5
1.3	Some auxiliary notation	8
1.4	Acknowledgments	10
2	Information from the graph	10
2.1	Quick review of graph homology	11
2.2	Information from the graph	12
3	Defining various spaces, all having a \mathcal{G}-action	13
3.1	Defining various spaces	14
3.2	Structure of X_Γ	18
3.3	Digression: what X_Γ looks like – an informal discussion	23

4	Re-constructing Kontsevich's characteristic classes	26
4.1	Defining Ω_Γ	28
4.2	Defining Kontsevich's characteristic classes	34
4.2.1	Cohomology push-forward	34
4.2.2	Defining Kontsevich's characteristic classes	36
5	Equivalence with the original definition	36
5.1	Čech to de Rham preliminary	37
5.2	Proof of Proposition 5.1	40
6	Some remarks about the condition in Theorem 1.2	45

1 Introduction

Given the following data:

- a smooth fiber bundle $E \xrightarrow{\pi} B$ whose fibers are homology spheres,
- a smooth section $s_\infty : B \rightarrow E$ and a trivialization t of π in a neighborhood U of $s_\infty(B)$,
- a vertical framing F on $E - s_\infty(B)$ which is standard (i.e. looks like the standard framing on \mathbb{R}^n near ∞) with respect to t in U ;

Kontsevich's characteristic classes are a collection of cohomology classes in $H^*(B; \mathbb{R})$, parameterized by some combinatorial data ("graph homology"). They were introduced by Kontsevich [8] and have been exploited by various authors thereafter; see [11] for a good introduction. In [17], Watanabe constructed smooth (trivialized near a section and framed, in the above sense) S^4 -bundles with non-trivial Kontsevich's characteristic classes, implying that as smooth fiber bundles (with fixed trivialization near a section) they are non-trivial, while as topological fiber bundles they are trivial.

We would like to understand why Kontsevich's characteristic classes are able to differentiate smooth fiber bundles that are topologically the same. These classes are constructed by considering the "configuration space bundles" associated to $E \xrightarrow{\pi} B$, which are obtained by doing a sequence of real (oriented) blow-up operations fiberwise, and then doing some sort of intersection in the total space of the configuration space bundle to get an intersection number. Since intersection theoretical invariants usually do not depend on the smooth structure, while the real blow-up operations do, it is plausible that different smooth structures on the original bundle $E \xrightarrow{\pi} B$ yield different topological structures on the induced configuration space bundles, and Kontsevich's characteristic classes only depend on the topological bundle structure of the configuration space bundles (together

with some information from the framing). The purpose of the present article is to make this statement precise and to give a detailed proof. The main theorem 1.2 says that the topological information from the 2-point configuration space bundle of $E \xrightarrow{\pi} B$, together with framing, determines Kontsevich's characteristic classes of $E \xrightarrow{\pi} B$.

Remark 1.1. The homotopy type of the real oriented blow up of a manifold X along a submanifold Y , $Bl_Y X$, does not depend on the smooth structure on X , since it is just $X - Y$; but the topological structure of $\partial Bl_Y X$ as a sphere bundle over Y and how $\partial Bl_Y X$ is attached to $X - Y$ do depend on the smooth structure in an essential way; and a framing on the normal bundle of Y can be used to capture this structure.

For example, the simplest Kontsevich's characteristic classes, when it is a number (Θ -graph invariants), can be viewed as the triple intersection number of a cohomology class (called "propagator class" in Section 4) in a space $\overline{C}_2(\pi)/\sim_F$, where $\overline{C}_2(\pi)$ is the total space of the 2-point configuration space bundle associated to $E \xrightarrow{\pi} B$, and \sim_F is an equivalence relation on $\partial^v \overline{C}_2(\pi)$ (the vertical boundary of $\overline{C}_2(\pi)$) – it uses the framing data F to "pinch" $\partial^v \overline{C}_2(\pi)$, making it lower-dimensional.

1.1 Statement of main result

Throughout Sections 1-5, let M be a closed smooth d -dimensional manifold whose R -homology groups are the same as that of the d -sphere, where $R = \mathbb{Z}$ or \mathbb{R} , and let ∞ be a fixed point in M . Let $\overline{C}_2(M, \infty)$ be the configuration space of 2 ordered, not equal and not ∞ , points in M , compactified by the Fulton-MacPherson compactification. More precisely, $\overline{C}_2(M, \infty)$ is obtained from $M \times M$ by first blowing up $\infty \times \infty$ and then blowing up the strict transforms of $\infty \times M$, $M \times \infty$ and the diagonal Δ . (All the blow-ups we use are oriented.) Denote by $f_+, f_- : \overline{C}_2(M, \infty) \rightarrow M$ the two forgetful maps lifting

$$M^2 \longrightarrow M, \quad f_+(x_1, x_2) = x_2, \quad f_-(x_1, x_2) = x_1,$$

respectively. So, $(f_-, f_+) : \overline{C}_2(M, \infty) \rightarrow M \times M$ is the blow down map.

Denote by $\text{Diff}_+(M)$ (resp. $\text{Homeo}_+(M)$) the group of orientation-preserving diffeomorphisms (resp. homeomorphisms) of M , with the Whitney (resp. compact-open) topology, and define

$$\begin{aligned} \text{Diff}_+(M, N_\infty) &:= \{g \in \text{Diff}_+(M) \mid \exists \text{ neighborhood } U \ni \infty \text{ such that } g|_U = \text{id}\} \\ \text{Homeo}_+(M, N_\infty) &:= \{g \in \text{Homeo}_+(M) \mid \exists \text{ neighborhood } U \ni \infty \text{ such that } g|_U = \text{id}\} \\ \mathcal{G} &:= \{(\tilde{g}, g) \in \text{Homeo}(\overline{C}_2(M, \infty)) \times \text{Homeo}_+(M, N_\infty) \mid g \circ f_\pm = f_\pm \circ \tilde{g}\}. \end{aligned}$$

By a smooth (M, ∞) -bundle $E \xrightarrow{\pi} B$ we mean a fiber bundle with typical fiber (M, ∞) and structure group $\text{Diff}_+(M, N_\infty)$. It has a canonical section s_∞ and a canonical germ of trivializations $t : B \times U \xrightarrow{\sim} \tilde{U}$ of some neighborhood $\tilde{U} \supset s_\infty(B)$ (U is some neighborhood of ∞ in M). Abusing notation we also denote the image $s_\infty(B)$ by s_∞ .

A framing F on a smooth (M, ∞) -bundle $E \xrightarrow{\pi} B$ is a continuous choice of basis for the vertical tangent space at every point of $E - s_\infty$, such that there are neighborhoods $E \supset \tilde{U} \supset s_\infty$, $M \supset U \supset \infty$, a diffeomorphism $(U, \infty) \approx ((\mathbb{R}^n - 0) \sqcup \infty, \infty)$, satisfying that $t^*(F|_{\tilde{U}})$ is the standard framing on \mathbb{R}^n under this diffeomorphism.

Let $\overline{C}_2(\pi) \rightarrow B$ be the associated $\overline{C}_2(M, \infty)$ -bundle, $f_+, f_- : \overline{C}_2(\pi) \rightarrow E$ the two forgetful maps and $\partial^v \overline{C}_2(\pi)$ consists of boundaries of every fiber. So

$$\partial^v \overline{C}_2(\pi) = (f_-, f_+)^{-1}(\Delta(\pi) \cup s_\infty \times_B E \cup E \times_B s_\infty),$$

where \times_B denotes fiber product over B and $\Delta(\pi) \subset E \times_B E$ is the fiberwise diagonal. The framing F induces a map $(f_-, f_+)^{-1}(\Delta(\pi)) \rightarrow S^{d-1}$ which at each point $x \in E - s_\infty$ maps $(f_-, f_+)^{-1}(x, x) = \mathcal{SN}_{\Delta(\pi)}^v E|_x \approx ST_x^v E$ to S^{d-1} using $F : T_x^v E \approx \mathbb{R}^d$; here $ST^v E$ denotes the sphere bundle of the vertical tangent bundle of E and $\mathcal{SN}_{\Delta(\pi)}^v E$ denotes the sphere bundle of the vertical normal bundle of $\Delta(\pi)$ in E . Using the trivialization t , this map can be extended to a map $\partial^v \overline{C}_2(\pi) \rightarrow S^{d-1}$ that we still denote by F , abusing notation; see for example [18, §2.3] for the detailed definition.

Kontsevich's invariants, as cohomology classes in B parameterized by graph homology, are defined for smooth framed (M, ∞) -bundles with smooth base B and $R = \mathbb{R}$. One of the by-products of this article is to extend Kontsevich's invariants to the case where B is any paracompact Hausdorff space and $R = \mathbb{Z}$. The main theorem is

Theorem 1.2 Let $E' \xrightarrow{\pi'} B'$, $E'' \xrightarrow{\pi''} B''$ be smooth (M, ∞) -fiber bundles and s'_∞, s''_∞ their canonical sections. Let F', F'' be framings on π', π'' , respectively. If there exist *continuous* maps

$$\tilde{h} : \overline{C}_2(\pi') \longrightarrow \overline{C}_2(\pi''), \quad h : E' \longrightarrow E'', \quad h_B : B' \longrightarrow B'', \quad h_S : S^{d-1} \xrightarrow{\text{homeomorphism}} S^{d-1}$$

such that the following diagrams commute

$$\begin{array}{ccc} \overline{C}_2(\pi') & \xrightarrow{\tilde{h}} & \overline{C}_2(\pi'') \\ f'_- \downarrow \parallel f'_+ & & f''_- \downarrow \parallel f''_+ \\ E' & \xrightarrow{h} & E'' \\ s'_\infty \uparrow \parallel \pi' & & s''_\infty \uparrow \parallel \pi'' \\ B' & \xrightarrow{h_B} & B'' \end{array} \quad \begin{array}{ccc} \partial^v \overline{C}_2(\pi') & \xrightarrow{\tilde{h}|_{\partial^v \overline{C}_2(\pi')}} & \partial^v \overline{C}_2(\pi'') \\ \downarrow F' & & \downarrow F'' \\ S^{d-1} & \xrightarrow{h_S} & S^{d-1} \end{array}$$

and for every point $b \in B$, \tilde{h}, h restrict to orientation-preserving homeomorphisms on the fibers over b , then the Kontsevich’s characteristic classes of π'' pull back to those of π' by h_B .

Remark 1.3. Without the commutativity condition of the diagram on the right, the assumptions of this theorem is the same as saying (\tilde{h}, h) is a \mathcal{G} -bundle map.

Remark 1.4. See Section 6 for some discussions about the condition in this theorem.

Remark 1.5. This theorem is a “uniqueness” statement: if two fiber bundles have the same “certain topological data”, then their Kontsevich’s invariants are the same. A natural question is whether there is an “existence” statement which abstract out the minimal amount of “certain topological data” needed to define Kontsevich’s invariants. (For example, this minimal data can be: a pair of topological fiber bundles $\overline{C}_2(\pi), E \rightarrow B$ together with bundle maps $f_{\pm} : \overline{C}_2(\pi) \rightarrow E$, a map $F : \overline{C}_2(\pi) \rightarrow S^{d-1}$, satisfying some conditions. These conditions are not trivial.) This question is not answered in the present article because it is not an immediate consequence of the present technique, and would be more suitable for a sequel paper. Section 6 is a possible proposal in this direction, but not the most optimal one and not fully proved either.

1.2 Outline of the proof

We re-construct Kontsevich’s characteristic classes in a way that all the definitions are made using only the topological bundle structure on $\overline{C}_2(\pi), \pi$ and the maps f_{\pm} , avoiding using the smooth structure in definitions. This will make Theorem 1.2 automatic. Sections 2-4 are devoted to this re-construction, which is really just a translation of the original construction. In Section 5 we show that the new definition is equivalent to the original one.

To make such a re-construction, the natural strategy is to translate the original construction using differential forms into the language of some topological cohomology theory—here we use Čech cochains—and thus avoid using the smooth structure. However, the cochains in all topological cohomology theories are rather cumbersome to work with, so we need to find the appropriate spaces so that we actually work with cohomology classes. The seemingly unmotivated definitions in Section 3 are for this purpose. This approach is similar to that of [10] and [7], but with some major difference; see Remark 1.6 below.

We describe in a bit more detail how the re-construction is done below. First, in this paragraph, we briefly recall how the original construction roughly goes. Let $E \xrightarrow{\pi} B$ be a smooth (M, ∞) -bundle with smooth, compact base B , and F a framing on π . Let Γ be a trivalent graph that is closed in graph homology; denote its vertex set and edge set by

$V(\Gamma), E(\Gamma)$, respectively. Denote by $\overline{C}_{V(\Gamma)}(\pi)$ the Fulton-MacPherson compactification of

$$C_{V(\Gamma)}(\pi) := \{(x_v \in E)_{v \in V(\Gamma)} \mid x_v \notin s_\infty, \pi(x_v) = \pi(x_w), x_v \neq x_w, \forall v, w \in V(\Gamma)\}.$$

It is a manifold with boundaries and corners. For every edge e of Γ , there is a forgetful map $f_e : \overline{C}_{V(\Gamma)}(\pi) \rightarrow \overline{C}_2(\pi)$ forgetting everything but the two points labeled by the vertices adjacent to e . Take a closed $(d-1)$ -form ω (called *propagator*) on $\overline{C}_2(\pi)$ satisfying $\omega|_{\partial^v \overline{C}_2(\pi)} = F^* \text{vol}$ for some form vol on S^{d-1} such that $\int_{S^{d-1}} \text{vol} = 1$. Then the desired characteristic class (with parameter Γ) is defined to be the class represented by the push-forward of $\bigwedge_e f_e^* \omega_e$ to B . This pushed-forward form is not automatically closed or independent of the choice of ω ; all the trouble here is that $\overline{C}_{V(\Gamma)}(\pi)$ has boundary. The codimension-1 boundary strata of $\overline{C}_{V(\Gamma)}(\pi)$ are in correspondence with subsets $A \subset \{\infty\} \sqcup V(\Gamma)$ having at least 2 elements. Denote by $\overline{\mathcal{S}}_A$ the closed boundary stratum corresponding to A ; it represents the configurations where the points with labels in A all coincide and “bubble off to a screen”. These boundary strata are divided into 4 types, and treated separately.

- (1) The subgraph Γ_A of Γ spanned by vertices in A has a zero- or univalent vertex, and is not of the form as in the 4-th type below. Then $\overline{\mathcal{S}}_A$ is contracted by the map to B and thus does not contribute.
- (2) Γ_A has a bivalent vertex, then there is an involution on $\overline{\mathcal{S}}_A$, making the contribution from $\overline{\mathcal{S}}_A$ cancel with itself.
- (3) $\infty \in A$ or $A = V(\Gamma)$. This case relies on the framing F , the trivialization of π near s_∞ , and that ω is defined to be compatible with F . A dimension count shows that $\overline{\mathcal{S}}_A$ does not contribute either.
- (4) Γ_A has two vertices connected by an edge, then these $\overline{\mathcal{S}}_A$ correspond to boundary terms of Γ in graph cohomology. Since Γ is closed, their total contribution is 0.

In our re-construction, we no longer use differential form propagators; instead, define the space $\overline{C}_2(\pi)/\sim_F$ obtained from $\overline{C}_2(\pi)$ by contracting $\partial^v \overline{C}_2(\pi)$ to S^{d-1} using the framing F , and the propagator can be naturally replaced by a *propagator class* $\Omega \in H^{d-1}(\overline{C}_2(\pi)/\sim_F)$. Denote by $q : \overline{C}_2(\pi) \rightarrow \overline{C}_2(\pi)/\sim_F$ the quotient map.

Our construction also treats the boundary strata of $\overline{C}_{V(\Gamma)}(\pi)$ type by type; each type is treated in the same spirit as the original arguments.

- (1) Instead of using $\overline{C}_{V(\Gamma)}(\pi)$, we use a “weaker” compactification of $C_{V(\Gamma)}(\pi)$, denoted by $\overline{C}_\Gamma(\pi)$. When marked points coincide, $\overline{C}_{V(\Gamma)}(\pi)$ records the collapsing directions of every pair of points, as well as the relative collapsing speed of each triple of points. But $\overline{C}_\Gamma(\pi)$ only records the collapsing directions of pairs of points whose labels in

$V(\Gamma)$ are connected by an edge of Γ ; $\overline{C}_\Gamma(\pi)$ also does not record the relative collapsing speed of triples of points. Indeed, $\overline{C}_\Gamma(\pi)$ is defined to be the closure of the image of

$$C_{V(\Gamma)}(\pi) \xrightarrow{(f_e)_{e \in E(\Gamma)}} \overline{C}_2(\pi)^{E(\Gamma)}.$$

We denote the image of each $\overline{\mathcal{S}}_A$ in $\overline{C}_\Gamma(\pi)$ by $\overline{\mathcal{S}}_A^\Gamma$. Those $\overline{\mathcal{S}}_A \subset \overline{C}_{V(\Gamma)}(M, \infty)$ for A of type 1 are “contracted”, i.e., $\overline{\mathcal{S}}_A^\Gamma \subset \overline{C}_\Gamma(\pi)$ has codimension 2 or higher.

- (2&4) The original arguments in these two cases tell us that each type 2 $\overline{\mathcal{S}}_A$ has an involution and type 4 $\overline{\mathcal{S}}_A$ ’s can be paired-up and cancel each other. Morally, we want to define a new space obtained from $\overline{C}_\Gamma(\pi)$ by gluing each type 2 $\overline{\mathcal{S}}_A^\Gamma$ to itself by the involution, and glue each type 4 pair $\overline{\mathcal{S}}_{A_1}^\Gamma, \overline{\mathcal{S}}_{A_2}^\Gamma$ to each other, thus they are no longer boundaries. However, the gluing procedure involves much technicality; so we instead just use the space

$$X_\Gamma(\pi) := \bigcup_{\sigma \in \tilde{S}_{E(\Gamma)}} \sigma \cdot \overline{C}_\Gamma(\pi) \subset \overline{C}_2(\pi)^{E(\Gamma)},$$

where $\tilde{S}_{E(\Gamma)}$ is a slight generalization of the permutation group of the set $E(\Gamma)$, which acts on $\overline{C}_2(\pi)^{E(\Gamma)}$ by permuting the factors. Taking the union of all the translates of $\overline{C}_\Gamma(\pi)$ by $\tilde{S}_{E(\Gamma)}$ makes the type 2 and type 4 boundary strata of them coincide and cancel with each other. We will explain in Section 3.3 that after removing a codimension 2 subset $T_2(\pi)$ from $X_\Gamma(\pi)$, it is (fiberwise) a “manifold with boundary and bindings” (see Figure 1 for what binding means), where the pages adjacent to a binding, when counted with sign, sum up to 0. Since $X_\Gamma(\pi)$ is a subspace of $\overline{C}_2(\pi)^{E(\Gamma)}$, for each edge e of Γ , we still have the forgetful map $f_e : X_\Gamma(\pi) \rightarrow \overline{C}_2(\pi)$.

- (3) Denote by $S(\pi) \subset X_\Gamma$ the union of all type 3 $\overline{\mathcal{S}}_A^\Gamma$. Then the boundary of $X_\Gamma(\pi) - T_2(\pi)$ is contained in $S(\pi)$. We show that the cup product $\Omega_\Gamma := \cup_{e \in E(\Gamma)} f_e^* q^* \Omega$ actually lands in the relative group $H^*(X_\Gamma(\pi), S(\pi))$.

Since $X_\Gamma(\pi)$ can be roughly thought of as a manifold with boundary $S(\pi)$ and bindings with pages summing up to 0, $H^*(X_\Gamma(\pi), S(\pi))$ is not trivial and Ω_Γ contains all the information we need. We push it forward to B (cohomology push-forward is defined in Section 4.2.1 using Leray-Serre spectral sequence) to obtain a class in $H^*(B)$, which is the desired Kontsevich’s characteristic class with parameter Γ .

Remark 1.6. Our approach above to modify $\overline{C}_{V(\Gamma)}(\pi)$ is similar to that of [10] and [7]: using a smaller configuration space to deal with boundary strata of type (1), and a gluing construction to deal with boundary strata of type (2) and (4). The main difference is about the construction of the “smaller configuration space”. In [10, 7], it is constructed by blowing up less diagonals in $M^{V(\Gamma)}$ (denote by M the 3-manifold whose configuration space is to be constructed), thus the “smaller configuration space” obtained is a smooth manifold, and




Figure 1: A binding with 4 adjacent pages




Figure 2: An illustration of X_Γ

when doing the gluing, how the corners glue together nicely is analyzed. Here, we construct the “smaller configuration space” in a much simpler way (the closure of the image of the forgetful maps), and define the glued space X_Γ in a very simple way as well. As a trade-off, the downside of this approach is that these spaces are not smooth, therefore we have to make an effort to show that the not-necessarily-smooth part are of at least codimension-2. Yet it is because the definition of $X_\Gamma(\pi)$ is so simple in our approach—essentially, only the topology of $\overline{C}_2(\pi)$ is used—that we are able to prove Theorem 1.2 which only involves the 2-point configuration space instead of the n -point configuration spaces for bigger n .

1.3 Some auxiliary notation

For a set S let $|S|$ denote the number of elements in S . For a set I and a space X , denote $X^I = \prod_{i \in I} X$; for $n \in \mathbb{Z}^{>0}$, denote $X^n = \underbrace{X \times X \dots \times X}_{n\text{-times}}$. Denote by $\Delta_{\text{big}} \subset X^I$ the big diagonal. For a map $f : X \rightarrow Y$, denote

$$f^I : X^I \rightarrow Y^I, \quad f^I((x_i)_{i \in I}) = (f(y_i))_{i \in I}.$$

By the real blow-up of a smooth manifold X along a submanifold Y we mean the oriented blow-up: replacing Y by the sphere normal bundle of Y in X .

We will be using Čech cochains and Čech cohomology. For a space X with an open cover \mathcal{U} , a family of supports Φ , and a coefficient ring R , we write $\check{C}_{\Phi, \mathcal{U}}^*(X; R)$ for the R -module of Čech cochains on X with respect to \mathcal{U} , valued in the constant sheaf $R \times X$, and has support in Φ . Write c instead of Φ (resp. omit Φ) if Φ is the collection of compact subsets (resp. closed subsets) of X . By a skew-symmetric Čech cochain $\alpha \in \check{C}_{\Phi, \mathcal{U}}^n(X; R)$ we mean α is such that

$$\alpha(U_0, \dots, U_i, U_{i+1}, \dots, U_n) = -\alpha(U_0, \dots, U_{i+1}, U_i, \dots, U_n)$$

for all $U_0, \dots, U_n \in \mathcal{U}$ and $0 \leq i < n$. If $Y \subset X$ is closed, we denote $\check{C}_{\Phi, \mathcal{U}}^*(X, A; R) = \check{C}_{\Phi, \mathcal{U}}^*(X; R|_{X-A})$, where $R|_{X-A}$ is the sheaf obtained by restricting $R \times X$ to $X - A$ and then extending it by 0 to X . So, $\check{C}_{\Phi, \mathcal{U}}^*(X, A; R)$ can be identified with a sub-module of $\check{C}_{\Phi, \mathcal{U}}^*(X; R)$: it consists of Čech cochains that vanish on A . Denote by $H_{\Phi, \mathcal{U}}^*(X; R)$ and $H_{\Phi, \mathcal{U}}^*(X, Y; R)$ the cohomology of $\check{C}_{\Phi, \mathcal{U}}^*(X; R)$, $\check{C}_{\Phi, \mathcal{U}}^*(X, A; R)$, respectively.

If \mathcal{U}' is a refinement of \mathcal{U} , then there is a well-defined map $H_{\Phi, \mathcal{U}}^*(X; R) \rightarrow H_{\Phi, \mathcal{U}'}^*(X; R)$ (resp. $H_{\Phi, \mathcal{U}}^*(X, Y; R) \rightarrow H_{\Phi, \mathcal{U}'}^*(X, Y; R)$), independent of the choice of refinements. Write $H_{\Phi}^*(X; R)$ (resp. $H_{\Phi}^*(X, Y; R)$) for the cohomology of X (resp. the pair (X, Y)) with R coefficients; it is the direct limit of $H_{\Phi, \mathcal{U}}^*(X; R)$ (resp. $H_{\Phi, \mathcal{U}}^*(X, Y; R)$) as \mathcal{U} gets more and more refined. In Sections 3 and 4, the coefficient ring R is assumed to be \mathbb{Z} or \mathbb{R} , and is often suppressed from notation; in Section 5, $R = \mathbb{R}$.

We follow [15, §2.4] for the definition of fiber bundles. When we say $p : B \rightarrow X$ is a fiber bundle with fiber Y and structure group G , we mean a fiber bundle as in [15, Definition 2.3], with given (maximal) coordinate functions that we do not explicitly mention. Given a G -action on some other space Y' (resp. and a G -equivariant map $f : Y' \rightarrow Y$), by *the associated bundle of p with fiber Y'* we mean a fiber bundle $p' : B' \rightarrow X$ with fiber Y' and structure group G (resp. and a G -bundle map (see [15, Definition 2.5] for definition) $\tilde{f} : B' \rightarrow B, p \circ \tilde{f} = p'$) built up by gluing coordinate charts in the same way as p . Then p' (resp. (p', \tilde{f})) is unique up to equivalence.

In Section 1.1 we defined a smooth (M, ∞) -bundle to be a fiber bundle with typical fiber (M, ∞) and structure group $\text{Diff}_+(M, N_\infty)$. If B is a smooth manifold, then it is the same as saying (by smooth approximation theorems we can make the transition maps smooth) $E \xrightarrow{\pi} B$ is a smooth submersion between smooth manifolds, with a smooth section $s_\infty : B \rightarrow E$, such that for all $b \in B$, $(\pi^{-1}(b), s_\infty(b))$ is diffeomorphic to (M, ∞) , together with a neighborhood $\tilde{U} \subset E$ of $s_\infty(B)$ and a trivialization $t : B \times (U, \infty) \xrightarrow{\sim} (\tilde{U}, s_\infty)$ where $U \subset M$ is some neighborhood of ∞ .

Throughout this article we assume the reader is familiar with the Fulton-MacPherson compactification – having the picture in their mind; see [5] or [11] for reference. We also

assume some familiarity with the original definition of Kontsevich's characteristic classes; see any one of [8][11][18, Section 2].

1.4 Acknowledgments

I thank Peter Kronheimer for many very helpful discussions throughout the course of the work. I thank Fabian Gundlach for helpful discussions regarding Example 6.1 and for providing the proof of Lemma 6.4. I thank the Society of Fellows at Harvard for support.

2 Information from the graph

Say a graph is *directed* if its edges have directions, *ordered* if both of its vertex set and edge set are ordered. If a vertex or edge is the i -th one in the ordering, we call i its *label*. Given such a graph Γ , we denote by $V(\Gamma)$ its vertex set and $E(\Gamma)$ its edge set. Denote by e_i^Γ the i -th edge of Γ , v_i^Γ the i -th vertex of Γ . Conversely, given an edge e or vertex v of Γ , we denote by $o_\Gamma(e), o_\Gamma(v)$ its label. (So $o_\Gamma(e_i^\Gamma) = o_\Gamma(v_i^\Gamma) = i$.) For an edge e of Γ , denote by $v_+(e), v_-(e)$ the output and input vertex connected to e , respectively.

Suppose Γ_1, Γ_2 are directed, ordered graphs, $\alpha : \Gamma_1 \rightarrow \Gamma_2$ an undirected, unordered graph isomorphism, then we denote by $\text{sgn}(\alpha, \text{vertex}), \text{sgn}(\alpha, \text{edge}) \in \{+1, -1\}$ the permutation signs of α on the set of vertices and edges, respectively, and denote

$$\begin{aligned} \text{sgn}(\alpha, \rightarrow) &= (-1)^{\text{number of edges whose direction is reversed by } \alpha}, \\ \text{sgn}_d(\alpha) &:= (d-1)\text{sgn}(\alpha, \text{edge}) + d(\text{sgn}(\alpha, \text{vertex}) + \text{sgn}(\alpha, \rightarrow)), \quad \text{for } d \in \mathbb{Z}. \end{aligned}$$

For a graph Γ , define $\text{Aut}^u(\Gamma)$ to be the group of automorphisms of Γ as an undirected, unlabeled graph. Define

$$|\text{Aut}^u(\Gamma)|_d^\pm = \sum_{\alpha \in \text{Aut}^u(\Gamma)} \text{sgn}_d(\alpha) \in \mathbb{Z}.$$

For a set I , denote by \tilde{S}_I the group of bijections from $\sqcup_{i \in I} \{i^+, i^-\}$ to itself satisfying that, if i^+ is mapped to j^+ or j^- , then i^- is also mapped to j^+ or j^- . There is an obvious map from \tilde{S}_I to S_I , the symmetry group of I as a set. For $\sigma \in \tilde{S}_I$, we denote by $\text{sgn}(\sigma)$ the sign of its image in S_I ; denote $\text{sgn}'(\sigma) := (-1)^{|\{i \in I \mid \sigma(i^+) = i^-\}|}$.

An element $\alpha \in \text{Aut}^u(\Gamma)$ consists of permutations $\alpha_V \in S_{V(\Gamma)}$ and $\alpha_E \in S_{E(\Gamma)}$, such that the adjacency between edges and vertices are preserved. Define

$$\psi_\Gamma : \text{Aut}^u(\Gamma) \longrightarrow \widetilde{S}_{E(\Gamma)}, \quad \psi_\Gamma(\alpha)(e^\pm) = \alpha_E(e)^\pm \text{ if } \alpha_V(v_+(e)) = \alpha_V(v_\pm(e)).$$


If Γ has no isolated vertex, this is an injective group homomorphism.

2.1 Quick review of graph homology

Let \mathfrak{G} be the free abelian group generated by directed, ordered graphs that are also non-empty, connected and such that every vertex is at least trivalent. Define two equivalence relations on \mathfrak{G} , \sim_{odd} and \sim_{even} , generated by the following: for Γ_1, Γ_2 directed, ordered graphs, if there exists an isomorphism $\phi : \Gamma_1 \rightarrow \Gamma_2$ as unoriented, unordered graphs, then

$$\Gamma_1 \sim_{\text{odd}} (\text{sgn}(\phi, \text{vertex}) \cdot \text{sgn}(\phi, \rightarrow)) \Gamma_2, \quad \Gamma_1 \sim_{\text{even}} \text{sgn}(\phi, \text{edge}) \Gamma_2.$$

For a directed, ordered graph Γ and an edge e of Γ , we define Γ/e to be the graph obtained from Γ by contracting e , with edge directions unchanged, vertices ordered in the same way as Γ , except for the new vertex, now put in the very front; edges ordered in the same way as Γ ; as below.



Define \mathbb{Z} -linear maps

$$\delta_{\text{odd}} : \mathfrak{G}/\sim_{\text{odd}} \longrightarrow \mathfrak{G}/\sim_{\text{odd}}, \quad \delta_{\text{even}} : \mathfrak{G}/\sim_{\text{even}} \longrightarrow \mathfrak{G}/\sim_{\text{even}}$$

induced by: for a directed, ordered graph Γ ,

$$\delta_{\text{odd}}(\Gamma) := \sum_{e \in E(\Gamma)} (-1)^{\text{or}(v_+(e)) - \text{or}(v_-(e))} \Gamma/e, \quad \delta_{\text{even}}(\Gamma) := \sum_{e \in E(\Gamma)} (-1)^{\text{or}(e)} \Gamma/e.$$

It can easily be seen that $\delta_{\text{odd}}^2, \delta_{\text{even}}^2 = 0$. Graph homology is defined to be the homology of $(\mathfrak{G}/\sim_{\text{odd}}, \delta_{\text{odd}})$ and $(\mathfrak{G}/\sim_{\text{even}}, \delta_{\text{even}})$.

The main takeaway we need is the following statement, which follows directly from definition: if $\Gamma_1 + \dots + \Gamma_n \in \mathfrak{G}$, $[\delta_{\text{odd}}(\sum_{i=1}^n \Gamma_i)]_{\sim_{\text{odd}}} = 0$ or $[\delta_{\text{even}}(\sum_{i=1}^n \Gamma_i)]_{\sim_{\text{even}}} = 0$ (“[]” denotes

taking equivalence class with respect to $\sim_{\text{odd}}, \sim_{\text{even}}$, respectively, then there exists an (often not unique) pairing between the edges of $\Gamma_1, \dots, \Gamma_n$, such that, if $e_{\Gamma_a, i}, e_{\Gamma_b, j}$ is a pair, then there is an undirected, unordered graph isomorphism

$$\alpha : \Gamma_a / e_i^{\Gamma_a} \longrightarrow \Gamma_b / e_j^{\Gamma_b}$$

such that

$$\text{sgn}(\alpha, \text{vertex}) \cdot \text{sgn}(\alpha, \rightarrow) \cdot (-1)^{o_{\Gamma_a}(v_+(e_i^{\Gamma_a})) - o_{\Gamma_a}(v_-(e_i^{\Gamma_a})) + o_{\Gamma_b}(v_+(e_j^{\Gamma_b})) - o_{\Gamma_b}(v_-(e_j^{\Gamma_b}))} = -1$$

or

$$\text{sgn}(\alpha, \text{edge}) \cdot (-1)^{o_{\Gamma_a}(e_i^{\Gamma_a}) + o_{\Gamma_b}(e_j^{\Gamma_b})} = -1,$$

respectively.

2.2 Information from the graph

Let $d \geq 3$ be an integer. Denote by $\bar{d} \in \{\text{odd}, \text{even}\}$ the parity of d . Let Γ be a connected, directed, ordered graph such that all vertices are at least trivalent, and $[\delta_{\bar{d}}(\Gamma)]_{\sim_{\bar{d}}} = 0$. (All the arguments in this paper can be modified to work if we assume Γ is a formal sum of graphs instead; for simplicity we focus on the case that Γ is a single graph.)

For $V' \subset V(\Gamma)$, we denote by $\Gamma_{V'}$ the subgraph of Γ spanned by the vertices in V' , and by Γ/V' the graph obtained from Γ by contracting $\Gamma_{V'}$ to one single vertex $[V']_v$. The order of edges in $\Gamma_{V'}, \Gamma/V'$ and the order of vertices in $\Gamma_{V'}$ are the same as in Γ . The order of vertices in Γ/V' is defined as: $[V']_v$ is the first one, and the rest are ordered as in Γ .

Suppose $A \subset \{\infty\} \sqcup V(\Gamma)$, $|A| \geq 2$. Define

$$\Gamma_A = \begin{cases} \Gamma_A, & \text{if } \infty \notin A, \\ \Gamma/(A-\infty), & \text{if } \infty \in A; \end{cases} \quad \Gamma/\Gamma_A = \begin{cases} \Gamma/A, & \text{if } \infty \notin A, \\ \Gamma_{A-\infty}, & \text{if } \infty \in A. \end{cases}$$

We say A is of

- type 1, if Γ_A has a zero-valent or univalent vertex, and $|A| \geq 3$ or Γ_A has no edge;
- type 2, if Γ_A has a bivalent vertex but no uni- or zero-valent vertex;
- type 3, if all vertices of Γ_A are at least trivalent,
- type 4, if Γ_A has exactly 2 vertices with 1 edge connecting them.

Notice that since all vertices of Γ are at least trivalent, $\infty \in A \implies A$ is of type 3.

Suppose A as above is of type 2. We fix v_A a bivalent vertex in A . Denote by $e_A^1, e_A^2 \in E(\Gamma)$ the two edges connected to v_A . Define $\sigma_A \in \tilde{S}_{E(\Gamma)}$ as follows:

$$\begin{cases} \sigma_A(e^\pm) = e^\pm & \text{if } e \neq e_A^1, e_A^2; \\ \sigma_A(e_A^{1\pm}) = e_A^{2\mp} & \text{if } e_A^1, e_A^2 \text{ both start or both end at } v_A; \\ \sigma_A(e_A^{1\pm}) = e_A^{2\pm} & \text{if one of } e_A^1, e_A^2 \text{ starts at } v_A, \text{ the other ends at } v_A. \end{cases}$$

Now we look at A of type 4 above. Since $[\delta_{\bar{d}}(\Gamma)]_{\sim_{\bar{d}}} = 0$, there exists a pairing (given Γ , let us choose it once and for all, and call it Γ -pairing) between type 4 A 's such that, if A_1, A_2 are paired (call them a Γ -pair), then there exists an undirected, unordered graph isomorphism $\alpha_{A_1 A_2} : \Gamma/A_1 \rightarrow \Gamma/A_2$, as in the last paragraph of Section 2.1. Denote by e_1, e_2 the edge in Γ between the two vertices in A_1, A_2 , respectively. We call e_1, e_2 a Γ -pair as well, and denote $\alpha_{A_1 A_2}$ also by $\alpha_{e_1 e_2}$. We define $\sigma_{e_1 e_2} \in \tilde{S}_{E(\Gamma)}$ to be:

$$\begin{cases} \sigma_{e_1 e_2}(e_1^\pm) = e_2^\pm; \\ \sigma_{e_1 e_2}(e_i^{\Gamma\pm}) = e_j^{\Gamma\pm}, & \text{if } \alpha_{A_1 A_2}(e_i^\Gamma) = e_j^\Gamma, \text{ preserving direction;} \\ \sigma_{e_1 e_2}(e_i^{\Gamma\pm}) = e_j^{\Gamma\mp}, & \text{if } \alpha_{A_1 A_2}(e_i^\Gamma) = e_j^\Gamma, \text{ reversing direction.} \end{cases}$$

Note that we implicitly identified edges in $E_\Gamma - \{e_1\}$ (resp. $E_\Gamma - \{e_2\}$) with edges in Γ/A_1 (resp. Γ/A_2). Evidently $\sigma_{e_2 e_1} = \sigma_{e_1 e_2}^{-1}$.

If Γ is trivalent, then $[A_1]_v, [A_2]_v$ are the only vertices of valency 4 in $\Gamma/A_1, \Gamma/A_2$, respectively. So $\alpha_{A_1 A_2}$ must map $[A_1]_v$ to $[A_2]_v$.

3 Defining various spaces, all having a \mathcal{G} -action

Let (M, ∞) be as in Section 1.1. Denote by $\Delta \subset M \times M$ the diagonal. Denote by $C_I(M) = M^I - \Delta_{\text{big}}$ the configuration space of distinct marked points on M labeled by I . If I is ordered, let $C_I(M)$ be oriented by the product orientation on M^I . Denote by $\overline{C}_I(M)$ its Fulton-MacPherson compactification [5]. It can be constructed from M^I by a sequence of real blow ups along various diagonals. For example, $\overline{C}_{\{1,2\}}(M)$ is the real blow up of $M \times M$ along Δ . The space $\overline{C}_I(M)$ has the structure of a smooth manifold with boundaries and corners. For $I' \subset I$ we have a smooth forgetful map $f_{I'} : \overline{C}_I(M) \rightarrow \overline{C}_{I'}(M)$ lifting the map

$$M^I \rightarrow M^{I'}, \quad (x_i)_{i \in I} \rightarrow (x_i)_{i \in I'}.$$

If I' has only one element i , we also denote $f_i = f_{I'}$.

Denote $C_I(M, \infty) = \{(x_i)_{i \in I} \in C_I(M) \mid x_i \neq \infty, \forall i\}$, oriented in the same way as $C_I(M)$. Denote by $\overline{C}_I(M, \infty)$ its Fulton-MacPherson compactification, which is defined to be the preimage of ∞ under the forgetful map $f_{\{*\}} : \overline{C}_{I \sqcup \{*\}}(M) \rightarrow M$. For simplicity we write $\overline{C}_n(M, \infty) := \overline{C}_{\{1, \dots, n\}}(M, \infty)$. For example, $\overline{C}_2(M, \infty)$ is described in Section 1.1. Define $\tau : \overline{C}_2(M, \infty) \rightarrow \overline{C}_2(M, \infty)$ to be the map swapping the two marked points.

For $A \subset \{\infty\} \sqcup I$, $|A| \geq 2$, we denote by $\mathring{S}_A \subset \overline{C}_I(M, \infty)$ the (open) boundary stratum corresponding to that the marked points with labels in A coincide. Denote by \overline{S}_A its closure in $\overline{C}_I(M, \infty)$.

3.1 Defining various spaces

Recall in Section 1.1 we defined

$$\mathcal{G} := \{(\tilde{g}, g) \in \text{Homeo}(\overline{C}_2(M, \infty)) \times \text{Homeo}_+(M, N_\infty) \mid g \circ f_\pm = f_\pm \circ \tilde{g}\}.$$

We call the action of \mathcal{G} on $\overline{C}_2(M, \infty)^I$ by acting simultaneously on every factor the *diagonal action*. All the actions of \mathcal{G} we talk about below are action by homeomorphisms.

Let Γ be a graph as in Section 2.2. Assume Γ is trivalent. For an edge e of Γ , we denote by $f_e : \overline{C}_{V(\Gamma)}(M, \infty) \rightarrow \overline{C}_2(M, \infty)$ the forgetful map lifting

$$M^{V(\Gamma)} \longrightarrow M^2, \quad (x_v)_{v \in V(\Gamma)} \longrightarrow (x_{v_-^\Gamma(e)}, x_{v_+^\Gamma(e)}).$$

Definition 3.1. Denote $f_\Gamma = (f_e)_{e \in E(\Gamma)} : \overline{C}_{V(\Gamma)}(M, \infty) \rightarrow \overline{C}_2(M, \infty)^{E(\Gamma)}$. Define

$$\begin{aligned} C_\Gamma(M, \infty) &\subset \overline{C}_\Gamma(M, \infty) \subset \overline{C}_2(M, \infty)^{E(\Gamma)} \\ \overline{C}_\Gamma(M, \infty) &= \text{image}(f_\Gamma), \quad C_\Gamma(M, \infty) = f_\Gamma(C_{V(\Gamma)}(M, \infty)). \end{aligned}$$

Since Γ is connected, $f_\Gamma|_{C_{V(\Gamma)}(M, \infty)}$ is an embedding. Thus it gives a diffeomorphism $C_{V(\Gamma)}(M, \infty) \rightarrow C_\Gamma(M, \infty)$. Since $V(\Gamma)$ is ordered, this also gives $C_\Gamma(M, \infty)$ an orientation. Since $C_\Gamma(M, \infty)$ can be written as

$$\{(z_e)_{e \in E(\Gamma)} \in \overline{C}_2(M, \infty)^{E(\Gamma)} \mid \forall e, e' \in E(\Gamma), \forall s, s' \in \{+, -\}, f_s(e) = f_{s'}(e') \text{ iff } v_s(e) = v_{s'}(e')\},$$

it is invariant under the diagonal \mathcal{G} -action.

Lemma 3.2 $\overline{C}_\Gamma(M, \infty)$ is the closure of $C_\Gamma(M, \infty)$ in $\overline{C}_2(M, \infty)^{E(\Gamma)}$.

Proof. Since $\overline{C}_{V(\Gamma)}(M, \infty)$ is compact, $f_\Gamma(\overline{C}_{V(\Gamma)}(M, \infty))$ is closed and thus contains the closure of $C_\Gamma(M, \infty)$. On the other hand, $C_{V(\Gamma)}(M, \infty)$ is dense in $\overline{C}_{V(\Gamma)}(M, \infty)$, so $C_\Gamma(M, \infty) = f_\Gamma(C_{V(\Gamma)}(M, \infty))$ is dense in $f_\Gamma(\overline{C}_{V(\Gamma)}(M, \infty))$. \square

Abusing notation, we denote by the projection $\overline{C}_\Gamma(M, \infty) \rightarrow \overline{C}_2(M, \infty)$ to the e -th coordinate still by f_e and denote the inclusion map $\overline{C}_\Gamma(M, \infty) \rightarrow \overline{C}_2(M, \infty)^{E(\Gamma)}$ still by f_Γ . It follows directly from Lemma 3.2 that $\overline{C}_\Gamma(M, \infty)$ is invariant under the diagonal action of \mathcal{G} on $\overline{C}_2(M, \infty)^{E(\Gamma)}$.

Definition 3.3. For I an ordered set, define an action ϕ of \tilde{S}_I on $\overline{C}_2(M, \infty)^I$ by diffeomorphisms: for $\sigma \in \tilde{S}_I$,

$$\phi(\sigma) : \overline{C}_2(M, \infty)^I \longrightarrow \overline{C}_2(M, \infty)^I, \quad \phi(\sigma)(z_i)_{i \in I} = \left(\begin{cases} z_i, & \text{if } \sigma(i^\pm) = i^\pm \\ \tau(z_i), & \text{if } \sigma(i^\pm) = i^\mp \end{cases} \right)_{i \in I}.$$

It is clear from definition that every $\phi(\sigma)$ is equivariant with respect to the diagonal action of \mathcal{G} on $\overline{C}_2(M, \infty)^I$.

Definition 3.4.

$$X_\Gamma := \bigcup_{\sigma \in \tilde{S}_{E(\Gamma)}} \phi(\sigma)(\overline{C}_\Gamma(M, \infty)) \subset \overline{C}_2(M, \infty)^{E(\Gamma)}$$

We still denote the inclusion map $X_\Gamma \rightarrow \overline{C}_2(M, \infty)^{E(\Gamma)}$ by f_Γ and its e -th factors by f_e .

It follows from the \mathcal{G} -invariance of $\overline{C}_\Gamma(M, \infty)$ and the \mathcal{G} -equivariance of each $\phi(\sigma)$ that X_Γ is invariant under the diagonal action of \mathcal{G} on $\overline{C}_2(M, \infty)^{E(\Gamma)}$. We therefore define the action of \mathcal{G} on X_Γ to be the restriction of the diagonal action. The maps f_e are clearly \mathcal{G} -equivariant.

It seems that X_Γ (and its bundle version) is the appropriate space that accommodates Kontsevich's characteristic classes. It is defined this way – taking the $|\tilde{S}_{E(\Gamma)}|$ copies of $\overline{C}_\Gamma(M, \infty)$ inside of $\overline{C}_2(M, \infty)^{E(\Gamma)}$ and taking their union – because this makes the type 2 and 4 boundary strata of the many copies of $\overline{C}_\Gamma(M, \infty)$ cancel with each other; see Figure 2. On the other hand, being a subspace of $\overline{C}_2(M, \infty)^{E(\Gamma)}$ automatically makes it (and all its subspaces) metrisable, as a topological space, which is needed for the various arguments regarding covering dimension and Čech cohomology below.

We first make a remark that in the definition of X_Γ as a union, the “main stratum” parts either coincide or do not intersect. This is the content of Lemma 3.5 below.

Recall $\text{Aut}^u(\Gamma)$ is the group of automorphisms of Γ as an undirected, unlabeled graph. An element $\alpha \in \text{Aut}^u(\Gamma)$ consists of permutations $\alpha_V \in S_{V(\Gamma)}$ and $\alpha_E \in S_{E(\Gamma)}$. Denote by $\gamma : \text{Aut}^u(\Gamma) \rightarrow \text{Diff}(\overline{C}_{V(\Gamma)}(M, \infty))$ the action of $\text{Aut}^u(\Gamma)$ on $\overline{C}_{V(\Gamma)}(M, \infty)$ by permuting marked points according to α_V , namely, $\gamma(\alpha)$ lifts the map

$$M^{V(\Gamma)} \rightarrow M^{V(\Gamma)}, \quad (x_v)_{v \in V(\Gamma)} \rightarrow (x_{\alpha_V(v)})_{v \in V(\Gamma)}.$$

Then (recall $\psi_\Gamma : \text{Aut}^u(\Gamma) \rightarrow \tilde{S}_{E(\Gamma)}$ defined right before Section 2.1)

$$f_\Gamma \circ \gamma(\alpha) = \phi(\psi_\Gamma(\alpha)) \circ f_\Gamma. \quad (1)$$

Lemma 3.5 For $\sigma \in \tilde{S}_{E(\Gamma)}$, if $\sigma \notin \text{image}(\psi_\Gamma)$, then $\phi(\sigma)(C_\Gamma(M, \infty)) \cap \overline{C}_\Gamma(M, \infty) = \emptyset$; if $\sigma = \psi_\Gamma(\alpha)$ for some $\alpha \in \text{Aut}^u(\Gamma)$, then $\phi(\sigma)(C_\Gamma(M, \infty)) = C_\Gamma(M, \infty)$ and $\phi(\sigma)$ changes its orientation by $\text{sgn}(\alpha, \text{vertex})^d$.

Proof. For simplicity we prove the lemma in the case Γ has no repeated edges; it can easily be generalized to the other cases as well. Given an element $z = (z_e)_{e \in E(\Gamma)} \in \overline{C}_2(M, \infty)^{E(\Gamma)}$, we define its *set of vertex positions* $\bigcup_e \{f_+(z_e), f_-(z_e)\} \subset M$. It does not change under the $\phi(\sigma)$ action which only permutes factors. If $x \in \overline{C}_{V(\Gamma)}(M, \infty)$, then $\{f_v(x)\}_{v \in V(\Gamma)}$ = the set of vertex positions of $f_\Gamma(x)$, which has exactly $|V(\Gamma)|$ (distinct) elements if $x \in C_{V(\Gamma)}(M, \infty)$ and less otherwise. Thus, if $x, y \in C_{V(\Gamma)}(M, \infty)$ are such that $f_\Gamma(x) = \phi(\sigma)f_\Gamma(y)$, then there is a permutation $\alpha_V \in S_{V(\Gamma)}$ such that $f_{\alpha_V(v)}(y) = f_v(x)$. Since Γ has no repeated edge, there is a unique $\alpha \in \text{Aut}^u(\Gamma)$ with α_V as given. So $\gamma(\alpha)(x) = y$. So,

$$f_\Gamma(x) = \phi(\sigma)f_\Gamma(y) = \phi(\sigma)(f_\Gamma(\gamma(\alpha)(x))) = \phi(\sigma) \circ \phi(\psi_\Gamma(\alpha))(f_\Gamma(x)) = \phi(\sigma\psi_\Gamma(\alpha))(f_\Gamma(x)).$$

Since $\phi(\sigma\psi_\Gamma(\alpha))$ acts by permuting factors of $\overline{C}_2(M, \infty)^{E(\Gamma)}$ composed with τ 's, and no two factors of $f_\Gamma(x)$ are the same even modulo τ , because Γ has no repeated edges, we must have $\sigma\psi_\Gamma(\alpha) = \text{id}$. So $\sigma = \psi_\Gamma(\alpha^{-1})$. This proves the first statement. The rest is immediate. \square

Definition 3.6. For $A \subset \{\infty\} \sqcup V(\Gamma)$, $|A| \geq 2$, Define

$$\begin{aligned} \mathcal{S}_A^\Gamma := \left\{ (z_e)_{e \in E(\Gamma)} \in \overline{C}_\Gamma(M, \infty) \subset \overline{C}_2(M, \infty)^{E(\Gamma)} \mid \exists x \in M, x = \infty \text{ iff } \infty \in A, \text{ such that} \right. \\ \left. \begin{aligned} &\forall e, e' \in E(\Gamma), \forall s, s' \in \{+, -\}, \text{ if } v_s(e) \in A, \text{ then } f_s(z_e) = x; \\ &\text{if } v_s(e) \neq v_{s'}(e') \text{ and } v_s(e) \notin A, \text{ then } \infty \neq f_s(z_e) \neq f_{s'}(z_{e'}). \end{aligned} \right\} \end{aligned} \quad (2)$$

Denote by $\overline{\mathcal{S}}_A^\Gamma$ the closure of \mathcal{S}_A^Γ . For an edge e of Γ , denote $\mathcal{S}_e^\Gamma := \mathcal{S}_{\{v_+^\Gamma(e), v_-^\Gamma(e)\}}^\Gamma$.

Intuitively, elements of \mathcal{S}_A^Γ are configurations of points such that those in A all coincide in M and no other two points coincide. Notice that $\bar{\mathcal{S}}_A = f_\Gamma^{-1}(\bar{\mathcal{S}}_A^\Gamma)$, $\mathring{\mathcal{S}}_A \subset f_\Gamma^{-1}(\mathcal{S}_A^\Gamma) \subset \bar{\mathcal{S}}_A$, and the inclusions are often strict.

Definition 3.7. For $A \subset \{\infty\} \sqcup V(\Gamma)$, $|A| \geq 2$, define $\mathring{\mathcal{S}}_A^\Gamma \subset \mathcal{S}_A^\Gamma$ to be the set of points such that locally $\bar{\mathcal{C}}_\Gamma(M, \infty)$ is a topological manifold with boundary, i.e.,

$$\mathcal{S}_A^\Gamma := \left\{ x \in \mathcal{S}_A^\Gamma \mid \exists U \subset \bar{\mathcal{C}}_\Gamma(M, \infty) \text{ open neighborhood of } x, \text{ and} \right. \\ \left. \text{homeomorphism } \nu: U \rightarrow \mathbb{R}^{d|V(\Gamma)|-1} \times \mathbb{R}^{\geq 0} \text{ such that } \nu(\mathcal{S}_A^\Gamma) = \mathbb{R}^{d|V(\Gamma)|-1} \times \{0\} \right\}.$$

Moreover, define

$$\mathring{\mathcal{S}}_A^\Gamma = \mathring{\mathcal{S}}_A^\Gamma - \bigcup_{\sigma \in \tilde{\mathcal{S}}_{E(\Gamma)}} \bigcup_{\substack{A' \subset \{\infty\} \sqcup V(\Gamma) \\ |A'| \geq 2}} \phi(\sigma)(\bar{\mathcal{S}}_{A'}^\Gamma - \mathring{\mathcal{S}}_{A'}^\Gamma).$$

Definition 3.8. Define the following subsets of X_Γ :

$$X_\Gamma^{\text{good}} = \bigcup_{\sigma \in \tilde{\mathcal{S}}_{E(\Gamma)}} \phi(\sigma) \left(C_\Gamma(M, \infty) \cup \bigcup_{A \text{ of type 2 or 4}} \mathring{\mathcal{S}}_A^\Gamma \right), \quad T_1 = X_\Gamma - X_\Gamma^{\text{good}}, \\ S = \bigcup_{\sigma \in \tilde{\mathcal{S}}_{E(\Gamma)}} \bigcup_{A \text{ of type 3}} \phi(\sigma)(\bar{\mathcal{S}}_A^\Gamma), \quad T_2 = \bigcup_{\sigma \in \tilde{\mathcal{S}}_{E(\Gamma)}} \phi(\sigma) \bigcup_A (\bar{\mathcal{S}}_A^\Gamma - \mathring{\mathcal{S}}_A^\Gamma).$$

Notice that if A is of type 1, then $\mathring{\mathcal{S}}_A^\Gamma = \emptyset$ since $\bar{\mathcal{S}}_A \subset \bar{\mathcal{C}}_{V(\Gamma)}(M, \infty)$ is contracted by f_Γ . So $T_1 \supset T_2 \supset T_1 - S$. Also T_2, T_1, S are invariant under the $\tilde{\mathcal{S}}_{E(\Gamma)}$ -action ϕ . And T_1, T_2 are closed, since every point in $\mathring{\mathcal{S}}_A^\Gamma$ (resp. \mathcal{S}_A^Γ) has a neighborhood lying in $C_\Gamma(M, \infty) \cup \mathring{\mathcal{S}}_A^\Gamma$ (resp. $C_\Gamma(M, \infty) \cup \mathcal{S}_A^\Gamma$). It is evident from the above three definitions that for every A , $\mathcal{S}_A^\Gamma, \bar{\mathcal{S}}_A^\Gamma, \mathring{\mathcal{S}}_A^\Gamma, \mathring{\mathcal{S}}_A^\Gamma \subset \bar{\mathcal{C}}_\Gamma(M, \infty)$ are invariant under the \mathcal{G} -action, and therefore S, T_1, T_2 are invariant under the \mathcal{G} -action too.

Recall the *covering dimension* of a topological space X is the biggest integer N satisfying: any open cover of X has a refinement \mathcal{U} such that if $U_0, \dots, U_{N+1} \in \mathcal{U}, U_i \neq U_j \forall i, j$, then $U_0 \cap \dots \cap U_{N+1} = \emptyset$. Below we write $\dim_t(X)$ for the covering dimension of X (“t” stands for “topological”). Notice that everything we have defined so far are subspaces of $\bar{\mathcal{C}}_2(M, \infty)^{E(\Gamma)}$, thus are all metrizable topological spaces.

In Section 3.2 below, we show that

$$\bullet \dim_t(T_2) \leq d|V(\Gamma)| - 2, \dim_t(T_1) \leq d|V(\Gamma)| - 1;$$

- $H^{d|V(\Gamma)|}(X_\Gamma, S; R)$ admits a \mathcal{G} -equivariant map to R . (Recall $R = \mathbb{Z}$ or \mathbb{R} is the coefficient ring; we take the trivial \mathcal{G} -action on R .) This statement is the goal of Section 3 and is all what we need to re-construct Kontsevich's classes. In all what follows we will call this map ρ .

3.2 Structure of X_Γ

We list some basic properties of the covering dimension, that will be used later. Let X, X' be non-empty metrisable spaces.

- If $Y \subset X$ is closed, then $\dim_t(Y) \leq \dim_t(X)$. This follows from definition.
- If $Y_1, \dots, Y_n \subset X$ are closed, $\dim_t(Y_i) \leq m, \forall i$, then $\dim_t(Y_1 \cup \dots \cup Y_n) \leq m$; see [12, Theorem 9-10].
- $\dim_t(X \times X') \leq \dim_t(X) + \dim_t(X')$; see [12, Theorem 12-14].

Lemma 3.9 Let $f : X \rightarrow Y$ be a smooth map between smooth manifolds (possibly with boundary and corners). Assume X is compact. Denote

$$A_r = \{x \in X \mid \text{rank}(d_x f) \leq r\}, \quad fA_r = f(A_r).$$

Then $\dim_t(fA_r) \leq r$. Specifically, $\dim_t(f(X)) \leq \dim(X)$.

Proof. This follows from two celebrated theorems. By [13, Corollary on Page 169], there exists countably many charts $(U_i \subset Y, \phi_i : U_i \rightarrow \mathbb{R}^n)_{i=1}^\infty$ of Y such that $fA_r \subset \bigcup_i U_i$ and the Hausdorff dimension of $\phi_i(U_i \cap fA_r)$ is at most r for all i . By [16], the covering dimension of a metrisable, separable space (which $\phi_i(U_i \cap fA_r)$ is) is no bigger than its Hausdorff dimension. So, $\dim_t(U_i \cap fA_r) \leq r$ for all i . Since X is compact and A_r is closed in X , fA_r is compact, and so it is covered by the charts U_1, \dots, U_m for some m . By shrinking each U_i a little bit, we have closed subsets $\{V_i \subset U_i \subset Y\}_{i=1}^m$ which still cover fA_r . And $\dim_t(V_i \cap fA_r) \leq r$ for all $i \leq m$. Since $V_i \cap fA_r$ are closed subsets of fA_r and fA_r is their union, $\dim_t(fA_r) \leq r$. \square

Corollary 3.10 $\dim_t(X_\Gamma) \leq d|V(\Gamma)|$, $\dim_t(S) \leq d|V(\Gamma)| - 1$.

Definition 3.11. Given a graph Γ' , define $\overline{V}_{\Gamma'} \subset (S^{d-1})^{E(\Gamma')}$ to be the image of the map

$$f_{\Gamma'}^{\mathbb{R}^d} = (f'_e)_{e \in E(\Gamma')} : \overline{C}_{V(\Gamma')}(\mathbb{R}^d) \rightarrow (S^{d-1})^{E(\Gamma')},$$

where $\overline{C}_{V(\Gamma')}(\mathbb{R}^d)$ is the Fulton-MacPherson space of configurations of $V(\Gamma')$ -marked points in \mathbb{R}^d , modulo translation and scaling (so $\dim(\overline{C}_{V(\Gamma')}(\mathbb{R}^d)) = d|V(\Gamma')| - d - 1$), and f'_e is the

unique map induced from

$$(\mathbb{R}^d)^{V(\Gamma')} - \Delta_{\text{big}} \longrightarrow S^{d-1}, \quad (x_v)_{v \in V(\Gamma')} \longrightarrow \frac{x_{v_+(e)} - x_{v_-(e)}}{|x_{v_+(e)} - x_{v_-(e)}|},$$

i.e., f'_e is the direction between the points marked by the vertices adjacent to e . Notice the $GL(d)$ action on \mathbb{R}^d induces $GL(d)$ -actions on $S^{d-1}, \overline{C}_{V(\Gamma')}(\mathbb{R}^d), \overline{V}_{\Gamma'}$.

Lemma 3.12 If $x, y \in C_{V(\Gamma')}(\mathbb{R}^d), x \neq y, f_{\Gamma'}^{\mathbb{R}^d}(x) = f_{\Gamma'}^{\mathbb{R}^d}(y)$, then $d_x f_{\Gamma'}^{\mathbb{R}^d}, d_y f_{\Gamma'}^{\mathbb{R}^d}$ are not injective.

Proof. Viewing $C_{V(\Gamma')}(\mathbb{R}^d)$ as the quotient of $(\mathbb{R}^d)^{V(\Gamma')} - \Delta_{\text{big}}$, let us take representatives $x' \in (\mathbb{R}^d)^{V(\Gamma')}$ of x and $y' \in (\mathbb{R}^d)^{V(\Gamma')}$ of y such that the marked point labeled by $v_1^{\Gamma'}$ is the origin and the marked point labeled by $v_2^{\Gamma'}$ has norm 1. It is easy to see that for all $0 < \lambda < 1$, if $\lambda x' + (1-\lambda)y' \in (\mathbb{R}^d)^{V(\Gamma')} - \Delta_{\text{big}}$, then $f_{\Gamma'}^{\mathbb{R}^d}([\lambda x' + (1-\lambda)y']) = f_{\Gamma'}^{\mathbb{R}^d}(x) = f_{\Gamma'}^{\mathbb{R}^d}(y)$. So the differential of $f_{\Gamma'}^{\mathbb{R}^d}$ at x or y is 0 in at least one direction. \square

For $A \subset \{\infty\} \sqcup V(\Gamma), |A| \geq 2$, recall the definitions of $\Gamma_A, \Gamma/\Gamma_A$ from Section 2.2. From the construction of Fulton-MacPherson compactification, we have

Fact 3.13 $\overline{\mathcal{S}}_A$ is a fiber bundle over $\overline{C}_{V(\Gamma/A)}(M, \infty)$ with fiber $\overline{C}_{V(\Gamma_A)}(\mathbb{R}^d)$ and structure group $SL(d)$.

Notice $\overline{\mathcal{S}}_A^\Gamma = f_\Gamma(\overline{\mathcal{S}}_A)$.

Corollary 3.14 $\overline{\mathcal{S}}_A^\Gamma$ is a fiber bundle over $\overline{C}_{\Gamma/A}(M, \infty)$ with fiber \overline{V}_{Γ_A} and structure group $SL(d)$. The map $f_\Gamma|_{\overline{\mathcal{S}}_A}$ is an $SL(d)$ -bundle map covering $f_{\Gamma/A} : \overline{C}_{V(\Gamma/A)}(M, \infty) \rightarrow \overline{C}_{\Gamma/A}(M, \infty)$, which is $f_{\Gamma_A}^{\mathbb{R}^d}$ on each fiber.

Notice that the covering dimension of the total space of a fiber bundle with compact base is no more than the sum of the covering dimensions of its fiber and base: over each chart, the bundle is a product, so this is true; and since covering dimension does not increase when taking finite unions of closed subsets, this is true for the whole total space.

Corollary 3.15 Suppose $A \subset \{\infty\} \sqcup V(\Gamma)$ is of type 1, then $\dim_t(\overline{\mathcal{S}}_A^\Gamma) \leq d|V(\Gamma)| - 2$.

Proof. Since Γ_A has a zero valent vertex v_0 or a uni-valent vertex v_1 adjacent to v'_1 , $f_{\Gamma_A}^{\mathbb{R}^d} : \overline{C}_{V(\Gamma_A)}(\mathbb{R}^d) \rightarrow \overline{V}_{\Gamma_A}$ factors through $f_{v_0} : \overline{C}_{V(\Gamma_A)}(\mathbb{R}^d) \rightarrow \overline{C}_{V(\Gamma_A) - \{v_0\}}(\mathbb{R}^d)$ which forgets the point labeled by v_0 , or through $f_{v_1} : \overline{C}_{V(\Gamma_A)}(\mathbb{R}^d) \rightarrow S^{d-1} \times \overline{C}_{V(\Gamma_A) - \{v_1\}}(\mathbb{R}^d)$ which forgets the distance between the point labeled by v_1, v'_1 , respectively. So, $\dim_t(\overline{V}_{\Gamma_A}) < \dim(\overline{C}_{V(\Gamma_A)}(\mathbb{R}^d))$. \square

Lemma 3.16 $\dim_t(\bar{\mathcal{S}}_A^\Gamma - \dot{\mathcal{S}}_A^\Gamma) \leq d|V(\Gamma)| - 2$ for all $A \subset \{\infty\} \sqcup V(\Gamma)$, $|A| \geq 2$.

Proof. Denote by $p_A: \bar{\mathcal{S}}_A \rightarrow \bar{C}_{V(\Gamma/A)}(M, \infty)$, $p_A^\Gamma: \bar{\mathcal{S}}_A^\Gamma \rightarrow \bar{C}_{\Gamma/A}(M, \infty)$ the fiber bundles in Fact 3.13 and Corollary 3.14, respectively. Denote

$$Z_A = \{x \in \bar{\mathcal{S}}_A \mid \text{rank } d_x(f_\Gamma|_{p_A^{-1}(p_A(x))}) < d|V(\Gamma_A)| - d - 1\}, \quad R_A = f_\Gamma(\bar{\mathcal{S}}_A - Z_A).$$

(Notice that $p_A^{-1}(p_A(x))$ is just the fiber of p_A containing x .) In other words, Z_A consists of points $x \in \bar{\mathcal{S}}_A$ at which $f_\Gamma|_{p_A^{-1}(p_A(x))}$ is not an immersion. $\bar{\mathcal{S}}_A^\Gamma - R_A$ is also a fiber bundle over $\bar{C}_{\Gamma/A}(M, \infty)$ with fiber

$$f_{\Gamma'}^{\mathbb{R}^d}(\{x \in \bar{C}_{V(\Gamma_A)}(\mathbb{R}^d) \mid \text{rank } f_{\Gamma_A}^{\mathbb{R}^d} < d|V(\Gamma_A)| - d - 1\}),$$

so $\dim_t(\bar{\mathcal{S}}_A^\Gamma - R_A) \leq d|V(\Gamma)| - 2$, and $\bar{\mathcal{S}}_A^\Gamma - R_A$ is closed in $\bar{\mathcal{S}}_A^\Gamma$. Since $\bar{\mathcal{S}}_A - \dot{\mathcal{S}}_A$ is covered by codimension-2 or higher strata of $\bar{C}_{V(\Gamma)}(M, \infty)$, $\dim_t(f_\Gamma(\bar{\mathcal{S}}_A - \dot{\mathcal{S}}_A)) \leq d|V(\Gamma)| - 2$. We claim that

$$\bar{\mathcal{S}}_A^\Gamma - (f_\Gamma(\bar{\mathcal{S}}_A - \dot{\mathcal{S}}_A) \cup (\bar{\mathcal{S}}_A^\Gamma - R_A)) \subset \dot{\mathcal{S}}_A^\Gamma. \quad (3)$$

If this is true, then $\bar{\mathcal{S}}_A^\Gamma - \dot{\mathcal{S}}_A^\Gamma$ is the union of two close subsets of $\dim_t \leq d|V(\Gamma)| - 2$ and we are done. Let x be in the LHS of (3). Since $x \in R_A$ and $x \notin f_\Gamma(\bar{\mathcal{S}}_A - \dot{\mathcal{S}}_A)$, by Lemma 3.12, $f_\Gamma^{-1}(x)$ consists of a single element $y \in \dot{\mathcal{S}}_A$. We next show that f_Γ is a homeomorphism onto its image in a neighborhood $U \subset \bar{C}_{V(\Gamma)}(M, \infty)$ of y . Since $p_A(\dot{\mathcal{S}}_A) = C_{V(\Gamma/A)}(M, \infty) = C_{\Gamma/A}(M, \infty)$, $f_\Gamma|_{\dot{\mathcal{S}}_A}$ is locally the product of $f_{\Gamma_A}^{\mathbb{R}^d}$ with a diffeomorphism. So $f_\Gamma|_{\dot{\mathcal{S}}_A}$ is an immersion at y . Since $f_\Gamma|_{C_{V(\Gamma)}(M, \infty)}$ is injective, f_Γ is injective in an open neighborhood $U \subset \bar{C}_{V(\Gamma)}(M, \infty)$ of y . Since f_Γ is a closed map ($\bar{C}_{V(\Gamma)}(M, \infty)$ is compact), $f_\Gamma|_U$ is a homeomorphism onto $f_\Gamma(U)$. Since $f_\Gamma(\bar{C}_{V(\Gamma)}(M, \infty) - U)$ is a closed subset of $\bar{C}_\Gamma(M, \infty)$ not containing x , there is a neighborhood $V \subset \bar{C}_\Gamma(M, \infty)$ of x such that $V \cap f_\Gamma(\bar{C}_{V(\Gamma)}(M, \infty) - U) = \emptyset$, so $V \subset f_\Gamma(U)$. This shows that $\bar{C}_\Gamma(M, \infty)$ has the structure of a topological manifold with boundary near x , completing the proof. \square

Corollary 3.17 $\dim_t(T_2) \leq d|V(\Gamma)| - 2$, $\dim_t(T_1) \leq d|V(\Gamma)| - 1$.

Lemma 3.18 Let Y be a compact metrizable space and $Y_2 \subset Y_1 \subset Y$ be closed subspaces such that $\dim_t(Y_1) \leq n - 1$, $\dim_t(Y_2) \leq n - 2$. Then, for every open cover \mathcal{U} of Y , there exists a refinement \mathcal{U}' of \mathcal{U} such that

- (*) there are open neighborhoods N_{Y_1} of Y_1 , N_{Y_2} of Y_2 , such that for all $U_0, \dots, U_n \in \mathcal{U}'$, pairwise distinct,

$$(U_0 \cap \dots \cap U_n) \cap N_{Y_1} = \emptyset, \quad (U_0 \cap \dots \cap U_{n-1}) \cap N_{Y_2} = \emptyset.$$

Hence, if $S \subset Y_1$ is a closed subset such that $Y_1 - Y_2 \subset S$, then there are canonical isomorphisms

$$H^n(Y, S) \approx H^n(Y, Y_1) \approx H_c^n(Y - Y_1).$$

Proof. The proof of [9, Lemma 21.2.1] goes through almost verbatim here and gives us the first statement. (We first use [12, Proposition 12-9 (1)(3)] where Y_1, Y_2 are plugged in as C_1, C_2 , and then use [12, Proposition 9-3].) The second statement easily follows using standard arguments in Čech cohomology. For the first isomorphism: the restriction map $\check{C}_{\mathcal{U}}^i(Y, Y_1) \rightarrow \check{C}_{\mathcal{U}}^i(Y, S)$ is an equality for all $i \geq n - 1$ and all open covers \mathcal{U} of Y satisfying (*), so $\varinjlim_{\mathcal{U} \text{ satisfying } (*)} H_{\mathcal{U}}^n(Y, Y_1) = \varinjlim_{\mathcal{U} \text{ satisfying } (*)} H_{\mathcal{U}}^n(Y, S)$; since every open cover of Y has a refinement satisfying (*), these two limits are equal to $H^n(Y, Y_1)$, $H^n(Y, S)$, respectively. For the second isomorphism: this follows from [2, Proposition 12-3]. Alternatively, let Φ be the collection of compact subsets of $Y - Y_1$, viewed as subsets of Y , and it is not hard to check directly that the natural restriction maps $H_{\Phi}^n(Y) \rightarrow H_c^n(Y - Y_1)$, $H_{\Phi}^n(Y) \rightarrow H^n(Y, Y_1)$ are isomorphisms. \square

Corollary 3.19 $H^{d|V(\Gamma)|}(X_{\Gamma}, S) \approx H^{d|V(\Gamma)|}(X_{\Gamma}, T_1) \approx H_c^{d|V(\Gamma)|}(X_{\Gamma} - T_1)$ via \mathcal{G} -equivariant isomorphisms.

To construct a map $H_c^{d|V(\Gamma)|}(X_{\Gamma} - T_1) \rightarrow R$, we next realize $X_{\Gamma} - T_1$ as the image of a proper map from an oriented topological manifold of dimension $d|V(\Gamma)|$. Recall at the end of Section 2.2 we defined, for every $A \subset \{\infty\} \sqcup V(\Gamma)$ of type 2, $\sigma_A \in \tilde{S}_{E(\Gamma)}$, and for every Γ -pair $A_1, A_2 \in E(\Gamma)$, $\sigma_{A_1 A_2} \in \tilde{S}_{E(\Gamma)}$. Thus, we have $\phi(\sigma_A), \phi(\sigma_{A_1 A_2}) : \overline{C}_2(M, \infty)^{E(\Gamma)} \rightarrow \overline{C}_2(M, \infty)^{E(\Gamma)}$.

Lemma 3.20 $\phi(\sigma_A)(\overline{S}_A^{\Gamma}) = \overline{S}_A^{\Gamma}$, $\phi(\sigma_{A_1 A_2})(\overline{S}_{A_1}^{\Gamma}) = \overline{S}_{A_2}^{\Gamma}$.

Proof. Let A be of type 2 with chosen bivalent vertex v_A , and vertices v_A^1, v_A^2 adjacent to it. There is a dense open subset $\mathring{S}'_A \subset \mathring{S}_A$ on which we can define an involution $\phi'_A : \mathring{S}'_A \rightarrow \mathring{S}'_A$ which fixes all other marked points and reflects the point labeled by v_A along the mid-point of the line segment between the points labeled by v_A^1, v_A^2 , on the screen that these marked points lie on. (This argument is in Kontsevich's original paper [8, Lemma 2.1]; see e.g. [18, Figure 16] for a nice picture. Notice here we use \mathring{S}'_A because ϕ'_A is not well-defined on \mathring{S}_A , due to cases when the new position of the point labeled by v_A coincides with other points.) Clearly $\phi(\sigma_A) \circ f_{\Gamma} = f_{\Gamma} \circ \phi'_A$. So

$$\phi(\sigma_A)(\overline{S}_A^{\Gamma}) = \phi(\sigma_A)(f_{\Gamma}(\overline{S}_A)) = \overline{\phi(\sigma_A)(f_{\Gamma}(\mathring{S}'_A))} = \overline{f_{\Gamma}(\phi'_A(\mathring{S}'_A))} = \overline{f_{\Gamma}(\mathring{S}'_A)} = \overline{S}_A^{\Gamma}.$$

Let A_1, A_2 be a Γ -pair. Since for A of type 4, \mathring{S}_A is an S^{d-1} -bundle over $C_{V(\Gamma/A)}(M, \infty)$ with fiber over x canonically identified with $ST_{f_{[A]_v}(x)}M$, we can define $\phi'_{A_1 A_2} : \mathring{S}_{A_1} \rightarrow \mathring{S}_{A_2}$

by lifting the map $C_{V(\Gamma/A_1)}(M) \rightarrow C_{V(\Gamma/A_2)}(M)$ switching marked points in the same way as $\alpha_{A_1A_2}$ (defined by the end of Section 2.2) maps vertices of Γ/A_1 to vertices of Γ/A_2 . Since $[A_1]_v$ is mapped to $[A_2]_v$, the fibers are canonically identified. By the definition of $\sigma_{A_1A_2}$, $\phi(\sigma_{A_1A_2}) \circ f_\Gamma = f_\Gamma \circ \phi'_{A_1A_2}$. So

$$\phi(\sigma_{A_1A_2})(\overline{\mathcal{S}_{A_1}^\Gamma}) = \phi(\sigma_{A_1A_2})(f_\Gamma(\overline{\mathcal{S}_{A_1}})) = \overline{\phi(\sigma_{A_1A_2})(f_\Gamma(\mathring{\mathcal{S}_{A_1}}))} = \overline{f_\Gamma(\phi'_{A_1A_2}(\mathring{\mathcal{S}_{A_1}}))} = \overline{f_\Gamma(\mathring{\mathcal{S}_{A_2}})} = \overline{\mathcal{S}_{A_2}^\Gamma}.$$

□

Corollary 3.21 $\phi(\sigma_A)(\mathring{\mathcal{S}}_A^\Gamma) = \mathring{\mathcal{S}}_A^\Gamma$, $\phi(\sigma_{e_1e_2})(\mathring{\mathcal{S}}_{e_1}^\Gamma) = \mathring{\mathcal{S}}_{e_2}^\Gamma$.

Denote

$$C'_\Gamma(M, \infty) := C_\Gamma(M, \infty) \cup \bigcup_{A \text{ of type 2 or 4}} \mathring{\mathcal{S}}_A^\Gamma \subset \overline{C}_\Gamma(M, \infty).$$

Then by the definition of $\mathring{\mathcal{S}}_A^\Gamma$, $C'_\Gamma(M, \infty)$ is a topological manifold with boundary.

Definition 3.22. Take $2^{|E(\Gamma)|}|E(\Gamma)|!$ copies of $C'_\Gamma(M, \infty)$, labeled by elements in $\tilde{S}_{E(\Gamma)}$. We write $C'_\Gamma(M, \infty)^{\textcircled{\sigma}}$ for the copy labeled by $\sigma \in \tilde{S}_{E(\Gamma)}$, and similarly write $(\mathring{\mathcal{S}}_A^\Gamma)^{\textcircled{\sigma}}$, $x^{\textcircled{\sigma}}$, etc., for its subspaces and elements. We orient $C'_\Gamma(M, \infty)^{\textcircled{\sigma}}$ by twisting the orientation on $C_\Gamma(M, \infty)$ by $(-1)^{(d-1)\text{sgn}(\sigma)+d\text{sgn}'(\sigma)}$. Define

$$\tilde{X}_\Gamma := \left(\bigsqcup_{\sigma \in \tilde{S}_{E(\Gamma)}} C'_\Gamma(M, \infty)^{\textcircled{\sigma}} \right) / \sim_\Gamma,$$

where \sim_Γ is the following equivalence relation (gluing boundary components pairwise):

- $\forall A \subset \{\infty\} \sqcup V(\Gamma)$ of type 2, $\forall x \in \mathring{\mathcal{S}}_A^\Gamma$, $\forall \sigma \in \tilde{S}_{E(\Gamma)}$, $x^{\textcircled{\sigma}} \sim_\Gamma (\phi(\sigma_A)(x))^{\textcircled{\sigma\sigma_A^{-1}}}$;
- $\forall A_1, A_2$ a Γ -pair, $\forall x \in \mathring{\mathcal{S}}_{A_1}^\Gamma$, $\forall \sigma \in \tilde{S}_{E(\Gamma)}$, $x^{\textcircled{\sigma}} \sim_\Gamma (\phi(\sigma_{A_1A_2})(x))^{\textcircled{\sigma\sigma_{A_1A_2}^{-1}}}$.

Moreover, define

$$\begin{aligned} \tilde{f} &= (\tilde{f}_e)_{e \in E(\Gamma)} : \tilde{X}_\Gamma \longrightarrow \overline{C}_2(M, \infty)^{E(\Gamma)}, \\ \tilde{f}|_{C'_\Gamma(M, \infty)^{\textcircled{\sigma}}} &= \phi(\sigma) \circ f_\Gamma. \end{aligned}$$

It is well-defined since $\phi(\sigma\sigma_A^{-1}) \circ \phi(\sigma_A) = \phi(\sigma)$, $\phi(\sigma\sigma_{A_1A_2}^{-1}) \circ \phi(\sigma_{A_1A_2}) = \phi(\sigma)$.

It can be easily seen that $\text{image}(\tilde{f}) = X_\Gamma - T_1$. It follows from Lemma 3.5 and the definition above that $\tilde{f}|_{\bigsqcup_{\sigma} C'_\Gamma(M, \infty)^{\textcircled{\sigma}}}$ is a covering map (onto its image) of degree $|\text{Aut}^u(\Gamma)|_d^\pm$. Since $\phi(\sigma)$ is \mathcal{G} -equivariant for all $\sigma \in \tilde{S}_{E(\Gamma)}$, the diagonal \mathcal{G} -action on $\overline{C}_2(M, \infty)^{E(\Gamma)}$ lifts to an action of \mathcal{G} on \tilde{X}_Γ , so that \tilde{f} is equivariant.

Lemma 3.23 \tilde{X}_Γ is an oriented topological manifold of dimension $d|V(\Gamma)|$.

Proof. That it is a topological manifold of dimension $d|V(\Gamma)|$ follows from that, $\phi(\sigma_A) : \mathring{S}_A^\Gamma \rightarrow \mathring{S}_A^\Gamma$ and $\phi(\sigma_{A_1 A_2}) : \mathring{S}_{e_1}^\Gamma \rightarrow \mathring{S}_{e_2}^\Gamma$ are homeomorphisms and \sim_Γ glues together these boundary components of $\bigsqcup_{\sigma \in \tilde{S}_{E(\Gamma)}} C'_\Gamma(M, \infty)^{\otimes \sigma}$ pairwise. It is not difficult to verify that $\phi(\sigma_A), \phi(\sigma_{A_1 A_2})$ are orientation-reversing, if $(\mathring{S}_A^\Gamma)^{\otimes \sigma}$'s are oriented as boundaries of $C'_\Gamma(M, \infty)$. \square

Lemma 3.24 $\tilde{f} : \tilde{X}_\Gamma \longrightarrow X_\Gamma - T_1$ is a proper map.

Proof. Let $K \subset X_\Gamma - T_1$ be compact. To show $\tilde{f}^{-1}(K)$ is compact, let $\{x_n\}_{n=1}^\infty$ be a sequence of points in $\tilde{f}^{-1}(K)$. There is a subsequence (still call it $\{x_n\}$) and some $\sigma \in \tilde{S}_{E(\Gamma)}$ such that $x_n \in C'_\Gamma(M, \infty)^{\otimes \sigma}$ for all n . After possibly passing to a subsequence, $\tilde{f}(x_n)$ converges to some $y \in \phi(\sigma)(\overline{C}_\Gamma(M, \infty)) \cap K$, since $\tilde{f}(x_n) \in \phi(\sigma)(C'_\Gamma(M, \infty))$ and $\overline{C}_\Gamma(M, \infty)$ is closed. But $y \notin \phi(\sigma)(\mathring{S}_A^\Gamma - \mathring{S}_A^\Gamma)$ for any A , since $\phi(\sigma)(\mathring{S}_A^\Gamma - \mathring{S}_A^\Gamma) \subset T_1$. Thus, $y \in \phi(\sigma)(C'_\Gamma(M, \infty))$. Since \tilde{f} maps $C'_\Gamma(M, \infty)^{\otimes \sigma}$ homeomorphically onto $\phi(\sigma)(C'_\Gamma(M, \infty))$, by the definition of $C'_\Gamma(M, \infty)$, $\{x_n\}$ converges to the unique element $x \in \tilde{f}^{-1}(y) \cap C'_\Gamma(M, \infty)^{\otimes \sigma}$. \square

Definition 3.25. Define $\rho : H^{d|V(\Gamma)|}(X_\Gamma, S; R) \longrightarrow R$ to be the composition

$$H^{d|V(\Gamma)|}(X_\Gamma, S) \xrightarrow{\text{Cr1 3.19}} H_c^{d|V(\Gamma)|}(X_\Gamma - T_1) \xrightarrow{\tilde{f}^*} H_c^{d|V(\Gamma)|}(\tilde{X}_\Gamma) \longrightarrow R,$$

where the last arrow is by cap product with the fundamental class of \tilde{X}_Γ (in the sense of Borel-Moore homology).

Since the \mathcal{G} -action on \tilde{X}_Γ is orientation-preserving, the last map is \mathcal{G} -equivariant (where R is equipped with the trivial action). So, \mathcal{G} acts on all the objects involved in this definition and all maps involved are equivariant, implying that ρ is \mathcal{G} -equivariant.

3.3 Digression: what X_Γ looks like – an informal discussion

This subsection can be skipped. It is here to justify Figure 2: after removing a codimension 2 subset from X_Γ , it looks like a manifold with boundary and bindings, where each binding component is connected to an even number of pages, summing up to 0 when counted with sign. This statement would also allow us to define ρ in a different (more canonical) way

than in Section 3.2. But working out everything precisely is quite technically involved, so we will only sketch such an approach in this subsection.

By Lemma 3.5,

$$X_\Gamma = \bigsqcup_{[\sigma] \in \tilde{S}_{E(\Gamma)} / \text{image}(\psi_\Gamma)} \phi(\sigma)(C_\Gamma(M, \infty)) \sqcup \bigcup_{\sigma \in \tilde{S}_{E(\Gamma)}} \phi(\sigma)(\overline{C}_\Gamma(M, \infty) - C_\Gamma(M, \infty)).$$

Denote the first term above by \mathring{X}_Γ .

First we define “binding points”.

Definition 3.26. For $A \subset \{\infty\} \sqcup V(\Gamma)$, $|A| \geq 2$, define $\mathring{\mathcal{S}}_A^\Gamma \subset \mathcal{S}_A^\Gamma$ to be the set of points p satisfying: there exists a neighborhood $U \subset \overline{C}_2(M, \infty)^{E(\Gamma)}$ of p , such that for every $\sigma \in \tilde{S}_{E(\Gamma)}$, either

- (1) $\phi(\sigma)(\overline{C}_\Gamma(M, \infty)) \cap U = \emptyset$, or
- (2) $p = \sigma(q)$ for some $A' \subset \{\infty\} \sqcup V(\Gamma)$, $|A'| \geq 2$, $q \in \mathcal{S}_{A'}^\Gamma$, and

- there is a homeomorphism

$$\nu : U \cap \phi(\sigma)(\overline{C}_\Gamma(M, \infty)) \longrightarrow \mathbb{R}^{N-1} \times \mathbb{R}^{\geq 0}, \text{ s.t. } \nu(U \cap \phi(\sigma)(\mathcal{S}_{A'}^\Gamma)) = \mathbb{R}^{N-1} \times \{0\}$$

(i.e., $\phi(\sigma)(\overline{C}_\Gamma(M, \infty))$ is a topological manifold with boundary near p);

- $U \cap \phi(\sigma)(\mathcal{S}_{A'}^\Gamma) = U \cap \mathcal{S}_A^\Gamma$.

For $p \in \mathring{\mathcal{S}}_A^\Gamma$, define the *signed count of pages at p* to be the signed count of elements in $\tilde{S}_{E(\Gamma)}$: those σ of case (1) above are counted with 0; those σ of case (2) above are counted with ± 1 : $+1$ if the boundary orientations of \mathcal{S}_A^Γ near p , as boundary of $\phi(\sigma)(\overline{C}_\Gamma(M, \infty))$ and as boundary of $\overline{C}_\Gamma(M, \infty)$, agree; -1 if they disagree.

Lemma 3.27 If $p \in \mathring{\mathcal{S}}_A^\Gamma$ where A is of type 2 or 4, then the signed count of pages at p is always 0.

Sketch of Proof. For type 2 A s, the pages come in pairs of opposite signs, see Lemma 3.20; for type 4 A s, the pages sum up to 0 because Γ is closed in graph homology. \square

We define S, T_1, T_2 verbatim as in Definition 3.8, just with $\mathring{\mathcal{S}}_A^\Gamma$ replaced by $\mathring{\mathcal{S}}_A^\Gamma$. The statements in the paragraph below Definition 3.8 still hold.

Lemma 3.28 There is a \mathcal{G} -equivariant surjective map $H_c^{d|V(\Gamma)|}(X_\Gamma - T_1; R) \rightarrow R$, where \mathcal{G} acts on R trivially.

Sketch of Proof. $X_\Gamma - T_1$ consists of two parts: \mathring{X}_Γ is an open subset of $X_\Gamma - T_1$ which is also an $d|V(\Gamma)|$ -dimensional (oriented) topological manifold, and $Y := X_\Gamma - T_1 - \mathring{X}_\Gamma$ is a closed subset of $X_\Gamma - T_1$ which is also an $(d|V(\Gamma)| - 1)$ -dimensional topological manifold (this follows from the definition of book binding points). It also follows from the definition of book binding points that $X_\Gamma - T_1$ is locally contractible. So we have the long exact sequence of compactly supported cohomology,

$$\cdots \longrightarrow H_c^{d|V(\Gamma)|-1}(Y) \xrightarrow{\delta} H_c^{d|V(\Gamma)|}(\mathring{X}_\Gamma) \longrightarrow H_c^{d|V(\Gamma)|}(X_\Gamma - T_1) \longrightarrow H_c^{d|V(\Gamma)|}(Y) \longrightarrow \cdots,$$

where the last term is 0. Denote by J_1, J_2 the set of connected components of Y and \mathring{X}_Γ , respectively, then

$$H_c^{d|V(\Gamma)|-1}(Y) \approx R^{\oplus J_1}, \quad H_c^{d|V(\Gamma)|}(\mathring{X}_\Gamma) \approx R^{\oplus J_2},$$

and δ is the coboundary map. So, by Lemma 3.27, the image of δ is contained in $\{(r_i)_{i \in J_2} \mid \sum_i r_i = 0\}$. Therefore, the map

$$H_c^{d|V(\Gamma)|}(\mathring{X}_\Gamma) \longrightarrow R, \quad (r_i)_{i \in J_2} \longrightarrow \sum_{i \in J_2} r_i$$

induces a surjective map from the quotient $H_c^{d|V(\Gamma)|}(\mathring{X}_\Gamma)/\text{image}(\delta) \approx H_c^{d|V(\Gamma)|}(X_\Gamma - T_1)$ to R , as desired. \square

Lemma 3.29 $\dim_t(T_2) \leq d|V(\Gamma)| - 2$.

Together with Lemma 3.18, Lemma 3.29 implies that $H^{d|V(\Gamma)|}(X_\Gamma, S) \approx H_c^{d|V(\Gamma)|}(X_\Gamma - T_1)$, and we can thus define $\rho : H^{d|V(\Gamma)|}(X_\Gamma, S) \rightarrow R$ using the map in Lemma 3.28. The rest of this subsection is devoted to the following

Sketch of Proof of Lemma 3.29. Recall T_2 consists of points in $X_\Gamma - \mathring{X}_\Gamma$ that are not binding points. So, we need to analyze, for $A, A' \in V(\Gamma) \sqcup \{\infty\}$ and $\sigma \in \tilde{S}_{E(\Gamma)}$, how \mathcal{S}_A^Γ and $\phi(\sigma)(\mathcal{S}_{A'}^\Gamma)$ intersect. Suppose A, A', σ are such that they do intersect. Then, by the same reasoning as in Lemma 3.5, there exists an unordered, unoriented graph isomorphism $\Gamma/\Gamma_{A'} \rightarrow \Gamma/\Gamma_A$ whose edge permutation is given by the restriction of σ to $E(\Gamma/\Gamma_{A'})$. So σ also restricts to a bijection $E(\Gamma_{A'}) \rightarrow E(\Gamma_A)$. Abusing notation we still denote by $\phi(\sigma)$ the map $(S^{d-1})^{E(\Gamma_{A'})} \rightarrow (S^{d-1})^{E(\Gamma_A)}$, permuting factors according to σ and composing with the antipodal map when there is a negative sign. Recall the notation “ $f_{\Gamma'}^{\mathbb{R}^d}$ ” in Definition 3.11. Denote

$$V_{A'}^\sigma = \phi(\sigma)(f_{\Gamma_{A'}}^{\mathbb{R}^d}(C_{V(\Gamma_{A'})}(\mathbb{R}^d))) \subset (S^{d-1})^{E(\Gamma_A)}, \quad V_A = f_{\Gamma_A}^{\mathbb{R}^d}(C_{V(\Gamma_A)}(\mathbb{R}^d)) \subset (S^{d-1})^{E(\Gamma_A)}.$$

Then, $\mathcal{S}_A^\Gamma \cap \phi(\sigma)(\mathcal{S}_{A'}^\Gamma)$ is a fiber bundle over $C_{\Gamma/A}(M, \infty)$ with fiber $V_A \cap V_{A'}^\sigma \subset (S^{d-1})^{E(\Gamma_A)}$.

Lemma 3.30 Let Γ' be a graph. Then $f_{\Gamma'}^{\mathbb{R}^d}(C_{V(\Gamma')}(\mathbb{R}^d)) \subset (S^{d-1})^{E(\Gamma')} \subset (\mathbb{R}^d)^{E(\Gamma')}$, where S^{d-1} is viewed as the unit sphere in \mathbb{R}^d , is semi-algebraic.

Proof. It is the image of the composition of a linear map $(\mathbb{R}^d)^{V(\Gamma)} - \Delta_{\text{big}} \rightarrow (\mathbb{R}^d)^{E(\Gamma')}$ with the projection map $(\mathbb{R}^d)^{E(\Gamma')} \rightarrow (S^{d-1})^{E(\Gamma')}$. Both of these maps' graphs are semi-algebraic. So the image is also semi-algebraic by Tarski–Seidenberg Theorem. \square

By the above theorem, $V_{A'}^\sigma$ and V_A are semi-algebraic. They are also open subsets of $(S^{d-1})^{E(\Gamma_A)}$. Let $Y_{A'}^\sigma, Y_A \subset (S^{d-1})^{E(\Gamma_A)}$ be minimal algebraic sets containing $V_{A'}^\sigma, V_A$, respectively. Then $V_{A'}^\sigma \subset Y_{A'}^\sigma, V_A \subset Y_A$ are open (in Euclidean topology), and the Krull dimensions of both Y_A and $Y_{A'}^\sigma$ are $d|V(\Gamma_A)| - d - 1$. Denote by $Z(A, A', \sigma)$ the union of irreducible components of $Y_{A'}^\sigma \cap Y_A$ whose Krull dimension is less than $d|V(\Gamma_A)| - d - 1$. Suppose $p \in V_A \cap V_{A'}^\sigma - Z(A, A', \sigma)$, then p is in some irreducible component Y of $Y_A \cap Y_{A'}^\sigma$ whose Krull dimension is $d|V(\Gamma_A)| - d - 1$, so Y must also be an irreducible component of both Y_A and $Y_{A'}^\sigma$. Since $p \notin Z(A, A', \sigma)$, there exists a neighborhood $U_p \subset (S^{d-1})^{E(\Gamma_A)}$ of p such that $V_A \cap U_p = Y \cap U_p = V_{A'}^\sigma \cap U_p$.

Now, define $\tilde{Z}(A, A', \sigma)$ to be the sub-fiber bundle of \mathcal{S}_A^Γ over $C_{\Gamma/A}(M, \infty)$ whose fibers are $Z(A, A', \sigma) \cap V_A$. For a given $A \subset V(\Gamma) \sqcup \{\infty\}$, define $\mathcal{S}_A^{\Gamma, \text{rmv}} = \bigcup_{\sigma, A'} \tilde{Z}(A, A', \sigma)$, where “rmv” stands for “remove”. It can be shown that the Krull dimension of an algebraic subset of \mathbb{R}^n equals to its covering dimension in Euclidean topology, so $\dim_t(\mathcal{S}_A^{\Gamma, \text{rmv}}) \leq d|V(\Gamma)| - 2$. By the conclusion of the previous paragraph, every point in $\bar{\mathcal{S}}_A^\Gamma$ which is not in (1) $\mathcal{S}_A^{\Gamma, \text{rmv}}$ or (2) $\mathcal{S}_A^\Gamma - \bar{\mathcal{S}}_A^\Gamma$ (as in Definition) or (3) the image of some codimension at least 2 stratum of $\bar{C}_{V(\Gamma)}(M, \infty)$ under f_Γ , is a binding point, by the definition of binding points. By Lemma 3.16, $\dim_t(\mathcal{S}_A^\Gamma - \bar{\mathcal{S}}_A^\Gamma) \leq d|V(\Gamma)| - 2$. So, the union of the above three sets has covering dimension at most $d|V(\Gamma)| - 2$. This completes the proof of the lemma. \square

4 Re-constructing Kontsevich's characteristic classes

Recall $R = \mathbb{Z}$ or \mathbb{R} and all cohomology in this section are with R -coefficients, which we omit. Let Γ be as in Section 2.2; assume also that Γ is trivalent. Let M be as in Section 1.1. Assume $d \geq 3$. Let $\pi: E \rightarrow B$ be a framed smooth (M, ∞) -fiber bundle as in Section

1.1. We assume B can be given a CW-structure. Using CW-approximation, Definition 4.10 and thus Corollary 4.11 generalize to cases where B is just a paracompact Hausdorff space.

Let

$$\begin{array}{cccccccccccc}
C_2(\pi) & \overline{C}_2(\pi) & \overline{C}_2^{E(\Gamma)}(\pi) & \overline{C}_\Gamma(\pi) & \mathcal{S}_A(\pi) & \mathcal{S}_A^\Gamma(\pi) & \overline{\mathcal{S}}_A^\Gamma(\pi) & X_\Gamma(\pi) & T_1(\pi) & T_2(\pi) & S(\pi) & \tilde{X}_\Gamma(\pi) \\
\downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow^{\pi_X} & \downarrow & \downarrow & \downarrow & \downarrow^{\pi_{\tilde{X}}} \\
B & B & B & B & B & B & B & B & B & B & B & B
\end{array}$$

be the associated bundles of π with fibers $C_2(M, \infty)$, $\overline{C}_2(M, \infty)$, $\overline{C}_2(M, \infty)^{E(\Gamma)}$, $\overline{C}_\Gamma(M, \infty)$, \mathcal{S}_A , \mathcal{S}_A^Γ , $\overline{\mathcal{S}}_A^\Gamma$, X_Γ , T_1 , T_2 , S , \tilde{X}_Γ , respectively. Correspondingly, the maps $f_\pm, f_\Gamma, \tilde{f}$ in Section 3.1 induce bundle maps. Abusing notation, we still denote them by $f_\pm, f_\Gamma, \tilde{f}$. Notice that $\overline{C}_2^{E(\Gamma)}(\pi)$ denotes the fiber product while $\overline{C}_2(\pi)^{E(\Gamma)}$ denotes the direct product of the total space, ignoring the fiber bundle structure.

Lemma 4.1 Under the condition of Theorem 1.2, (\tilde{h}, h) induces \mathcal{G} -bundle maps between the π' and π'' version of all the bundles above.

Proof. By the existence of \tilde{h} , h is a \mathcal{G} -bundle map. Since every space defined in Section 3.1 has induced \mathcal{G} -action and the maps between them defined in Section 3.1 are all \mathcal{G} -equivariant,

$$\tilde{h}^{E(\Gamma)} : \overline{C}_2(\pi')^{E(\Gamma)} \longrightarrow \overline{C}_2(\pi'')^{E(\Gamma)}$$

restricts to \mathcal{G} -bundle maps

$$\begin{aligned}
\tilde{h}^{E(\Gamma)} : \overline{C}_2^{E(\Gamma)}(\pi') &\rightarrow \overline{C}_2^{E(\Gamma)}(\pi''), \quad \tilde{h}_\Gamma : \overline{C}_\Gamma(\pi') \rightarrow \overline{C}_\Gamma(\pi''), \\
\tilde{h}_X : X_\Gamma(\pi') &\rightarrow X_\Gamma(\pi''), \quad \tilde{h}_S : S(\pi') \rightarrow S(\pi''), \quad \text{etc..}
\end{aligned}$$

□

The framing on π induces a map $F : \partial^v \overline{C}_2(\pi) \rightarrow S^{d-1}$, as in [18, §2.4.3] (it is called $p(\tau_E)$ there). Define \sim_F to be the following equivalence relation on $\overline{C}_2(\pi)$:

$$x \sim_F y \quad \text{if} \quad x = y \in C_2(\pi) = \overline{C}_2(\pi) - \partial^v \overline{C}_2(\pi) \quad \text{or} \quad F(x) = F(y), \quad x, y \in \partial^v \overline{C}_2(\pi).$$

Denote by $q : \overline{C}_2(\pi) \rightarrow \overline{C}_2(\pi)/\sim_F$ the quotient map by \sim_F , where the target is equipped with the quotient topology. Let us denote $S_\pi^{d-1} = q(\partial^v \overline{C}_2(\pi))$ and denote $(\overline{C}_2(\pi)/\sim_F) - S_\pi^{d-1}$ still by $C_2(\pi)$, then $\overline{C}_2(\pi)/\sim_F = C_2(\pi) \sqcup S_\pi^{d-1}$. It is not hard to see $\overline{C}_2(\pi)/\sim_F$ is Hausdorff and S_π^{d-1} is a deformation retract of some neighborhood of it (hint: since

$\overline{C}_2(\pi)$ restricted to each cell of B is a manifold with compact boundary, using a cell-by-cell construction we can find a collar neighborhood of $\partial^v \overline{C}_2(\pi)$ in $\overline{C}_2(\pi)$. Notice that the orientation on M specifies an orientation on S_π^{d-1} .

Definition 4.2. The *propagator class* $\Omega(\pi) \in H^{d-1}(\overline{C}_2(\pi)/\sim_F; R)$ is the unique class satisfying that $\Omega(\pi)|_{S_\pi^{d-1}} \in H^{d-1}(S_\pi^{d-1}; R)$ is the Poincaré dual of the point class.

The existence and uniqueness of such a class follows from the exact sequence

$$H^{d-1}(\overline{C}_2(\pi)/\sim_F, S_\pi^{d-1}) \rightarrow H^{d-1}(\overline{C}_2(\pi)/\sim_F) \rightarrow H^{d-1}(S_\pi^{d-1}) \rightarrow H^d(\overline{C}_2(\pi)/\sim_F, S_\pi^{d-1})$$

and the vanishing of its first and last terms: for $n = d, d-1$,

$$H^n(\overline{C}_2(\pi)/\sim_F, S_\pi^{d-1}) \approx H^n((\overline{C}_2(\pi)/\sim_F)/S_\pi^{d-1}) = H^n(\overline{C}_2(\pi)/\partial \overline{C}_2(\pi)) \approx H^n(\overline{C}_2(\pi), \partial \overline{C}_2(\pi))$$

and the vanishing of the last term follows from the proof of [18, Lemma 2.10]. The definition of propagator class here is completely analogous to that in [10].

We next show that $\Omega(\pi)$ gives us a class $\Omega_\Gamma(\pi) \in H^{|E(\Gamma)|(d-1)}(X_\Gamma(\pi), S(\pi))$.

4.1 Defining Ω_Γ

Define

$$\iota : X_\Gamma(\pi) \subset \overline{C}_2^{E(\Gamma)}(\pi) \subset \overline{C}_2(\pi)^{E(\Gamma)} \xrightarrow{q, \dots, q} (\overline{C}_2(\pi)/\sim_F)^{E(\Gamma)}.$$

We remark that $\overline{C}_2(\pi)^{E(\Gamma)}$ is the direct product and $\overline{C}_2^{E(\Gamma)}(\pi)$ is the fiber product. For $I \subset E(\Gamma)$, denote by $p_I : (\overline{C}_2(\pi)/\sim_F)^{E(\Gamma)} \rightarrow (\overline{C}_2(\pi)/\sim_F)^I$ the projection to the I -factors. If $I = \{e\}$ we also denote $p_e = p_I$. For $\sigma \in \tilde{S}_{E(\Gamma)}$ and $A \subset \{\infty\} \sqcup V(\Gamma)$, define $I_{\sigma, A} = \sigma(E(\Gamma_A)) \subset E(\Gamma)$, where we implicitly identify edges of Γ_A also as edges of Γ and abuse notation to still write σ for its image in $S_{E(\Gamma)}$.

Lemma 4.3 Suppose $A \subset \{\infty\} \sqcup V(\Gamma)$ and $\sigma \in \tilde{S}_{E(\Gamma)}$. Denote

$$\overline{V}_A^\sigma = p_{I_{\sigma, A}}(\iota(\phi(\sigma)(\overline{S}_A^\Gamma(\pi)))) \subset (\overline{C}_2(\pi)/\sim_F)^{I_{\sigma, A}},$$

then $\overline{V}_A^\sigma \subset (S_\pi^{d-1})^{I_{\sigma, A}}$ and it is the image of $\overline{C}_{V(\Gamma_A)}(\mathbb{R}^d)$ under a smooth map.

Proof. This is a consequence of the framing F on π (in the case $\infty \notin A$) and the trivialization of π near $s_\infty(B)$ (in the case $\infty \in A$). For simplicity we only consider the case $\sigma = \text{id}$; the other cases follow easily. First assume $\infty \notin A$. Recall $\overline{V}_{\Gamma_A} \subset (S^{d-1})^{E(\Gamma_A)}$ as

in Definition 3.11. Since the framing F identifies the vertical tangent space of E at each point not in $s_\infty(B)$ with \mathbb{R}^d ,

$$\mathcal{S}_A^\Gamma(\pi) \approx C_{\Gamma/A}(\pi) \times \bar{V}_{\Gamma_A} \subset C_2(\pi)^{E(\Gamma/A)} \times (S_\pi^{d-1})^{E(\Gamma_A)} \subset (\bar{C}_2(\pi)/\sim_F)^{E(\Gamma/A)} \times (\bar{C}_2(\pi)/\sim_F)^{E(\Gamma_A)},$$

and $(p_{E(\Gamma_A)} \circ \iota)(\mathcal{S}_A^\Gamma(\pi)) = \bar{V}_{\Gamma_A}$. Since \bar{V}_{Γ_A} is closed, $(p_{E(\Gamma_A)} \circ \iota)(\bar{\mathcal{S}}_A^\Gamma(\pi)) = \bar{V}_{\Gamma_A}$ as well.

Now assume $\infty \in A$. Since π is trivialized near $s_\infty(B)$, the vertical tangent spaces of E at points in $s_\infty(B)$ are all identified with $T_\infty M$. So

$$\bar{\mathcal{S}}_A(\pi) \approx \bar{C}_{V(\Gamma/A)}(\pi) \times \bar{C}_{V(\Gamma_A)}(T_\infty M).$$

And $(p_{E(\Gamma_A)} \circ \iota)(\bar{\mathcal{S}}_A^\Gamma(\pi)) = (p_{E(\Gamma_A)} \circ \iota \circ f_\Gamma)(\bar{\mathcal{S}}_A(\pi))$; the map $p_{E(\Gamma_A)} \circ \iota \circ f_\Gamma$ factors through the projection to the second factor above. \square

By Lemma 4.6 and Lemma 4.7 below (these are two somewhat technical lemmas whose statements and proofs are postponed to below Proposition 4.5, due to their lengths), the cup product

$$\begin{aligned} \cup : H^{d-1}(\bar{C}_2(\pi)/\sim_F)^{\otimes E(\Gamma)} &\longrightarrow H^{|E(\Gamma)|(d-1)}((\bar{C}_2(\pi)/\sim_F)^{E(\Gamma)}), \\ \otimes_{i=1}^{|E(\Gamma)|} \sigma_{e_i^\Gamma} &\longrightarrow p_{e_1^\Gamma}^* \sigma_{e_1^\Gamma} \cup \dots \cup p_{e_{|E(\Gamma)|}^\Gamma}^* \sigma_{e_{|E(\Gamma)|}^\Gamma} \end{aligned}$$

factors through $H^{|E(\Gamma)|(d-1)}((\bar{C}_2(\pi)/\sim_F)^{E(\Gamma)}, \bigcup_{\sigma, A} p_{I_{\sigma, A}}^{-1}(\bar{V}_A^\sigma))$, where the union ranges through all $\sigma \in \tilde{S}_{E(\Gamma)}$ and $A \subset \{\infty\} \sqcup V(\Gamma)$ of type 3 (same below).

Definition 4.4.

$$\Omega'_\Gamma(\pi) := p_{e_1^\Gamma}^* \Omega(\pi) \cup \dots \cup p_{e_{|E(\Gamma)|}^\Gamma}^* \Omega(\pi) \in H^{|E(\Gamma)|(d-1)}((\bar{C}_2(\pi)/\sim_F)^{E(\Gamma)}, \bigcup_{\sigma, A} p_{I_{\sigma, A}}^{-1}(\bar{V}_A^\sigma))$$

$$\Omega_\Gamma(\pi) := \iota^* \Omega_\Gamma(\pi) \in H^{|E(\Gamma)|(d-1)}(X_\Gamma(\pi), S(\pi))$$

Proposition 4.5 Under the assumptions of Theorem 1.2, let $\tilde{h}_X : X_\Gamma(\pi') \rightarrow X_\Gamma(\pi'')$ be the \mathcal{G} -bundle map as in Lemma 4.1, then $\tilde{h}_X^*(\Omega_\Gamma(\pi'')) = \Omega_\Gamma(\pi')$.

Proof. By the second commutative diagram in Theorem 1.2, \tilde{h} induces

$$\tilde{h}_q : \bar{C}_2(\pi')/\sim_{F'} \longrightarrow \bar{C}_2(\pi'')/\sim_{F''}$$

which commutes with q and restricts to the homeomorphism $h_S : S_{\pi'}^{d-1} \rightarrow S_{\pi''}^{d-1}$. It follows that $\tilde{h}_q^* \Omega(\pi'') = \Omega(\pi')$. Since ι commutes with \tilde{h}_X and $(\tilde{h}_q)^{E(\Gamma)}$, by the naturality statement in Lemma 4.6 below, $\tilde{h}_X^*(\Omega_\Gamma(\pi'')) = \Omega_\Gamma(\pi')$. \square

Lemma 4.6 Let $Y = Y_1 \times \dots \times Y_n$ be a product of paracompact Hausdorff spaces. Let $s, r \in \mathbb{Z}^{>0}$. Suppose for all $i = 1, \dots, r$, we have $I_i = \{a_1^i, \dots, a_{m_i}^i\} \subset \{1, \dots, n\}$ and closed subset $V_i \subset Y_{a_1^i} \times \dots \times Y_{a_{m_i}^i}$ satisfying the following condition: every open cover of Y has a refinement of the form

$$\mathcal{U}_1 \times \dots \times \mathcal{U}_n := \{U_1 \times \dots \times U_n \mid U_j \in \mathcal{U}_j\}, \text{ where } \mathcal{U}_j \text{ is an open cover of } Y_j$$

such that

(†) for all $i = 1, \dots, r$, $s'_1, \dots, s'_{m_i} = s$ or $s-1$, with at most one of them being $s-1$,

$$\left((U_{a_1^i}^0 \cap \dots \cap U_{a_1^i}^{s'_1}) \times \dots \times (U_{a_{m_i}^i}^0 \cap \dots \cap U_{a_{m_i}^i}^{s'_{m_i}}) \right) \cap V_i = \emptyset$$

where $U_{a_j^i}^0, \dots, U_{a_j^i}^{s'_j}$ are any pairwise distinct elements of $\mathcal{U}_{a_j^i}$, for every $j = 1, \dots, m_i$.

Denote $p_{I_i} : Y \rightarrow Y_{a_1^i} \times \dots \times Y_{a_{m_i}^i}$ the projection. Then there is a map

$$\Xi : H^s(Y_1) \otimes \dots \otimes H^s(Y_n) \longrightarrow H^{sn}(Y, \bigcup_{i=1}^r p_{I_i}^{-1}(V_i))$$

such that Ξ composed with the restriction to $H^{sn}(Y)$ is the cup product. And Ξ is natural in the following sense: if $Y' = Y'_1 \times \dots \times Y'_n$, $\{V'_i \subset Y'_{a_1^i} \times \dots \times Y'_{a_{m_i}^i}\}_{i=1}^r$ satisfy the same condition as $Y, \{Y_i\}, \{V_i\}$ above, with Ξ' the corresponding map, and there are continuous maps $\{f_i : Y'_i \rightarrow Y_i\}_{i=1}^r$ such that $(f_{a_1^i} \times \dots \times f_{a_{m_i}^i})(V'_i) \subset V_i$ for all i , then $\Xi' \circ (f_1^* \otimes \dots \otimes f_n^*) = (f_1 \times \dots \times f_n)^* \circ \Xi$.

Proof. First we define Ξ . Given $\sigma_1 \in H^s(Y_1), \dots, \sigma_n \in H^s(Y_n)$, take an open cover $\mathcal{U} = \mathcal{U}_1 \times \dots \times \mathcal{U}_n$ of Y satisfying (†) and such that for all i , σ_i is represented by some skew-symmetric Čech cochain $\alpha_i \in \check{C}_{\mathcal{U}_i}^s(Y_i)$ (see, e.g., [3] for the equivalence between usual and skew-symmetric Čech cohomology; in particular, every (usual) Čech cohomology class has a skew-symmetric cochain representative). Denote by $p_i : Y \rightarrow Y_i$ projection to the i -th factor. Define

$$\alpha := p_1^* \alpha_1 \cup \dots \cup p_n^* \alpha_n \in \check{C}_{\mathcal{U}}^{sn}(Y). \quad (4)$$

We claim $\alpha|_{\bigcup_{i=1}^r p_{I_i}^{-1}(V_i)} = 0$. Suppose $x \in p_{I_i}^{-1}(V_i)$ for some i and

$$x \in (U_1^0 \times \dots \times U_n^0) \cap (U_1^1 \times \dots \times U_n^1) \cap \dots \cap (U_1^{sn} \times \dots \times U_n^{sn}), \quad U_i^j \in \mathcal{U}_i,$$

then

$$\alpha((U_1^0 \times \dots \times U_n^0) \cap (U_1^1 \times \dots \times U_n^1) \cap \dots \cap (U_1^{sn} \times \dots \times U_n^{sn}))(x)$$

$$= \alpha_1(U_1^0 \cap \dots \cap U_1^s)(p_1(x)) \cdot \dots \cdot \alpha_n(U_n^{s(n-1)} \cap \dots \cap U_n^{sn})(p_n(x))$$

contains factor

$$\alpha_{a_1^i}(U_{a_1^i}^{s(a_1^i-1)} \cap \dots \cap U_{a_1^i}^{sa_1^i})(p_{a_1^i}(x)) \cdot \dots \cdot \alpha_{a_{m_i}^i}(U_{a_{m_i}^i}^{s(a_{m_i}^i-1)} \cap \dots \cap U_{a_{m_i}^i}^{sa_{m_i}^i})(p_{a_{m_i}^i}(x)). \quad (5)$$

Since $\alpha_1, \dots, \alpha_n$ are skew-symmetric, for (5) to be non-zero, $U_{a_j^i}^{s(a_j^i-1)}, \dots, U_{a_j^i}^{sa_j^i}$ must be pairwise different for all $j = 1, \dots, m_i$. But $p_{I_i}(x) \in V_i$. So by (\dagger) , (5) $\neq 0$ is impossible. This proves $\alpha|_{\bigcup_{i=1}^r p_{I_i}^{-1}(V_i)} = 0$, so $\alpha \in \check{C}_{\mathcal{U}}^{sn}(Y, \bigcup_{i=1}^r p_{I_i}^{-1}(V_i))$. We define

$$\Xi(\sigma_1, \dots, \sigma_n) = \text{the direct image of } [\alpha] \text{ in } H^{sn}(Y, \bigcup_{i=1}^r p_{I_i}^{-1}(V_i)).$$

Then Ξ is clearly linear.

Next we show that $[\alpha]$ above does not depend on the choices of α_i . Suppose $\alpha_i, \alpha'_i \in \check{C}_{\mathcal{U}_i}^s(Y_i)$ are both skew-symmetric cocycles, $[\alpha_i] = [\alpha'_i] = \sigma_i$, then there exists a skew-symmetric $\tilde{\alpha}_i \in \check{C}_{\mathcal{U}_i}^{d-1}(Y_i)$, $d\tilde{\alpha}_i = \alpha_i - \alpha'_i$. Define

$$\tilde{\alpha} := p_1^* \alpha_1 \cup \dots \cup p_{i-1}^* \alpha_{i-1} \cup p_i^* \tilde{\alpha}_i \cup p_{i+1}^* \alpha_{i+1} \cup \dots \cup p_n^* \alpha_n \in \check{C}_{\mathcal{U}}^{sn-1}(Y).$$

By the same argument as above, $\tilde{\alpha}$ vanishes on $\bigcup_{i=1}^r p_{I_i}^{-1}(V_i)$, so $\tilde{\alpha} \in \check{C}_{\mathcal{U}}^{sn-1}(Y, \bigcup_{i=1}^r p_{I_i}^{-1}(V_i))$. Since all α_j are cocycles,

$$d\tilde{\alpha} = (-1)^i (p_1^* \alpha_1 \cup \dots \cup d(p_i^* \tilde{\alpha}_i) \cup \dots \cup p_n^* \alpha_n) = (-1)^i (p_1^* \alpha_1 \cup \dots \cup p_i^* (\alpha_i - \alpha'_i) \cup \dots \cup p_n^* \alpha_n).$$

Therefore $[\alpha]$ does not depend on the choice of α_i .

We then show that $\Xi(\sigma_1, \dots, \sigma_n)$ does not depend on the choices of \mathcal{U}_i . Suppose $\mathcal{U}' = \mathcal{U}'_1 \times \dots \times \mathcal{U}'_n$ also satisfy (\dagger) and for all i , \mathcal{U}'_i is a refinement of \mathcal{U}_i . Let $\{\mu_i : \mathcal{U}'_i \rightarrow \mathcal{U}_i\}_{i=1}^n$ be some refinement maps. Denote $\mu = (\mu_1, \dots, \mu_n) : \mathcal{U}' \rightarrow \mathcal{U}$. Then for $U_j^i \in \mathcal{U}'_j$,

$$\begin{aligned} & \mu^* \alpha((U_1^0 \times \dots \times U_n^0) \cap (U_1^1 \times \dots \times U_n^1) \cap \dots \cap (U_1^{sn} \times \dots \times U_n^{sn}))(x) \\ &= \alpha((\mu_1(U_1^0) \times \dots \times \mu_n(U_n^0)) \cap (\mu_1(U_1^1) \times \dots \times \mu_n(U_n^1)) \cap \dots \cap (\mu_1(U_1^{sn}) \times \dots \times \mu_n(U_n^{sn})))(x) \\ &= \alpha_1(\mu_1(U_1^0) \cap \dots \cap \mu_1(U_1^s))(p_1(x)) \cdot \dots \cdot \alpha_n(\mu_n(U_n^{s(n-1)}) \cap \dots \cap \mu_n(U_n^{sn}))(p_n(x)) \\ &= \mu_1^* \alpha_1(U_1^0 \cap \dots \cap U_1^s)(p_1(x)) \cdot \dots \cdot \mu_n^* \alpha_n(U_n^{s(n-1)} \cap \dots \cap U_n^{sn})(p_n(x)). \end{aligned}$$

This says $\mu^* \alpha = p_1^*(\mu_1^* \alpha_1) \cup \dots \cup p_n^*(\mu_n^* \alpha_n) \in \check{C}_{\mathcal{U}'}^{sn}(Y, \bigcup_{i=1}^r p_{I_i}^{-1}(V_i))$. So, if we define $\Xi(\sigma_1, \dots, \sigma_n)$ using \mathcal{U}' , then it is the direct image of $\mu^*[\alpha] = [\mu^* \alpha] \in H_{\mathcal{U}'}^{sn}(Y, \bigcup_{i=1}^r p_{I_i}^{-1}(V_i))$ which is the same as the direct image of $[\alpha]$. Now, if we assume \mathcal{U}' satisfies (\dagger) but not

necessarily a refinement of \mathcal{U} , then by the assumption of the lemma we can find a common refinement of $\mathcal{U}, \mathcal{U}'$ that also satisfies (\dagger) . This implies that $\Xi(\sigma_1, \dots, \sigma_n)$ is independent of the choice of \mathcal{U} , and thus Ξ is well-defined.

Naturality follows immediately from the definition of Ξ : let $Y', \{Y'_i\}, \{f_i\}, \{V'_i\}$ be as in the lemma and $\mathcal{U}, \alpha, \alpha_i$ as in the first paragraph of the proof, then $\mathcal{U}' := f_1^{-1}\mathcal{U}_1 \times \dots \times f_n^{-1}\mathcal{U}_n$ satisfies (\dagger) . Take $\alpha'_i = f_i^*(\alpha_i) \in \check{C}_{\mathcal{U}'}^s(Y')$, then α' defined using (4) is the same as $(f_1 \times \dots \times f_n)^*(\alpha)$ and the conclusion follows. \square

Lemma 4.7 Every open cover of $(\overline{C}_2(\pi)/\sim_F)^{E(\Gamma)}$ has a refinement of the form

$$\mathcal{U}_{e_1^\Gamma} \times \dots \times \mathcal{U}_{e_{|E(\Gamma)|}^\Gamma}, \text{ where each } \mathcal{U}_{e_i^\Gamma} \text{ is an open cover of } \overline{C}_2(\pi)/\sim_F$$

such that

(*) for all A of type 3, $\sigma \in \tilde{S}_{E(\Gamma)}$, $s'_e = d-1$ or $d-2$ with at most one being $d-2$,

$$\overline{V}_A^\sigma \cap \prod_{e \in I_{\sigma,A}} (U_e^0 \cap \dots \cap U_e^{s'_e}) = \emptyset$$

where $U_e^0, \dots, U_e^{s'_e}$ are any pairwise distinct elements of \mathcal{U}_e , for every $e \in I_{\sigma,A}$.

Proof. Since A is of type 3, $|V(\Gamma_A)| \leq 2/3E(\Gamma_A)$. Since $d \geq 3$, $(2d/3) \leq d-1$, and

$$\dim \overline{C}_{V(\Gamma_A)}(\mathbb{R}^d) = d|V(\Gamma_A)| - d - 1 \leq (2d/3)|E(\Gamma_A)| - (d+1) \leq \dim((S^{d-1})^{E(\Gamma_A)}) - 4.$$

So by Lemma 4.3 \overline{V}_A^σ is the image of a smooth map $f_{\sigma,A} : \overline{C}_{V(\Gamma_A)} \rightarrow (S^{d-1})^{I_{\sigma,A}}$, where the domain is of codimension at least 4 with respect to the target. (Indeed, codimension 2 would suffice for this lemma.)

Fix a metric D on $\overline{C}_2(\pi)/\sim_F$. By Lebesgue's Number Lemma, it suffices to show that for any $\epsilon > 0$, there exist $(\mathcal{U}_e)_{e \in E(\Gamma)}$ such that for all e and $U \in \mathcal{U}_e$, $\text{diameter}_D(U) < \epsilon$, satisfying (*).

We first reduce the lemma to the following statement: for all $\epsilon > 0$, there exist $(\mathcal{U}_e)_{e \in E(\Gamma)}$ open covers of S_π^{d-1} such that for all $U \in \bigcup_e \mathcal{U}_e$, $\text{diameter}_D(U) < \epsilon/4$, satisfying (*). Suppose this is true. We can enlarge each $U \in \mathcal{U}_e$ a little bit to get an open subset $l(U)$ of $\overline{C}_2(\pi)/\sim_F$, still of diameter less than ϵ . (For example, take $l(U) = \{x \in C_2(\pi) \mid D(x, U) < \epsilon/2\} \cup U$.) Denote $l(\mathcal{U}_e) = \{l(U)\}_{U \in \mathcal{U}_e}$. Since $\overline{C}_2(\pi)/\sim_F - \bigcup_{U \in \mathcal{U}_e} l(U)$ is compact, we can cover it by finitely many open subsets of $C_2(\pi)$ of diameter less than ϵ ; call this collection of open sets \mathcal{U}'_e . Then for every e , $l(\mathcal{U}_e) \cup \mathcal{U}'_e$ is an open cover of $\overline{C}_2(\pi)/\sim_F$ of diameter less than ϵ . Since for every A , \overline{V}_A^σ is contained in $(S_\pi^{d-1})^{I_{\sigma,A}}$ whereas elements in \mathcal{U}'_e do not

intersect S_π^{d-1} , that $(*)$ is satisfied by $(l(\mathcal{U}_e) \cup \mathcal{U}'_e)_{e \in E(\Gamma)}$ follows from that it is satisfied by $(\mathcal{U}_e)_{e \in E(\Gamma)}$.

We then reduce the statement at the beginning of last paragraph to the following: for any $\epsilon > 0$ there exist triangulations $(T_e)_{e \in E(\Gamma)}$ of S^{d-1} , compatible with the smooth structure on S^{d-1} , such that the diameter of each simplex is less than $\epsilon/4$, satisfying

(**) for all σ, A , and all collections of simplices $(S_e : \text{a simplex in } T_e)_{e \in I_{\sigma,A}}$ such that S_e is of dimension 0 or 1, at most one of dimension 1, $\bar{V}_A^\sigma \cap \prod_{e \in I_{\sigma,A}} \bar{S}_e = \emptyset$.

Suppose this is true, we can take \mathcal{U}_e to be obtained from T_e by slightly enlarging each top-dimensional simplex S to an open neighborhood $U(S)$ of its closure, so that,

(1) (for a non-top dimensional simplex S , still denote by $U(S) = \bigcap_{S'} U(S')$ where S' runs through the top-dimensional simplices that S is a face of,)

$$\bar{V}_A^\sigma \cap \prod_{e \in I_{\sigma,A}} U(S_e) = \emptyset \text{ whenever } \bar{V}_A^\sigma \cap \prod_{e \in I_{\sigma,A}} \bar{S}_e = \emptyset;$$

(2) for any finite collection, S_1, \dots, S_k , of simplices in T_e , $S_1 \cap \dots \cap S_k = \emptyset \implies U(S_1) \cap \dots \cap U(S_k) = \emptyset$.

(These are clearly easy to satisfy. For (2), for every S in T_e , take $U(S)$ contained in the union of the stars of the barycenters of its faces in the barycentric subdivision of T . For (1), take $U(S)$ contained in the ϵ' -neighborhood of S , where ϵ' is the minimal distance between some \bar{V}_A^σ and the union of products of closed simplices \bar{V}_A^σ does not intersect.) We also require $U(S)$ to be contained in the $\epsilon/8$ -neighborhood of S . Then $\{\mathcal{U}_e\}$ satisfies $(*)$.

It remains to prove the statement at the beginning of the last paragraph. Take arbitrary triangulations $(T_e^0)_{e \in E(\Gamma)}$ of S^{d-1} with diameter smaller than $\epsilon/8$, we perturb them simplex by simplex to satisfy (**). The point is that every time we perturb a simplex away from \bar{V}_A^σ , we do it so slightly that no new unwanted intersection appears. The rest of the proof consists of technical details of this.

Below by “distance” we mean the restriction of D to S_π^{d-1} and its products. By a triangulation T of S^{d-1} we mean a homeomorphism $T : |K_T| \rightarrow S^{d-1}$, where K_T is a finite simplicial complex and $|K_T|$ its realization. Given $\mathcal{T} = (T_e)_{e \in E(\Gamma)}$ a tuple of triangulations of S^{d-1} , denote

$$\epsilon(\mathcal{T}) = \text{minimal distance between } \prod_{e \in I_{\sigma,A}} \bar{S}_e \text{ and } \bar{V}_A^\sigma,$$

where the “minimal” is taken over all σ, A , $(\bar{S}_e)_{e \in I_{\sigma,A}}$ such that $\bar{V}_A^\sigma \cap \prod_{e \in I_{\sigma,A}} \bar{S}_e = \emptyset$. There are only finitely many of them. And $\epsilon(\mathcal{T}) > 0$. Denote by $\mathcal{J}_1(\mathcal{T})$ (resp. $\mathcal{J}_0(\mathcal{T})$) the set of

tuples $(\sigma, A, (x_e)_{e \in I_{\sigma, A}})$ such that x_e is the image of a 0- or 1-simplex of T_e , exactly one of them (resp. none of them) being a 1-simplex, and $\bar{V}_A^\sigma \cap \prod_{e \in I_{\sigma, A}} x_e \neq \emptyset$. Then $\mathcal{J}_0(\mathcal{T}), \mathcal{J}_1(\mathcal{T})$ are finite.

Now take arbitrary triangulations $\mathcal{T}^0 = (T_e^0)_{e \in E(\Gamma)}$ such that the diameter of any simplex is smaller than $\epsilon/8$. Denote $\epsilon' = \epsilon/(8N(\mathcal{T}^0))$. For the rest of the paragraph, for all \mathcal{T} , define $\epsilon'(\mathcal{T}) = \min\{\epsilon', \epsilon(\mathcal{T})\}$. Take an element $(\sigma, A, (x_e)_{e \in I_{\sigma, A}}) \in \mathcal{J}_0(\mathcal{T})$. Since \bar{V}_A^σ cannot cover a neighborhood of $\prod_{e \in I_{\sigma, A}} x_e$, we can find $\prod_{e \in I_{\sigma, A}} x'_e$ in the $\epsilon'(\mathcal{T}^0)/2$ -neighborhood of it that does not meet \bar{V}_A^σ . For each e we can perturb the triangulation map T_e^0 in a neighborhood of $(T_e^0)^{-1}(x_e)$ to get T_e^1 , in such a way that $T_e^1((T_e^0)^{-1}(x_e)) = x'_e$, and the distance between old and new images of any point is less than $\epsilon'(\mathcal{T}^0)/|E(\Gamma)|$. For $e \notin I_{\sigma, A}$, define $T_e^1 = T_e^0$. Define $\mathcal{T}^1 = (T_e^1)_{e \in E(\Gamma)}$. Then $|\mathcal{J}_0(\mathcal{T}^1)| < |\mathcal{J}_0(\mathcal{T}^0)|$, $|\mathcal{J}_1(\mathcal{T}^1)| \leq |\mathcal{J}_1(\mathcal{T}^0)|$. We do this one by one for all elements of $\mathcal{J}_0(\mathcal{T}^0)$, getting new triangulations $\mathcal{T}^k = (T_e^k)_{e \in E(\Gamma)}$ by the end. Now take $(\sigma, A, (x_e)_{e \in I_{\sigma, A}}) \in \mathcal{J}_1(\mathcal{T}^k)$ where $e_* \in I_{\sigma, A}$ is such that x_{e_*} is the (image of a) 1-simplex. Let $p : \bar{V}_A^\sigma \subset (S^{d-1})^{I_{\sigma, A}} \rightarrow (S^{d-1})^{I_{\sigma, A} - e_*}$ be the projection forgetting the e_* -factor. Recall $f_{\sigma, A} : \bar{C}_{V(\Gamma_A)}(\mathbb{R}^d) \rightarrow (S^{d-1})^{I_{\sigma, A}}$ defined at the beginning of this proof. By possibly perturbing $(x_e)_{e \in I_{\sigma, A} - e_*}$ in the same way as above, we can assume that $(x_e)_{e \in I_{\sigma, A} - e_*}$ is a regular value of $p \circ f_{\sigma, A}$. So $\bar{V}_A^\sigma \cap p^{-1}((x_e)_{e \in I_{\sigma, A} - e_*}) \subset S^{d-1}$ is the smooth image of a manifold of dimension less than $d-2$. So we can perturb $T_{e_*}^k$ in a neighborhood of $(T_{e_*}^k)^{-1}(x_{e_*})$, fixing some neighborhoods of the 0-simplices (since $\mathcal{J}_0 = \emptyset$ now), to get a new triangulation $T_{e_*}^{k+1}$, so that $T_{e_*}^{k+1}((T_{e_*}^k)^{-1}(x_{e_*}))$ does not intersect $\bar{V}_A^\sigma \cap p^{-1}((x_e)_{e \in I_{\sigma, A} - e_*})$, and the new and old image of any point has distance less than $\epsilon'(\mathcal{T}^k)/|E(\Gamma)|$. For $e \in I_{\sigma, A}$ where x_e is not perturbed, or $e \notin I_{\sigma, A}$, define $T_e^{k+1} = T_e^k$. Define $\mathcal{T}^{k+1} = (T_e^{k+1})_{e \in E(\Gamma)}$. Then $|\mathcal{J}_1(\mathcal{T}^{k+1})| < |\mathcal{J}_1(\mathcal{T}^k)|$. Keep doing this one by one for all elements in $\mathcal{J}_1(\mathcal{T}^k)$. By the end we obtain a tuple of triangulations $\mathcal{T}^{|\mathcal{J}_0(\mathcal{T})|+|\mathcal{J}_1(\mathcal{T})|}$ satisfying (**). \square

4.2 Defining Kontsevich's characteristic classes

To “push forward” $\Omega_\Gamma(\pi)$ to a cohomology class on the base B , the Leray-Serre spectral sequence is a convenient tool to formulate it. We follow [6] for the definition of Leray-Serre spectral sequence. First we make a general definition.

4.2.1 Cohomology push-forward

Suppose B is a CW complex and $X \xrightarrow{\pi} B$ a fiber bundle with fiber F . Denote by B_p the p -skeleton of B and $X_p = \pi^{-1}(B_p)$. Suppose $k_0 > 0$ is such that $H^k(F) = 0$ for all $k > k_0$.

Then, for any integers n and $p < n - k_0$, in the Leray-Serre spectral sequence for $X_p \xrightarrow{\pi} B_p$, $E_2^{a,b} = 0$ for all $a + b = n$, so $H^n(X_p) = 0$.

The Leray-Serre spectral sequence for $X \xrightarrow{\pi} B$ tells us the following (cf. [6, Theorem 5.15]; note here we use the local coefficient version; cf. [14]): suppose $n \geq k_0 \in \mathbb{Z}$,

- $H^n(X)$ has a filtration by subgroups $F_p^n = \ker(H^n(X) \rightarrow H^n(X_{p-1}))$ and $E_\infty^{p,n-p} \approx F_p^n / F_{p+1}^n$;
- $E_2^{p,q} \approx H^p(B; H^q(F))$, where the latter is understood as cohomology with local coefficients;
- $d_r : E_r^{p,q} \rightarrow E_r^{p+r, q+1-r}$.

Suppose $r \geq 2$. Since $H^k(F) = 0$ for all $k > k_0$,

$$E_2^{n-k_0-r, k_0+r-1} \approx H^{n-k_0-r}(B, H^{k_0+r-1}(F)) = \{0\}.$$

Since $E_r^{p,q}$ is obtained from $E_2^{p,q}$ by taking subgroups and quotients, $E_r^{n-k_0-r, k_0+r-1} = \{0\}$ for all $r \geq 2$. Therefore, all the d_r 's mapping into $E_r^{n-k_0, k_0}$ vanish and $E_\infty^{n-k_0, k_0}$ is a subgroup of $E_2^{n-k_0, k_0} \approx H^{n-k_0}(B; H^{k_0}(F))$. Since $H^n(X_{n-k_0-1}) = 0$, $H^n(X) = F_{n-k_0}^n$. This identifies a map (which we denote by π_*)

$$\pi_* : H^n(X) \longrightarrow H^{n-k_0}(B; H^{k_0}(F)). \quad (6)$$

Definition 4.8. We call π_* *cohomology push-forward* of the fiber bundle $X \xrightarrow{\pi} B$.

By the naturality of Leray-Serre spectral sequence [6, page 537-538], π_* does not depend on the choice of the CW structure on B , and is natural: suppose $X' \xrightarrow{\pi'} B'$ is another fiber bundle with fiber F' such that $H^k(F') = 0$ for all $k > k_0$ and $(\tilde{f} : X' \rightarrow X, f : B' \rightarrow B)$ is a bundle map (so $H^{k_0}(F') \approx f^* H^{k_0}(F)$ as local systems over B'), then $f^* \circ \pi_* = \pi'_* \circ \tilde{f}^*$. Using CW approximation, π_* can be generalized to the case where B is an arbitrary space.

The above procedure can be generalized to the relative version: $X \xrightarrow{\pi} B$ has a subbundle $Y \xrightarrow{\pi} B$ with fiber $A \subset F$. Replacing F with the pair (F, A) and X with (X, Y) everywhere, everything goes through without change.

Remark 4.9. An explicit description of π_* can be obtained by carefully unwinding the definition; we follow [6] for the construction of Leray-Serre spectral sequences and specifically, what we do below comes from the diagram on page 526 and Φ in the proof of Theorem 5.3 in [6]. Denote $p_0 = n - k_0$ for simplicity. Given an element $\sigma \in H^n(X)$, first restrict it to $H^n(X_{p_0})$; the image will lie in the image of $H^n(X_{p_0}, X_{p_0-1})$. Take such a preimage, say σ' .

For each p_0 -cell $e : (D^{p_0}, \partial D^{p_0}) \rightarrow (X_{p_0}, X_{p_0-1})$ (D^{p_0} being the standard p_0 -dimensional ball), $e^* \sigma' \in H^n(e^*(X), (e|_{\partial D^{p_0}})^*(X))$. Since $e^* \pi$ is trivializable, Künneth formula gives

$$H^n(e^*(X), (e|_{\partial D^{p_0}})^*(X)) \approx H^{p_0}(D^{p_0}, \partial D^{p_0}) \otimes H^{k_0}(F_b),$$

where $F_b = \pi^{-1}(e(b))$ is the fiber over an arbitrarily fixed point $b \in \mathring{D}^{p_0}$. Other fibers over D^{p_0} are identified with F_b via the trivialization. Notice that the Künneth isomorphism does not depend on the choice of the trivialization of $e^* \pi$: the Künneth map is determined by the projection map $e^*(X) \rightarrow F_b$ (note that ∂D^{p_0} is not involved here); any two trivializations give homotopic projection maps since D^{p_0} is contractible. Let us denote by $\sigma(e) \in H^{k_0}(F_b)$ the image of $e^* \sigma'$ under the Künneth map, where $H^{p_0}(D^{p_0}, \partial D^{p_0}) \approx R$ identified using the canonical orientation on D^{p_0} . Then $\{e \rightarrow \sigma(e)\}_e$, where e ranges through all p_0 -cells of B , gives a cellular cochain on B with coefficients in the local system $H^{k_0}(F)$. It is a cocycle and represents a cohomology class in $H^{p_0}(B; H^{k_0}(F))$, which is $\pi_*(\sigma)$. The relative version is similar.

4.2.2 Defining Kontsevich's characteristic classes

Applying the construction above to the fiber bundle $(X_\Gamma(\pi), S(\pi)) \xrightarrow{\pi_X} B$ with $k_0 = d|V(\Gamma)|$, we get

$$\pi_{X*} : H^{|E(\Gamma)|(d-1)}(X_\Gamma(\pi), S(\pi)) \longrightarrow H^{|E(\Gamma)|(d-1)-d|V(\Gamma)|}(B; H^{d|V(\Gamma)|}(X_\Gamma, S)). \quad (7)$$

The map $\rho : H^{d|V(\Gamma)|}(X_\Gamma, S) \rightarrow R$ in Definition 3.25 induces the corresponding map of local systems on B . So we get an induced map

$$H^*(B; H^{d|V(\Gamma)|}(X_\Gamma, S)) \longrightarrow H^*(B; R). \quad (8)$$

Definition 4.10. Define $K_{\Gamma, \pi, F} \in H^{|E(\Gamma)|(d-1)-d|V(\Gamma)|}(B; R)$, to be the image of $\Omega_\Gamma(\pi)$ under (7) and (8).

The corollary below is a direct consequence of Proposition 4.5 and the naturality of cohomology push-forward.

Corollary 4.11 Under the assumptions of Theorem 1.2, $K_{\Gamma, \pi', F'} = h_B^* K_{\Gamma, \pi'', F''}$.

5 Equivalence with the original definition

In this section the coefficient ring $R = \mathbb{R}$. All open covers are assumed to be locally finite. The goal of this section is to prove the following

Proposition 5.1 Suppose $(E \xrightarrow{\pi} B, F)$ is a framed smooth (M, ∞) bundle over a smooth manifold B , then $K_{\Gamma, \pi, F}$ defined above agrees, up to scaling by a constant depending only on Γ , with the usual definition of Kontsevich's characteristic classes for $(E \xrightarrow{\pi} B, F)$; see, for example, [18] for definition.

5.1 Čech to de Rham preliminary

First we state some general facts translating Čech to de Rham cohomology. Let Y be a smooth manifold. Denote by \mathcal{A}_Y^q the sheaf of differential q -form germs on Y , \mathcal{Z}_Y^q the subsheaf of closed q -form germs and $A^q(Y)$ the space of global q -forms on Y . Let \mathcal{U} be an open cover of Y . Let $\underline{l} = \{l_U : Y \rightarrow \mathbb{R}^{\geq 0}\}_{U \in \mathcal{U}}$ be a partition of unity subordinate to \mathcal{U} . For any $p, q \in \mathbb{Z}^{\geq 0}$ define

$$h_{p,q}^l : \check{C}_{\mathcal{U}}^p(Y; \mathcal{A}_Y^q) \longrightarrow \check{C}_{\mathcal{U}}^{p-1}(Y; \mathcal{A}_Y^{q+1})$$

$$h_{p,q}^l(\sigma)(U_0 \cap \dots \cap U_{p-1}) = (-1)^p \sum_{U \in \mathcal{U}} d(l_U \cdot \sigma(U \cap U_0 \cap \dots \cap U_{p-1})).$$

By $l_U \cdot \sigma(U \cap \dots \cap U_{p-1})$ we mean a form on $U_0 \cap \dots \cap U_{p-1}$ which is given by this formula on $U \cap U_0 \cap \dots \cap U_{p-1}$ and 0 elsewhere; it is smooth since l_U vanishes in a neighborhood of ∂U . Clearly $\text{image}(h_{p,q}^l) \subset \check{C}_{\mathcal{U}}^{p-1}(Y; \mathcal{Z}_Y^{q+1})$. For each p , define

$$h^l : \check{C}_{\mathcal{U}}^p(Y; \mathbb{R}) \longrightarrow \check{C}_{\mathcal{U}}^0(Y; \mathcal{Z}_Y^p), \quad h^l(\sigma) = (-1)^p (h_{1,p-1}^l \circ h_{2,p-2}^l \dots \circ h_{p,0}^l)(\sigma).$$

By [1, Proposition 9.8], if $\sigma \in \check{C}_{\mathcal{U}}^p(Y; \mathbb{R})$ is a Čech cocycle, then $h^l(\sigma)$ is a global closed form; if \mathcal{U} is such that any finite intersection of elements has trivial cohomology, then $\sigma \rightarrow h^l(\sigma)$ induces the canonical isomorphism between $\check{H}^p(Y; \mathbb{R})$ and $H_{\text{de Rham}}^p(Y)$. If Φ is a family of supports on Y , the arguments in [1, Section 8] still go through if all differential forms are assumed to have supports in Φ (i.e., $\check{C}_{\mathcal{U}}^p$ is replaced with its subspace of Φ -supported cochains $\check{C}_{\mathcal{U}, \Phi}^p(Y; \mathcal{A}_Y^q)$); so h^l still induces the canonical isomorphism between $\check{H}_{\Phi}^p(Y; \mathbb{R})$ and $H_{\text{de Rham}, \Phi}^p(Y)$, where $H_{\text{de Rham}, \Phi}^p(Y)$ is the cohomology of the cochain complex of Φ -supported differential forms on Y .

Suppose $\mathcal{U}, \mathcal{U}'$ are two open covers of Y and $\mu : \mathcal{U} \rightarrow \mathcal{U}'$ is a refinement. Let $\underline{l} = \{l_U\}_{U \in \mathcal{U}}$ be a partition of unity subordinate to \mathcal{U} , then

$$\mu_* \underline{l} := \left\{ l_{U'} := \sum_{\substack{U \in \mathcal{U} \\ \mu(U) = U'}} l_U : Y \longrightarrow \mathbb{R} \right\}_{U' \in \mathcal{U}'}$$

is a partition of unity subordinate to \mathcal{U}' . It is easy to check that $h_{p,q}^{\mu_* \underline{l}}(\mu^* \sigma) = \mu^*(h_{p,q}^{\underline{l}}(\sigma))$ for all p, q and $\sigma \in \check{C}_{\mathcal{U}'}^p(Y; \mathcal{A}_Y^q)$.

Define cup product

$$\cup : \check{C}_{\mathcal{U}}^{p_1}(Y; \mathcal{A}_Y^{q_1}) \otimes \check{C}_{\mathcal{U}}^{p_2}(Y; \mathcal{A}_Y^{q_2}) \longrightarrow \check{C}_{\mathcal{U}}^{p_1+p_2}(Y; \mathcal{A}_Y^{q_1+q_2})$$

$$(\sigma_1 \cup \sigma_2)(U_0 \cap \dots \cap U_{p_1+p_2}) = (-1)^{q_1 p_2} \sigma_1(U_0 \cap \dots \cap U_{p_1}) \wedge \sigma_2(U_{p_1} \cap \dots \cap U_{p_1+p_2}),$$

where the two forms on the RHS are restricted to $U_0 \cap \dots \cap U_{p_1+p_2}$. For simplicity we omit the notation for restriction; same below. When restricted to $\check{C}_{\mathcal{U}}^{p_1}(Y; \mathcal{Z}_Y^0) \otimes \check{C}_{\mathcal{U}}^{p_2}(Y; \mathcal{Z}_Y^0)$ this is the usual cup product for Čech cochains.

Lemma 5.2 Let $Y = Y_1 \times \dots \times Y_m$ be a product of smooth manifolds. Denote by $\pi_i : Y \rightarrow Y_i$ the projection to the i -th factor. For every $i = 1, \dots, m$, let \mathcal{U}_i be an open cover of Y_i and $\underline{l}_i = \{l_U\}_{U \in \mathcal{U}_i}$ be a partition of unity. Denote $\mathcal{U} = \mathcal{U}_1 \times \dots \times \mathcal{U}_m$ the product open cover of Y , then

$$\underline{l} = \{l_{U_1 \times \dots \times U_m} := (l_{U_1} \circ \pi_1) \cdot \dots \cdot (l_{U_m} \circ \pi_m) : Y \longrightarrow \mathbb{R}\}_{U_1 \in \mathcal{U}_1, \dots, U_m \in \mathcal{U}_m}$$

is a partition of unity of Y subordinate to \mathcal{U} . Let $p = \sum_{i=1}^m p_i$, $q = \sum_{i=1}^m q_i$ be non-negative integers. Let $\sigma_i \in \check{C}_{\mathcal{U}_i}^{p_i}(Y_i; \mathcal{Z}_{Y_i}^{q_i})$ be Čech cocycles. Define

$$\sigma = \pi_1^* \sigma_1 \cup \dots \cup \pi_m^* \sigma_m \in \check{C}_{\mathcal{U}}^p(Y; \mathcal{Z}_Y^q).$$

Then,

1. if $m = 2$ (this is only for simplicity),

$$h_{p,q}^{\underline{l}}(\sigma) = \begin{cases} \pi_1^* h_{p_1, q_1}^{l_1}(\sigma_1) \cup \pi_2^*(\sigma_2), & \text{if } p_1 > 0, \\ \pi_1^* \sigma_1 \cup \pi_2^* h_{p_2, q_2}^{l_2}(\sigma_2), & \text{if } p_1 = 0; \end{cases}$$

2. if $q_i = 0$ for all i , $h^{\underline{l}}(\sigma) = \pi_1^* h^{l_1}(\sigma_1) \wedge \dots \wedge \pi_m^* h^{l_m}(\sigma_m)$.

Proof. This is direct computation. For (1), when $p_1 > 0$,

$$\begin{aligned} & h_{p,q}^{\underline{l}}(\sigma)((U_1^0 \times U_2^0) \cap \dots \cap (U_1^{p-1} \times U_2^{p-1})) \\ &= (-1)^p \sum_{U_1 \in \mathcal{U}_1, U_2 \in \mathcal{U}_2} ((\pi_1^* dl_{U_1})(l_{U_2} \circ \pi_2) + (l_{U_1} \circ \pi_1)(\pi_2^* dl_{U_2})) \wedge \\ & \quad (-1)^{q_1 p_2} \pi_1^* \sigma_1(U_1 \cap U_1^0 \cap \dots \cap U_1^{p_1-1}) \wedge \pi_2^* \sigma_2(U_2^{p_1-1} \cap \dots \cap U_2^{p-1}) \\ &= (-1)^{p+q_1 p_2} \left(\left(\sum_{U_1} (\pi_1^* dl_{U_1}) \wedge \pi_1^* \sigma_1(U_1 \cap \dots \cap U_1^{p_1-1}) \right) \wedge \left(\sum_{U_2} (l_{U_2} \circ \pi_2) \pi_2^* \sigma_2(U_2^{p_1-1} \cap \dots \cap U_2^{p-1}) \right) \right) \\ & \quad + \left(\sum_{U_1} (-1)^{q_1} (l_{U_1} \circ \pi_1) \pi_1^* \sigma_1(U_1 \cap \dots \cap U_1^{p_1-1}) \right) \wedge \underbrace{\left(\sum_{U_2} (\pi_2^* dl_{U_2}) \wedge \pi_2^* \sigma_2(U_2^{p_1-1} \cap \dots \cap U_2^{p-1}) \right)}_{=0} \end{aligned}$$

$$\begin{aligned}
&= (-1)^{p+p_1+q_1p_2} \pi_1^* h_{p_1, q_1}^{l_1}(\sigma_1)(U_1^0 \cap \dots \cap U_1^{p_1-1}) \wedge \pi_2^* \sigma_2(U_2^{p_1-1} \cap \dots \cap U_2^{p-1}) \\
&= (\pi_1^* h_{p_1, q_1}^{l_1}(\sigma_1) \cup \pi_2^* \sigma_2)((U_1^0 \times U_2^0) \cap \dots \cap (U_1^{p-1} \times U_2^{p-1})).
\end{aligned}$$

When $p_1 = 0$,

$$\begin{aligned}
&h_{p,q}^l(\sigma)((U_1^0 \times U_2^0) \cap \dots \cap (U_1^{p-1} \times U_2^{p-1})) \\
&= (-1)^{p+q_1p_2} \sum_{U_1 \in \mathcal{U}_1, U_2 \in \mathcal{U}_2} ((\pi_1^* dl_{U_1})(l_{U_2} \circ \pi_2) + (l_{U_1} \circ \pi_1)(\pi_2^* dl_{U_2})) \wedge \pi_1^* \sigma_1(U_1) \wedge \pi_2^* \sigma_2(U_2 \cap U_2^0 \cap \dots \cap U_2^{p-1}) \\
&= (-1)^{p+q_1p_2} \left(\underbrace{\left(\sum_{U_1} (\pi_1^* dl_{U_1}) \wedge \pi_1^* \sigma_1(Y_1) \right)}_{=0} \wedge \left(\sum_{U_2} (l_{U_2} \circ \pi_2) \pi_2^* \sigma_2(U_2 \cap U_2^0 \cap \dots \cap U_2^{p-1}) \right) \right) \\
&\quad + \left(\sum_{U_1} (-1)^{q_1} (l_{U_1} \circ \pi_1) \pi_1^* \sigma_1(Y_1) \right) \wedge \left(\sum_{U_2} (\pi_2^* dl_{U_2}) \wedge \pi_2^* \sigma_2(U_2 \cap U_2^0 \cap \dots \cap U_2^{p-1}) \right) \\
&= (-1)^{p+q_1+p_2+q_1p_2} \pi_1^* \sigma_1(Y_1) \wedge \pi_2^* h_{p_2, q_2}^{l_2}(\sigma_2)(U_2 \cap U_2^0 \cap \dots \cap U_2^{p-1}) \\
&= (\pi_1^* \sigma_1 \cup \pi_2^* h_{p_2, q_2}^{l_2}(\sigma_2))((U_1^0 \times U_2^0) \cap \dots \cap (U_1^{p-1} \times U_2^{p-1}));
\end{aligned}$$

notice that we can write $\sigma_1(Y_1)$ because σ_1 is a degree 0 Čech cocycle.

For (2), in the case $m = 2$,

$$\begin{aligned}
h^l(\sigma) &= (-1)^p (h_{1,p-1}^l \circ \dots \circ h_{p,0}^l)(\pi_1^* \sigma_1 \cup \pi_2^* \sigma_2) \\
&= (-1)^p (h_{1,p-1}^l \circ \dots \circ h_{p_2,p-p_2}^l)(\pi_1^*(h_{1,p_1-1}^{l_1} \circ \dots \circ h_{p_1,0}^{l_1})(\sigma_1) \cup \pi_2^* \sigma_2) \\
&= (-1)^{p+p_1} (h_{1,p-1}^l \circ \dots \circ h_{p_2,p-p_2}^l)(\pi_1^* h^{l_1}(\sigma_1) \cup \pi_2^* \sigma_2) \\
&= (-1)^{p+p_1} (\pi_1^* h^{l_1}(\sigma_1) \cup \pi_2^*(h_{1,p_2-1}^{l_2} \circ \dots \circ h_{p_2,0}^{l_2})\sigma_2) \\
&= \pi_1^* h^{l_1}(\sigma_1) \cup \pi_2^* h^{l_2}(\sigma_2) = \pi_1^* h^{l_1}(\sigma_1) \wedge \pi_2^* h^{l_2}(\sigma_2).
\end{aligned}$$

The general case follows by induction. □

We next check that h is natural. Let $f : X \rightarrow Y$ be a smooth map between smooth manifolds. For an open cover \mathcal{U} on Y with partition of unity $\underline{l} = \{l_U\}_U$, $f^*(\mathcal{U}) := \{f^{-1}(U)\}_{U \in \mathcal{U}}$ is an open cover of X and $f^*\underline{l} := \{l_{f^{-1}(U)} := l_U \circ f : X \rightarrow \mathbb{R}^{\geq 0}\}_{U \in \mathcal{U}}$ is a partition of unity. We have the pull-back map $f^* : \check{C}_{\mathcal{U}}^p(Y; \mathcal{Z}_Y^q) \rightarrow \check{C}_{f^*\mathcal{U}}^p(X; \mathcal{Z}_X^q)$, and

$$\begin{aligned}
(f^* h^l \sigma)(f^{-1}(U_0) \cap \dots \cap f^{-1}(U_{p-1})) &= \sum_{U \in \mathcal{U}} (-1)^p d((l_U \circ f) \wedge f^*(\sigma(U \cap U_0 \cap \dots \cap U_{p-1}))) \\
&= \sum_{U \in \mathcal{U}} (-1)^p d(l_{f^{-1}(U)} \cdot (f^* \sigma)(f^{-1}(U) \cap f^{-1}(U_0) \cap \dots \cap f^{-1}(U_{p-1})))
\end{aligned}$$

$$= (h^{f^* \underline{l}} f^* \sigma)(f^{-1}(U_0) \cap \dots \cap f^{-1}(U_{p-1})).$$

Notice that everything in this subsection works for cohomology with supports as well.

5.2 Proof of Proposition 5.1

Let \mathcal{U}' be an open cover of $\overline{C}_2(\pi)/\sim_F$ such that there exists $\check{\omega} \in \check{C}_{\mathcal{U}'}^{d-1}(\overline{C}_2(\pi)/\sim_F)$ a cocycle representative of Ω . We will make a careful choice of an open cover and a partition of unity on $\overline{C}_2(\pi)$. Note the following commutative square, where \hat{i}, ι are inclusion maps,

$$\begin{array}{ccc} \partial \overline{C}_2(\pi) & \xrightarrow{\hat{i}} & \overline{C}_2(\pi) \\ \downarrow F & & \downarrow q \\ S^{d-1} & \xrightarrow{\iota} & \overline{C}_2(\pi)/\sim_F. \end{array}$$

Lemma 5.3 There exist an open cover \mathcal{U} of $\overline{C}_2(\pi)$ such that all intersections of its elements are contractible, a refining map $\mu : \mathcal{U} \rightarrow q^* \mathcal{U}'$, and partitions of unity \underline{l} on $\overline{C}_2(\pi)$ subordinate to \mathcal{U} , \underline{l}_S on S^{d-1} subordinate to $\iota^* \mathcal{U}'$, such that $\hat{i}^*(\mu_* \underline{l}) = F^* \underline{l}_S$ (they are both subordinate to $\hat{i}^* q^* \mathcal{U}' = F^* \iota^* \mathcal{U}'$ on $\partial \overline{C}_2(\pi)$).

Proof. By the way F is defined, it can be extended to a smooth map $F : N_\partial \rightarrow S^{d-1}$, where $N_\partial \subset \overline{C}_2(\pi)$ is a neighborhood of $\partial \overline{C}_2(\pi)$. Still denote by $\hat{i} : N_\partial \rightarrow \overline{C}_2(\pi)$ the inclusion. Let \underline{l}_S be a partition of unity on S^{d-1} subordinate to $\iota^* \mathcal{U}'$. Then $F^* \underline{l}_S =: \{F^* \underline{l}_S(U)\}_{U \in \mathcal{U}'}$, where $F^* \underline{l}_S(U)$ is supported on $q^{-1}(U) \cap N_\partial$, is a partition of unity on N_∂ subordinate to $\hat{i}^* q^* \mathcal{U}'$. Our next goal is to find a partition of unity \underline{l}' on $\overline{C}_2(\pi)$ subordinate to $q^* \mathcal{U}'$, such that $\hat{i}^* \underline{l}' = F^* \underline{l}_S$.

We can find a smooth function $g : \overline{C}_2(\pi) \rightarrow \mathbb{R}^{\geq 0}$ supported in N_∂ such that $g|_{\partial \overline{C}_2(\pi)} \equiv 1$: let $K \subset C_2(\pi)$ be a compact subset containing $C_2(\pi) - N_\partial$, and let $N'_\partial \subset \overline{C}_2(\pi) - K$ be a neighborhood of $\partial \overline{C}_2(\pi)$; let $\{V_i \subset \overline{C}_2(\pi) - N'_\partial\}_{i \in I}$ be an open cover of K ; let $\{g_i\}_{i \in I} \sqcup \{g'\}$, where g_i is supported in V_i and g' is supported in $C_2(\pi) - K$, be a partition of unity on $C_2(\pi)$ subordinate to $\{V_i\}_i \sqcup \{C_2(\pi) - K\}$, then $(\sum_i g_i)|_K \equiv 1$ and $(\sum_i g_i)|_{N'_\partial \cap C_2(\pi)} \equiv 0$; so $g := 1 - \sum_i g_i$, extended by 0 to $\partial \overline{C}_2(\pi)$, satisfies the requirement. For each $U \in \mathcal{U}'$, define $h_U = g \cdot F^* \underline{l}_S(U) : N_\partial \rightarrow \mathbb{R}$, then it can be smoothly extended to the entire $\overline{C}_2(\pi)$, taking value 0 out of N_∂ ; so $h_U|_{\partial \overline{C}_2(\pi)} = F^* \underline{l}_S(U)|_{\partial \overline{C}_2(\pi)}$, h_U is supported in $q^{-1}(U)$ and $(\sum_U h_U)|_{N'_\partial} \equiv 1$.

Let $K' \subset C_2(\pi)$ be a compact subset containing $C_2(\pi) - N'_\partial$. Let $\{G_U \subset q^{-1}(U) \cap C_2(\pi)\}_{U \in \mathcal{U}'}$ be compact subsets that still cover K' . For each $U \in \mathcal{U}'$, take $\phi_U : \overline{C}_2(\pi) \rightarrow$

$\mathbb{R}^{\geq 0}$ that is supported in $q^{-1}(U) \cap C_2(\pi)$ and $\phi_U|_{G_U} \equiv 1$. Then $\sum_{U \in \mathcal{U}'} (\phi_U + h_U)$, as a function on $\overline{C}_2(\pi)$, is positive everywhere and equals to 1 on $\partial \overline{C}_2(\pi)$. Define $l'_U = (\phi_U + h_U) / (\sum_U (\phi_U + h_U))$. Then $l' := \{l'_U\}_{U \in \mathcal{U}'}$ is a partition of unity as required.

Next we define \mathcal{U} . Fix a Riemannian metric on $\overline{C}_2(\pi)$ (which is a smooth manifold with boundary and corners). For every $U \in \mathcal{U}'$, let $\{V_i^U \subset q^{-1}(U) \subset \overline{C}_2(\pi)\}_{i \in I_U}$ be a finite collection of geodesically convex open subsets, such that $\text{supp}(l'_U) \subset \bigcup_{i \in I_U} V_i^U$. Then $\mathcal{U} := \{V_i^U\}_{U \in \mathcal{U}', i \in I_U}$ is an open cover of $\overline{C}_2(\pi)$ whose intersections are all contractible. Define $\mu : \mathcal{U} \rightarrow q^*\mathcal{U}'$, $\mu(V_i^U) = q^{-1}(U)$; it is a refinement map.

Now, we construct a partition of unity \underline{l} on $\overline{C}_2(\pi)$ subordinate to \mathcal{U} such that $\mu_* \underline{l} = l'$. Using the same argument used to find the g_i s above, for every $U \in \mathcal{U}'$ we can find smooth functions $\psi_i^U : \overline{C}_2(\pi) \rightarrow \mathbb{R}^{\geq 0}$ supported in V_i^U , for every $i \in I_U$, such that $(\sum_{i \in I_U} \psi_i^U)|_{\text{supp}(l'_U)} \equiv 1$. Define $l_i^U = \psi_i^U \cdot l'_U$. Then $\sum_{i \in I_U} l_i^U = l'_U$. So $\underline{l} := \{l_i^U\}_{U \in \mathcal{U}', i \in I_U}$ is a partition of unity as required. \square

Corollary 5.4 $h^l(\mu^* q^* \tilde{\omega})$ is a closed $(d-1)$ -form on $\overline{C}_2(\pi)$ such that $\hat{\iota}^* h^l(\mu^* q^* \tilde{\omega}) = F^* \alpha$ for some closed form $\alpha \in A^{d-1}(S^{d-1})$, $\int_{S^{d-1}} \alpha = 1$. In other words, $h^l(\mu^* q^* \tilde{\omega})$ is a propagator.

Proof. That it is a closed $(d-1)$ -form is clear. And

$$\hat{\iota}^* h^l(\mu^* q^* \tilde{\omega}) = \hat{\iota}^* h^{\mu_* l}(\mu^* q^* \tilde{\omega}) = h^{\hat{\iota}^* \mu_* l}(\hat{\iota}^* \mu^* q^* \tilde{\omega}) = h^{F^* l_S}(F^* \iota^* \tilde{\omega}) = F^* h^{l_S}(\iota^* \tilde{\omega}).$$

Since $[\tilde{\omega}] = \Omega \in H^{d-1}(\overline{C}_2(\pi)/\sim_F)$ and h^{l_S} induces the canonical isomorphism between Čech and de Rham cohomology, $[h^{l_S}(\iota^* \tilde{\omega})] \in H^{d-1}(S^{d-1}; \mathbb{R})$ is the Poincaré dual of the point class by the definition of Ω . So $\int_{S^{d-1}} h^{l_S}(\iota^* \tilde{\omega}) = 1$. \square

Now, let $\{\mathcal{U}'_e\}_{e \in E(\Gamma)}$ be a collection of open covers of $\overline{C}_2(\pi)/\sim_F$ given by Lemma 4.7. For every e , applying the above argument with \mathcal{U}' replaced by \mathcal{U}'_e , we get an open cover \mathcal{U}_e with refinement map $\mu_e : \mathcal{U}_e \rightarrow q^*\mathcal{U}'_e$ and partition of unity \underline{l}^e on $\overline{C}_2(\pi)$ subordinate to \mathcal{U}_e , such that $h^{\underline{l}^e}(\mu^* q^* \tilde{\omega})$ is a propagator. Denote $\omega_e = h^{\underline{l}^e}(\mu^* q^* \tilde{\omega})$. Denote by $\text{pr}_e : \overline{C}_2(\pi)^{E(\Gamma)} \rightarrow \overline{C}_2(\pi)$ the projection to the e -th factor. (Recall $\overline{C}_2^{E(\Gamma)}(\pi)$ is the fiber product and $\overline{C}_2(\pi)^{E(\Gamma)}$ is the direct product of the total space.) Denote

$$\tilde{\mathcal{U}} := \text{pr}_{e_1}^* \mathcal{U}_{e_1} \times \dots \times \text{pr}_{e_{|E(\Gamma)|}}^* \mathcal{U}_{e_{|E(\Gamma)|}}^\Gamma$$

the product open cover on $\overline{C}_2(\pi)^{E(\Gamma)}$. Define (that it is supported away from $S(\pi)$ follows from the choice of $\{\mathcal{U}'_e\}$ given by Lemma 4.7)

$$\tilde{\omega}'_\Gamma := \text{pr}_{e_1}^* \mu^* q^* \tilde{\omega} \cup \text{pr}_{e_2}^* \mu^* q^* \tilde{\omega} \cup \dots \cup \text{pr}_{e_{|E(\Gamma)|}}^* \mu^* q^* \tilde{\omega} \in \check{C}_{\tilde{\mathcal{U}}}^{(|E(\Gamma)|)(d-1)}(\overline{C}_2(\pi)^{E(\Gamma)}, S(\pi));$$

its restriction to X_Γ represents the class $\Omega_\Gamma(\pi) \in H^{|E(\Gamma)|(d-1)}(X_\Gamma(\pi), S(\pi))$. Define

$$\omega_\Gamma = \text{pr}_{e_1^\Gamma}^* \omega_{e_1^\Gamma} \wedge \dots \wedge \text{pr}_{e_{|E(\Gamma)|}^\Gamma}^* \omega_{e_{|E(\Gamma)|}^\Gamma} \in A^{|E(\Gamma)|(d-1)}(\overline{C}_2(\pi)^{E(\pi)});$$

then the push-forward of $\omega_\Gamma|_{C_\Gamma(\pi)}$ to B represents Kontsevich's class in the usual definition. By Lemma 5.2, $\omega_\Gamma = h^{\tilde{l}} \tilde{\omega}'_\Gamma$, where \tilde{l} is the partition of unity on $\overline{C}_2(\pi)^{E(\Gamma)}$ subordinate to $\tilde{\mathcal{U}}$ given by taking “product” of the \tilde{l}^e s as in Lemma 5.2. Below we denote the restriction of ω_Γ to $\overline{C}_2^{E(\Gamma)}(\pi)$ still by ω_Γ .

Fix a triangulation on B and denote by B_p the p -skeleton of B with respect to this triangulation. Denote

$$p_0 = |E(\Gamma)|(d-1) - d|V(\Gamma)| = \deg \Omega_\Gamma(\pi) - \dim X_\Gamma.$$

Recall that $f = (f_e)_{e \in E(\Gamma)} : \overline{C}_{V(\Gamma)}(\pi) \rightarrow \overline{C}_2^{E(\Gamma)}(\pi)$ is the forgetful map, and ϕ is the factor-permuting action of $\tilde{S}_{E(\Gamma)}$ on $\overline{C}_2^{E(\Gamma)}(\pi)$. Denote by

$$\pi_V : \overline{C}_{V(\Gamma)}(\pi) \longrightarrow B, \quad \hat{\pi} : \overline{C}_2^{E(\Gamma)}(\pi) \longrightarrow B, \quad \pi_X : X_\Gamma(\pi) \longrightarrow B$$

the bundle projection maps (the first two were both denoted by π ; here we want to distinguish them to avoid confusion). Define

$$\mathring{X}_\Gamma = \bigcup_{\sigma \in \tilde{S}_{E(\Gamma)}} \phi(\sigma)(C_\Gamma(M, \infty)), \quad T = X_\Gamma - \mathring{X}_\Gamma \subset \overline{C}_2(M, \infty),$$

and denote $\mathring{X}_\Gamma \pi, T(\pi) \subset \overline{C}_2^{E(\Gamma)}(\pi)$ the bundle version of them. For all p , denote $X_p = \hat{\pi}^{-1}(B_p) \cap X_\Gamma(\pi)$.

Define $s(\sigma) = (d-1)\text{sgn}(\sigma) + d\text{sgn}'(\sigma)$. For a differential form $\alpha \in A^m(\overline{C}_2^{E(\Gamma)}(\pi))$, define $\pi_*^s(\alpha)$ to be the degree $(m - d|V(\Gamma)|)$ simplicial cochain on B that sends a dimension $m - d|V(\Gamma)|$ simplex Δ to

$$\sum_{\sigma \in \tilde{S}_{E(\Gamma)}} (-1)^{s(\sigma)} \int_{\pi_V^{-1}(\Delta)} (\phi(\sigma) \circ f)^* \alpha = \sum_{\sigma \in \tilde{S}_{E(\Gamma)}} (-1)^{s(\sigma)} \int_{\pi_V^{-1}(\Delta)} f^* \phi(\sigma)^* \alpha.$$

Then, since $\phi(\sigma)^* \omega_\Gamma = (-1)^s \omega_\Gamma$, $[\pi_*^s \omega_\Gamma] \in H^{p_0}(B)$ is $2^{|E(\Gamma)|} |E(\Gamma)|!$ times Kontsevich's class in the usual definition.

Recall $\tilde{X}_\Gamma(\pi)$ is the bundle version of Definition 3.22, and $\tilde{f} : \tilde{X}_\Gamma(\pi) \rightarrow \overline{C}_2^{E(\Gamma)}(\pi)$. In Definition 3.22, by choosing a collar neighborhood when gluing the copies of $C'_\Gamma(M, \infty)$,

\tilde{X}_Γ can be given a smooth structure so that $\tilde{f} : \tilde{X}_\Gamma(\pi) \rightarrow \overline{C}_2^{E(\Gamma)}(\pi)$ is piecewise smooth (smooth away from $\tilde{f}^{-1}(T(\pi))$); so pulling back a differential form by \tilde{f} is still well defined (the result would be a piecewise-smooth form), and the usual differential form push-forward $(\pi_{\tilde{X}})_*$ is also well defined for these forms (by integrating along each piece of the fiber and summing up). It follows from definition that $\pi_*^s(\alpha) = (\pi_{\tilde{X}})_* \tilde{f}^* \alpha$,

Lemma 5.5 Suppose $\alpha_1, \alpha_2 \in A^*(\overline{C}_2^{E(\Gamma)}(\pi))$ are such that there exists $\tilde{\alpha} \in A^{*-1}(\overline{C}_2^{E(\Gamma)}(\pi))$ supported away from $S(\pi)$ and $d\tilde{\alpha} = \alpha_1 - \alpha_2$, then $\pi_*^s \alpha_1 - \pi_*^s \alpha_2$ is a coboundary.

Proof. For a simplex Δ in B ,

$$\begin{aligned} \sum_{\sigma \in \tilde{S}_{E(\Gamma)}} (-1)^{s(\sigma)} \int_{\pi_V^{-1}(\Delta)} (\phi(\sigma) \circ f)^*(\alpha_1 - \alpha_2) &= \sum_{\sigma \in \tilde{S}_{E(\Gamma)}} (-1)^{s(\sigma)} \int_{\partial \pi_V^{-1}(\Delta)} (\phi(\sigma) \circ f)^* \tilde{\alpha} \\ &= \sum_{\sigma \in \tilde{S}_{E(\Gamma)}} (-1)^{s(\sigma)} \int_{\pi_V^{-1}(\partial \Delta)} (\phi(\sigma) \circ f)^* \tilde{\alpha} + \sum_{\sigma \in \tilde{S}_{E(\Gamma)}} (-1)^{s(\sigma)} \sum_A \int_{\pi_V^{-1}(\Delta) \cap \mathcal{S}_A} (\phi(\sigma) \circ f)^* \tilde{\alpha}, \end{aligned}$$

where $A \subset V(\Gamma) \cup \{\infty\}$, $|A| \geq 2$. The second term vanishes: for A of type 1, $f(\mathcal{S}_A)$ has smaller dimension; for A of type 2 or 4, they cancel with each other when summed over σ ; for A of type 3, because $\tilde{\alpha}$ vanishes on $S(\pi)$ by assumption. So,

$$(\pi_*^s \alpha_1 - \pi_*^s \alpha_2)(\Delta) = (\pi_*^s \tilde{\alpha})(\partial \Delta) = (\delta \pi_*^s \tilde{\alpha})(\Delta).$$

□

By Lemma 3.18, we can find an open cover $\hat{\mathcal{U}}$ of $\overline{C}_2(\pi)^{E(\Gamma)}$ refining $\tilde{\mathcal{U}}$, such that there exists a neighborhood N of $(T(\pi) \cap X_{p_0}) \cup X_{p_0-1}$ in $\overline{C}_2(\pi)^{E(\Gamma)}$, satisfying

$$U_0 \cap \dots \cap U_{|E(\Gamma)|(d-1)} \cap N = \emptyset, \quad \forall U_0 \neq \dots \neq U_{|E(\Gamma)|(d-1)} \in \hat{\mathcal{U}}.$$

Let $\hat{\mu} : \hat{\mathcal{U}} \rightarrow \tilde{\mathcal{U}}$ be a refinement map. Then $\hat{\mu}^* \tilde{\omega}'_\Gamma$ is supported away from N . Define $\tilde{\omega}_\Gamma = (\hat{\mu}^* \tilde{\omega}'_\Gamma)|_{X_\Gamma(\pi)}$.

Let \hat{l} be a partition of unity on $\overline{C}_2(\pi)^{E(\Gamma)}$ subordinate to $\hat{\mathcal{U}}$; then $\hat{\mu}_* \hat{l}$ is a partition of unity subordinate to $\tilde{\mathcal{U}}$. Define $\bar{\omega}_\Gamma := h^{\hat{\mu}_* \hat{l}}(\tilde{\omega}'_\Gamma) = h^{\hat{l}}(\hat{\mu}^* \tilde{\omega}'_\Gamma)$. Then $\bar{\omega}_\Gamma|_{X_\Gamma(\pi)} = h^{\hat{l}|_{X_\Gamma(\pi)}}(\tilde{\omega}_\Gamma)$. Since all the intersections of elements of $\tilde{\mathcal{U}}$ are contractible (since the same is true for each \mathcal{U}_e and $\tilde{\mathcal{U}}$ is their product), both $h^{\hat{\mu}_* \hat{l}}$ and $h^{\hat{l}}$ induce the isomorphism between Čech and de Rham cohomology (here we let the family of supports be the collection of compact subsets in $\overline{C}_2(\pi)^{E(\pi)}$ that do not intersect $S(\pi)$), so

$$[\bar{\omega}_\Gamma] = [\tilde{\omega}'_\Gamma] = [\omega_\Gamma] \in H^*(\overline{C}_2(\pi)^{E(\Gamma)}, S(\pi)).$$

Denote the restriction of $\bar{\omega}_\Gamma$ to $\bar{C}_2^{E(\Gamma)}(\pi)$ still by $\bar{\omega}_\Gamma$, then, pulling back the above equation by restriction, $[\bar{\omega}_\Gamma] = [\omega_\Gamma] \in H^*(\bar{C}_2^{E(\Gamma)}(\pi), S(\pi))$. So, by Lemma 5.5, $\pi_*^s(\bar{\omega}_\Gamma)$ also represents $2^{|E(\Gamma)|} |E(\Gamma)|!$ times Kontsevich's class in the usual definition.

We apply Remark 4.9 to the situation here. For each p_0 -simplex Δ^{p_0} in B , $\check{\omega}_\Gamma|_{\pi_X^{-1}(\Delta^{p_0})}$ is supported away from $(T(\pi) \cap \pi_X^{-1}(\Delta^{p_0})) \cup \pi_X^{-1}(\partial\Delta^{p_0})$. So $\check{\omega}_\Gamma|_{\pi_X^{-1}(\Delta^{p_0})}$ is a cocycle in

$$\check{C}_{\check{U}|_{\pi_X^{-1}(\Delta^{p_0})}}^{|E(\Gamma)|(d-1)}(X_\Gamma(\pi)|_{\Delta^{p_0}}, X_\Gamma(\pi)|_{\partial\Delta^{p_0}} \cup T(\pi)|_{\Delta^{p_0}}).$$

By Künneth formula,

$$H^{|E(\Gamma)|(d-1)}(X_\Gamma(\pi)|_{\Delta^{p_0}}, X_\Gamma(\pi)|_{\partial\Delta^{p_0}} \cup T(\pi)|_{\Delta^{p_0}}) \approx H^{p_0}(\Delta^{p_0}, \partial\Delta^{p_0}) \otimes H^{d|V(\Gamma)|}(X_{\Gamma,b}, T_b),$$

where $b \in \Delta^{p_0}$ is an arbitrary point and $(X_{\Gamma,b}, T_b)$ is the fiber of $(X_\Gamma(\pi), T(\pi))$ over b . Let $\chi'(\Delta^{p_0}) \in H^{d|V(\Gamma)|}(X_{\Gamma,b}, T_b)$ be such that $[\check{\omega}_\Gamma|_{\pi_X^{-1}(\Delta^{p_0})}] \approx 1 \otimes \chi'(\Delta^{p_0})$ under the Künneth isomorphism, where $1 \in \mathbb{R} \approx H^{p_0}(\Delta^{p_0}, \partial\Delta^{p_0})$ (notice the identification uses the orientation of B). Then $\{\Delta^{p_0} \rightarrow \chi'(\Delta^{p_0})\}_{\Delta^{p_0}}$ is a simplicial cocycle on B with coefficients in the local system $H^{d|V(\Gamma)|}(X_\Gamma, T)$. Its restriction to $H^{d|V(\Gamma)|}(X_\Gamma, T_1)$ represents $(\pi_X)_* \Omega_\Gamma(\pi)$. Let $\chi(\Delta^{p_0}) = \tilde{f}^* \chi'(\Delta^{p_0}) \in H_c^{d|V(\Gamma)|}(\tilde{X}_{\Gamma,b}) \approx \mathbb{R}$ (the last identification uses the orientation on \tilde{X}_Γ), then $\{\Delta^{p_0} \rightarrow \chi(\Delta^{p_0})\}_{\Delta^{p_0}}$ is a representative of $K_{\Gamma,\pi,F}$ by the definition of $K_{\Gamma,\pi,F}$ (7),(8).

Note the bundle map

$$\begin{array}{ccccc} \tilde{X}_\Gamma(\pi) & \xrightarrow{\tilde{f}} & X_\Gamma(\pi) - T_1(\pi) & \hookrightarrow & X_\Gamma(\pi) \\ & \searrow \pi_{\tilde{X}} & \downarrow \pi_X & \swarrow \pi_X & \\ & & B & & \end{array}.$$

Since $\check{\omega}_\Gamma = 0$ in a neighborhood of $T(\pi) \cap X_{p_0}$, $\tilde{f}^* \check{\omega}_\Gamma|_{\pi_X^{-1}(B_{p_0})}$ is compactly supported. For each p_0 -simplex Δ^{p_0} of B , by the naturality of the Künneth formula, $[\tilde{f}^* \check{\omega}_\Gamma|_{\pi_X^{-1}(\Delta^{p_0})}] \approx 1 \otimes \tilde{f}^* \chi'(\Delta^{p_0}) = 1 \otimes \chi(\Delta^{p_0})$ under the Künneth isomorphism

$$H_c^{|E(\Gamma)|(d-1)}(\tilde{X}_\Gamma(\pi)|_{\Delta^{p_0}}, \tilde{X}_\Gamma(\pi)|_{\partial\Delta^{p_0}}) \approx H^{p_0}(\Delta^{p_0}, \partial\Delta^{p_0}) \otimes H_c^{d|V(\Gamma)|}(\tilde{X}_{\Gamma,b}).$$

Since $\bar{\omega}_\Gamma = h^{\hat{L}}(\hat{\mu}^* \check{\omega}'_\Gamma)$ and $\hat{\mu}^* \check{\omega}'_\Gamma$ is supported away from N , $\bar{\omega}_\Gamma$ is also supported away from N . So $[\bar{\omega}_\Gamma] = [\check{\omega}'_\Gamma] \in H^*(\bar{C}_2(\pi)^{E(\Gamma)}, N)$, and $\tilde{f}^* \bar{\omega}_\Gamma$ is a smooth form. For every simplex Δ^{p_0} in B , pulling back the cohomology equality to $\tilde{X}_\Gamma(\pi)$ and restricting to Δ^{p_0} in B ,

$$[\tilde{f}^* \bar{\omega}_\Gamma|_{\pi_X^{-1}(\Delta^{p_0})}] = [\tilde{f}^* \check{\omega}_\Gamma|_{\pi_X^{-1}(\Delta^{p_0})}] = 1 \otimes \chi(\Delta^{p_0}) \in H_c^{|E(\Gamma)|(d-1)}(\tilde{X}_\Gamma(\pi)|_{\Delta^{p_0}}, \tilde{X}_\Gamma(\pi)|_{\partial\Delta^{p_0}})$$

$$\approx H^{p_0}(\Delta^{p_0}, \partial\Delta^{p_0}) \otimes H_c^{d|V(\Gamma)|}(\tilde{X}_{\Gamma,b}) \approx \mathbb{R} \otimes \mathbb{R}.$$

Therefore, $\int_{\pi_{\tilde{X}}^{-1}(\Delta^{p_0})} \tilde{f}^* \tilde{\omega}_{\Gamma} = \chi(\Delta^{p_0})$. This completes the proof of Proposition 5.1.

6 Some remarks about the condition in Theorem 1.2

Theorem 1.2 can potentially be formulated in a different way.

For open subsets $U, V \subset \mathbb{R}^d$ and homeomorphism $f : U \rightarrow V$, say f is *almost smooth* if the map $(f, f) : U \times U \rightarrow V \times V$ lifts to a continuous map $\tilde{f} : Bl_{\Delta}(U \times U) \rightarrow Bl_{\Delta}(V \times V)$, where Δ denotes the diagonal in $U \times U$ and $V \times V$, respectively, and Bl_{Δ} denotes real oriented blow-up along Δ . We can define an *almost smooth manifold* to be a topological manifold together with a maximal collection of charts where the transition maps are almost smooth. So, if M is an almost smooth manifold, then $Bl_{\Delta}(M \times M)$ is well-defined. The corresponding automorphism group $\text{Aut}^{as}(M)$ in this category consists of homeomorphisms $f : M \rightarrow M$ such that $(f, f) : M \times M \rightarrow M \times M$ lifts to $Bl_{\Delta}(M \times M) \rightarrow Bl_{\Delta}(M \times M)$. Denote by $\pi : Bl_{\Delta}(M \times M) \rightarrow M \times M$ the blow down map.

Given two real vector spaces T_1, T_2 and a linear isomorphism $f : T_1 \rightarrow T_2$, since $f(\lambda v) = \lambda f(v)$ for all $v \in T_1 - 0$, $\lambda \in \mathbb{R} - 0$, f induces a homeomorphism $ST_1 \rightarrow ST_2$, where $ST_i = (T_i - 0)/\text{scaling}$ denotes the unit sphere in T_i .

Suppose M is a d -dimensional almost smooth manifold. Define a *framing* F on M to be a continuous map $F : \partial Bl_{\Delta}(M \times M) \rightarrow S^{d-1}$ such that for every $x \in M$, $F|_{\pi^{-1}(x,x)}$ satisfies: if $\phi : \mathbb{R}^d \supset U \xrightarrow{\sim} N \subset M$, $\phi(0) = x$ is a chart of M near x , then $F|_{\pi^{-1}(x,x)} := ST_x U \rightarrow S^{d-1}$ is a homeomorphism induced from a linear map $T_x U \rightarrow \mathbb{R}^d$. By Proposition 6.3 below, if this condition is satisfied for one chart ϕ , then it is satisfied for all charts.

Suppose M is a d -dimensional almost smooth manifold and $\infty \in M$ a fixed point. Then the group \mathcal{G} defined in Section 1.1 can be similarly defined here:

$$\mathcal{G} := \{g \in \text{Aut}^{as}(M) \mid \exists \text{ neighborhood } N \ni \infty \text{ such that } g|_N = \text{id}\}.$$

I expect that the $K_{\Gamma,\pi,F}$ constructed in Section 3.4 can be generalized to the case where $E \xrightarrow{\pi} B$ is an (M, ∞) -bundle with group \mathcal{G} and F is a vertical framing on π in some appropriate sense (a vertical framing on E away from s_{∞} , and “standard” near s_{∞}), and Theorem 1.2 can then be rephrased as the naturality of $K_{\Gamma,\pi,F}$ in the category of vertically framed almost smooth fiber bundles.

The almost smooth condition is actually quite strong. I do not know at the time of writing if an almost smooth manifold (respectively, an almost smooth bundle) necessarily has a unique smooth structure. We close this section with three auxiliary observations. Example 6.1 below shows that almost smoothness does not imply smoothness. Proposition 6.2 below shows that almost smooth implies quasi-conformal. Proposition 6.3 below shows that an almost smooth map induces a linear map between tangent bundles, modulo scaling by a positive smooth function.

Example 6.1. Let B_ϵ^n be the standard ball of radius $\epsilon < 1/(2e)$ in \mathbb{R}^n . Define

$$f : B_\epsilon^n \rightarrow \mathbb{R}^n, \quad f(x) = -2 \log(|x|) \cdot x,$$

then f maps B_ϵ^n homeomorphically onto its image, and f is almost smooth, but not continuously differentiable at 0. See [19] for a detailed proof – the point being that the function $-2 \log |x|$ approaches ∞ slow enough as $x \rightarrow 0$.

The following definition of quasi-conformal of a map is copied from [4]. A homeomorphism $f : U \rightarrow \mathbb{R}^d$ from an open subset U to its image is k *quasiconformal* if for all $x \in U$

$$H_f(x) = \limsup_{r \rightarrow 0} \frac{\max\{|f(y) - f(x)| \mid |y - x| = r\}}{\min\{|f(y) - f(x)| \mid |y - x| = r\}} \leq k.$$

f is *quasiconformal* if it is k quasiconformal for some $k \geq 1$.

Proposition 6.2 Let $U \subset \mathbb{R}^d$ be open and $f : U \rightarrow \mathbb{R}^d$ be an almost smooth homeomorphism to its image. Then for every compact subset $K \subset U$, f is quasiconformal on K .

Proof. First notice that f being almost smooth implies: for every point $x \in U$ there is a map $f'_x : S^{d-1} \rightarrow S^{d-1}$ such that for a sequence of pairs of points $\{(x_n, y_n) \in U \times U\}_{n=1}^\infty$,

$$\lim_{n \rightarrow \infty} (x_n, y_n) \rightarrow (x, x), \quad \lim_{n \rightarrow \infty} \frac{y_n - x_n}{|y_n - x_n|} = v \implies \lim_{n \rightarrow \infty} \frac{f(y_n) - f(x_n)}{|f(y_n) - f(x_n)|} = f'_x(v).$$

Suppose f is not quasiconformal on some K . Then for every $k > 0$ there exists $x_k \in K$ such that, for all $\epsilon > 0$, there exist $y_k^b, y_k^s \in U$ (the superscripts stand for “big” and “small”) satisfying

$$|y_k^b - x_k| = |y_k^s - x_k| < \epsilon, \quad \frac{|f(y_k^b) - f(x_k)|}{|f(y_k^s) - f(x_k)|} > k.$$

Let k range over $\mathbb{Z}^{>0}$. Since K is compact, by possibly passing to a subsequence, we can assume $x_k \rightarrow x$ as $k \rightarrow \infty$ for some $x \in K$. Plugging in $\epsilon = 1/k$ above, for every k we

get a tuple of points x_k, y_k^b, y_k^s , all limit to x as $k \rightarrow \infty$. For each k , denote by S_k the sphere centered at x_k on which y_k^s, y_k^b lie. Define $z_k \in S_k$ to be the midpoint of the shortest geodesic between y_k^b, y_k^s on S_k . This implies that the angle between the vectors $y_k^b - x_k$ and $y_k^b - z_k$ is at least $\pi/4$; same for the vectors $z_k - y_k^s$ and $z_k - x_k$. Since

$$\frac{|f(y_k^b) - f(x_k)|}{|f(z_k) - f(x_k)|} \cdot \frac{|f(z_k) - f(x_k)|}{|f(y_k^s) - f(x_k)|} > k,$$

one of the factors must be bigger than \sqrt{k} . In the case it is the first factor, define $z_k^b = y_k^b$, $z_k^s = z_k$; in the case it is the second factor, define $z_k^b = z_k$, $z_k^s = y_k^s$. Now we have a sequence of tuples (x_k, z_k^b, z_k^s) , all limit to x as $k \rightarrow \infty$, and

$$\lim_{k \rightarrow \infty} \frac{f(z_k^b) - f(x_k)}{|f(z_k^b) - f(x_k)|} = \lim_{k \rightarrow \infty} \frac{f(z_k^b) - f(z_k^s)}{|f(z_k^b) - f(z_k^s)|},$$

because in the triangle $(f(z_k^b), f(z_k^s), f(x_k))$, the length between the edges $z_k^s x_k$ and $z_k^b x_k$ goes to 0, implying that the angle between the edges $z_k^b z_k^s$ and $z_k^b x_k$ goes to 0. Now, since f is almost smooth and thus so is f^{-1} ,

$$\begin{aligned} \lim_{k \rightarrow \infty} \frac{z_k^b - x_k}{|z_k^b - x_k|} &= (f^{-1})'_{f(x)} \left(\lim_{k \rightarrow \infty} \frac{f(z_k^b) - f(x_k)}{|f(z_k^b) - f(x_k)|} \right) \\ &= (f^{-1})'_{f(x)} \left(\lim_{k \rightarrow \infty} \frac{f(z_k^b) - f(z_k^s)}{|f(z_k^b) - f(z_k^s)|} \right) = \lim_{k \rightarrow \infty} \frac{z_k^b - z_k^s}{|z_k^b - z_k^s|}, \end{aligned}$$

which contradicts that the angle between the two vectors is at least $\pi/4$. \square

The converse to Proposition 6.2 is not true. For example, $f : \mathbb{R}^2 \rightarrow \mathbb{R}^2, f(x) = |x|^{-1/2} \cdot x$ is quasiconformal but not almost smooth.

Proposition 6.3 Let U_1, U_2 be open subsets of \mathbb{R}^d and $f : U_1 \rightarrow U_2$ be an almost smooth homeomorphism. For every point $x \in U$, denote $\tilde{f}_x : ST_x U_1 \rightarrow ST_{f(x)} U_2$ the homeomorphism given by restricting \tilde{f} to $\pi^{-1}(x, x)$, where ST denotes the unit sphere in the tangent space. Then \tilde{f}_x is induced from a linear isomorphism $T_x U_1 \rightarrow T_x U_2$.

Proof. Given a directed, ordered graph Γ and $U \subset \mathbb{R}^d$ an open subset, define

$$g_\Gamma^U : U^{V(\Gamma)} - \Delta_{\text{big}} \rightarrow (U \times U - \Delta)^{E(\Gamma)}, \quad (x_v)_{v \in V(\Gamma)} \longrightarrow ((x_{v_-^\Gamma(e)}, x_{v_+^\Gamma(e)}))_{e \in E(\Gamma)};$$

then, by the construction of Fulton-MacPherson compactification, g_Γ^U extends to a map

$$\bar{g}_\Gamma^U : \bar{C}_{V(\Gamma)}(U) \longrightarrow Bl_\Delta(U \times U)^{E(\Gamma)},$$

where $\overline{C}_{V(\Gamma)}(U)$ is the Fulton-MacPherson configuration space of $V(\Gamma)$ -labeled marked points in U . Similar to the proof of Lemma 3.2, $\text{image}(\bar{g}_\Gamma^U)$ is the closure of $\text{image}(g_\Gamma^U)$ in $Bl_\Delta(U \times U)^{E(\Gamma)}$. Let $\pi : Bl_\Delta(U \times U)^{E(\Gamma)} \rightarrow (U \times U)^{E(\Gamma)}$ be the blow-down map. For a point $x \in U$, $\pi^{-1}((x, x))_{e \in E(\Gamma)} = (ST_x U)^{E(\Gamma)} = (S^{d-1})^{E(\Gamma)}$ and, by the construction of Fulton-MacPherson compactification,

$$\pi^{-1}(((x, x))_{e \in E(\Gamma)}) \cap \text{image}(\bar{g}_\Gamma^U) = \overline{V}_\Gamma \subset (S^{d-1})^{E(\Gamma)};$$

recall \overline{V}_Γ is defined in Definition 3.11. Now plug in U_1, U_2 for U . From the definition of g_Γ^U , $(f, f)^{E(\Gamma)} \circ g_\Gamma^{U_1} = g_\Gamma^{U_2} \circ f^{V(\Gamma)}$, so $\text{image}(g_\Gamma^{U_2}) = (f, f)^{E(\Gamma)}(\text{image}(g_\Gamma^{U_1}))$. Passing to their closures in $Bl_\Delta(U_i \times U_i)^{E(\Gamma)}$, we have $\text{image}(\bar{g}_\Gamma^{U_2}) = \tilde{f}^{E(\Gamma)}(\text{image}(\bar{g}_\Gamma^{U_1}))$. Therefore, for each $x \in U_1$,

$$(\tilde{f}_x)^{E(\Gamma)} : (S^{d-1})^{E(\Gamma)} = (ST_x U_1)^{E(\Gamma)} \longrightarrow (ST_{f(x)} U_2)^{E(\Gamma)} = (S^{d-1})^{E(\Gamma)}$$

maps \overline{V}_Γ to \overline{V}_Γ . The conclusion of the proposition follows from Lemma 6.4 below. \square

Lemma 6.4 Suppose $f : S^{d-1} \rightarrow S^{d-1}$ is a homeomorphism such that for any directed, ordered graph Γ , \overline{V}_Γ is invariant under $f^{E(\Gamma)} := (f, \dots, f) : (S^{d-1})^{E(\Gamma)} \rightarrow (S^{d-1})^{E(\Gamma)}$, then f is induced by a map $F \in GL(d)$.

Proof. (This proof is given by Fabian Gundlach.) The strategy is to define an increasing sequence of subsets $\{A_n \subset S^{d-1}\}_{n=1}^\infty$, $A_n \subset A_{n+1}$, such that $\cup_n A_n$ is dense in S^{d-1} , and show that for each n , $f|_{A_n}$ is induced by a map $F_n \in GL(d)$. For each $n \in \mathbb{Z}^{>0}$, let Γ_n be the complete graph with $(n+1)^d$ vertices labeled by elements in $L_n := \{0, \dots, n\}^d$. Then putting the vertex labeled by (m_1, \dots, m_d) at $(m_1, \dots, m_d) \in \mathbb{R}^d$ gives an element in $C_{V(\Gamma_n)}(\mathbb{R}^d)$. Since \overline{V}_{Γ_n} is invariant under $f^{E(\Gamma_n)}$, there is an element $x = (x_{\underline{m}} \in \mathbb{R}^d)_{\underline{m} \in L_n} \in (\mathbb{R}^d)^{V(\Gamma_n)}$ such that for any $\underline{m}_1 \neq \underline{m}_2 \in L_n$, $x_{\underline{m}_1} \neq x_{\underline{m}_2}$ and

$$\frac{x_{\underline{m}_1} - x_{\underline{m}_2}}{|x_{\underline{m}_1} - x_{\underline{m}_2}|} = f\left(\frac{\underline{m}_1 - \underline{m}_2}{|\underline{m}_1 - \underline{m}_2|}\right). \quad (9)$$

Denote $e_j = (0, \dots, 0, 1, 0, \dots, 0) \in L_n$ where 1 is at the j -th place. We also view e_j as an element in S^{d-1} . For all $\underline{m} \in L_n$ and $j \in \{1, \dots, d\}$, $x_{\underline{m}+e_j} - x_{\underline{m}}$ has direction $f(e_j)$ by (9). We next show that $|x_{\underline{m}+e_j} - x_{\underline{m}}|$ does not depend on \underline{m} either. For $k \neq j$, since $x_{\underline{m} \pm e_k} - x_{\underline{m}}$ is parallel to $x_{\underline{m}+e_j \pm e_k} - x_{\underline{m}+e_j}$, the points $x_{\underline{m}}, x_{\underline{m}+e_j}, x_{\underline{m} \pm e_k}, x_{\underline{m}+e_j \pm e_k}$ form a parallelogram, so we must have $x_{\underline{m} \pm e_k + e_j} - x_{\underline{m} \pm e_k} = x_{\underline{m}+e_j} - x_{\underline{m}}$. Therefore, plugging in k for j , we have $x_{\underline{m}+e_k} - x_{\underline{m}} = x_{\underline{m}-e_j+e_k} - x_{\underline{m}-e_j}$ as well. So, in the two triangles $(x_{\underline{m}}, x_{\underline{m}+e_j}, x_{\underline{m}+e_k})$ and $(x_{\underline{m}-e_j}, x_{\underline{m}}, x_{\underline{m}-e_j+e_k})$, one of the pairs of corresponding edges are equal as vectors. The other two pairs of corresponding edges are both parallel by (9), so they must both be equal. This shows $x_{\underline{m}+e_j} - x_{\underline{m}} = x_{\underline{m}} - x_{\underline{m}-e_j}$. Therefore, $x_{\underline{m}+e_j} - x_{\underline{m}}$ does not depend on the choice of \underline{m} . Without loss of generality we can assume $x_{(0, \dots, 0)} = (0, \dots, 0) \in \mathbb{R}^d$.

Then $x_{\underline{m}+e_j} - x_{\underline{m}} = x_{e_j}$ for all \underline{m}, j . So, $x_{\underline{m}_1+\underline{m}_2} = x_{\underline{m}_1} + x_{\underline{m}_2}$ for all $\underline{m}_1, \underline{m}_2$. This shows that the map $F_n \in GL(d)$ defined by “ $\forall i \ F_n(e_i) = x_{e_i}$ ” maps \underline{m} to $x_{\underline{m}}$ for all $\underline{m} \in L_n$. For $F \in GL(d)$, denote by $\hat{F} : S^{d-1} \rightarrow S^{d-1}$ the homeomorphism induced by F . Now define

$$A_n = \left\{ \frac{\underline{m}_1 - \underline{m}_2}{|\underline{m}_1 - \underline{m}_2|} \right\}_{\underline{m}_1 \neq \underline{m}_2 \in L_n} \subset S^{d-1},$$

then $f|_{A_n} = \hat{F}_n|_{A_n}$. On the other hand, the condition $\hat{F}_{A_n} = f|_{A_n}$ uniquely determines F up to scaling: since $\{e_i\}_{i=1}^d \subset A_n$, there exists $(0 \neq \lambda_i \in \mathbb{R})_{i=1}^d$ such that $F_n(e_i) = \lambda_i f(e_i)$; since $e_i + e_j \in A_n$, the direction of $F(e_i + e_j) = \lambda_i f(e_i) + \lambda_j f(e_j)$ is determined by f , so λ_i/λ_j is determined. Therefore, for different n , F_n differ only by scaling. This determines a map $F \in GL(d)$ up to scaling, satisfying $\hat{F}|_{\cup_n A_n} = f|_{\cup_n A_n}$. Since $\cup_n A_n$ is dense in S^{d-1} , $\hat{F} = f$. \square

References

- [1] R. Bott and L. Tu, *Differential Forms in Algebraic Topology*, Springer, 1982
- [2] G. E. Bredon, *Sheaf Theory*, Springer, GTM 170, 1997
- [3] B. Conrad, lecture notes, math.stanford.edu/~conrad/papers/cech.pdf
- [4] S. Donaldson and D. Sullivan, *Quasiconformal 4-manifolds*, Acta Math., 163 (1989), 181-252
- [5] W. Fulton and R. MacPherson, *A Compactification of Configuration Spaces*, Annals of Mathematics Vol. 139, No.1 (Jan., 1994), pp. 183-225
- [6] A. Hatcher, *Algebraic Topology, Chapter 5: Spectral Sequences*
- [7] R. Koytcheff, *Bott-Taubes-Vassiliev cohomology classes by cut-and-paste topology*, International Journal of Mathematics Vol. 30, No. 10, 1950047 (2019)
- [8] M. Kontsevich, *Feynman Diagrams and Low-Dimensional Topology*, from: “First European Congress of Mathematics, Vol II (Paris, 1992)”, Progr. Math. 120, Birkhäuser (1994) 97
- [9] P. Kronheimer and T. Mrowka, *Monopoles and Three-Manifolds*, Cambridge University Press, 2007
- [10] G. Kuperberg and D. Thurston, *Perturbative 3-manifold invariants by cut-and-paste topology*, arxiv/9912167

- [11] C. Lescop, *Invariants of links and 3-manifolds from graph configurations*, arxiv/2001.09929
- [12] K. Nagami, *Dimension Theory*, Academic Press, 1970
- [13] A. Sard, *Hausdorff Measure of Critical Images on Banach Manifolds*, American Journal of Mathematics Vol. 87, No. 1 (Jan., 1965), pp. 158-174
- [14] N. Steenrod, *Homology with local coefficients*, Annals of Mathematics, Second Series, Vol. 44, No. 4 (Oct., 1943), pp. 610-627
- [15] N. Steenrod, *The Topology of Fiber Bundles*, Princeton University Press, 1951
- [16] E. Szpilrajn, *La dimension et la mesure*, Fundamenta Mathematicae 28 (1937), 81-89
- [17] T. Watanabe, *Some exotic nontrivial elements of the rational homotopy groups of $\text{Diff}(S^4)$* , arxiv/1812.02448
- [18] T. Watanabe, *Addendum to: Some exotic nontrivial elements of the rational homotopy groups of $\text{Diff}(S^4)$ (homological interpretation)*, arxiv/2109.01609
- [19] mathoverflow.net/q/436642, Xujia's answer

xujiachen@math.harvard.edu