

Bayesian Data Analysis, class 2a

Andrew Gelman

Chapter 2. Single-parameter models (part 1)

2. Single-parameter models (part 1)

- ▶ The basics:
 - ▶ Data model (thus, likelihood)
 - ▶ Prior density
 - ▶ Posterior density
- ▶ Work with analytic conjugate forms
- ▶ Likelihood comes before the prior; why?

2. Single-parameter models (part 1)

- ▶ The basics:
 - ▶ Data model (thus, likelihood)
 - ▶ Prior density
 - ▶ Posterior density
- ▶ Work with analytic conjugate forms
- ▶ Likelihood comes before the prior; why?

2. Single-parameter models (part 1)

- ▶ The basics:
 - ▶ Data model (thus, likelihood)
 - ▶ Prior density
 - ▶ Posterior density
- ▶ Work with analytic conjugate forms
- ▶ Likelihood comes before the prior; why?

2. Single-parameter models (part 1)

- ▶ The basics:
 - ▶ Data model (thus, likelihood)
 - ▶ Prior density
 - ▶ Posterior density
- ▶ Work with analytic conjugate forms
- ▶ Likelihood comes before the prior; why?

2. Single-parameter models (part 1)

- ▶ The basics:
 - ▶ Data model (thus, likelihood)
 - ▶ Prior density
 - ▶ Posterior density
- ▶ Work with analytic conjugate forms
- ▶ Likelihood comes before the prior; why?

2. Single-parameter models (part 1)

- ▶ The basics:
 - ▶ Data model (thus, likelihood)
 - ▶ Prior density
 - ▶ Posterior density
- ▶ Work with analytic conjugate forms
- ▶ Likelihood comes before the prior; why?

2. Single-parameter models (part 1)

- ▶ The basics:
 - ▶ Data model (thus, likelihood)
 - ▶ Prior density
 - ▶ Posterior density
- ▶ Work with analytic conjugate forms
- ▶ Likelihood comes before the prior; why?

2.1. Estimating a probability from binomial data




Figure 2.1 *Unnormalized posterior density for binomial parameter θ , based on uniform prior distribution and y successes out of n trials. Curves displayed for several values of n and y .*

2.1. Estimating a probability from binomial data

- ▶ Bayes' original example
 - ▶ $\frac{y}{n}$, $\frac{y+1}{n+2}$, and other alternatives
 - ▶ Consider some examples
-
- ▶ Dependence of θ and n

2.1. Estimating a probability from binomial data

- ▶ Bayes' original example
- ▶ $\frac{y}{n}$, $\frac{y+1}{n+2}$, and other alternatives
- ▶ Consider some examples
 - ▶ Election polls
 - ▶ Sampling of financial records
 - ▶ Rare diseases
- ▶ Dependence of θ and n

2.1. Estimating a probability from binomial data

- ▶ Bayes' original example
- ▶ $\frac{y}{n}$, $\frac{y+1}{n+2}$, and other alternatives
- ▶ Consider some examples
 - ▶ Opinion polls
 - ▶ Sampling of financial records
 - ▶ Rare diseases
- ▶ Dependence of θ and n

2.1. Estimating a probability from binomial data

- ▶ Bayes' original example
- ▶ $\frac{y}{n}$, $\frac{y+1}{n+2}$, and other alternatives
- ▶ Consider some examples
 - ▶ Opinion polls
 - ▶ Sampling of financial records
 - ▶ Rare diseases
- ▶ Dependence of θ and n

2.1. Estimating a probability from binomial data

- ▶ Bayes' original example
- ▶ $\frac{y}{n}$, $\frac{y+1}{n+2}$, and other alternatives
- ▶ Consider some examples
 - ▶ Opinion polls
 - ▶ Sampling of financial records
 - ▶ Rare diseases
- ▶ Dependence of θ and n

2.1. Estimating a probability from binomial data

- ▶ Bayes' original example
- ▶ $\frac{y}{n}$, $\frac{y+1}{n+2}$, and other alternatives
- ▶ Consider some examples
 - ▶ Opinion polls
 - ▶ Sampling of financial records
 - ▶ Rare diseases
- ▶ Dependence of θ and n

2.1. Estimating a probability from binomial data

- ▶ Bayes' original example
- ▶ $\frac{y}{n}$, $\frac{y+1}{n+2}$, and other alternatives
- ▶ Consider some examples
 - ▶ Opinion polls
 - ▶ Sampling of financial records
 - ▶ Rare diseases
- ▶ Dependence of θ and n

2.1. Estimating a probability from binomial data

- ▶ Bayes' original example
- ▶ $\frac{y}{n}$, $\frac{y+1}{n+2}$, and other alternatives
- ▶ Consider some examples
 - ▶ Opinion polls
 - ▶ Sampling of financial records
 - ▶ Rare diseases
- ▶ Dependence of θ and n

2.2. Posterior as compromise between data and prior information

- ▶ The posterior variance is on average smaller than the prior variance
- ▶ When does it happen that posterior variance is *larger* than the prior variance?
 - Bad luck
 - Bad model
- ▶ Research problem: when do models have “warning lights”?

2.2. Posterior as compromise between data and prior information

- ▶ The posterior variance is on average smaller than the prior variance
- ▶ When does it happen that posterior variance is *larger* than the prior variance?
 - ▶ Bad luck
 - ▶ Bad model
- ▶ Research problem: when do models have “warning lights”?

2.2. Posterior as compromise between data and prior information

- ▶ The posterior variance is on average smaller than the prior variance
- ▶ When does it happen that posterior variance is *larger* than the prior variance?
 - ▶ Bad luck
 - ▶ Bad model
- ▶ Research problem: when do models have “warning lights”?

2.2. Posterior as compromise between data and prior information

- ▶ The posterior variance is on average smaller than the prior variance
- ▶ When does it happen that posterior variance is *larger* than the prior variance?
 - ▶ Bad luck
 - ▶ Bad model
- ▶ Research problem: when do models have “warning lights”?

2.2. Posterior as compromise between data and prior information

- ▶ The posterior variance is on average smaller than the prior variance
- ▶ When does it happen that posterior variance is *larger* than the prior variance?
 - ▶ Bad luck
 - ▶ Bad model
- ▶ Research problem: when do models have “warning lights”?

2.2. Posterior as compromise between data and prior information

- ▶ The posterior variance is on average smaller than the prior variance
- ▶ When does it happen that posterior variance is *larger* than the prior variance?
 - ▶ Bad luck
 - ▶ Bad model
- ▶ Research problem: when do models have “warning lights”?

2.3. Summarizing posterior inference

- ▶ Simulation draws
- ▶ Simulations + analytics
- ▶ Posterior intervals




Figure 2.2 *Hypothetical posterior density for which the 95% central interval and 95% highest posterior density region dramatically differ: (a) central posterior interval, (b) highest posterior density region.*

2.3. Summarizing posterior inference

- ▶ Simulation draws
- ▶ Simulations + analytics
- ▶ Posterior intervals




Figure 2.2 *Hypothetical posterior density for which the 95% central interval and 95% highest posterior density region dramatically differ: (a) central posterior interval, (b) highest posterior density region.*

2.3. Summarizing posterior inference

- ▶ Simulation draws
- ▶ Simulations + analytics
- ▶ Posterior intervals




Figure 2.2 *Hypothetical posterior density for which the 95% central interval and 95% highest posterior density region dramatically differ: (a) central posterior interval, (b) highest posterior density region.*

2.3. Summarizing posterior inference

- ▶ Simulation draws
- ▶ Simulations + analytics
- ▶ Posterior intervals






Figure 2.2 *Hypothetical posterior density for which the 95% central interval and 95% highest posterior density region dramatically differ: (a) central posterior interval, (b) highest posterior density region.*

Central and shortest intervals



Shortest posterior intervals



Posterior intervals: choices and goals

- ▶ Central intervals
- ▶ Highest posterior density intervals
- ▶ Nested or non-nested?
- ▶ Asymmetry and multiple modes
- ▶ Multidimensional contours
- ▶ The purpose of interval estimation

Posterior intervals: choices and goals

- ▶ Central intervals
- ▶ Highest posterior density intervals
- ▶ Nested or non-nested?
- ▶ Asymmetry and multiple modes
- ▶ Multidimensional contours
- ▶ The purpose of interval estimation

Posterior intervals: choices and goals

- ▶ Central intervals
- ▶ Highest posterior density intervals
- ▶ Nested or non-nested?
- ▶ Asymmetry and multiple modes
- ▶ Multidimensional contours
- ▶ The purpose of interval estimation

Posterior intervals: choices and goals

- ▶ Central intervals
- ▶ Highest posterior density intervals
- ▶ Nested or non-nested?
- ▶ Asymmetry and multiple modes
- ▶ Multidimensional contours
- ▶ The purpose of interval estimation

Posterior intervals: choices and goals

- ▶ Central intervals
- ▶ Highest posterior density intervals
- ▶ Nested or non-nested?
- ▶ Asymmetry and multiple modes
- ▶ Multidimensional contours
- ▶ The purpose of interval estimation

Posterior intervals: choices and goals

- ▶ Central intervals
- ▶ Highest posterior density intervals
- ▶ Nested or non-nested?
- ▶ Asymmetry and multiple modes
- ▶ Multidimensional contours
- ▶ The purpose of interval estimation
 - ▶ Accepting what's in the interval
 - ▶ Rejecting what's outside the interval

Posterior intervals: choices and goals

- ▶ Central intervals
- ▶ Highest posterior density intervals
- ▶ Nested or non-nested?
- ▶ Asymmetry and multiple modes
- ▶ Multidimensional contours
- ▶ The purpose of interval estimation
 - ▶ Accepting what's in the interval
 - ▶ Rejecting what's outside the interval

Posterior intervals: choices and goals

- ▶ Central intervals
- ▶ Highest posterior density intervals
- ▶ Nested or non-nested?
- ▶ Asymmetry and multiple modes
- ▶ Multidimensional contours
- ▶ The purpose of interval estimation
 - ▶ Accepting what's in the interval
 - ▶ Rejecting what's outside the interval

Posterior intervals: choices and goals

- ▶ Central intervals
- ▶ Highest posterior density intervals
- ▶ Nested or non-nested?
- ▶ Asymmetry and multiple modes
- ▶ Multidimensional contours
- ▶ The purpose of interval estimation
 - ▶ Accepting what's in the interval
 - ▶ Rejecting what's outside the interval

Constructing shortest posterior intervals from simulations

- ▶ Empirical Spin
- ▶ Smoothing
- ▶ Bootstrapping
- ▶ Computational tradeoffs

Constructing shortest posterior intervals from simulations

- ▶ Empirical Spin
- ▶ Smoothing
- ▶ Bootstrapping
- ▶ Computational tradeoffs

Constructing shortest posterior intervals from simulations

- ▶ Empirical Spin
- ▶ Smoothing
- ▶ Bootstrapping
- ▶ Computational tradeoffs


Constructing shortest posterior intervals from simulations

- ▶ Empirical Spin
- ▶ Smoothing
- ▶ Bootstrapping
- ▶ Computational tradeoffs

Constructing shortest posterior intervals from simulations

- ▶ Empirical Spin
- ▶ Smoothing
- ▶ Bootstrapping
- ▶ Computational tradeoffs

Sampling variability of the empirical shortest posterior interval



2.4. Informative prior distributions

- ▶ Interpretations
 - ▶ Population
 - ▶ State of knowledge
 - ▶ Software defaults (statisticians in a fix)
- ▶ Binomial model:
- ▶ Conjugate and non-conjugate priors

2.4. Informative prior distributions

- ▶ Interpretations

- ▶ Population
- ▶ State of knowledge
- ▶ Software defaults (statistician in a box)

- ▶ Binomial model:

- ▶ Conjugate and non-conjugate priors

2.4. Informative prior distributions

- ▶ Interpretations
 - ▶ Population
 - ▶ State of knowledge
 - ▶ Software defaults (statistician in a box)
- ▶ Binomial model:
- ▶ Conjugate and non-conjugate priors

2.4. Informative prior distributions

- ▶ Interpretations
 - ▶ Population
 - ▶ State of knowledge
 - ▶ Software defaults (statistician in a box)
- ▶ Binomial model:
 - ▶ $\text{Beta}(\alpha, \beta)$ prior is equivalent to $\alpha + \beta$ data points
 - ▶ Or not? $\beta \sim 2$ data points?
- ▶ Conjugate and non-conjugate priors

2.4. Informative prior distributions

- ▶ Interpretations
 - ▶ Population
 - ▶ State of knowledge
 - ▶ Software defaults (statistician in a box)
- ▶ Binomial model:
 - ▶ Beta(α, β) prior as equivalent to $\alpha + \beta$ data points
 - ▶ Or $\alpha + \beta - 2$ data points?
- ▶ Conjugate and non-conjugate priors

2.4. Informative prior distributions

- ▶ Interpretations
 - ▶ Population
 - ▶ State of knowledge
 - ▶ Software defaults (statistician in a box)
- ▶ Binomial model:
 - ▶ $\text{Beta}(\alpha, \beta)$ prior as equivalent to $\alpha + \beta$ data points
 - ▶ Or $\alpha + \beta - 2$ data points?
- ▶ Conjugate and non-conjugate priors

2.4. Informative prior distributions

- ▶ Interpretations
 - ▶ Population
 - ▶ State of knowledge
 - ▶ Software defaults (statistician in a box)
- ▶ Binomial model:
 - ▶ $\text{Beta}(\alpha, \beta)$ prior as equivalent to $\alpha + \beta$ data points
 - ▶ Or $\alpha + \beta - 2$ data points?
- ▶ Conjugate and non-conjugate priors

2.4. Informative prior distributions


- ▶ Interpretations
 - ▶ Population
 - ▶ State of knowledge
 - ▶ Software defaults (statistician in a box)
- ▶ Binomial model:
 - ▶ Beta(α, β) prior as equivalent to $\alpha + \beta$ data points
 - ▶ Or $\alpha + \beta - 2$ data points?
- ▶ Conjugate and non-conjugate priors

2.4. Informative prior distributions

- ▶ Interpretations
 - ▶ Population
 - ▶ State of knowledge
 - ▶ Software defaults (statistician in a box)
- ▶ Binomial model:
 - ▶ $\text{Beta}(\alpha, \beta)$ prior as equivalent to $\alpha + \beta$ data points
 - ▶ Or $\alpha + \beta - 2$ data points?
- ▶ Conjugate and non-conjugate priors


Probability of a girl birth

- In general population, $\Pr(\text{girl birth}) = 0.485$




Probability of a girl birth

- ▶ In general population, $\Pr(\text{girl birth}) = 0.485$



Variation!



Example: probability of a girl birth given placenta previa

- ▶ Data: 437 girls out of 980 births, $y/n = 437/980 = 0.446$
- ▶ Consider different $\text{Beta}(\alpha, \beta)$ priors

$\frac{\alpha}{\alpha + \beta}$	$\alpha + \beta$	Posterior median of θ	95% posterior interval for θ
0.500	2	0.446	[0.415, 0.477]
0.485	2	0.446	[0.415, 0.477]
0.485	5	0.446	[0.415, 0.477]
0.485	10	0.446	[0.415, 0.477]
0.485	20	0.447	[0.416, 0.478]
0.485	100	0.450	[0.420, 0.479]
0.485	200	0.453	[0.424, 0.481]

Example: probability of a girl birth given placenta previa

- ▶ Data: 437 girls out of 980 births, $y/n = 437/980 = 0.446$
- ▶ Consider different $\text{Beta}(\alpha, \beta)$ priors

$\frac{\alpha}{\alpha + \beta}$	$\alpha + \beta$	Posterior median of θ	95% posterior interval for θ
0.500	2	0.446	[0.415, 0.477]
0.485	2	0.446	[0.415, 0.477]
0.485	5	0.446	[0.415, 0.477]
0.485	10	0.446	[0.415, 0.477]
0.485	20	0.447	[0.416, 0.478]
0.485	100	0.450	[0.420, 0.479]
0.485	200	0.453	[0.424, 0.481]

Example: probability of a girl birth given placenta previa

- ▶ Data: 437 girls out of 980 births, $y/n = 437/980 = 0.446$
- ▶ Consider different $\text{Beta}(\alpha, \beta)$ priors

$\frac{\alpha}{\alpha + \beta}$	$\alpha + \beta$	Posterior median of θ	95% posterior interval for θ
0.500	2	0.446	[0.415, 0.477]
0.485	2	0.446	[0.415, 0.477]
0.485	5	0.446	[0.415, 0.477]
0.485	10	0.446	[0.415, 0.477]
0.485	20	0.447	[0.416, 0.478]
0.485	100	0.450	[0.420, 0.479]
0.485	200	0.453	[0.424, 0.481]

Constructing a prior distribution

- ▶ p = probability of a girl birth given placenta previa
- ▶ Suppose we believe p should be between 0.47 and 0.50
 - ▶ Beta(α, β) prior with mean 0.485 and sd 0.01
 - ▶ Mean $= \frac{\alpha}{\alpha + \beta} = 0.485$ and $\text{var} = \frac{\alpha\beta}{(\alpha + \beta)^2} = 0.01^2$
 - ▶ Solve to get $\alpha = 2940$ (actually, 2940.75) and $\beta = 1212.5$,
 $\beta = 1207.5$
- ▶ Combine with $n = 980$ data points

Constructing a prior distribution

- ▶ p = probability of a girl birth given placenta previa
- ▶ Suppose we believe p should be between 0.47 and 0.50
 - ▶ Beta(α, β) prior with mean 0.485 and sd 0.01
 - ▶ Need $\frac{\alpha}{\alpha+\beta} = 0.485$ and $\frac{\alpha\beta}{(\alpha+\beta)^2(\alpha+\beta+1)} = 0.01^2$
 - ▶ Solve to get $\alpha \approx \beta = 2540$ (actually, 2540.75) and $\alpha = 1212.5$, $\beta = 1257.5$
- ▶ Combine with $n = 980$ data points

Constructing a prior distribution

- ▶ p = probability of a girl birth given placenta previa
- ▶ Suppose we believe p should be between 0.47 and 0.50
 - ▶ Beta(α, β) prior with mean 0.485 and sd 0.01
 - ▶ Need $\frac{\alpha}{\alpha+\beta} = 0.485$ and $\frac{\alpha\beta}{(\alpha+\beta)^2(\alpha+\beta+1)} = 0.01^2$
 - ▶ Solve to get $\alpha + \beta = 2500$ (actually, 2496.75) and $\alpha = 1212.5$, $\beta = 1287.5$
- ▶ Combine with $n = 980$ data points

Constructing a prior distribution

- ▶ p = probability of a girl birth given placenta previa
- ▶ Suppose we believe p should be between 0.47 and 0.50
 - ▶ $\text{Beta}(\alpha, \beta)$ prior with mean 0.485 and sd 0.01
 - ▶ Need $\frac{\alpha}{\alpha+\beta} = 0.485$ and $\frac{\alpha\beta}{(\alpha+\beta)^2(\alpha+\beta+1)} = 0.01^2$
 - ▶ Solve to get $\alpha + \beta = 2500$ (actually, 2496.75) and $\alpha = 1212.5$, $\beta = 1287.5$
- ▶ Combine with $n = 980$ data points

Constructing a prior distribution

- ▶ p = probability of a girl birth given placenta previa
- ▶ Suppose we believe p should be between 0.47 and 0.50
 - ▶ $\text{Beta}(\alpha, \beta)$ prior with mean 0.485 and sd 0.01
 - ▶ Need $\frac{\alpha}{\alpha+\beta} = 0.485$ and $\frac{\alpha\beta}{(\alpha+\beta)^2(\alpha+\beta+1)} = 0.01^2$
 - ▶ Solve to get $\alpha + \beta = 2500$ (actually, 2496.75) and $\alpha = 1212.5$, $\beta = 1287.5$
- ▶ Combine with $n = 980$ data points

Constructing a prior distribution

- ▶ p = probability of a girl birth given placenta previa
- ▶ Suppose we believe p should be between 0.47 and 0.50
 - ▶ $\text{Beta}(\alpha, \beta)$ prior with mean 0.485 and sd 0.01
 - ▶ Need $\frac{\alpha}{\alpha+\beta} = 0.485$ and $\frac{\alpha\beta}{(\alpha+\beta)^2(\alpha+\beta+1)} = 0.01^2$
 - ▶ Solve to get $\alpha + \beta = 2500$ (actually, 2496.75) and $\alpha = 1212.5$, $\beta = 1287.5$
- ▶ Combine with $n = 980$ data points

Constructing a prior distribution

- ▶ p = probability of a girl birth given placenta previa
- ▶ Suppose we believe p should be between 0.47 and 0.50
 - ▶ $\text{Beta}(\alpha, \beta)$ prior with mean 0.485 and sd 0.01
 - ▶ Need $\frac{\alpha}{\alpha+\beta} = 0.485$ and $\frac{\alpha\beta}{(\alpha+\beta)^2(\alpha+\beta+1)} = 0.01^2$
 - ▶ Solve to get $\alpha + \beta = 2500$ (actually, 2496.75) and $\alpha = 1212.5$, $\beta = 1287.5$
- ▶ Combine with $n = 980$ data points