

Bayesian Data Analysis, class 1a

Andrew Gelman

Introductory examples:
Soccer ratings and Population toxicokinetics

Bayesian data analysis: (1) Modeling

- ▶ “Generative models”
 - ▶ Data are a realization from a (multivariate) probability distribution
 - ▶ Data vector y , probability model $p(y|\theta)$, parameter vector θ
- ▶ Prior distributions

Bayesian data analysis: (1) Modeling

- ▶ “Generative models”
 - ▶ Data are a realization from a (multivariate) probability distribution
 - ▶ Data vector y , probability model $p(y|\theta)$, parameter vector θ
- ▶ Prior distributions

Bayesian data analysis: (1) Modeling

- ▶ “Generative models”
 - ▶ Data are a realization from a (multivariate) probability distribution
 - ▶ Data vector y , probability model $p(y|\theta)$, parameter vector θ
- ▶ Prior distributions

Interpretation: y is a realization from a probability distribution, $p(y|\theta)$, which depends on the parameter vector θ . We want to learn about θ from the data y .

Bayesian data analysis: (1) Modeling

- ▶ “Generative models”
 - ▶ Data are a realization from a (multivariate) probability distribution
 - ▶ Data vector y , probability model $p(y|\theta)$, parameter vector θ
- ▶ Prior distributions
 - ▶ In Bayes inference, the parameter vector θ is a realization from a prior distribution, $p(\theta|\phi)$
 - ▶ Vector of hyperparameters ϕ is specified or itself modeled

Bayesian data analysis: (1) Modeling

- ▶ “Generative models”
 - ▶ Data are a realization from a (multivariate) probability distribution
 - ▶ Data vector y , probability model $p(y|\theta)$, parameter vector θ
- ▶ Prior distributions
 - ▶ In Bayes inference, the parameter vector θ is a realization from a prior distribution, $p(\theta|\phi)$
 - ▶ Vector of hyperparameters ϕ is specified or itself modeled

Bayesian data analysis: (1) Modeling

- ▶ “Generative models”
 - ▶ Data are a realization from a (multivariate) probability distribution
 - ▶ Data vector y , probability model $p(y|\theta)$, parameter vector θ
- ▶ Prior distributions
 - ▶ In Bayes inference, the parameter vector θ is a realization from a prior distribution, $p(\theta|\phi)$
 - ▶ Vector of hyperparameters ϕ is specified or itself modeled

Bayesian data analysis: (1) Modeling

- ▶ “Generative models”
 - ▶ Data are a realization from a (multivariate) probability distribution
 - ▶ Data vector y , probability model $p(y|\theta)$, parameter vector θ
- ▶ Prior distributions
 - ▶ In Bayes inference, the parameter vector θ is a realization from a prior distribution, $p(\theta|\phi)$
 - ▶ Vector of hyperparameters ϕ is specified or itself modeled

Bayesian data analysis: (2) Inference

- ▶ Inference is represented by a matrix of posterior simulations
 - ▶ 1000 simulations of 90 parameters: a 1000×90 matrix
- ▶ Postprocessing

Bayesian data analysis: (2) Inference

- ▶ Inference is represented by a matrix of posterior simulations
 - ▶ 1000 simulations of 90 parameters: a 1000×90 matrix
- ▶ Postprocessing

Bayesian data analysis: (2) Inference

- ▶ Inference is represented by a matrix of posterior simulations
 - ▶ 1000 simulations of 90 parameters: a 1000×90 matrix
- ▶ Postprocessing
 - ▶ Inference for qoi's
 - ▶ Decision analysis

Bayesian data analysis: (2) Inference

- ▶ Inference is represented by a matrix of posterior simulations
 - ▶ 1000 simulations of 90 parameters: a 1000×90 matrix
- ▶ Postprocessing
 - ▶ Inference for qoi's
 - ▶ Decision analysis

Bayesian data analysis: (2) Inference

- ▶ Inference is represented by a matrix of posterior simulations
 - ▶ 1000 simulations of 90 parameters: a 1000×90 matrix
- ▶ Postprocessing
 - ▶ Inference for qoi's
 - ▶ Decision analysis

Bayesian data analysis: (2) Inference

- ▶ Inference is represented by a matrix of posterior simulations
 - ▶ 1000 simulations of 90 parameters: a 1000×90 matrix
- ▶ Postprocessing
 - ▶ Inference for qoi's
 - ▶ Decision analysis

Bayesian data analysis: (3) Model checking/improvement

- ▶ Do the inferences make sense?
- ▶ Are the model's predictions consistent with the data?
- ▶ *Not*: Is the model true?
- ▶ *Not*: What is $\Pr(\text{model is true})$?
- ▶ *Not*: Can we “reject” the model?
- ▶ Expanding the model
- ▶ Including more data

Bayesian data analysis: (3) Model checking/improvement

- ▶ Do the inferences make sense?
- ▶ Are the model's predictions consistent with the data?
- ▶ *Not*: Is the model true?
- ▶ *Not*: What is $\Pr(\text{model is true})$?
- ▶ *Not*: Can we “reject” the model?
- ▶ Expanding the model
- ▶ Including more data

Bayesian data analysis: (3) Model checking/improvement

- ▶ Do the inferences make sense?
- ▶ Are the model's predictions consistent with the data?
- ▶ *Not*: Is the model true?
- ▶ *Not*: What is $\Pr(\text{model is true})$?
- ▶ *Not*: Can we “reject” the model?
- ▶ Expanding the model
- ▶ Including more data

Bayesian data analysis: (3) Model checking/improvement

- ▶ Do the inferences make sense?
- ▶ Are the model's predictions consistent with the data?
- ▶ *Not*: Is the model true?
- ▶ *Not*: What is $\Pr(\text{model is true})$?
- ▶ *Not*: Can we “reject” the model?
- ▶ Expanding the model
- ▶ Including more data

Bayesian data analysis: (3) Model checking/improvement

- ▶ Do the inferences make sense?
- ▶ Are the model's predictions consistent with the data?
- ▶ *Not*: Is the model true?
- ▶ *Not*: What is $\Pr(\text{model is true})$?
- ▶ *Not*: Can we “reject” the model?
- ▶ Expanding the model
- ▶ Including more data

Bayesian data analysis: (3) Model checking/improvement

- ▶ Do the inferences make sense?
- ▶ Are the model's predictions consistent with the data?
- ▶ *Not*: Is the model true?
- ▶ *Not*: What is $\Pr(\text{model is true})$?
- ▶ *Not*: Can we “reject” the model?
- ▶ Expanding the model
- ▶ Including more data


Bayesian data analysis: (3) Model checking/improvement

- ▶ Do the inferences make sense?
- ▶ Are the model's predictions consistent with the data?
- ▶ *Not*: Is the model true?
- ▶ *Not*: What is $\Pr(\text{model is true})$?
- ▶ *Not*: Can we “reject” the model?
- ▶ Expanding the model
- ▶ Including more data

Bayesian data analysis: (3) Model checking/improvement

- ▶ Do the inferences make sense?
- ▶ Are the model's predictions consistent with the data?
- ▶ *Not*: Is the model true?
- ▶ *Not*: What is $\Pr(\text{model is true})$?
- ▶ *Not*: Can we “reject” the model?
- ▶ Expanding the model
- ▶ Including more data

Stan goes to the World Cup



The model

- ▶ Fit data on signed square roots:

$$y_{ij} = \sqrt{\text{score differential when team } i \text{ plays team } j}$$

- ▶ Model $y_{ij} \sim N(a_i - a_j, \sigma_y^2)$
- ▶ a_i and a_j are “ability parameters”
- ▶ σ_y is a scale parameter
- ▶ To allow for outliers, use t_7 instead of normal
- ▶ Prior info on abilities:

$$a_i \sim N(\mu + b * \text{prior.score}_i, \sigma_a^2)$$

- ▶ We can set $\mu = 0$
- ▶ No further prior info in model

The model

- ▶ Fit data on signed square roots:

$$y_{ij} = \sqrt{\text{score differential when team } i \text{ plays team } j}$$

- ▶ Model $y_{ij} \sim N(a_i - a_j, \sigma_y^2)$
- ▶ a_i and a_j are “ability parameters”
- ▶ σ_y is a scale parameter
- ▶ To allow for outliers, use t_7 instead of normal
- ▶ Prior info on abilities:

$$a_i \sim N(\mu + b * \text{prior.score}_i, \sigma_a^2)$$

- ▶ We can set $\mu = 0$
- ▶ No further prior info in model

The model

- ▶ Fit data on signed square roots:

$$y_{ij} = \sqrt{\text{score differential when team } i \text{ plays team } j}$$

- ▶ Model $y_{ij} \sim N(a_i - a_j, \sigma_y^2)$
- ▶ a_i and a_j are “ability parameters”
- ▶ σ_y is a scale parameter
- ▶ To allow for outliers, use t_7 instead of normal
- ▶ Prior info on abilities:

$$a_i \sim N(\mu + b * \text{prior.score}_i, \sigma_a^2)$$

- ▶ We can set $\mu = 0$
- ▶ No further prior info in model

The model

- ▶ Fit data on signed square roots:

$$y_{ij} = \sqrt{\text{score differential when team } i \text{ plays team } j}$$

- ▶ Model $y_{ij} \sim N(a_i - a_j, \sigma_y^2)$
- ▶ a_i and a_j are “ability parameters”
- ▶ σ_y is a scale parameter
- ▶ To allow for outliers, use t_7 instead of normal
- ▶ Prior info on abilities:

$$a_i \sim N(\mu + b * \text{prior.score}_i, \sigma_a^2)$$

- ▶ We can set $\mu = 0$
- ▶ No further prior info in model

The model

- ▶ Fit data on signed square roots:

$$y_{ij} = \sqrt{\text{score differential when team } i \text{ plays team } j}$$

- ▶ Model $y_{ij} \sim N(a_i - a_j, \sigma_y^2)$
- ▶ a_i and a_j are “ability parameters”
- ▶ σ_y is a scale parameter
- ▶ To allow for outliers, use t_7 instead of normal
- ▶ Prior info on abilities:

$$a_i \sim N(\mu + b * \text{prior.score}_i, \sigma_a^2)$$

- ▶ We can set $\mu = 0$
- ▶ No further prior info in model

The model

- ▶ Fit data on signed square roots:

$$y_{ij} = \sqrt{\text{score differential when team } i \text{ plays team } j}$$

- ▶ Model $y_{ij} \sim N(a_i - a_j, \sigma_y^2)$
- ▶ a_i and a_j are “ability parameters”
- ▶ σ_y is a scale parameter
- ▶ To allow for outliers, use t_7 instead of normal
- ▶ Prior info on abilities:

$$a_i \sim N(\mu + b * \text{prior.score}_i, \sigma_a^2)$$

- ▶ We can set $\mu = 0$
- ▶ No further prior info in model

The model

- ▶ Fit data on signed square roots:

$$y_{ij} = \sqrt{\text{score differential when team } i \text{ plays team } j}$$

- ▶ Model $y_{ij} \sim N(a_i - a_j, \sigma_y^2)$
- ▶ a_i and a_j are “ability parameters”
- ▶ σ_y is a scale parameter
- ▶ To allow for outliers, use t_7 instead of normal
- ▶ Prior info on abilities:

$$a_i \sim N(\mu + b * \text{prior.score}_i, \sigma_a^2)$$

- ▶ We can set $\mu = 0$
- ▶ No further prior info in model

The model

- ▶ Fit data on signed square roots:

$$y_{ij} = \sqrt{\text{score differential when team } i \text{ plays team } j}$$

- ▶ Model $y_{ij} \sim N(a_i - a_j, \sigma_y^2)$
- ▶ a_i and a_j are “ability parameters”
- ▶ σ_y is a scale parameter
- ▶ To allow for outliers, use t_7 instead of normal
- ▶ Prior info on abilities:

$$a_i \sim N(\mu + b * \text{prior.score}_i, \sigma_a^2)$$

- ▶ We can set $\mu = 0$
- ▶ No further prior info in model

The model

- ▶ Fit data on signed square roots:

$$y_{ij} = \sqrt{\text{score differential when team } i \text{ plays team } j}$$

- ▶ Model $y_{ij} \sim N(a_i - a_j, \sigma_y^2)$
- ▶ a_i and a_j are “ability parameters”
- ▶ σ_y is a scale parameter
- ▶ To allow for outliers, use t_7 instead of normal
- ▶ Prior info on abilities:

$$a_i \sim N(\mu + b * \text{prior.score}_i, \sigma_a^2)$$

- ▶ We can set $\mu = 0$
- ▶ No further prior info in model

Stan model (part 1)

```
data {  
  int nteams;  
  int ngames;  
  vector[nteam] prior_score;  
  int team1[ngames];  
  int team2[ngames];  
  vector[ngames] score1;  
  vector[ngames] score2;  
  real df;  
}  
transformed data {  
  vector[ngames] dif;  
  vector[ngames] sqrt_dif;  
  dif <- score1 - score2;  
  for (i in 1:ngames)  
    sqrt_dif[i] <- (step(dif[i])-.5)*sqrt(fabs(dif[i]));  
}
```

Stan model (part 2)

```
parameters {  
  real b;  
  real<lower=0> sigma_a;  
  real<lower=0> sigma_y;  
  vector[nteams] eta_a;  
}  
transformed parameters {  
  vector[nteams] a;  
  a <- b*prior_score + sigma_a*eta_a;  
}  
model {  
  eta_a ~ normal(0,1);  
  for (i in 1:ngames)  
    sqrt_dif[i] ~ student_t(df, a[team1[i]]-a[team2[i]],sigma_y)  
}
```

Fitting the model

- ▶ Go into R
- ▶ Read in the data
- ▶ Fit the Stan model
- ▶ Check convergence
- ▶ Graph the estimated team abilities
- ▶ Re-fit without prior information
- ▶ Compare to model with prior information

Fitting the model

- ▶ Go into R
- ▶ Read in the data
- ▶ Fit the Stan model
- ▶ Check convergence
- ▶ Graph the estimated team abilities
- ▶ Re-fit without prior information
- ▶ Compare to model with prior information

Fitting the model

- ▶ Go into R
- ▶ Read in the data
- ▶ Fit the Stan model
- ▶ Check convergence
- ▶ Graph the estimated team abilities
- ▶ Re-fit without prior information
- ▶ Compare to model with prior information

Fitting the model

- ▶ Go into R
- ▶ Read in the data
- ▶ Fit the Stan model
- ▶ Check convergence
- ▶ Graph the estimated team abilities
- ▶ Re-fit without prior information
- ▶ Compare to model with prior information

Fitting the model

- ▶ Go into R
- ▶ Read in the data
- ▶ Fit the Stan model
- ▶ Check convergence
- ▶ Graph the estimated team abilities
- ▶ Re-fit without prior information
- ▶ Compare to model with prior information

Fitting the model

- ▶ Go into R
- ▶ Read in the data
- ▶ Fit the Stan model
- ▶ Check convergence
- ▶ Graph the estimated team abilities
- ▶ Re-fit without prior information
- ▶ Compare to model with prior information

Fitting the model

- ▶ Go into R
- ▶ Read in the data
- ▶ Fit the Stan model
- ▶ Check convergence
- ▶ Graph the estimated team abilities
- ▶ Re-fit without prior information
- ▶ Compare to model with prior information

Fitting the model

- ▶ Go into R
- ▶ Read in the data
- ▶ Fit the Stan model
- ▶ Check convergence
- ▶ Graph the estimated team abilities
- ▶ Re-fit without prior information
- ▶ Compare to model with prior information

Checking model fit

- ▶ Still inside R
- ▶ For each game, plot actual score differential and 95% predictive intervals
 - ▶ Not cross-validated but no big deal in this case because n is large
- ▶ The predictions don't fit the data!!
- ▶ Redoing the predictive intervals
- ▶ Re-plot, still a problem!

Checking model fit

- ▶ Still inside R
- ▶ For each game, plot actual score differential and 95% predictive intervals
 - ▶ Not cross-validated but no big deal in this case because n is large
- ▶ The predictions don't fit the data!!
- ▶ Redoing the predictive intervals
- ▶ Re-plot, still a problem!

Checking model fit

- ▶ Still inside R
- ▶ For each game, plot actual score differential and 95% predictive intervals
 - ▶ Not cross-validated but no big deal in this case because n is large
- ▶ The predictions don't fit the data!!
- ▶ Redoing the predictive intervals
- ▶ Re-plot, still a problem!

Checking model fit

- ▶ Still inside R
- ▶ For each game, plot actual score differential and 95% predictive intervals
 - ▶ Not cross-validated but no big deal in this case because n is large
- ▶ The predictions don't fit the data!!
- ▶ Redoing the predictive intervals
- ▶ Re-plot, still a problem!

Checking model fit

- ▶ Still inside R
- ▶ For each game, plot actual score differential and 95% predictive intervals
 - ▶ Not cross-validated but no big deal in this case because n is large
- ▶ The predictions don't fit the data!!
- ▶ Redoing the predictive intervals
- ▶ Re-plot, still a problem!

Checking model fit

- ▶ Still inside R
- ▶ For each game, plot actual score differential and 95% predictive intervals
 - ▶ Not cross-validated but no big deal in this case because n is large
- ▶ The predictions don't fit the data!!
- ▶ Redoing the predictive intervals
- ▶ Re-plot, still a problem!

Checking model fit

- ▶ Still inside R
- ▶ For each game, plot actual score differential and 95% predictive intervals
 - ▶ Not cross-validated but no big deal in this case because n is large
- ▶ The predictions don't fit the data!!
- ▶ Redoing the predictive intervals
- ▶ Re-plot, still a problem!

I found the bug!

- ▶ Still inside R
- ▶ Re-fit the model on the original scale
- ▶ Display the estimated team abilities
- ▶ Updated plot of data with predictive intervals—now it's ok!
- ▶ Go back and find the bug in the square-root-scale model
- ▶ Re-fit the debugged model

I found the bug!

- ▶ Still inside R
- ▶ Re-fit the model on the original scale
- ▶ Display the estimated team abilities
- ▶ Updated plot of data with predictive intervals—now it's ok!
- ▶ Go back and find the bug in the square-root-scale model
- ▶ Re-fit the debugged model

I found the bug!

- ▶ Still inside R
- ▶ Re-fit the model on the original scale
- ▶ Display the estimated team abilities
- ▶ Updated plot of data with predictive intervals—now it's ok!
- ▶ Go back and find the bug in the square-root-scale model
- ▶ Re-fit the debugged model

I found the bug!

- ▶ Still inside R
- ▶ Re-fit the model on the original scale
- ▶ Display the estimated team abilities
- ▶ Updated plot of data with predictive intervals—now it's ok!
- ▶ Go back and find the bug in the square-root-scale model
- ▶ Re-fit the debugged model

I found the bug!

- ▶ Still inside R
- ▶ Re-fit the model on the original scale
- ▶ Display the estimated team abilities
- ▶ Updated plot of data with predictive intervals—now it's ok!
- ▶ Go back and find the bug in the square-root-scale model
- ▶ Re-fit the debugged model

I found the bug!

- ▶ Still inside R
- ▶ Re-fit the model on the original scale
- ▶ Display the estimated team abilities
- ▶ Updated plot of data with predictive intervals—now it's ok!
- ▶ Go back and find the bug in the square-root-scale model
- ▶ Re-fit the debugged model

I found the bug!

- ▶ Still inside R
- ▶ Re-fit the model on the original scale
- ▶ Display the estimated team abilities
- ▶ Updated plot of data with predictive intervals—now it's ok!
- ▶ Go back and find the bug in the square-root-scale model
- ▶ Re-fit the debugged model

Structure of the course

- ▶ Textbook: *Bayesian Data Analysis*, third edition, by Gelman, [Carlin], Stern, Dunson, Vehtari, and [Rubin]
- ▶ All other course material is at <http://www.stat.columbia.edu/~gelman/bda.course/>
- ▶ In class, we'll go through the book, section by section:

Structure of the course

- ▶ Textbook: *Bayesian Data Analysis*, third edition, by Gelman, [Carlin], Stern, Dunson, Vehtari, and [Rubin]
- ▶ All other course material is at <http://www.stat.columbia.edu/~gelman/bda.course/>
- ▶ In class, we'll go through the book, section by section:

▶ Chapter 1: Introduction
▶ Chapter 2: Data
▶ Chapter 3: Probability
▶ Chapter 4: Decision Theory

Structure of the course

- ▶ Textbook: *Bayesian Data Analysis*, third edition, by Gelman, [Carlin], Stern, Dunson, Vehtari, and [Rubin]
- ▶ All other course material is at <http://www.stat.columbia.edu/~gelman/bda.course/>
- ▶ In class, we'll go through the book, section by section:
 - ▶ Stories
 - ▶ Tips on what's tricky
 - ▶ Computer exercises

Structure of the course

- ▶ Textbook: *Bayesian Data Analysis*, third edition, by Gelman, [Carlin], Stern, Dunson, Vehtari, and [Rubin]
- ▶ All other course material is at <http://www.stat.columbia.edu/~gelman/bda.course/>
- ▶ In class, we'll go through the book, section by section:
 - ▶ Stories
 - ▶ Tips on what's tricky
 - ▶ Connections and open problems

Structure of the course

- ▶ Textbook: *Bayesian Data Analysis*, third edition, by Gelman, [Carlin], Stern, Dunson, Vehtari, and [Rubin]
- ▶ All other course material is at <http://www.stat.columbia.edu/~gelman/bda.course/>
- ▶ In class, we'll go through the book, section by section:
 - ▶ Stories
 - ▶ Tips on what's tricky
 - ▶ Connections and open problems

Structure of the course

- ▶ Textbook: *Bayesian Data Analysis*, third edition, by Gelman, [Carlin], Stern, Dunson, Vehtari, and [Rubin]
- ▶ All other course material is at <http://www.stat.columbia.edu/~gelman/bda.course/>
- ▶ In class, we'll go through the book, section by section:
 - ▶ Stories
 - ▶ Tips on what's tricky
 - ▶ Connections and open problems

Structure of the course

- ▶ Textbook: *Bayesian Data Analysis*, third edition, by Gelman, [Carlin], Stern, Dunson, Vehtari, and [Rubin]
- ▶ All other course material is at <http://www.stat.columbia.edu/~gelman/bda.course/>
- ▶ In class, we'll go through the book, section by section:
 - ▶ Stories
 - ▶ Tips on what's tricky
 - ▶ Connections and open problems

Class assignments

- ▶ Homeworks due at beginning of class 2b, 3b, ...
 - ▶ Assignments and data at <http://www.stat.columbia.edu/~gelman/bda.course/>
 - ▶ Each week: one theory problem, one computing problem, one applied problem
 - ▶ Hand in hard copies, no email
- ▶ Just-in-time-teaching assignments due one hour before every class
- ▶ Final exam

Class assignments

- ▶ Homeworks due at beginning of class 2b, 3b, ...
 - ▶ Assignments and data at <http://www.stat.columbia.edu/~gelman/bda.course/>
 - ▶ Each week: one theory problem, one computing problem, one applied problem
 - ▶ Hand in hard copies, no email
- ▶ Just-in-time-teaching assignments due one hour before every class
- ▶ Final exam

Class assignments

- ▶ Homeworks due at beginning of class 2b, 3b, ...
 - ▶ Assignments and data at <http://www.stat.columbia.edu/~gelman/bda.course/>
 - ▶ Each week: one theory problem, one computing problem, one applied problem
 - ▶ Hand in hard copies, no email
- ▶ Just-in-time-teaching assignments due one hour before every class
- ▶ Final exam

Class assignments

- ▶ Homeworks due at beginning of class 2b, 3b, ...
 - ▶ Assignments and data at <http://www.stat.columbia.edu/~gelman/bda.course/>
 - ▶ Each week: one theory problem, one computing problem, one applied problem
 - ▶ Hand in hard copies, no email
- ▶ Just-in-time-teaching assignments due one hour before every class
- ▶ Final exam

Class assignments

- ▶ Homeworks due at beginning of class 2b, 3b, ...
 - ▶ Assignments and data at <http://www.stat.columbia.edu/~gelman/bda.course/>
 - ▶ Each week: one theory problem, one computing problem, one applied problem
 - ▶ Hand in hard copies, no email
- ▶ Just-in-time-teaching assignments due one hour before every class
 - ▶ Testing background knowledge
 - ▶ Making sure you do the reading and are prepared for class
- ▶ Final exam

Class assignments

- ▶ Homeworks due at beginning of class 2b, 3b, ...
 - ▶ Assignments and data at <http://www.stat.columbia.edu/~gelman/bda.course/>
 - ▶ Each week: one theory problem, one computing problem, one applied problem
 - ▶ Hand in hard copies, no email
- ▶ Just-in-time-teaching assignments due one hour before every class
 - ▶ Testing background knowledge
 - ▶ Making sure you do the reading and are prepared for class
- ▶ Final exam

Class assignments

- ▶ Homeworks due at beginning of class 2b, 3b, ...
 - ▶ Assignments and data at <http://www.stat.columbia.edu/~gelman/bda.course/>
 - ▶ Each week: one theory problem, one computing problem, one applied problem
 - ▶ Hand in hard copies, no email
- ▶ Just-in-time-teaching assignments due one hour before every class
 - ▶ Testing background knowledge
 - ▶ Making sure you do the reading and are prepared for class
- ▶ Final exam

Class assignments

- ▶ Homeworks due at beginning of class 2b, 3b, ...
 - ▶ Assignments and data at <http://www.stat.columbia.edu/~gelman/bda.course/>
 - ▶ Each week: one theory problem, one computing problem, one applied problem
 - ▶ Hand in hard copies, no email
- ▶ Just-in-time-teaching assignments due one hour before every class
 - ▶ Testing background knowledge
 - ▶ Making sure you do the reading and are prepared for class
- ▶ Final exam
 - ▶ Take home
 - ▶ Comparable to homework problems

Class assignments

- ▶ Homeworks due at beginning of class 2b, 3b, ...
 - ▶ Assignments and data at <http://www.stat.columbia.edu/~gelman/bda.course/>
 - ▶ Each week: one theory problem, one computing problem, one applied problem
 - ▶ Hand in hard copies, no email
- ▶ Just-in-time-teaching assignments due one hour before every class
 - ▶ Testing background knowledge
 - ▶ Making sure you do the reading and are prepared for class
- ▶ Final exam
 - ▶ Take home
 - ▶ Comparable to homework problems

Class assignments

- ▶ Homeworks due at beginning of class 2b, 3b, ...
 - ▶ Assignments and data at <http://www.stat.columbia.edu/~gelman/bda.course/>
 - ▶ Each week: one theory problem, one computing problem, one applied problem
 - ▶ Hand in hard copies, no email
- ▶ Just-in-time-teaching assignments due one hour before every class
 - ▶ Testing background knowledge
 - ▶ Making sure you do the reading and are prepared for class
- ▶ Final exam
 - ▶ Take home
 - ▶ Comparable to homework problems

Class assignments

- ▶ Homeworks due at beginning of class 2b, 3b, ...
 - ▶ Assignments and data at <http://www.stat.columbia.edu/~gelman/bda.course/>
 - ▶ Each week: one theory problem, one computing problem, one applied problem
 - ▶ Hand in hard copies, no email
- ▶ Just-in-time-teaching assignments due one hour before every class
 - ▶ Testing background knowledge
 - ▶ Making sure you do the reading and are prepared for class
- ▶ Final exam
 - ▶ Take home
 - ▶ Comparable to homework problems

Your first jitt!

- ▶ Pull it up on the screen
- ▶ Spend 3 minutes on it
- ▶ Then we (briefly) discuss

Your first jitt!

- ▶ Pull it up on the screen
- ▶ Spend 3 minutes on it
- ▶ Then we (briefly) discuss

Your first jitt!

- ▶ Pull it up on the screen
- ▶ Spend 3 minutes on it
- ▶ Then we (briefly) discuss

Your first jitt!

- ▶ Pull it up on the screen
- ▶ Spend 3 minutes on it
- ▶ Then we (briefly) discuss

An example from toxicology

- ▶ Central story: 4-compartment model of toxicokinetics of perchloroethylene
- ▶ Bayesian inference combines prior information and data
- ▶ Unresolved questions
- ▶ How the model all fits together

An example from toxicology

- ▶ Central story: 4-compartment model of toxicokinetics of perchloroethylene
- ▶ Bayesian inference combines prior information and data
- ▶ Unresolved questions
- ▶ How the model all fits together

An example from toxicology

- ▶ Central story: 4-compartment model of toxicokinetics of perchloroethylene
- ▶ Bayesian inference combines prior information and data
- ▶ Unresolved questions
- ▶ How the model all fits together

An example from toxicology

- ▶ Central story: 4-compartment model of toxicokinetics of perchloroethylene
- ▶ Bayesian inference combines prior information and data
- ▶ Unresolved questions
- ▶ How the model all fits together

An example from toxicology

- ▶ Central story: 4-compartment model of toxicokinetics of perchloroethylene
- ▶ Bayesian inference combines prior information and data
- ▶ Unresolved questions
- ▶ How the model all fits together

Toxicokinetics of perchloroethylene

- ▶ Goal:
 - ▶ How much PERC is metabolized at low doses
 - ▶ Population distribution
- ▶ Experimental data: Expose 6 healthy volunteers to PERC for four hours, then measure concentrations in blood and air for 2 weeks
- ▶ 4-compartment model, metabolism in liver
- ▶ Our analysis:

Toxicokinetics of perchloroethylene

- ▶ Goal:
 - ▶ How much PERC is metabolized at low doses
 - ▶ Population distribution
- ▶ Experimental data: Expose 6 healthy volunteers to PERC for four hours, then measure concentrations in blood and air for 2 weeks
- ▶ 4-compartment model, metabolism in liver
- ▶ Our analysis:

Toxicokinetics of perchloroethylene

- ▶ Goal:
 - ▶ How much PERC is metabolized at low doses
 - ▶ Population distribution
- ▶ Experimental data: Expose 6 healthy volunteers to PERC for four hours, then measure concentrations in blood and air for 2 weeks
- ▶ 4-compartment model, metabolism in liver
- ▶ Our analysis:

Toxicokinetics of perchloroethylene

- ▶ Goal:
 - ▶ How much PERC is metabolized at low doses
 - ▶ Population distribution
- ▶ Experimental data: Expose 6 healthy volunteers to PERC for four hours, then measure concentrations in blood and air for 2 weeks
- ▶ 4-compartment model, metabolism in liver
- ▶ Our analysis:

Toxicokinetics of perchloroethylene

- ▶ Goal:
 - ▶ How much PERC is metabolized at low doses
 - ▶ Population distribution
- ▶ Experimental data: Expose 6 healthy volunteers to PERC for four hours, then measure concentrations in blood and air for 2 weeks
- ▶ 4-compartment model, metabolism in liver
- ▶ Our analysis:
 - ▶ We use a simple compartmental model with four compartments
 - ▶ We use a simple model for metabolism, which we treat as a random effect

Toxicokinetics of perchloroethylene

- ▶ Goal:
 - ▶ How much PERC is metabolized at low doses
 - ▶ Population distribution
- ▶ Experimental data: Expose 6 healthy volunteers to PERC for four hours, then measure concentrations in blood and air for 2 weeks
- ▶ 4-compartment model, metabolism in liver
- ▶ Our analysis:
 - ▶ Simple data-fitting did not work
 - ▶ Use Bayes to combine data and prior info within model

Toxicokinetics of perchloroethylene

- ▶ Goal:
 - ▶ How much PERC is metabolized at low doses
 - ▶ Population distribution
- ▶ Experimental data: Expose 6 healthy volunteers to PERC for four hours, then measure concentrations in blood and air for 2 weeks
- ▶ 4-compartment model, metabolism in liver
- ▶ Our analysis:
 - ▶ Simple data-fitting did not work
 - ▶ Use Bayes to combine data and prior info within model


Toxicokinetics of perchloroethylene

- ▶ Goal:
 - ▶ How much PERC is metabolized at low doses
 - ▶ Population distribution
- ▶ Experimental data: Expose 6 healthy volunteers to PERC for four hours, then measure concentrations in blood and air for 2 weeks
- ▶ 4-compartment model, metabolism in liver
- ▶ Our analysis:
 - ▶ Simple data-fitting did not work
 - ▶ Use Bayes to combine data and prior info within model



Toxicokinetics of perchloroethylene

- ▶ Goal:
 - ▶ How much PERC is metabolized at low doses
 - ▶ Population distribution
- ▶ Experimental data: Expose 6 healthy volunteers to PERC for four hours, then measure concentrations in blood and air for 2 weeks
- ▶ 4-compartment model, metabolism in liver
- ▶ Our analysis:
 - ▶ Simple data-fitting did not work
 - ▶ Use Bayes to combine data and prior info within model

4-compartment model



Some data



Connections to general Bayesian principles

- ▶ Sometimes the model comes first, based on substantive considerations (toxicology, economics, ...)
- ▶ Sometimes the model is chosen based on the data collection (traditional statistics of surveys and experiments)
- ▶ Other times the data come first (other statistics examples)
- ▶ Usually it's a mix

Connections to general Bayesian principles

- ▶ Sometimes the model comes first, based on substantive considerations (toxicology, economics, . . .)
- ▶ Sometimes the model is chosen based on the data collection (traditional statistics of surveys and experiments)
- ▶ Other times the data come first (other statistics examples)
- ▶ Usually it's a mix

Connections to general Bayesian principles

- ▶ Sometimes the model comes first, based on substantive considerations (toxicology, economics, . . .)
- ▶ Sometimes the model is chosen based on the data collection (traditional statistics of surveys and experiments)
- ▶ Other times the data come first (other statistics examples)
- ▶ Usually it's a mix
 - ▶ Discuss an example of prediction
 - ▶ Discuss an example of causal inference

Connections to general Bayesian principles

- ▶ Sometimes the model comes first, based on substantive considerations (toxicology, economics, . . .)
- ▶ Sometimes the model is chosen based on the data collection (traditional statistics of surveys and experiments)
- ▶ Other times the data come first (other statistics examples)
- ▶ Usually it's a mix
 - ▶ Discuss an example of prediction
 - ▶ Discuss an example of causal inference

Connections to general Bayesian principles

- ▶ Sometimes the model comes first, based on substantive considerations (toxicology, economics, . . .)
- ▶ Sometimes the model is chosen based on the data collection (traditional statistics of surveys and experiments)
- ▶ Other times the data come first (other statistics examples)
- ▶ Usually it's a mix
 - ▶ Discuss an example of prediction
 - ▶ Discuss an example of causal inference

Connections to general Bayesian principles

- ▶ Sometimes the model comes first, based on substantive considerations (toxicology, economics, . . .)
- ▶ Sometimes the model is chosen based on the data collection (traditional statistics of surveys and experiments)
- ▶ Other times the data come first (other statistics examples)
- ▶ Usually it's a mix
 - ▶ Discuss an example of prediction
 - ▶ Discuss an example of causal inference

Simple statistical ideas did not work

- ▶ Fitting 4-compartment model directly to data
- ▶ Assisted model fit
- ▶ 1 or 2-compartment model
- ▶ Simulation from prior distribution

Simple statistical ideas did not work

- ▶ Fitting 4-compartment model directly to data
- ▶ Assisted model fit
- ▶ 1 or 2-compartment model
- ▶ Simulation from prior distribution

Simple statistical ideas did not work

- ▶ Fitting 4-compartment model directly to data
- ▶ Assisted model fit
- ▶ 1 or 2-compartment model
- ▶ Simulation from prior distribution

Simple statistical ideas did not work

- ▶ Fitting 4-compartment model directly to data
- ▶ Assisted model fit
- ▶ 1 or 2-compartment model
- ▶ Simulation from prior distribution

Simple statistical ideas did not work

- ▶ Fitting 4-compartment model directly to data
- ▶ Assisted model fit
- ▶ 1 or 2-compartment model
- ▶ Simulation from prior distribution

Simple statistical ideas that did not work:

Fitting 4-compartment model directly to data

- ▶ Nonlinear least squares
- ▶ Fitting to each person separately:
 - ▶ One fit to every 30 data points, 15 parameters
 - ▶ “15 by 15”
- ▶ Pooling data and estimating parameters for “the standard man”

Simple statistical ideas that did not work:

Fitting 4-compartment model directly to data

- ▶ Nonlinear least squares
- ▶ Fitting to each person separately:
 - ▶ Unstable: approx 30 data points, 15 param
 - ▶ “8 kg liver”
- ▶ Pooling data and estimating parameters for “the standard man”

Simple statistical ideas that did not work:

Fitting 4-compartment model directly to data

- ▶ Nonlinear least squares
- ▶ Fitting to each person separately:
 - ▶ Unstable: approx 30 data points, 15 param
 - ▶ “8 kg liver”
- ▶ Pooling data and estimating parameters for “the standard man”

Simple statistical ideas that did not work:

Fitting 4-compartment model directly to data

- ▶ Nonlinear least squares
- ▶ Fitting to each person separately:
 - ▶ Unstable: approx 30 data points, 15 param
 - ▶ “8 kg liver”
- ▶ Pooling data and estimating parameters for “the standard man”

Simple statistical ideas that did not work:

Fitting 4-compartment model directly to data

- ▶ Nonlinear least squares
- ▶ Fitting to each person separately:
 - ▶ Unstable: approx 30 data points, 15 param
 - ▶ “8 kg liver”
- ▶ Pooling data and estimating parameters for “the standard man”
 - ▶ Not useful for our goal of population inference

Simple statistical ideas that did not work:

Fitting 4-compartment model directly to data

- ▶ Nonlinear least squares
- ▶ Fitting to each person separately:
 - ▶ Unstable: approx 30 data points, 15 param
 - ▶ “8 kg liver”
- ▶ Pooling data and estimating parameters for “the standard man”
 - ▶ Not useful for our goal of population inference

Simple statistical ideas that did not work:

Fitting 4-compartment model directly to data

- ▶ Nonlinear least squares
- ▶ Fitting to each person separately:
 - ▶ Unstable: approx 30 data points, 15 param
 - ▶ “8 kg liver”
- ▶ Pooling data and estimating parameters for “the standard man”
 - ▶ Not useful for our goal of population inference

Simple statistical ideas that did not work: Assisted model fit

- ▶ Set some parameters to fixed values (from the pharmacology literature)
- ▶ Estimate the other parameters
- ▶ Results:

Simple statistical ideas that did not work:

Assisted model fit

- ▶ Set some parameters to fixed values (from the pharmacology literature)
- ▶ Estimate the other parameters
- ▶ Results:

▶ The model fit the data well

▶ But it was not clear how well the model fit the data

▶ The model fit the data well

Simple statistical ideas that did not work: Assisted model fit

- ▶ Set some parameters to fixed values (from the pharmacology literature)
- ▶ Estimate the other parameters
- ▶ Results:
 - ▶ Couldn't fit the data well
 - ▶ Difficult to get fixed values for PERC-specific parameters such as equilibrium concentration ratios

Simple statistical ideas that did not work: Assisted model fit

- ▶ Set some parameters to fixed values (from the pharmacology literature)
- ▶ Estimate the other parameters
- ▶ Results:
 - ▶ Couldn't fit the data well
 - ▶ Difficult to get fixed values for PERC-specific parameters such as equilibrium concentration ratios

Simple statistical ideas that did not work:

Assisted model fit

- ▶ Set some parameters to fixed values (from the pharmacology literature)
- ▶ Estimate the other parameters
- ▶ Results:
 - ▶ Couldn't fit the data well
 - ▶ Difficult to get fixed values for PERC-specific parameters such as equilibrium concentration ratios

Simple statistical ideas that did not work:

Assisted model fit

- ▶ Set some parameters to fixed values (from the pharmacology literature)
- ▶ Estimate the other parameters
- ▶ Results:
 - ▶ Couldn't fit the data well
 - ▶ Difficult to get fixed values for PERC-specific parameters such as equilibrium concentration ratios

Simple statistical ideas that did not work:

1 or 2-compartment model

- ▶ Simpler model can be estimated easily and robustly
- ▶ Does not fit the data well
 - Most of the ITCU leaves in a few hours, but some stays in the body after a week or more
- ▶ Not realistic for low-dose extrapolation

Simple statistical ideas that did not work: 1 or 2-compartment model

- ▶ Simpler model can be estimated easily and robustly
- ▶ Does not fit the data well
 - ▶ Most of the PERC leaves in a few hours, but some stays in the body after a week or more
- ▶ Not realistic for low-dose extrapolation

Simple statistical ideas that did not work: 1 or 2-compartment model

- ▶ Simpler model can be estimated easily and robustly
- ▶ Does not fit the data well
 - ▶ Most of the PERC leaves in a few hours, but some stays in the body after a week or more
 - ▶ Not realistic for low-dose extrapolation

Simple statistical ideas that did not work: 1 or 2-compartment model

- ▶ Simpler model can be estimated easily and robustly
- ▶ Does not fit the data well
 - ▶ Most of the PERC leaves in a few hours, but some stays in the body after a week or more
- ▶ Not realistic for low-dose extrapolation

Simple statistical ideas that did not work: 1 or 2-compartment model

- ▶ Simpler model can be estimated easily and robustly
- ▶ Does not fit the data well
 - ▶ Most of the PERC leaves in a few hours, but some stays in the body after a week or more
- ▶ Not realistic for low-dose extrapolation

Simple statistical ideas that did not work: Simulation from prior distribution

- ▶ Get prior information on parameters from pharmacology literature
- ▶ Try to fit data within these prior constraints
- ▶ Does not fit the data well
- ▶ Difficult to get good prior information for PERC-specific parameters such as equilibrium concentration ratios

Simple statistical ideas that did not work: Simulation from prior distribution

- ▶ Get prior information on parameters from pharmacology literature
- ▶ Try to fit data within these prior constraints
- ▶ Does not fit the data well
- ▶ Difficult to get good prior information for PERC-specific parameters such as equilibrium concentration ratios

Simple statistical ideas that did not work: Simulation from prior distribution

- ▶ Get prior information on parameters from pharmacology literature
- ▶ Try to fit data within these prior constraints
- ▶ Does not fit the data well
- ▶ Difficult to get good prior information for PERC-specific parameters such as equilibrium concentration ratios

Simple statistical ideas that did not work: Simulation from prior distribution

- ▶ Get prior information on parameters from pharmacology literature
- ▶ Try to fit data within these prior constraints
- ▶ Does not fit the data well
- ▶ Difficult to get good prior information for PERC-specific parameters such as equilibrium concentration ratios

Simple statistical ideas that did not work: Simulation from prior distribution

- ▶ Get prior information on parameters from pharmacology literature
- ▶ Try to fit data within these prior constraints
- ▶ Does not fit the data well
- ▶ Difficult to get good prior information for PERC-specific parameters such as equilibrium concentration ratios

Bayesian inference

- ▶ 4-compartment model
 - ▶ 15 parameters for each person
 - ▶ Prior information
-
- ▶ Posterior simulation: random walk through parameter space
 - ▶ Inference for parameters and predictions
 - ▶ Model checking

Bayesian inference

- ▶ 4-compartment model
- ▶ 15 parameters for each person
- ▶ Prior information
 - ▶ Strong for some parameters (e.g., volume of blood)
 - ▶ Weak for others (e.g., Michaelis-Menten const)
 - ▶ Model includes uncertainty and variation
- ▶ Posterior simulation: random walk through parameter space
- ▶ Inference for parameters and predictions
- ▶ Model checking

Bayesian inference

- ▶ 4-compartment model
- ▶ 15 parameters for each person
- ▶ Prior information
 - ▶ Strong for some parameters (e.g., volume of liver)
 - ▶ Weak for others (e.g., Michaelis-Menten coef)
 - ▶ Model includes uncertainty and variation
- ▶ Posterior simulation: random walk through parameter space
- ▶ Inference for parameters and predictions
- ▶ Model checking

Bayesian inference

- ▶ 4-compartment model
- ▶ 15 parameters for each person
- ▶ Prior information
 - ▶ Strong for some parameters (e.g., volume of liver)
 - ▶ Weak for others (e.g., Michaelis-Menten coef)
 - ▶ Model includes uncertainty and variation
- ▶ Posterior simulation: random walk through parameter space
- ▶ Inference for parameters and predictions
- ▶ Model checking

Bayesian inference

- ▶ 4-compartment model
- ▶ 15 parameters for each person
- ▶ Prior information
 - ▶ Strong for some parameters (e.g., volume of liver)
 - ▶ Weak for others (e.g., Michaelis-Menten coef)
 - ▶ Model includes uncertainty and variation
- ▶ Posterior simulation: random walk through parameter space
- ▶ Inference for parameters and predictions
- ▶ Model checking

Bayesian inference

- ▶ 4-compartment model
- ▶ 15 parameters for each person
- ▶ Prior information
 - ▶ Strong for some parameters (e.g., volume of liver)
 - ▶ Weak for others (e.g., Michaelis-Menten coef)
 - ▶ Model includes uncertainty and variation
- ▶ Posterior simulation: random walk through parameter space
- ▶ Inference for parameters and predictions
- ▶ Model checking

Bayesian inference

- ▶ 4-compartment model
- ▶ 15 parameters for each person
- ▶ Prior information
 - ▶ Strong for some parameters (e.g., volume of liver)
 - ▶ Weak for others (e.g., Michaelis-Menten coef)
 - ▶ Model includes uncertainty and variation
- ▶ Posterior simulation: random walk through parameter space
- ▶ Inference for parameters and predictions
- ▶ Model checking

Bayesian inference

- ▶ 4-compartment model
- ▶ 15 parameters for each person
- ▶ Prior information
 - ▶ Strong for some parameters (e.g., volume of liver)
 - ▶ Weak for others (e.g., Michaelis-Menten coef)
 - ▶ Model includes uncertainty and variation
- ▶ Posterior simulation: random walk through parameter space
- ▶ Inference for parameters and predictions
- ▶ Model checking

Bayesian inference

- ▶ 4-compartment model
- ▶ 15 parameters for each person
- ▶ Prior information
 - ▶ Strong for some parameters (e.g., volume of liver)
 - ▶ Weak for others (e.g., Michaelis-Menten coef)
 - ▶ Model includes uncertainty and variation
- ▶ Posterior simulation: random walk through parameter space
- ▶ Inference for parameters and predictions
- ▶ Model checking

Bayesian inference

- ▶ 4-compartment model
- ▶ 15 parameters for each person
- ▶ Prior information
 - ▶ Strong for some parameters (e.g., volume of liver)
 - ▶ Weak for others (e.g., Michaelis-Menten coef)
 - ▶ Model includes uncertainty and variation
- ▶ Posterior simulation: random walk through parameter space
- ▶ Inference for parameters and predictions
- ▶ Model checking

Hierarchical prior distributions

- ▶ Prior distribution for a rate parameter in the metabolism, θ_j for person j
 - ▶ $\log \theta_j \sim N(\mu, \tau^2)$
 - ▶ $\mu \sim N(\log 16, (\log 10)^2)$
 - ▶ $\tau \sim \log 2$
- ▶ Large uncertainty, small variation
- ▶ Can learn about μ using data from several people
- ▶ Can't do this without a hierarchical model
- ▶ Transformations and prior correlations (why transformations are particularly important for Bayesians)

Hierarchical prior distributions

- ▶ Prior distribution for a rate parameter in the metabolism, θ_j for person j
 - ▶ $\log \theta_j \sim N(\mu, \tau^2)$
 - ▶ $\mu \sim N(\log 16, (\log 10)^2)$
 - ▶ $\tau \approx \log 2$
- ▶ Large uncertainty, small variation
- ▶ Can learn about μ using data from several people
- ▶ Can't do this without a hierarchical model
- ▶ Transformations and prior correlations (why transformations are particularly important for Bayesians)

Hierarchical prior distributions

- ▶ Prior distribution for a rate parameter in the metabolism, θ_j for person j
 - ▶ $\log \theta_j \sim N(\mu, \tau^2)$
 - ▶ $\mu \sim N(\log 16, (\log 10)^2)$
 - ▶ $\tau \approx \log 2$
- ▶ Large uncertainty, small variation
- ▶ Can learn about μ using data from several people
- ▶ Can't do this without a hierarchical model
- ▶ Transformations and prior correlations (why transformations are particularly important for Bayesians)

Hierarchical prior distributions

- ▶ Prior distribution for a rate parameter in the metabolism, θ_j for person j
 - ▶ $\log \theta_j \sim N(\mu, \tau^2)$
 - ▶ $\mu \sim N(\log 16, (\log 10)^2)$
 - ▶ $\tau \approx \log 2$
- ▶ Large uncertainty, small variation
- ▶ Can learn about μ using data from several people
- ▶ Can't do this without a hierarchical model
- ▶ Transformations and prior correlations (why transformations are particularly important for Bayesians)

Hierarchical prior distributions

- ▶ Prior distribution for a rate parameter in the metabolism, θ_j for person j
 - ▶ $\log \theta_j \sim N(\mu, \tau^2)$
 - ▶ $\mu \sim N(\log 16, (\log 10)^2)$
 - ▶ $\tau \approx \log 2$
- ▶ Large uncertainty, small variation
- ▶ Can learn about μ using data from several people
- ▶ Can't do this without a hierarchical model
- ▶ Transformations and prior correlations (why transformations are particularly important for Bayesians)

Hierarchical prior distributions

- ▶ Prior distribution for a rate parameter in the metabolism, θ_j for person j
 - ▶ $\log \theta_j \sim N(\mu, \tau^2)$
 - ▶ $\mu \sim N(\log 16, (\log 10)^2)$
 - ▶ $\tau \approx \log 2$
- ▶ Large uncertainty, small variation
- ▶ Can learn about μ using data from several people
- ▶ Can't do this without a hierarchical model
- ▶ Transformations and prior correlations (why transformations are particularly important for Bayesians)

Hierarchical prior distributions

- ▶ Prior distribution for a rate parameter in the metabolism, θ_j for person j
 - ▶ $\log \theta_j \sim N(\mu, \tau^2)$
 - ▶ $\mu \sim N(\log 16, (\log 10)^2)$
 - ▶ $\tau \approx \log 2$
- ▶ Large uncertainty, small variation
- ▶ Can learn about μ using data from several people
- ▶ Can't do this without a hierarchical model
- ▶ Transformations and prior correlations (why transformations are particularly important for Bayesians)

Hierarchical prior distributions

- ▶ Prior distribution for a rate parameter in the metabolism, θ_j for person j
 - ▶ $\log \theta_j \sim N(\mu, \tau^2)$
 - ▶ $\mu \sim N(\log 16, (\log 10)^2)$
 - ▶ $\tau \approx \log 2$
- ▶ Large uncertainty, small variation
- ▶ Can learn about μ using data from several people
- ▶ Can't do this without a hierarchical model
- ▶ Transformations and prior correlations (why transformations are particularly important for Bayesians)

Hierarchical prior distributions

- ▶ Prior distribution for a rate parameter in the metabolism, θ_j for person j
 - ▶ $\log \theta_j \sim N(\mu, \tau^2)$
 - ▶ $\mu \sim N(\log 16, (\log 10)^2)$
 - ▶ $\tau \approx \log 2$
- ▶ Large uncertainty, small variation
- ▶ Can learn about μ using data from several people
- ▶ Can't do this without a hierarchical model
- ▶ Transformations and prior correlations (why transformations are particularly important for Bayesians)

Hierarchical prior distributions

<i>Parameter</i>	<i>Population prior</i>		
Ventilation/perfusion ratio (VPR)	$1.6(\times \div 1.3)$ $\times \div 1.3$		
Blood flow, well-perfused tissues (Fwp)	$.47(\times \div 1.17)$ $\times \div 1.17$	Partition coeff, blood/air (Pba)	$12(\times \div 1.5)$ $\times \div 1.3$
Blood flow, poorly perfused tissues (Fpp)	$.20(\times \div 1.22)$ $\times \div 1.22$	Partition coeff, well-perfused (Pwp)	$4.8(\times \div 1.5)$ $\times \div 1.3$
Blood flow, fat (Ff)	$.07(\times \div 1.27)$ $\times \div 1.27$	Partition coeff, poorly perfused (Ppp)	$1.6(\times \div 1.5)$ $\times \div 1.3$
Blood flow, liver (Fl)	$.25(\times \div 1.15)$ $\times \div 1.15$	Partition coeff, fat (Pf)	$125(\times \div 1.5)$ $\times \div 1.3$
Volume, well-perfused tissues (Vwp)	$.27(\times \div 1.36)$ $\times \div 1.36$	Partition coeff, liver (Pl)	$4.8(\times \div 1.5)$ $\times \div 1.3$
Volume, poorly perfused tissues (Vpp)	$.55(\times \div 1.17)$ $\times \div 1.17$	Max metabolic rate in liver (VMI)	$.042(\times \div 10)$ $\times \div 2$
Volume, liver (VI)	$.033(\times \div 1.1)$ $\times \div 1.1$	K_m in liver (KMI)	$16(\times \div 10)$ $\times \div 1.5$

What we did

- ▶ Set up a hierarchical prior distribution with uncertainty and population variation for a 4-compartment model
- ▶ Fit the model to data (much computation)
- ▶ Checked inferences about parameters to see that they made sense
- ▶ Re-ran model under hypothetical low-dose, high-dose exposures

What we did

- ▶ Set up a hierarchical prior distribution with uncertainty and population variation for a 4-compartment model
- ▶ Fit the model to data (much computation)
- ▶ Checked inferences about parameters to see that they made sense
- ▶ Re-ran model under hypothetical low-dose, high-dose exposures

What we did

- ▶ Set up a hierarchical prior distribution with uncertainty and population variation for a 4-compartment model
- ▶ Fit the model to data (much computation)
- ▶ Checked inferences about parameters to see that they made sense
- ▶ Re-ran model under hypothetical low-dose, high-dose exposures

What we did

- ▶ Set up a hierarchical prior distribution with uncertainty and population variation for a 4-compartment model
- ▶ Fit the model to data (much computation)
- ▶ Checked inferences about parameters to see that they made sense
- ▶ Re-ran model under hypothetical low-dose, high-dose exposures

What we did

- ▶ Set up a hierarchical prior distribution with uncertainty and population variation for a 4-compartment model
- ▶ Fit the model to data (much computation)
- ▶ Checked inferences about parameters to see that they made sense
- ▶ Re-ran model under hypothetical low-dose, high-dose exposures

Fitting and using the model

- ▶ Use Gibbs sampler and Metropolis algorithm to take a random walk through parameter space
- ▶ Computationally intensive
 - ▶ Each step requires evaluation of the numerical differential equation solver
- ▶ Check inferences: Do they make sense?
- ▶ Re-run the model several times to simulate what would happen under different conditions

Fitting and using the model

- ▶ Use Gibbs sampler and Metropolis algorithm to take a random walk through parameter space
- ▶ Computationally intensive
 - ▶ Each step requires evaluation of the numerical differential equation solver
- ▶ Check inferences: Do they make sense?
- ▶ Re-run the model several times to simulate what would happen under different conditions

Fitting and using the model

- ▶ Use Gibbs sampler and Metropolis algorithm to take a random walk through parameter space
- ▶ Computationally intensive
 - ▶ Each step requires evaluation of the numerical differential equation solver
- ▶ Check inferences: Do they make sense?
- ▶ Re-run the model several times to simulate what would happen under different conditions

Fitting and using the model

- ▶ Use Gibbs sampler and Metropolis algorithm to take a random walk through parameter space
- ▶ Computationally intensive
 - ▶ Each step requires evaluation of the numerical differential equation solver
- ▶ Check inferences: Do they make sense?
- ▶ Re-run the model several times to simulate what would happen under different conditions


Fitting and using the model

- ▶ Use Gibbs sampler and Metropolis algorithm to take a random walk through parameter space
- ▶ Computationally intensive
 - ▶ Each step requires evaluation of the numerical differential equation solver
- ▶ Check inferences: Do they make sense?
- ▶ Re-run the model several times to simulate what would happen under different conditions


Fitting and using the model

- ▶ Use Gibbs sampler and Metropolis algorithm to take a random walk through parameter space
- ▶ Computationally intensive
 - ▶ Each step requires evaluation of the numerical differential equation solver
- ▶ Check inferences: Do they make sense?
- ▶ Re-run the model several times to simulate what would happen under different conditions


Inference for 6 individuals




Inference for the population




Prediction of data from a new study





Sensitivity to priors



(a)



(b)



Putting it all together

- (a) Physiological pharmacokinetic model
- (b) Hierarchical population model
- (c) Prior information
- (d) Experimental data
- (e) Bayesian inference
- (f) Computation
- (g) Model checking

► We need all of these!

Putting it all together

- (a) Physiological pharmacokinetic model
- (b) Hierarchical population model
- (c) Prior information
- (d) Experimental data
- (e) Bayesian inference
- (f) Computation
- (g) Model checking

► We need all of these!

Putting it all together

- (a) Physiological pharmacokinetic model
- (b) Hierarchical population model
- (c) Prior information
- (d) Experimental data
- (e) Bayesian inference
- (f) Computation
- (g) Model checking

► We need all of these!

Putting it all together

- (a) Physiological pharmacokinetic model
- (b) Hierarchical population model
- (c) Prior information
- (d) Experimental data
- (e) Bayesian inference
- (f) Computation
- (g) Model checking

► We need all of these!

Putting it all together

- (a) Physiological pharmacokinetic model
- (b) Hierarchical population model
- (c) Prior information
- (d) Experimental data
- (e) Bayesian inference
- (f) Computation
- (g) Model checking

► We need all of these!

Putting it all together

- (a) Physiological pharmacokinetic model
- (b) Hierarchical population model
- (c) Prior information
- (d) Experimental data
- (e) Bayesian inference
- (f) Computation
- (g) Model checking

► We need all of these!

Putting it all together

- (a) Physiological pharmacokinetic model
- (b) Hierarchical population model
- (c) Prior information
- (d) Experimental data
- (e) Bayesian inference
- (f) Computation
- (g) Model checking

► We need all of these!

Putting it all together

- (a) Physiological pharmacokinetic model
- (b) Hierarchical population model
- (c) Prior information
- (d) Experimental data
- (e) Bayesian inference
- (f) Computation
- (g) Model checking

► We need all of these!

Putting it all together

- (a) Physiological pharmacokinetic model
- (b) Hierarchical population model
- (c) Prior information
- (d) Experimental data
- (e) Bayesian inference
- (f) Computation
- (g) Model checking

► We need all of these!

(a) Physiological pharmacokinetic model

- ▶ Without a physiological model, there is no good way to get prior information on the parameters
- ▶ We need physiological parameters (not just curve-fitting of the data) to efficiently combine information across different people

(a) Physiological pharmacokinetic model

- ▶ Without a physiological model, there is no good way to get prior information on the parameters
- ▶ We need physiological parameters (not just curve-fitting of the data) to efficiently combine information across different people

(a) Physiological pharmacokinetic model

- ▶ Without a physiological model, there is no good way to get prior information on the parameters
- ▶ We need physiological parameters (not just curve-fitting of the data) to efficiently combine information across different people

(b) Hierarchical population model

- ▶ Without a population model, there generally are not enough data to estimate the parameters separately for each individual
- ▶ And there is too much variation among bodies (even among healthy young male volunteers) to pool all the data together and estimate common parameters

(b) Hierarchical population model

- ▶ Without a population model, there generally are not enough data to estimate the parameters separately for each individual
- ▶ And there is too much variation among bodies (even among healthy young male volunteers) to pool all the data together and estimate common parameters

(b) Hierarchical population model

- ▶ Without a population model, there generally are not enough data to estimate the parameters separately for each individual
- ▶ And there is too much variation among bodies (even among healthy young male volunteers) to pool all the data together and estimate common parameters

- (c) Prior information
- (d) Experimental data

- ▶ We need prior information. Otherwise, our estimates don't make sense (the 8 kg liver)
- ▶ We need experimental data to learn about perchloroethylene in particular

- (c) Prior information
- (d) Experimental data

- ▶ We need prior information. Otherwise, our estimates don't make sense (the 8 kg liver)
- ▶ We need experimental data to learn about perchloroethylene in particular

- (c) Prior information
- (d) Experimental data

- ▶ We need prior information. Otherwise, our estimates don't make sense (the 8 kg liver)
- ▶ We need experimental data to learn about perchloroethylene in particular

(e) Bayesian inference

- ▶ Using Bayesian inference, we can find parameters that are consistent with both prior information and data, if such agreement is possible
- ▶ Automatically includes uncertainty and variability, so inferences can be plugged in directly to risk assessment and decision analysis

(e) Bayesian inference

- ▶ Using Bayesian inference, we can find parameters that are consistent with both prior information and data, if such agreement is possible
- ▶ Automatically includes uncertainty and variability, so inferences can be plugged in directly to risk assessment and decision analysis

(e) Bayesian inference

- ▶ Using Bayesian inference, we can find parameters that are consistent with both prior information and data, if such agreement is possible
- ▶ Automatically includes uncertainty and variability, so inferences can be plugged in directly to risk assessment and decision analysis

(f) Computation

- ▶ Our models are big. Least squares, maximum likelihood, etc., are not enough
- ▶ Old-fashioned differential-equation solver is still sitting inside the model
- ▶ Our computers are never fast enough. We want more, more, more!

(f) Computation

- ▶ Our models are big. Least squares, maximum likelihood, etc., are not enough
- ▶ Old-fashioned differential-equation solver is still sitting inside the model
- ▶ Our computers are never fast enough. We want more, more, more!

(f) Computation

- ▶ Our models are big. Least squares, maximum likelihood, etc., are not enough
- ▶ Old-fashioned differential-equation solver is still sitting inside the model
- ▶ Our computers are never fast enough. We want more, more, more!

(f) Computation

- ▶ Our models are big. Least squares, maximum likelihood, etc., are not enough
- ▶ Old-fashioned differential-equation solver is still sitting inside the model
- ▶ Our computers are never fast enough. We want more, more, more!

(g) Model checking

- ▶ Check inferences about parameters
 - ▶ Do they make sense?
 - ▶ Are they consistent with prior distributions
- ▶ Check fit to data
- ▶ Check predictions on new data

(g) Model checking

- ▶ Check inferences about parameters
 - ▶ Do they make sense?
 - ▶ Are they consistent with prior distributions
- ▶ Check fit to data
- ▶ Check predictions on new data

(g) Model checking

- ▶ Check inferences about parameters
 - ▶ Do they make sense?
 - ▶ Are they consistent with prior distributions
- ▶ Check fit to data
- ▶ Check predictions on new data

(g) Model checking

- ▶ Check inferences about parameters
 - ▶ Do they make sense?
 - ▶ Are they consistent with prior distributions
- ▶ Check fit to data
- ▶ Check predictions on new data

(g) Model checking

- ▶ Check inferences about parameters
 - ▶ Do they make sense?
 - ▶ Are they consistent with prior distributions
- ▶ Check fit to data
- ▶ Check predictions on new data

Using Bayesian ideas to improve existing analyses

- ▶ Regularization (for example, avoiding estimates on the boundary of parameter space)
- ▶ Accounting for uncertainty (especially for decisions)
- ▶ Checking model fit
- ▶ Using models to combine different sources of information (partial pooling)
- ▶ Better dialogue with subject-matter experts (more windows into the model and data)

Using Bayesian ideas to improve existing analyses

- ▶ Regularization (for example, avoiding estimates on the boundary of parameter space)
- ▶ Accounting for uncertainty (especially for decisions)
- ▶ Checking model fit
- ▶ Using models to combine different sources of information (partial pooling)
- ▶ Better dialogue with subject-matter experts (more windows into the model and data)

Using Bayesian ideas to improve existing analyses

- ▶ Regularization (for example, avoiding estimates on the boundary of parameter space)
- ▶ Accounting for uncertainty (especially for decisions)
- ▶ Checking model fit
- ▶ Using models to combine different sources of information (partial pooling)
- ▶ Better dialogue with subject-matter experts (more windows into the model and data)

Using Bayesian ideas to improve existing analyses

- ▶ Regularization (for example, avoiding estimates on the boundary of parameter space)
- ▶ Accounting for uncertainty (especially for decisions)
- ▶ Checking model fit
- ▶ Using models to combine different sources of information (partial pooling)
- ▶ Better dialogue with subject-matter experts (more windows into the model and data)

Using Bayesian ideas to improve existing analyses

- ▶ Regularization (for example, avoiding estimates on the boundary of parameter space)
- ▶ Accounting for uncertainty (especially for decisions)
- ▶ Checking model fit
- ▶ Using models to combine different sources of information (partial pooling)
- ▶ Better dialogue with subject-matter experts (more windows into the model and data)

Using Bayesian ideas to improve existing analyses

- ▶ Regularization (for example, avoiding estimates on the boundary of parameter space)
- ▶ Accounting for uncertainty (especially for decisions)
- ▶ Checking model fit
- ▶ Using models to combine different sources of information (partial pooling)
- ▶ Better dialogue with subject-matter experts (more windows into the model and data)

Summary of example

- ▶ Population pharmacokinetic models have many moving parts
- ▶ Complexity in one place can make it *easier*, not harder, to add information in other places

Summary of example

- ▶ Population pharmacokinetic models have many moving parts
- ▶ Complexity in one place can make it *easier*, not harder, to add information in other places

Summary of example

- ▶ Population pharmacokinetic models have many moving parts
- ▶ Complexity in one place can make it *easier*, not harder, to add information in other places