

Bcl-6 controlled T_{FH} polarization and memory: the known unknowns

Hai Qi, Dan Liu, Weiwei Ma, Yifeng Wang and Hu Yan

Upon antigenic activation *in vivo*, naïve CD4 T cells can differentiate into one of several helper (Th) subsets under the control of lineage-specifying transcription factors to tailor immune responses against different types of pathogens. Follicular T-helper (T_{FH}) cells are a recently defined subset that is controlled by Bcl-6 and specializes in promoting B cell-mediated humoral immunity. T_{FH} cells exhibit unique spatiotemporal and functional features, but it is not settled as to how Bcl-6 promotes the T_{FH} development, how T_{FH} cells relate to other Th subsets, and how T_{FH} cells relate to memory. Here we review recent advances and crucial gaps in our understanding of Bcl-6-controlled T_{FH} development and function.

Addresses

Tsinghua-Peking Center for Life Sciences, Laboratory of Dynamic Immunobiology, School of Medicine, School of Life Sciences, Tsinghua University, Beijing 100084, PR China

Corresponding author: Qi, Hai (qihai@biomed.tsinghua.edu.cn)

Current Opinion in Immunology 2014, 28:34–41

This review comes from a themed issue on **Lymphocyte activation and effector functions**

Edited by **Claude-Agnès Reynaud** and **Stuart Tangye**

For a complete overview see the [Issue](#) and the [Editorial](#)

Available online 28th February 2014

0952-7915/\$ – see front matter, © 2014 Elsevier Ltd. All rights reserved.

<http://dx.doi.org/10.1016/j.coi.2014.01.016>

Introduction

The most distinct feature of T_{FH} cells is the follicular site of helper actions [1,2]. As neutralizing antibodies re-emerge as the hope for anti-viral vaccines [3] and dysregulated T_{FH} biology is increasingly implicated in diseases [4], how T_{FH} cells develop, acquire localization features, and function to regulate the quality and magnitude of humoral immunity has become not only a fascinating basic question but also a key area of potential translational research [5,6]. As such, the T_{FH} subset has been the main topic of more than 30 review articles since 2011. Not to be excessively repetitive, we synthesize recent results on mapping T_{FH} development and its relationship with memory formation. We highlight conceptual roadblocks on our way forward to better understand the essence of a Bcl-6-controlled T_{FH} program.

GC T_{FH} cells and T_{FH}-associated molecules


Whereas histologically defined, GC-associated Th cells are *bona fide* GC T_{FH} cells, CXCR5⁺ Th cells at the T–B border

can engage in cognate interactions with antigen-specific B cells [7[•],8[•]], promote B cell proliferation and differentiation [9,10[•]], and are considered pre-GC T_{FH} cells [1]. These cells contain precursors that subsequently become GC T_{FH} cells (Figure 1). GC accumulation of T_{FH} cells, as identified by the CXCR5⁺PD-1⁺ phenotype [11] or by dynamic visualization in the GC structure [9], requires cognate interactions with antigen-specific B cells [12,13[•]], in part reflecting their dependence on continuous antigen presentation inside the follicle and GCs [14,15[•]]. CXCR5⁺PD-1⁺ T cells fail to develop without T cell-intrinsic Bcl-6 expression, and Bcl-6 overexpression promotes the T_{FH} phenotype [16[•],17[•],18[•]]. By histology, Yu *et al.* showed a complete absence of Bcl-6^{−/−} Th cells from GCs formed in Bcl-6^{+/+};Bcl-6^{−/−} mixed bone-marrow chimera [18[•]]. These data establish that the absence of Bcl-6^{−/−} GC T_{FH} cells is a primary T-cell defect and not secondary to a lack of GC B cells or the organized GC structure. Given the complexity of reciprocal regulation between T and B cells [19], such tests in mixed chimera are crucial for precisely assigning gene functions involved in T_{FH} and GC development, a point illustrated again by the recent discovery that distinct biochemical mechanisms are responsible for Bcl-6-controlled T_{FH} development and GC B cell differentiation [20[•]]. Based on known functions, major T_{FH}-associated molecules can be categorized into three functional modules (Figure 2), with the positioning and interaction modules together specifying where and how helper signals are delivered to B cells. In addition to Bcl-6, c-Maf [21[•]], IRF-4 [22] and BATF [23] may also contribute to T_{FH} programming. However, these other transcriptional factors have not been tested in systems where GCs can form with ‘help’ provided by wildtype T_{FH} cells.

Bcl-6 induction and pre-GC T_{FH} cells at the T–B border

Bcl-6 is upregulated in T cells within 30 minutes upon activation *in vitro* [24]. T cells substantially increase Bcl-6 protein expression from 2 to 4 days following activation *in vivo* [7[•],8[•],13[•],25[•]], and essentially all divided T cells at day 2 highly express Bcl-6 [25[•]]. Cells with a CXCR5⁺PD-1⁺ phenotype also rapidly appear, exhibiting kinetics identical to or slightly slower than that of Bcl-6 [7[•],8[•],25[•]]. Early-arising Bcl-6⁺CXCR5⁺ cells are located in the T–B border and interfollicular regions, poised to migrate into the follicle [7[•],8[•]]. The appearance of these pre-GC T_{FH} cells requires DC- but not B cell-mediated antigen presentation [8[•],13[•],26[•]], and Bcl-6 upregulation is promoted by IL-6 [27–29] and IL-12 [30–32]. Because ICOS signaling is required for T_{FH} development and GC formation [33]

Figure 1




Spatiotemporal development of the T_{FH} subset and its relationship to other lineages and memory. Upon activation, naïve T cells rapidly upregulate Bcl-6 and CXCR5 and become pre-GC T_{FH} cells at the T-B border. Varied biases toward Th1/Th2/Th17 potentials likely exist among individual cells at this stage, and such T cells can productively interact with cognate B cells here (not depicted). CXCR5 and ICOS upregulation combined with CCR7 and PSGL-1 downregulation facilitates follicular re-localization, for which the ICOSL molecule expressed by bystander B cells is crucial (red shade in the follicle). Some pre-GC T_{FH} cells at the T-B border may not enter the follicle but encounter milieu that favors Blimp-1 induction and drives the terminal effector differentiation. The entry of T_{FH} cells into the GC requires their SAP expression and productive interactions with cognate B cells. Besides location, functional polarization in other aspects of GC T_{FH} cells is not well characterized. T cells may exit the secondary lymphoid organ from both T-B border and the follicle, eventually joining the general circulation to become blood T_{FH} cells. GC T_{FH} cells can exit from GCs and join other GCs. They might also enter general circulation and constitute a numerical minority of the blood T_{FH} compartment. Blood T_{FH} cells are more quiescent and memory-competent, and their relationship with T_{CM} cells that survey lymphoid organs requires further investigation. A provocative possibility is that T_{CM} cells are essentially derived from cells along the T_{FH} developmental pathway. Upon re-challenge, such memory cells can give rise to not only non-polarized or Th1-/Th2-/Th17-biased T_{FH} cells but also terminally differentiated and Th1/Th2/Th17-polarized effectors. Alternatively, blood T_{FH} cells may be memory cells committed to the T_{FH} pathway and cannot generate terminal Th1/Th2/Th17 effectors in a recall response. Dotted lines and question marks indicate uncertainties. Time axis at the top indicates the approximate time frame in which T_{FH} cells of a given stage appear before the next stage in the lymphoid organ, thus giving the overlap. The time axis does not imply a schedule for T_{FH} cells to exit from a give location into the blood. The progressive localization from the T-B border into the GC is likely a feature shared by follicular NKT cells and follicular T regulatory cells (not depicted).

and ICOS-deficient T cells fail to generate Bcl-6⁺CXCR5⁺ cells [13^{••}], ICOS co-stimulation is thought to specifically induce Bcl-6 [1], although evidence for a direct signaling connection is lacking.

How Bcl-6 promotes pre-GC T_{FH} development is not clear. A very high level of Bcl-6 overexpression in human T cells led to upregulation of CXCR5, PD-1, SAP, ICOS,

and CD40L in five days [21[•]], implying that Bcl-6 programs all functional modules characteristic of T_{FH} cells. However, this kinetics is slower than the appearance of pre-GC T_{FH} cells *in vivo*. Bcl-6 was shown to promote CXCR5 transcription [17[•]], possibly through suppression of microRNA species including the mir-17~92 family that might target CXCR5 [18[•]]. Following *Listeria* infection, antigen-specific Bcl-6^{-/-} T cells failed

Figure 2



Modular characteristics of T_{FH} cells. The 'positioning' module facilitates T cell migration into the follicle and promotes T cell co-localization with cognate B cells. ICOS plays a key role in this module, and the ligand it requires in regulating follicular recruitment of T cells is on bystander B cells (see Figure 1). The 'interaction' module controls the strength of physical interactions with cognate B cells, which likely impinge on the efficiency with which not only surface-bound but also soluble 'helper-signal' molecules are delivered to cognate B cells across the synapse. Bcl-6 is required for activated T cells to develop into T_{FH} cells and for cognate B cells to differentiate into the GC state. Precise functions of ICOS and PD-1 in the context of antigen-specific interactions between T_{FH} and GC B cells are not yet clarified.

to express CXCR5 [34^{••}], but *Bcl-6*^{-/-} OT-II T cells normally expressed CXCR5 3 days after OVA immunization in CFA [35^{••}], suggesting requirement of Bcl-6 for CXCR5 expression is context-dependent. New evidence indicates that mir-17~92 does not inhibit but actually promote T_{FH} development and GC formation [36^{••}, 37^{••}], making this microRNA cluster an unlikely link between Bcl-6 and CXCR5. Very recently, Achaete-Scute homologue 2 (Ascl-2), a new basic helix-loop-helix transcription factor, was found to bind directly to the *Cxcr5* locus, trigger CXCR5 upregulation, and initiate the T_{FH} development [38^{••}].

Transition from pre-GC T_{FH} cells to GC T_{FH} cells

All recently primed T cells move to the T-B border but do not necessarily move into the follicle (Figure 1). When activated T cells were blocked from exiting the lymphoid organ, it became evident that the GC T_{FH} cell compartment constitutes a numerical minority of all activated T cells of the same specificity [39]. It is not yet clear how the GC T_{FH} cell compartment is selectively established from the pool of all activated T cells that have ever arrived at the T-B border [19, 40], although the strength of TCR signals appears to be an important factor [39, 41].

Does Bcl-6 commit T cells to the unique follicular location and/or functions?

Although Bcl-6 does not seem to regulate CXCR5 directly, it could control other factors that promote follicular localization, a possibility that remains to be vigorously tested. It is also possible that, between individual pre-GC T_{FH} cells at the T-B border and their clonal progenies eventually in the GC T_{FH} compartment at a later time, Bcl-6 instructionally polarizes biological features of T_{FH} cells [1] so as to fulfill a particular set of helper functions inside the follicle or GCs. Somewhat contrary to this notion, Bcl-6 protein levels on individual CXCR5⁺PD-1⁺ T cells were found to peak around day 3 after immunization at the T-B border and decrease thereafter, leading to some GC T_{FH} cells with undetectable Bcl-6 expression [7^{••}]. Studies with an IRES-driven Bcl-6 reporter line revealed the same peak time but not the subsequent declining phase [35^{••}]. This difference may result from the complex transcriptional and post-translational regulation of Bcl-6 protein expression, extensively characterized in B but not yet in T cells [42].

Importantly, we cannot yet define T_{FH} function properties unique to the follicular or GC phase, as the entire helper signal module does not distinguish GC from pre-GC T_{FH}

cells (Figure 2). PD-1 is involved in regulating plasma cell differentiation [43,44], but cellular mechanisms remain to be clarified. IL-21 is arguably the most specific TFH-associated helper signal molecule [45,46], although it is not controlled by Bcl-6 but c-Maf [21*,47]. Experiments with an IL-21 reporter mouse strain showed no sign of IL-21 polarization, as a comparable fraction of CXCR5⁺PD-1⁺ TFH cells expressed this cytokine from the pre-GC to the GC stage [48**]. In fact, the magnitude of GC formation remains the most frequently used readout to evaluate TFH functions in general, even though it depends more on 'help' delivered before GC formation. Until we come to grasp how TFH cells function to help B cells inside the follicle and GCs, possibly in a manner that is distinct from that at the T-B border, we may not be able to define TFH functional polarization in substance, that is, other than location.

ICOS-controlled follicular localization of TFH cells and bystander B cells

CXCR5 expression is necessary but not sufficient for activated T cells to localize into follicle, even when CCR7 is downregulated [49]. Another key factor is ICOS. Our recent work demonstrates that ICOS exerts direct motility control on T cells [50**]. This effect does not involve classical co-stimulation but hinges on ICOS engagement by ICOSL on the follicular bystander B cells. The resultant p110δ-mediated signaling promotes pseudopod formation and persistent T cell migration, which is necessary for CXCR5⁺ T cells to move efficiently from the T-B border into the follicle [50**]. Interestingly, the TFH- and GC-promoting effect of mir-17~92 is in part due to its suppression of Phlpp2, a phosphatase that antagonizes ICOS-triggered PI3K signaling and inhibits follicular localization of T cells [36**], further highlighting the importance of direct ICOS-mediated motility control. Given its crucial role, ICOS transcription is under redundant control by RNA-binding protein Roquin-1 and Roquin-2 [51,52]. Because enforced Bcl-6 overexpression cannot correct the deficient TFH recruitment and GC formation in the absence of ICOS [50**], Bcl-6 alone is insufficient to instruct a pre-GC to GC TFH transition, at least in terms of follicular localization (Figure 1). Given the role of bystander B cells in regulating TFH development, studies involving gene ablation in the entire B or T cell compartment should not always be interpreted solely in the context of antigen presentation and cognate interactions.

Bcl-6 and other Th fates

Bcl-6 may antagonize Th1, Th2, and Th17 development by suppressing T-bet, GATA-3, and ROR-γ expression and activities [17*,18*,53], implying a zero-sum competition between the TFH fate and the other lineages. However, TFH cells sorted from immunized animals still exhibit positive histone marks at T-bet, GATA-3 and ROR-γ loci [54**]. Th1 and Th2 cytokines are needed for regulating isotype-switching inside GCs and are indeed expressed by TFH cells [48**,55]. IFN-γ may even

promote TFH development under certain conditions [56]. In Peyer's Patches, IL-17-producing cells are TFH-like and induce IgA-producing GC B cells [57]. T cell-specific Bcl-6 ablation does not cause exaggerated Th1, Th2, or Th17 development [58,59]. In the strictest sense, therefore, the TFH fate is not a parallel alternative to the other effector lineages (Figure 1).

Bcl-6, Blimp-1 and memory competence of TFH cells

From either the T-B border or the follicle, recently activated T cells can exit the lymphoid tissue to eventually join the circulation (Figure 1). Cells with a TFH-like phenotype are found in the blood, and their numbers and helper qualities correlate with ongoing immune responses [60*,61*,62*,63*]. These cells are of a more quiescent phenotype (e.g. low in PD-1 expression) compared to TFH cells in the lymphoid organ, suggestive of memory formation [62*,63*].

In B cells, Bcl-6 is required to maintain the highly proliferative GC state, which arguably represents a differentiation stage in transition to humoral memory [5*]. In T cells, Bcl-6 is essential for the memory competence, in part because it helps maintain T cell proliferative capacity and suppresses apoptosis [59,64,65]. Interestingly, central memory T (TCM) cells identified during a *Listeria* infection express CXCR5, and their development requires Bcl-6, intact B cell compartment, and ICOS signaling triggered by B cell-derived ICOSL [34**]. These are the same set of requirements for generating GC TFH cells. CXCR5⁺ TCM-like memory CD4 T cells have been identified in other systems [66*,67,68]. An intriguing possibility is that TFH cells at the T-B border and inside the follicle are a shared precursor pool for GC TFH cells and TCM cells (Figure 1). Consistent with this scenario, IL-2-driven upregulation of Blimp-1, which antagonizes Bcl-6 [69], promotes terminal effector T cell differentiation, suppresses TCM development, and inhibits TFH formation [16*,70,71*,72–74]. Chemokine receptors other than CXCR5 may guide recently activated T cells to milieu particularly suited for terminal effector differentiation, as in the case of CXCR3 for Th1 differentiation [75].

TFH-polarized memory cells or TCM cells capable of TFH development

Given the relatedness of TFH and TCM cells, it is challenging to vigorously establish whether an immune response produces memory T cells with TFH functional polarization, at least until the latter concept can be better defined as discussed above. Compared to terminally differentiated effector cells, which are enriched in the non-TFH compartment and lack memory or TFH potential owing to high levels of Blimp-1, TCM cells undergo secondary expansion of much higher magnitudes. Thus, even if secondary TFH development and GC formation is only coincidental and proportional to the recall response, a TFH 'bias' can be

detected in the T_{CM} -as compared to the effector-mediated recall. Indeed, when T_{FH} and non- T_{FH} cells isolated late during influenza infection were recalled in naïve hosts after a period of 'rest' [48**], a huge difference in proliferative capacity was observed, reflecting inherent differences in the memory competence between the seeding populations. Such difference in the proliferative capacity between T_{FH} and non- T_{FH} cells after transfer was reported in two studies of LCMV infection [66*,71*] and in one [35**] but not another [76] model of protein immunization. Optical-highlighting experiments demonstrate that GC T_{FH} cells could exit GCs [77*], and some GC T_{FH} cells downmodulate Bcl-6 [7**] and increase IL-7R α expression [7**,71*], making it possible that even GC T_{FH} cells are not terminal but maintain memory competence. Future studies have to determine whether the same or different biochemical mechanisms underlie the Bcl-6 requirement for T_{FH} and T_{CM} formation and to test how B cells, bystander or cognate, may help maintain joint or separate GC T_{FH} and T_{CM} competence.

Conclusions

Moving forward, we must clarify two key concepts for T_{FH} cells, functional polarization and memory potential. To help B cells to form GCs or differentiate into memory or plasma cells, do T_{FH} cells work in different manners or by different mechanisms inside follicle or GCs as compared to any Th cells at the T-B border? Regarding memory, it seems ' T_{FH} -ness' could be important for T_{CM} formation. Alternatively, is the T_{CM} competence somehow important for B helper functions of T_{FH} cells? Or, are some aspects (e.g. cell cycling, quiescence, metabolism) of T cell biology potentially controlled by Bcl-6 important for both? It is provocatively possible that a single cell and its progenies may traverse the pre-GC and GC T_{FH} compartments before becoming T_{CM} cells in a primary response, and can still give rise to Th1/Th2/Th17/ T_{FH} cells in a recall response (Figure 1). The crux of these issues remains to understand what Bcl-6 does. Solving the puzzle should also help to identify the most inclusive aspect of T_{FH} biology that may unify follicular NKT cells and regulatory T cells with the conventional helper [78]. Is it reliance on B cells, follicular passage, functional polarization, memory competence or still just that, Bcl-6?

Acknowledgements

This work is supported in part by the National Natural Science Foundation of China (grant nos. 81330070 and 81361120397), the Ministry of Science and Technology '973' program (grant no. 2014CB542501), and the Tsinghua-Peking Center for Life Sciences.

References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Crotty S: **Follicular helper CD4 T cells (T_{FH})**. *Annu Rev Immunol* 2011, **29**:621-663.
2. Vinuesa CG, Cyster JG: **How T cells earn the follicular rite of passage**. *Immunity* 2011, **35**:671-680.
3. Corti D, Lanzavecchia A: **Broadly neutralizing antiviral antibodies**. *Annu Rev Immunol* 2013, **31**:705-742.
4. Tangye SG, Ma CS, Brink R, Deenick EK: **The good, the bad and the ugly – T_{FH} cells in human health and disease**. *Nat Rev Immunol* 2013, **13**:412-426.
5. Tarlinton D, Good-Jacobson K: **Diversity among memory B cells: origin, consequences, and utility**. *Science* 2013, **341**:1205-1211.
- A thorough and authoritative review on humoral memory and a must-read for contemplating how T_{FH} cells may function beyond promoting GC formation.
6. Streeck H, D'Souza MP, Littman DR, Crotty S: **Harnessing CD4⁺ T cell responses in HIV vaccine development**. *Nat Med* 2013, **19**:143-149.
7. Kitano M, Moriyama S, Ando Y, Hikida M, Mori Y, Kurosaki T, Okada T: **Bcl6 protein expression shapes pre-germinal center B cell dynamics and follicular helper T cell heterogeneity**. *Immunity* 2011, **34**:961-972.
- By static and dynamic imaging aided with a Bcl-6-YFP fusion reporter, this study thoroughly characterized Bcl-6 expression kinetics in T cells following activation *in vivo*, traced spatiotemporal T_{FH} development from the T-B border to the GC stage, and identified Bcl-6⁺ GC T_{FH} cells. This study also showed IL-7R α upregulation on some of these T_{FH} cells, hinting at their potential memory competence. Data in this study and Refs. [8**,25*] show the frequency of Bcl-6⁺CXCR5⁺ cells peaks very early around day 3 after immunization.
8. Kerfoot SM, Yaari G, Patel JR, Johnson KL, Gonzalez DG, Kleinstein SH, Haberman AM: **Germinal center B cell and T follicular helper cell development initiates in the interfollicular zone**. *Immunity* 2011, **34**:947-960.
- This study demonstrates Bcl-6 expression and other T_{FH} -associated markers are upregulated early during an immune response when they are still at the T-B border or the interfollicular region. Curiously, a majority of antigen-reactive T cells appear PD-1⁺Bcl-6⁺ between day 1 and day 2. Also see annotation for Refs. [7**,25*]. Together with Refs. [13**,26*], this study further demonstrates the early appearance of T_{FH} cells requires only DC priming but not antigen presentation by cognate B cells.
9. Qi H, Cannons JL, Klauschen F, Schwartzberg PL, Germain RN: **SAP-controlled T-B cell interactions underlie germinal centre formation**. *Nature* 2008, **455**:764-769.
10. Lee SK, Rigby RJ, Zotos D, Tsai LM, Kawamoto S, Marshall JL, Ramiscal RR, Chan TD, Gatto D, Brink R *et al.*: **B cell priming for extrafollicular antibody responses requires Bcl-6 expression by T cells**. *J Exp Med* 2011, **208**:1377-1388.
- This study demonstrates T cell help for the extrafollicular B cell response also depends on Bcl-6.
11. Haynes NM, Allen CD, Lesley R, Ansel KM, Killeen N, Cyster JG: **Role of CXCR5 and CCR7 in follicular Th cell positioning and appearance of a programmed cell death gene-1^{high} germinal center-associated subpopulation**. *J Immunol* 2007, **179**:5099-5108.
12. Salek-Ardakani S, Choi YS, Rafii-El-Idrissi Benhnia M, Flynn R, Arens R, Shoenberger S, Crotty S, Croft M: **B cell-specific expression of B7-2 is required for follicular Th cell function in response to vaccinia virus**. *J Immunol* 2011, **186**:5294-5303.
13. Choi YS, Kageyama R, Eto D, Escobar TC, Johnston RJ, Monticelli L, Lao C, Crotty S: **ICOS receptor instructs T follicular helper cell versus effector cell differentiation via induction of the transcriptional repressor Bcl6**. *Immunity* 2011, **34**:1-15.
- Together with Refs. [8**,26*], this study demonstrates that the early appearance of T_{FH} cells requires only DC priming but not antigen presentation by cognate B cells. Based on kinetic analyses of T_{FH} differentiation in the absence of ICOS signaling at different temporal stages of an immune response, this study was the first to propose a coherent cellular model ('ICOS \rightarrow Bcl-6 \rightarrow CXCR5') for how the ICOS molecule might regulate T_{FH} development and GC formation, even though more recent work challenges such a linear instructional model.
14. Deenick EK, Chan A, Ma CS, Gatto D, Schwartzberg PL, Brink R, Tangye SG: **Follicular helper T cell differentiation requires continuous antigen presentation that is independent of unique B cell signaling**. *Immunity* 2010, **33**:241-253.

15. Baumjohann D, Preite S, Rebaldi A, Ronchi F, Ansel KM, Lanzavecchia A, Sallusto F: **Persistent antigen and germinal center B cells sustain T follicular helper cell responses and phenotype.** *Immunity* 2013, **38**:596-605.

By carefully correlating the magnitude of GC and T_{FH} responses and by testing effects of acute GC B cell deletion on T_{FH} cells, this study confirms and extends previous findings that GC B cells are required for maintaining GC T_{FH} cells and that the pool size of GC T_{FH} cells is dictated by the number of GC B cells. The authors therefore argue that T_{FH} cells represent a transient state instead of a stable lineage.

16. Johnston RJ, Poholek AC, DiToro D, Yusuf I, Eto D, Barnett B, Dent AL, Craft J, Crotty S: **Bcl6 and Blimp-1 are reciprocal and antagonistic regulators of T follicular helper cell differentiation.** *Science* 2009, **325**:1006-1010.

Together with Refs. [17*,18*], this study provides the first evidence that Bcl-6 is required for T_{FH} development. It also demonstrates that Blimp-1 suppresses T_{FH} development.

17. Nurieva RI, Chung Y, Martinez GJ, Yang XO, Tanaka S, Matskevitch TD, Wang YH, Dong C: **Bcl6 mediates the development of T follicular helper cells.** *Science* 2009, **325**:1001-1005.

Together with Refs. [16*,18*], this study provides the first evidence that Bcl-6 is required for T_{FH} development. In addition, this study suggests that Bcl-6 may direct T_{FH} development by suppressing Th1/Th2/Th17 lineages, a point of ongoing debate.

18. Yu D, Rao S, Tsai LM, Lee SK, He Y, Sutcliffe EL, Srivastava M, Linterman M, Zheng L, Simpson N et al.: **The transcriptional repressor Bcl-6 directs T follicular helper cell lineage commitment.** *Immunity* 2009, **31**:457-468.

Together with Refs. [16*,17*], this study provides the first evidence that Bcl-6 is required for T_{FH} development. In addition, this study suggests that Bcl-6 may direct T_{FH} development by suppressing alternative Th1/Th2/Th17 lineages and promote the T_{FH} signature including CXCR5 expression by suppressing microRNA species. Both points are of ongoing debate.

19. Qi H: **From SAP-less T cells to helpless B cells and back: dynamic T-B cell interactions underlie germinal center development and function.** *Immunol Rev* 2012, **247**:24-35.
20. Huang C, Hatzl K, Melnick A: **Lineage-specific functions of Bcl-6 in immunity and inflammation are mediated by distinct biochemical mechanisms.** *Nat Immunol* 2013, **14**:380-388.

This study demonstrates that Bcl-6 promotes T_{FH} development and GC B cell differentiation by distinct mechanisms that differentially depend on Bcl-6 BTB domain interactions with other co-repressors.

21. Kroenke MA, Eto D, Locci M, Cho M, Davidson T, Haddad EK, Crotty S: **Bcl6 and Maf cooperate to instruct human follicular helper CD4 T cell differentiation.** *J Immunol* 2012, **188**:3734-3744.

This study provides *in vitro* evidence that Bcl-6 and Maf may directly instruct a T_{FH} phenotype in human T cells. Of particular interest, a very high level of Bcl-6 overexpression leads to upregulation of many T_{FH}-associated markers 5 days after retroviral transduction.

22. Bollig N, Brustle A, Kellner K, Ackermann W, Abass E, Raifer H, Camara B, Brendel C, Giel G, Bothur E et al.: **Transcription factor IRF4 determines germinal center formation through follicular T-helper cell differentiation.** *Proc Natl Acad Sci U S A* 2012, **109**:8664-8669.
23. Ise W, Kohyama M, Schraml BU, Zhang T, Schwer B, Basu U, Alt FW, Tang J, Oltz EM, Murphy TL et al.: **The transcription factor BATF controls the global regulators of class-switch recombination in both B cells and T cells.** *Nat Immunol* 2011, **12**:536-543.
24. Fukuda T, Miki T, Yoshida T, Hatano M, Ohashi K, Hirokawa S, Tokuhisa T: **The murine BCL6 gene is induced in activated lymphocytes as an immediate early gene.** *Oncogene* 1995, **11**:1657-1663.
25. Baumjohann D, Okada T, Ansel KM: **Cutting Edge: distinct waves of BCL6 expression during T follicular helper cell development.** *J Immunol* 2011, **187**:2089-2092.

By tracing cell divisions and Bcl-6 expression, this study provides definitive evidence that Bcl-6 upregulation is as early as the first division and as extensive as on all those divided 48 h after immunization. Combined with data from Refs. [7*,8*,13**], it suggests some progenies of these cells downregulate Bcl-6 by day 4.

26. Goenka R, Barnett LG, Silver JS, O'Neill PJ, Hunter CA, Cancro MP, Lauffer TM: **Cutting edge: dendritic cell-restricted antigen presentation initiates the follicular helper T cell program but cannot complete ultimate effector differentiation.** *J Immunol* 2011, **187**:1091-1095.

Together with Refs. [8**,13**], this study demonstrates that the early appearance of T_{FH} cells requires only DC priming but not antigen presentation by cognate B cells.

27. Choi YS, Eto D, Yang JA, Lao C, Crotty S: **Cutting edge: STAT1 is required for IL-6-mediated Bcl6 induction for early follicular helper cell differentiation.** *J Immunol* 2013, **190**:3049-3053.
28. Harker JA, Lewis GM, Mack L, Zuniga EI: **Late interleukin-6 escalates T follicular helper cell responses and controls a chronic viral infection.** *Science* 2011, **334**:825-829.
29. Petrovas C, Yamamoto T, Gerner MY, Boswell KL, Wloka K, Smith EC, Ambrozak DR, Sandler NG, Timmer KJ, Sun X et al.: **CD4 T follicular helper cell dynamics during SIV infection.** *J Clin Invest* 2012, **122**:3281-3294.
30. Nakayama S, Kanno Y, Takahashi H, Jankovic D, Lu KT, Johnson TA, Sun HW, Vahedi G, Hakim O, Hannon R et al.: **Early Th1 cell differentiation is marked by a Tfh cell-like transition.** *Immunity* 2011, **35**:919-931.
31. Ma CS, Suryani S, Avery DT, Chan A, Nanan R, Santner-Nanan B, Deenick EK, Tangye SG: **Early commitment of naive human CD4+ T cells to the T follicular helper (T_{FH}) cell lineage is induced by IL-12.** *Immunol Cell Biol* 2009, **87**:590-600.
32. Schmitt N, Bustamante J, Bourdery L, Bentebibel SE, Boisson-Dupuis S, Hamlin F, Tran MV, Blankenship D, Pascual V, Savino DA et al.: **IL-12 receptor β 1 deficiency alters *in vivo* T follicular helper cell response in humans.** *Blood* 2013, **121**:3375-3385.
33. Gigoux M, Shang J, Pak Y, Xu M, Choe J, Mak TW, Suh WK: **Inducible costimulator promotes helper T-cell differentiation through phosphoinositide 3-kinase.** *Proc Natl Acad Sci U S A* 2009, **106**:20371-20376.
34. Pepper M, Pagan AJ, Igyarto BZ, Taylor JJ, Jenkins MK: **Opposing signals from the Bcl6 transcription factor and the interleukin-2 receptor generate T helper 1 central and effector memory cells.** *Immunity* 2011, **35**:583-595.

By tracking memory formation from endogenous CD4+ T cells specific to *Listeria* antigen, this study shows an early bifurcation of Th1-biased effector memory cells and apparently uncommitted central memory cells that exhibit many T_{FH} characteristics, including CXCR5 expression and dependence on Bcl-6, ICOS, and an intact B cell compartment. This study raises the crucial question as to how T_{FH} cells are related to T_{CM} cells in general.

35. Liu X, Yan X, Zhong B, Nurieva RI, Wang A, Wang X, Martin-Orozco N, Wang Y, Chang SH, Esplugues E et al.: **Bcl6 expression specifies the T follicular helper cell program *in vivo*.** *J Exp Med* 2012, **209**:1841-1852.

A 'knock-in' IRES-driven RFP reporter for Bcl-6 was used in this study to characterize Bcl-6 expression and T_{FH} development *in vivo*. This study also provides genetic evidence that CXCR5 upregulation *per se* does not require Bcl-6 following protein antigen immunization in CFA. Memory formation by T_{FH} cells is also reported.

36. Kang SG, Liu WH, Lu P, Jin HY, Lim HW, Shepherd J, Fremgen D, Verdin E, Oldstone MB, Qi H et al.: **MicroRNAs of the miR-17~92 family are critical regulators of T_{FH} differentiation.** *Nat Immunol* 2013, **14**:849-857.

A comprehensive study of the entire miR-17~92 microRNA family during T_{FH} development. It demonstrates miR-17~92 is required for T_{FH} and GC responses because it protects PI3K activities downstream of ICOS from suppression by PTEN and Phlpp2, a well-known and a new target of miR-17~92, respectively.

37. Baumjohann D, Kageyama R, Clingan JM, Morar MM, Patel S, de Kouchkovsky D, Bannard O, Bluestone JA, Matloubian M, Ansel KM et al.: **The microRNA cluster miR-17~92 promotes T_{FH} cell differentiation and represses subset-inappropriate gene expression.** *Nat Immunol* 2013, **14**:840-848.

Another study of the role of miR-17~92 in T_{FH} development and GC formation, reaching similar functional conclusions as Ref. [36**]. Mechanistically, this study suggests that increased expression of PTEN and additional 'T_{FH}-inappropriate' genes such *Rora* is what impairs T_{FH} development in the absence of miR-17~92.

38. Liu X, Chen X, Zhong B, Wang YA, Wang X, Chu F, Nurieva R, ●● Yan X, Chen P, van der Flier L *et al.*: **Transcription factor Achaete-Scute homologue 2 initiates T follicular helper cell development.** *Nature* 2014, <http://dx.doi.org/10.1038/nature12910>.
This study identifies a new transcription factor that controls CXCR5 and initiates TFH development.
39. Fazilleau N, McHeyzer-Williams LJ, Rosen H, McHeyzer-Williams MG: **The function of follicular helper T cells is regulated by the strength of T cell antigen receptor binding.** *Nat Immunol* 2009, **10**:375-384.
40. Qi H, Chen X, Chu C, Lu P, Xu H, Yan J: **Follicular T-helper cells: controlled localization and cellular interactions.** *Immunol Cell Biol* 2013, **92**:28-33.
41. Tubo NJ, Pagan AJ, Taylor JJ, Nelson RW, Linehan JL, Ertelt JM, Huseby ES, Way SS, Jenkins MK: **Single naive CD4⁺ T cells from a diverse repertoire produce different effector cell types during infection.** *Cell* 2013, **153**:785-796.
42. Basso K, Dalla-Favera R: **Roles of BCL6 in normal and transformed germinal center B cells.** *Immunol Rev* 2012, **247**:172-183.
43. Good-Jacobson KL, Szumilas CG, Chen L, Sharpe AH, Tomayko MM, Shlomchik MJ: **PD-1 regulates germinal center B cell survival and the formation and affinity of long-lived plasma cells.** *Nat Immunol* 2010, **11**:535-542.
44. Kawamoto S, Tran TH, Maruya M, Suzuki K, Doi Y, Tsutsui Y, Kato LM, Fagarasan S: **The inhibitory receptor PD-1 regulates IgA selection and bacterial composition in the gut.** *Science* 2012, **336**:485-489.
45. Zotos D, Coquet JM, Zhang Y, Light A, D'Costa K, Kallies A, Corcoran LM, Godfrey DI, Toellner KM, Smyth MJ *et al.*: **IL-21 regulates germinal center B cell differentiation and proliferation through a B cell-intrinsic mechanism.** *J Exp Med* 2010, **207**:365-378.
46. Linterman MA, Beaton L, Yu D, Ramiscal RR, Srivastava M, Hogan JJ, Verma NK, Smyth MJ, Rigby RJ, Vinuesa CG: **IL-21 acts directly on B cells to regulate Bcl-6 expression and germinal center responses.** *J Exp Med* 2010, **207**:353-363.
47. Hiramatsu Y, Suto A, Kashiwakuma D, Kanari H, Kagami S, Ikeda K, Hirose K, Watanabe N, Grusby MJ, Iwamoto I *et al.*: **c-Maf activates the promoter and enhancer of the IL-21 gene, and TGF-beta inhibits c-Maf-induced IL-21 production in CD4⁺ T cells.** *J Leukoc Biol* 2010, **87**:703-712.
48. Luthje K, Kallies A, Shimohakamada Y, Belz GT, Light A, ●● Tarlinton DM, Nutt SL: **The development and fate of follicular helper T cells defined by an IL-21 reporter mouse.** *Nat Immunol* 2012, **13**:491-498.
This study reports the first IL-21 reporter and comprehensive analyses of TFH development and memory formation in an influenza infection model. It reveals that a relatively stable percentage of activated T cells produce IL-21, from the pre-GC to the GC stage. It provides the clearest evidence so far that TFH cells are cycling and non-terminal and that the biggest difference between a recall response by TFH cells and non-TFH effector cells after 'rest' is the proliferative capacity.
49. Ansel KM, McHeyzer-Williams LJ, Ngo VN, McHeyzer-Williams MG, Cyster JG: **In vivo-activated CD4 T cells upregulate CXC chemokine receptor 5 and reprogram their response to lymphoid chemokines.** *J Exp Med* 1999, **190**:1123-1130.
50. Xu H, Li X, Liu D, Li J, Zhang X, Chen X, Hou S, Peng L, Xu C, Liu W ●● *et al.*: **Follicular T-helper cell recruitment governed by bystander B cells and ICOS-driven motility.** *Nature* 2013, **496**:523-527.
This study uncovers a new mechanism for how ICOS regulate TFH development via motility control. It shows ICOS engaged by ICOSL expressed on follicular bystander B cells optimize T cell pseudopod formation and power T cell migration from the T-B border into the follicle. This new layer of positioning control does not replace follicle-bound directionality provided by a combination of chemokine receptors including CXCR5. This study also shows Bcl-6 overexpression cannot correct TFH deficiency due to a lack of ICOS.

51. Vogel KU, Edelmann SL, Jeltsch KM, Bertossi A, Heger K, Heinz GA, Zoller J, Warth SC, Hoefig KP, Lohs C *et al.*: **Roquin paralogs 1 and 2 redundantly repress the Icos and Ox40 costimulator mRNAs and control follicular helper T cell differentiation.** *Immunity* 2013, **38**:655-668.
52. Pratama A, Ramiscal RR, Silva DG, Das SK, Athanasopoulos V, Fitch J, Botelho NK, Chang PP, Hu X, Hogan JJ *et al.*: **Roquin-2 shares functions with its paralog Roquin-1 in the repression of mRNAs controlling T follicular helper cells and systemic inflammation.** *Immunity* 2013, **38**:669-680.
53. Mondal A, Sawant D, Dent AL: **Transcriptional repressor BCL6 controls Th17 responses by controlling gene expression in both T cells and macrophages.** *J Immunol* 2010, **184**:4123-4130.
54. Lu KT, Kanno Y, Cannons JL, Handon R, Bible P, Elkahoul AG, ●● Anderson SM, Wei L, Sun H, O'Shea JJ *et al.*: **Functional and epigenetic studies reveal multistep differentiation and plasticity of in vitro-generated and in vivo-derived follicular T helper cells.** *Immunity* 2011, **35**:622-632.
A comprehensive study of Th plasticity shows that CXCR5⁺PD-1⁺ TFH cells derived *in vitro* and TFH cells at GC time points *in vivo* possess positive histone marks on *T-bet*, *Gata-3*, and *Rorc* loci. It also shows that Th1/Th2/Th17 cells *in vitro* remain poised to express Bcl-6 and Blimp-1. These data are consistent with a high degree of plasticity of TFH cells whose lineage is defined by Bcl-6; they are also consistent with the possibility that Th1/Th2/Th17-biased cells are competent to become memory or terminally differentiated effectors, as respectively conditioned by Bcl-6 and Blimp-1.
55. Vijayanand P, Seumois G, Simpson LJ, Abdul-Wajid S, Baumjohann D, Panduro M, Huang X, Interlandi J, Djuretic VM, Brown DR *et al.*: **Interleukin-4 production by follicular helper T cells requires the conserved IL4 enhancer hypersensitivity site V.** *Immunity* 2012, **36**:175-187.
56. Lee SK, Silva DG, Martin JL, Pratama A, Hu X, Chang PP, Walters G, Vinuesa CG: **Interferon-γ excess leads to pathogenic accumulation of follicular helper T cells and germinal centers.** *Immunity* 2012, **37**:880-892.
57. Hirota K, Turner JE, Villa M, Duarte JH, Demengeot J, Steinmetz OM, Stockinger B: **Plasticity of Th17 cells in Peyer's patches is responsible for the induction of T cell-dependent IgA responses.** *Nat Immunol* 2013, **14**:372-379.
58. Kaji T, Ishige A, Hikida M, Taka J, Hijikata A, Kubo M, Nagashima T, Takahashi Y, Kurosaki T, Okada M *et al.*: **Distinct cellular pathways select germline-encoded and somatically mutated antibodies into immunological memory.** *J Exp Med* 2012, **209**:2079-2080.
59. Hollister K, Kusam S, Wu H, Clegg N, Mondal A, Sawant DV, Dent AL: **Insights into the role of Bcl6 in follicular Th cells using a new conditional mutant mouse model.** *J Immunol* 2013, **191**:3705-3710.
60. Morita R, Schmitt N, Bentebibel SE, Ranganathan R, Bourdery L, ● Zurewski G, Foucat E, Dullaers M, Oh S, Sabzghabaei N *et al.*: **Human blood CXCR5⁺CD4⁺ T cells are counterparts of T follicular cells and contain specific subsets that differentially support antibody secretion.** *Immunity* 2011, **34**:108-121.
See annotation for [63*].
61. Bentebibel SE, Lopez S, Obermoser G, Schmitt N, Mueller C, ● Harrod C, Flano E, Mejias A, Albrecht RA, Blankenship D *et al.*: **Induction of ICOS⁺CXCR3⁺CXCR5⁺ T_H cells correlates with antibody responses to influenza vaccination.** *Sci Transl Med* 2013, **5**:176ra132.
See annotation for [63*].
62. Locci M, Havenar-Daughton C, Landais E, Wu J, Kroenke MA, ● Arlehamn CL, Su LF, Cubas R, Davis MM, Sette A *et al.*: **Human circulating PD-1⁺CXCR3⁺CXCR5⁺ memory T_H cells are highly functional and correlate with broadly neutralizing HIV antibody responses.** *Immunity* 2013, **39**:758-769.
See annotation for [63*].
63. He J, Tsai LM, Leong YA, Hu X, Ma CS, Chevalier N, Sun X, ● Vandenberg K, Rockman S, Ding Y *et al.*: **Circulating precursor CCR7^{lo}PD-1^{hi} CXCR5⁺ CD4⁺ T cells indicate T_H cell activity and promote antibody responses upon antigen reexposure.** *Immunity* 2013, **39**:770-781.

Together with Refs. [60*,61*,62*], these studies demonstrate blood CXCR5⁺ Th cells are likely derived from T_{FH} cells in the lymphoid organ at various stages during ongoing immune responses. These cells maintain various levels of B-cell helper activity and are competent to form memory.

64. Ichii H, Sakamoto A, Arima M, Hatano M, Kuroda Y, Tokuhisa T: **Bcl6 is essential for the generation of long-term memory CD4⁺ T cells.** *Int Immunol* 2007, **19**:427-433.
 65. Ichii H, Sakamoto A, Hatano M, Okada S, Toyama H, Taki S, Arima M, Kuroda Y, Tokuhisa T: **Role for Bcl-6 in the generation and maintenance of memory CD8⁺ T cells.** *Nat Immunol* 2002, **3**:558-563.
 66. Hale JS, Youngblood B, Latner DR, Mohammed AU, Ye L, Akondy RS, Wu T, Iyer SS, Ahmed R: **Distinct memory CD4⁺ T cells with commitment to T follicular helper- and T helper 1-cell lineages are generated after acute viral infection.** *Immunity* 2013, **38**:805-817.
- An extensive study of memory formation during LCMV infection in mice, which emphasizes distinct lineage-committed Th1 and T_{FH} memory cells. See also annotation to Ref. [71*].
67. MacLeod MK, David A, McKee AS, Crawford F, Kappler JW, Marrack P: **Memory CD4 T cells that express CXCR5 provide accelerated help to B cells.** *J Immunol* 2011, **186**:2889-2890.
 68. Chevalier N, Jarrossay D, Ho E, Avery DT, Ma CS, Yu D, Sallusto F, Tangye SG, Mackay CR: **CXCR5 expressing human central memory CD4 T cells and their relevance for humoral immune responses.** *J Immunol* 2011, **186**:5556-5560.
 69. Crotty S, Johnston RJ, Schoenberger SP: **Effectors and memories: Bcl-6 and Blimp-1 in T and B lymphocyte differentiation.** *Nat Immunol* 2010, **11**:114-120.
 70. Johnston RJ, Choi Y, Diamond JA, Yang JA, Crotty S: **STAT5 is a potent negative regulator of T_{FH} cell differentiation.** *J Exp Med* 2012, **209**:243-250.
 71. Choi YS, Yang JA, Yusuf I, Johnston RJ, Greenbaum J, Peters B, Crotty S: **Bcl6 expressing follicular helper CD4 T cells are fate committed early and have the capacity to form memory.** *J Immunol* 2013, **190**:4014-4026.
- Another extensive study of memory formation during LCMV infection in mice, which emphasizes distinct lineage-committed Th1 and T_{FH} memory cells. See also annotation to Ref. [66*].
72. Nurieva RI, Podd A, Chen Y, Alekseev AM, Yu M, Qi X, Huang H, Wen R, Wang J, Li HS et al.: **STAT5 negatively regulates T follicular helper (Tfh) cell generation and function.** *J Biol Chem* 2012, **287**:11234-11239.
 73. Oestreich KJ, Mohn SE, Weinmann AS: **Molecular mechanisms that control the expression and activity of Bcl-6 in T_H1 cells to regulate flexibility with a T_{FH}-like gene profile.** *Nat Immunol* 2012, **13**:405-411.
 74. Ballesteros-Tato A, Leon B, Graf BA, Moquin A, Adams PS, Lund FE, Randall TD: **Interleukin-2 inhibits germinal center formation by limiting T follicular helper cell differentiation.** *Immunity* 2012, **36**:847-856.
 75. Groom JR, Richmond J, Murooka TT, Sorensen EW, Sung JH, Bankert K, von Andrian UH, Moon JJ, Mempel TR, Luster AD: **CXCR3 chemokine receptor-ligand interactions in the lymph node optimize CD4⁺ T helper 1 cell differentiation.** *Immunity* 2012, **37**:1091-1100.
 76. Weber JP, Fuhrmann F, Hutloff A: **T-follicular helper cells survive as long-term memory cells.** *Eur J Immunol* 2012, **42**:1981-1988.
 77. Shulman Z, Gittlin AD, Targ S, Jankovic M, Pasqual G, Nussenzweig MC, Victora GD: **T follicular helper cell dynamics in germinal centers.** *Science* 2013, **341**:673-677.
- Using photo-activation and lineage tracing, this study provides definitive evidence that GC T_{FH} cells can exit original GCs and join other GCs. One implication is that GC T_{FH} cells might have the potential to enter the blood and forming memory.
78. Linterman MA, Liston A, Vinuesa CG: **T-follicular helper cell differentiation and the co-option of this pathway by non-helper cells.** *Immunol Rev* 2012, **247**:143-159.