



RICE

TOMO
DACHI

MA
CHI

Fabrication and Optimization of Lenses using Two-photon Polymerization Technology

Presenter: Hsu-Ting (Sophie) Kuo

Supervisor: Dr. Tomasz Tkaczyk

Mentor: Jiawei Lu


Overview

Miniature objective lens

- Ideal for **weight** and **size-sensitive** applications, such as endoscopic microscopy

Research objective: An optical system for two-photon fluorescence imaging

- Four **commercial lenses** + three **3D printed lenses**
- Input numerical aperture (NA) is 0.2
- Output NA is 0.56



Current method and challenges

Diamond turning technique

- Uses a diamond tool to cut the required surface geometry of the lens

Drawbacks

1. Time-consuming
2. Expensive
3. Difficult to fabricate free-form lenses
4. Manual alignment and assembly are required




Fig 1: Diamond turning (Photonics Media, 2018)

Fabrication technique

Two-photon polymerization technology

- Uses a **femtosecond pulse laser** in the near-infrared region
- Sequentially illuminate the **photosensitive resin**
 - Laser beam focused by a objective lens triggers the polymerization process
 - 150 nm feature sizes and $10\text{-}15\text{ nm}$ surface roughness

Advantages

1. Fast
2. Cheap
3. Fabricates lenses with automatic alignment features






Fig 2: Two photon polymerization (Bunea et al., 2021)

Zemax design

Right Half for testing


- The right half can be tested in air since it still has diffraction-limited performance
- ➔ Enables finalization of all printing parameters with either 25X or 10X objectives



Zemax design

Right Half for testing



- The right half can be tested in air since it still has diffraction-limited performance
- Enables finalization of all printing parameters with either 25X or 10X objectives



Generating 3D printing files

- The general MATLAB code for generating grayscale images

```
% initiate some parameters  
step_size = 1; % unit is micron  
  
x = -1000:1:1000;  
  
[X,Y] = meshgrid(x,x);  
  
% lens parameters  
R = 1496;  
r = 616;  
sag_n = sqrt(R^2-r^2);  
ET = 667;  
  
% sphere  
Z = sqrt(R^2 - (X.^2 + Y.^2));  
Z(Z<sag_n) = sag_n;  
Z(Z>sag_n) = Z(Z>sag_n) + ET;  
Z = Z - sag_n;  
  
% convert to 16 bit  
scale = 65535/(max(Z(:))-min(Z(:)));  
Z_new = (Z-min(Z(:)))*scale;  
  
% save images  
imwrite(uint16(Z_new), 'FirstLens.png');
```



The output grayscale image

Printing parameters

- **Slicing:** separates the object into many layers
- **Hatching:** differentiates the materials and areas with parallel lines





Fig 4. Slicing and hatching (Saraswat et al., 2020)

The first several trials gave us some estimation of the printing time:

1. 25X objective, 1-micron slicing, 0.2-micron hatching: **16-17 hours**
2. 10X objective, 1.6-micron slicing, 0.2-micron hatching: **12-13 hours**
3. 10X objective, 1.6-micron slicing, 0.6-micron hatching: **3-4 hours**

Printing parameters

- Printing setup for 25X objectives



Design Parameters

Source Path	FirstLens.png
Size X	2001 μm
Size Y	2001 μm
Size Z	132.71 μm
Center X	0 μm
Center Y	0 μm
2GL Calibration Profile	Select file...
Tiling X	— 1 +
Tiling Y	— 1 +
Interpolate	<input checked="" type="checkbox"/>
Display Color	[Color Swatch]
Preview Downsampling	— 1 +

Compensation Image


Enable	<input type="checkbox"/>
Structure Parameters	
Slice Distance	1 μm
Hatch Distance	0.2 μm
Hatch Direction	Y
Base Slice Count	0
Interface Offset	-1.5 μm
Exposure Parameters	
Scan Speed	200000 $\mu\text{m/s}$
Sparse Threshold	-1
Sparse Slice	<input type="checkbox"/>

Splitting Parameters

Enabled	<input checked="" type="checkbox"/>
Field Size X	420 μm
Field Size Y	420 μm
Field Overlap	60 μm
2GL Stitching	<input checked="" type="checkbox"/>
Field Offset Auto	<input checked="" type="checkbox"/>
Meander	<input type="checkbox"/>
Shear Angle	15 °

Characterization of the lens

Zygo white light interferometer – 10X



Problems encountered

1. Curvature deviation
2. Stitching defects

Proposed solutions

1. $R_{new} = \frac{R_{measure} - R_{design}}{2} + R_{design}$
2. Optimize laser power

Conclusions

What I learned over the past four weeks


- Observed how to design a lens in Zemax
- Participated in the fabrication process (grayscale image generation + 3D printing)
- Fundamental knowledge of optical components, such as lenses and fibers

Future work

- Solve previously encountered problems
- Finish the design and fabrication of the second lens
- Assemble the first and the second lenses

Acknowledgement:

Modern Optical Instrumentation and Bio-imaging Laboratory



This research was conducted as part of the 2022 MACHI Program at Rice University. Special thanks to members of the Tkacyzk team for their guidance and support and to Prof. Junichiro Kono, the Taipei Economic and Cultural Office in Houston (TECO-Houston), the sponsors, and everyone who makes this program possible.