

수학으로부터 인류를 자유롭게 하라
Free Humankind from Mathematics

Basic Algebra


Chap.2 Sets



Data Structures in Math


- (1) Sets
- (2) Sequences
- (3) Vectors
- (4) Matrices
- (3) Trees
- ⋮

Characteristics of Data Structures

- 
- (1) Sets: unordered objects
 - (2) Sequences: ordered objects with specific patterns
 - (3) Vectors: ordered numbers
 - (4) Matrices: numbers arranged in rectangular grids
 - (3) Trees: nodes connected by edges

Data Structures in Math

Operations on Data Structures



2.1 Definition and Notations of Sets

What's Sets?

a collection of **distinct** and **well-defined** things(or elements)

distinct: 서로 같지 않은

well-defined: doesn't change from person to person

things: 서로 같은 종류의 object들

don't have to be numbers

e.g. natural numbers, letters, rectangulars, images, persons

2.1 Definition and Notations of Sets

Notations**Enumerating Elements(Roster Form)**

Set = {element₁, element₂, ..., element_n}

$$A = \{1, 2, 3, 4\}$$

$$B = \left\{ \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \right\}$$

$$C = \{red, green, blue\}$$

$$D = \{free, guarantee, help, prices, winner, chance\}$$

Set Builder

Set = {element | element's condition}

$$A = \{x \mid 1 \leq x \in \mathbb{N} \leq 4\}$$


$$B = \{x \mid x \text{는 } \mathbb{R}^{2 \times 2} \text{의 standard basis vector}\}$$

$$C = \{x \mid x \text{는 빛의 삼원색}\}$$

$$D = \{x \mid x \text{는 스팸메일에 자주 등장하는 단어}\}$$

Notations

Venn Diagram



2.1 Definition and Notations of Sets

Universal/Empty Sets**Empty Sets**

$\emptyset = \{\}$ → 아무런 원도 갖고 있지 않은 거

Universal Sets

U : 가능한 모든 원소들의 집합

ex.1) $U = \{x \mid (x \text{ is a } 200 \times 200 \text{ images})\}$

$A = \{x \mid (x \text{ is a } 200 \times 200 \text{ images}) \wedge (x \text{ contains humans in it})\}$

$B = \{x \mid (x \text{ is a } 200 \times 200 \text{ images}) \wedge (x \text{ contains dogs in it})\}$

$C = \{x \mid (x \text{ is a } 200 \times 200 \text{ images}) \wedge (x \text{ contains humans and dogs in it})\}$

ex.2) $U = \{x \mid (x \text{ is an English word})\}$

$A = \{x \mid (x \text{ is a frequently occurred word in spams})\}$

Algebra

Common Number Sets

Natural Numbers(자연수)

$$\mathbb{N} = \{1, 2, 3, \dots\} = \{x \mid (x \text{는 자연수})\}$$

Whole Number

$$\mathbb{W} = \{0, 1, 2, \dots\} = \{x \mid (x \text{는 } 0 \vee x \text{는 자연수})\} \rightarrow 0 \text{을 포함한 자연수}$$

Integers(정수)

$$\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\} = \{x \mid (x \text{는 정수})\}$$

Rational Numbers(유리수)

$$\begin{aligned} \mathbb{Q} &= \left\{\frac{1}{2}, \frac{1}{3}, \frac{2}{3}, \frac{101}{100}, \dots\right\} = \{x \mid (x \text{는 유리수})\} \\ &= \{x \mid x = \frac{p}{q}, (p, q \text{ are integers } \wedge q \neq 0)\} \end{aligned}$$

Irrational Numbers(무리수)

$$\mathbb{I} = \{\pi, e, \sqrt{2}, \dots\} = \{x \mid (x \text{는 무리수})\} = \{x \mid \neg(x \text{는 유리수})\}$$

Real Numbers(실수)

$$\mathbb{R} = \{x \mid (x \text{는 실수})\} = \{x \mid (x \text{는 유리수 } \vee x \text{는 무리수})\}$$

Complex Numbers(복소수)

$$\mathbb{C} = \{x \mid (x \text{는 복소수})\} = \{a + j \cdot b \mid (a, b \text{는 실수})\}$$

Algebra

Coordinate Spaces

Coord. Plane, Space

$$\mathbb{R}^2 = \{(x, y) \mid x, y \text{는 실수}\}$$

$$\mathbb{R}^3 = \{(x, y, z) \mid x, y, z \text{는 실수}\}$$

Higher Dimensional Spaces

$$\mathbb{R}^n = \{(x_1, x_2, \dots, x_n) \mid \forall x_i \text{는 실수}\}$$

Algebra


Functions

Domain: a **set of** departure of a function

Codomain: a **set of** destination of a function


Range(Image): The image of f is then **the subset of** Y consisting of only those elements y of Y such that there is at least one x in X with $f(x) = y$.

Function: a binary relation between two sets that associates to each element of the first set exactly one element of the second set.



Algebra

Lines and Planes



Line: a set of points whose coordinates satisfy a given linear equation


$$y = ax + b$$

$$L = \{(x, y) \mid y = ax + b\}$$

평면

Plane: a set of all points \mathbf{r} such that, $\mathbf{n} \cdot (\mathbf{r} - \mathbf{r}_0) = 0$

$$P = \{(x, y, z) \mid \mathbf{n} \cdot (\mathbf{r} - \mathbf{r}_0) = 0\}$$



Algebra

Intersections


Intersection of L and P = the set of all points that lie on both L and P

No intersection $\implies S = \emptyset$

Point intersection $\implies S = \{(x_s, y_s, z_s)\}$

Lines intersection $\implies S = \{\mathbf{r} \mid (\mathbf{r} \text{ is on } L)\}$

→ 객체가 큰 교정이 된다



Algebra**Solution Sets****Solution Set of Equations** $f(x) = 0$

solution set of the equation = the set of all x 's that satisfy the equation

$$(x - 2)(x + 3) = 0 \rightarrow S = \{2, -3\}$$

$$\sin(x) = 0 \rightarrow S = \{\dots, -2\pi, -\pi, 0, \pi, 2\pi, \dots\} = \{x \mid x = k\pi, k \text{ is an integer}\}$$

Solution Set of Inequalities $f(x) < 0$

solution set of the inequality = the set of all x 's that satisfy the inequality

$$(x - 2)(x + 3) < 0 \rightarrow S = \{x \mid -3 < x < 2\}$$

Linear Algebra

Vector Space

a **set of** objects called vectors

Linear Subspace

W is a subset of V , then W is a linear subspace of V if under the operations of V , W is a vector space over K .

Linear Span

the set of linear combinations of elements of S .

Basis

a **set B** is a basis if its elements are linearly independent and every element of V is a linear combination of elements of B .

Spectrum of a Matrix

the set of its eigenvalues.

Eigenspace

The set of all eigenvectors of T corresponding to the same eigenvalue, together with the zero vector, is called an eigenspace

\Rightarrow 집합이 선형 대수에서도 많이 사용되구나..!

Probability and Statistics

Sample Space **the set of** all possible outcomes or results of that experiment.

Event A subset of the sample space is an event

Random Process A stochastic or random process can be defined as **a collection of random variables** that is indexed by some mathematical set

Statistical Model A statistical model is a mathematical model that embodies **a set of** statistical assumptions concerning the generation of sample data

Cardinality of Sets

or cardinal number

$$|A| = (\# \text{ elements})$$

ex.1) $A = \{0, 1\} \longrightarrow |A| = 2 \rightarrow \text{cardinal number} \rightarrow 2$

ex.2) $B = \{a, b, c\} \longrightarrow |B| = 3$

ex.3) $C = \{x \mid (x \text{ is an 1-digit integer})\} \longrightarrow |C| = 10 \rightarrow 0, 1, 2, \dots, 9$

ex.4) $D = \{x \mid (x \text{ is an alphabet})\} \longrightarrow |D| = 26$

원소를 하나만 가지는 집합

Cardinality of Empty Set $|\emptyset| = 0$

Singleton Set $|A| = 1$

Equivalent Sets $|A| = |B|$

\hookrightarrow Cardinality가 같은 Set 의미

Finite/Infinite Sets

Finite Sets


원소의 개수가 한정되어 있는 집합

$$|A| = 0 \text{ or } n$$

⇒ 유한하게 한정됨.

Encoding of Elements

주로 컴퓨터의 연산을 위해, 원소들을 index에 대응시키는 과정




Finite/Infinite Sets

Infinite Sets 무한 집합

원소의 개수가 무한한 집합

$$|A| = \infty$$

ex) $\mathbb{N}, \mathbb{W}, \mathbb{Z}, \mathbb{Q}, \mathbb{I}, \mathbb{R}, \mathbb{C}$



Countably Infinite Sets

원소들을 index에 대응시킬 수 있는 무한집합

ex) $\mathbb{N}, \mathbb{W}, \mathbb{Z}, \mathbb{Q}$

encoding이 가능함

어떻게 countable?

→ $\{ \dots, -3, -2, -1, 0, 1, 2, 3, \dots \}$

순서를 바꿔보자!

⇒ $\{ 0, 1, -1, 2, -2, 3, -3, \dots \}$

↓ ↓ ↓ ↓ ↓
1 2 3 4 5 ...

수학으로부터 인류를 자유롭게 하라

Uncountably Infinite Sets

원소들을 index에 대응시킬 수 없는 무한집합

ex) \mathbb{R}, \mathbb{C}

2.4 Inclusion and Exclusion

Inclusion/Exclusion of Elements

원소들은 어떤 집합에 포함될 수도, 포함되지 않을 수 있다.


$$\text{(원소 } a \text{가 집합 } A \text{에 포함됨)} = (a \in A)$$

$$\text{(원소 } a \text{가 집합 } A \text{에 포함되지 않은)} = (a \notin A)$$

ex) $A = \{a, b, c, d\}$

$$a \in A, b \in A, c \in A, d \in A$$

$$e \notin A, f \notin A, g \notin A, h \notin A$$



$$a \notin A$$

$$b \in A$$

2.4 Inclusion and Exclusion


Equal Sets

집합 A 의 모든 원소가 집합 B 에 포함되고 반대도 성립할 때, A, B 는 서로 같은 집합이다.

$$A = B \iff [\underbrace{(\forall a \in A)}_{\text{U}} \underbrace{\in B} \wedge [\underbrace{(\forall b \in B)}_{\text{U}} \underbrace{\in A}]]$$

$$A = \{a, b, c\}$$

$$B = \{a, b, c\}$$



Inclusion/Exclusion of Sets


Subset

집합 A 의 모든 원소가 집합 B 에 포함될 때, A 는 B 의 subset이라 한다.

$$A \subseteq B \longleftrightarrow (\forall a \in A) a \in B$$

Cardinality

$$A \subseteq B \longrightarrow |A| \leq |B|$$




Superset


집합 B 의 모든 원소가 집합 A 에 포함될 때, A 는 B 의 superset이라 한다.

$$A \supseteq B \longleftrightarrow (\forall b \in B) b \in A$$

$$A \supseteq B \longrightarrow |A| \geq |B|$$



Inclusion/Exclusion of Sets



$$\begin{aligned} A &\subseteq B, B \subseteq C, A \subseteq C \\ B &\supseteq A, C \supseteq B, C \supseteq A \end{aligned}$$

2.4 Inclusion and Exclusion

Inclusion/Exclusion of Sets

Example

$$A = \{a, b, c, d\}$$

$$\emptyset \subseteq A$$

$$\{a\} \subseteq A, \{b\} \subseteq A, \{c\} \subseteq A, \{d\} \subseteq A$$

$$\{a, b\} \subseteq A, \{a, c\} \subseteq A, \{a, d\} \subseteq A, \\ \{b, c\} \subseteq A, \{b, d\} \subseteq A, \{c, d\} \subseteq A$$

$$\{a, b, c\} \subseteq A, \{a, b, d\} \subseteq A, \\ \{a, c, d\} \subseteq A, \{b, c, d\} \subseteq A$$

$$\{a, b, c, d\} \subseteq A \longrightarrow A \subseteq A$$

A is a subset of itself.

$$\{a, b, c, d\} \supseteq A \longrightarrow A \supseteq A$$

$$\{a, b, c, d, e\} \supseteq A$$

$$\{a, b, c, d, f\} \supseteq A$$

$$\{a, b, c, d, e, f\} \supseteq A$$

2.4 Inclusion and Exclusion

Proper Subsets/Supersets


집부분집합

Proper Subsets

집합 A, B 에 대해 A 가 B 의 subset이지만 완전히 같지는 않을 때, A 는 B 의 proper subset이라 한다.
 적어도 B 의 원소 중 하나는 A 에 포함되지 않아야 한다.

$$A \subset B \iff [(\forall a \in A) \in B] \wedge [A \neq B]$$

$$A \subset B \longrightarrow |A| < |B|$$




Proper Supersets

집합 A, B 에 대해 A 가 B 의 superset이지만 완전히 같지는 않을 때, A 는 B 의 proper superset이라 한다.
 적어도 A 의 원소 중 하나는 B 에 포함되지 않아야 한다.

$$A \supset B \iff [(\forall b \in B) \in A] \wedge [A \neq B]$$

$$A \supseteq B \longrightarrow |A| \geq |B|$$



2.4 Inclusion and Exclusion

Proper Subsets/Supersets**Example.1**

$$A = \{a, b, c, d\}$$

$$A \not\supset A$$

$$\emptyset \subset A$$

$$\{a\} \subset A, \{b\} \subset A, \{c\} \subset A, \{d\} \subset A$$

$$\{a, b\} \subset A, \{a, c\} \subset A, \{a, d\} \subset A, \\ \{b, c\} \subset A, \{b, d\} \subset A, \{c, d\} \subset A$$

$$\{a, b, c\} \subset A, \{a, b, d\} \subset A, \\ \{a, c, d\} \subset A, \{b, c, d\} \subset A$$

A is not a proper subset of A .

A is not a proper superset of A .


$$\{a, b, c, d, e\} \supset A$$

$$\{a, b, c, d, f\} \supset A$$

$$\{a, b, c, d, e, f\} \supset A$$

Proper Subsets/Supersets

Example.2



2.4 Inclusion and Exclusion

Proper Subsets/Supersets

ex.3)

$$\begin{aligned}
 X &= \{x \mid (x \text{는 } 4\text{의 배수})\} & Y &= \{x \mid (x \text{는 } 8\text{의 배수})\} \\
 &= \{4, 8, 12, 16, 20, \dots\} & &= \{8, 16, 24, 32, 40, \dots\}
 \end{aligned}$$

$$X \supset Y, Y \subset X$$



ex.4)

$$\begin{aligned}
 A &= \{2, 4\} & B &= \{x \mid (x \text{는 } 12\text{의 약수})\} \\
 & & &= \{1, 2, 3, 4, 6, 12\}
 \end{aligned}$$

$$A \subset B, B \supset A$$

ex.5)

$$A = \{3, 5, 7, 9\} \quad B = \{x \mid \alpha \leq x \in \mathbb{N} \leq \beta\}$$

$$A \subseteq B \text{이기 위해, } \alpha \leq 3, \beta \geq 9$$

증가하고 있는
집합들의 나열?


Increasing/Decreasing Sequences of Sets

Increasing Sequences of Sets

$$A_1 \subseteq A_2 \subseteq \dots \subseteq A_n$$

$$A_k \subseteq A_{k+1}, 1 \leq k \in \mathbb{N} \leq n-1$$

$$A_k \subseteq A_{k+1}, 1 \leq k \in \mathbb{N} \leq n-1 \longrightarrow |A_k| \leq |A_{k+1}|$$




Decreasing Sequences of Sets

$$A_1 \supseteq A_2 \supseteq \dots \supseteq A_n$$

$$A_k \supseteq A_{k+1}, 1 \leq k \in \mathbb{N} \leq n-1$$

$$A_k \supseteq A_{k+1}, 1 \leq k \in \mathbb{N} \leq n-1 \longrightarrow |A_k| \geq |A_{k+1}|$$



Increasing/Decreasing Sequences of Sets

Example.1

$$A_i = \{x \mid (x \text{는 } 2^i \text{의 배수})\}$$

$$A_1 = \{x \mid (x \text{는 } 2 \text{의 배수})\} = \{2, 4, 6, 8, 10, \dots\}$$

$$A_2 = \{x \mid (x \text{는 } 4 \text{의 배수})\} = \{4, 8, 12, 16, 20, \dots\}$$


$$A_3 = \{x \mid (x \text{는 } 8 \text{의 배수})\} = \{8, 16, 24, 32, 40, \dots\}$$

$$\implies A_1 \supset A_2 \supset A_3 \supset \dots$$

$$\implies A_1, A_2, A_3, \dots : \text{decreasing sequence}$$

Increasing/Decreasing Sequences of Sets

Example.2 The key idea of residual networks



Unary/Binary Operations

Operations on Sets

일정한 규칙을 통해 새로운 집합을 만들어내는 과정

Unary Operations

$$f: A \longrightarrow B$$

$\sqrt{4} = 2 \rightarrow$ 이걸 집합 하나 가지고
새로운 집합을 만들어내는
연산

- power set of sets
- complement of sets

Binary Operations

$$f: A \times B \longrightarrow C$$

- Intersection of sets
- union of sets
- set difference
- symmetric difference
- Cartesian product of sets

\rightarrow 두개의 집합 이용해서
새로운 집합 만들어내는거

Unary Operations - Power Sets

Power Sets

집합 A 의 모든 subset들의 집합 $\mathcal{P}(A)$

모든 원소들은 “집합”

Power Set and Cardinality

ex.1)

$$A = \{0, 1\}$$

Subsets $\Rightarrow \emptyset, \{0\}, \{1\}, \{0, 1\}$

$$\emptyset \in P(A)$$

$$\{0\} \in P(A)$$

$$\{1\} \in P(A)$$

$$\{0, 1\} \in P(A)$$

$$\mathcal{P}(A) = \{\emptyset, \{0\}, \{1\}, \{0, 1\}\}$$

$$\mathcal{P}(A) = \{X \mid X \subseteq A\}$$

$$|\mathcal{P}(A)| = 2^{|A|}$$

이 집합의
~하트

ex.2)

$$B = \{a, b, c\}$$

Subsets $\Rightarrow \emptyset, \{a\}, \{b\}, \{c\}, \{a, b\}, \{a, c\}, \{b, c\}, \{a, b, c\}$

$$\emptyset \in P(B)$$

$$\{a\} \in P(B)$$

$$\{b\} \in P(B)$$

$$\{c\} \in P(B)$$

$$\{a, b\} \in P(B)$$

$$\{b, c\} \in P(B)$$

$$\{a, c\} \in P(B)$$

$$\{a, b, c\} \in P(B)$$

$$\mathcal{P}(B) = \{\emptyset, \{a\}, \{b\}, \{c\}, \{a, b\}, \{a, c\}, \{b, c\}, \{a, b, c\}\}$$

Unary Operations - Power Sets

Power Sets and Binary Numbers

포함 | 비포함 → 2개의 경우.
↑
B = {a, b, c}

| Decimal | Bin. Num. | | | Subset |
|---------|-----------|--|--|-------------|
| 0 | 000 | | | \emptyset |
| 1 | 001 | | | {c} |
| 2 | 010 | | | {b} |
| 3 | 011 | | | {b, c} |
| 4 | 100 | | | {a} |
| 5 | 101 | | | {a, c} |
| 6 | 110 | | | {a, b} |
| 7 | 111 | | | {a, b, c} |

*Symmetric Table

Unary Operations - Power Sets

Note!

1. Power set의 원소들은 “집합”
2. \emptyset 과 A 는 $\mathcal{P}(A)$ 의 원소
3. $\mathcal{P}(A)$ 의 원소들은 binary number로 인코딩이 가능하다.

Unary Operations - Power Sets

ex.1)

$$A = \left\{ \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \right\}$$

$$\longrightarrow \mathcal{P}(A) = \left\{ \emptyset, \left\{ \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \right\}, \left\{ \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \right\}, \left\{ \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \right\} \right\}$$

ex.2)

$$B = \left\{ \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} \right\}$$

$$\longrightarrow \mathcal{P}(B) = \left\{ \emptyset, \left\{ \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} \right\}, \left\{ \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} \right\}, \left\{ \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} \right\}, \left\{ \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} \right\}, \left\{ \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} \right\}, \left\{ \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} \right\}, \left\{ \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} \right\} \right\}$$

ex.3)


$$C = \{R, G, B\}$$

$$\longrightarrow \mathcal{P}(C) = \{ \emptyset, \{R\}, \{G\}, \{B\}, \{R, G\}, \{R, B\}, \{G, B\}, \{R, G, B\} \}$$

Unary Operations - Complements

A 에 포함되지 않은 원소들을 모은 집합을 A 의 complement이라 하고, A^c 로 표현한다.


$$A^c = \{x \mid x \notin A\}$$



$$|A| + |A^c| = |U|$$

Cardinality

$$|A^c| = |U| - |A|$$




Unary Operations - Complements

ex.1)

$$U = \{x \mid 5 \leq (x \in \mathbb{N}) \leq 10\}$$

$$A = \{7, 8\}$$

$$B = \{8, 9, 10\}$$




$$A^c = \{5, 6, 9, 10\}$$

$$B^c = \{5, 6, 7\}$$

ex.2)

$$U = \left\{ \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \right\}$$

$$A = \left\{ \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \right\} \quad B = \left\{ \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \right\}$$



$$A^c = \left\{ \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \right\}$$

$$B^c = \left\{ \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} \right\}$$



Binary Operations - Intersections and Unions

집합 A, B 에 모두 포함되는 원소들을 모든 집합을 A 와 B 의 intersection(교집합)이라고 부르고, $A \cap B$ 로 나타낸다.

$$A \cap B = \{x \mid (x \in A) \wedge (x \in B)\} \quad \text{A} \cap B \text{는 가끔 } AB \text{로 표현하기도 한다.}$$

집합 A 또는(or) B 에 포함되는 원소들을 모든 집합을 A 와 B 의 union(합집합)이라고 부르고, $A \cup B$ 로 나타낸다.

$$A \cup B = \{x \mid (x \in A) \vee (x \in B)\}$$



Binary Operations - Intersections and Unions

Cardinality

$$|A \cup B| = |A| + |B| - |A \cap B|$$

$$|A \cap B| \leq |A|, \quad |A \cap B| \leq |B|$$

$$|A \cup B| \geq |A|, \quad |A \cup B| \geq |B|$$



Special Cases

$$A \subseteq B \longrightarrow A \cup B = B, \quad A \cap B = A$$

$$|A \cap B| = |A|, \quad |A \cup B| = |B|$$

$$A \supseteq B \longrightarrow A \cup B = A, \quad A \cap B = B$$

$$|A \cap B| = |B|, \quad |A \cup B| = |A|$$




Binary Operations - Intersections and Unions

General Intersections

$$\bigcap_{i=1}^n A_i = A_1 \cap A_2 \cap \dots \cap A_n$$


$$\bigcap_{i=1}^n A_i = A_1 A_2 \dots A_n$$



Binary Operations - Intersections and Unions

General Unions

$$\bigcup_{i=1}^n A_i = A_1 \cup A_2 \cup \dots \cup A_n$$




Binary Operations - Intersections and Unions

Special Cases(2)

$$A_k \subseteq A_{k+1}, \quad 1 \leq k \in \mathbb{N} \leq n-1 \quad \longrightarrow \quad \bigcap_{i=1}^n A_i = A_1, \quad \bigcup_{i=1}^n A_i = A_n$$

$$\left| \bigcup_{i=1}^n A_i \right| = |A_n|, \quad \left| \bigcap_{i=1}^n A_i \right| = |A_1|$$

$$\left| \bigcup_{i=1}^{k-1} A_i \right| \leq \left| \bigcup_{i=1}^k A_i \right|, \quad \left| \bigcap_{i=1}^{k-1} A_i \right| = \left| \bigcap_{i=1}^k A_i \right| = |A_1|$$



Binary Operations - Intersections and Unions

Special Cases(2)

$$A_k \supseteq A_{k+1}, \quad 1 \leq k \in \mathbb{N} \leq n-1 \quad \longrightarrow \quad \bigcap_{i=1}^n A_i = A_n, \quad \bigcup_{i=1}^n A_i = A_1$$



$$\left| \bigcup_{i=1}^n A_i \right| = |A_1|, \quad \left| \bigcap_{i=1}^n A_i \right| = |A_n|$$

$$\left| \bigcup_{i=1}^{k-1} A_i \right| = \left| \bigcup_{i=1}^k A_i \right| = |A_1|, \quad \left| \bigcap_{i=1}^{k-1} A_i \right| \geq \left| \bigcap_{i=1}^k A_i \right|$$

Binary Operations - Intersections and Unions

The Algebraic Properties

Commutative Law

$$A \cup B = B \cup A$$

$$A \cap B = B \cap A$$

Associative Law

$$(A \cup B) \cup C = A \cup (B \cup C)$$

$$(A \cap B) \cap C = A \cap (B \cap C)$$


Distributive Law

$$A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$$

$$A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$$

$$A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$$

$$A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$$



Binary Operations - Intersections and Unions

Identities


$$(1) A \cup \emptyset = A$$

$$(2) A \cap \emptyset = \emptyset$$

$$(3) A \cup U = U$$

$$(4) A \cap U = A$$

$$(5) A \cup A^c = U$$

$$(6) A \cap A^c = \emptyset$$

$$(7) (A^c)^c = A$$

Binary Operations - Intersections and Unions

De Morgan's Law


$$(A \cap B)^c = A^c \cup B^c$$

$$(A \cup B)^c = A^c \cap B^c$$

Examples

$$(A^c \cap B)^c = [(A^c)^c \cup B^c] = A \cup B^c$$

$$(A \cup B^c)^c = [A^c \cap (B^c)^c] = A^c \cap B$$




Binary Operations - Set Differences

집합 A, B 에 대해 A 에는 포함되고, B 에는 포함되지 않은 원소들을 모은 집합을 $A - B$ 로 나타내고, set difference(차집합)이라고 부른다.

$$A - B = \{x \mid (x \in A) \wedge (x \notin B)\}$$

$A - B$ 는 $A \setminus B$ 로 표현하기도 한다.



Binary Operations - Set Differences

Computation Exercises

$$(1) \quad A - B = A \cap B^c$$

$$(2) \quad B - A = B \cap A^c$$

$$(3) \quad A - B = A - (A \cap B) \\ = (A \cup B) - B$$

$$(4) \quad B - A = B - (A \cap B) \\ = (A \cup B) - A$$

$$A - B = A \cap B^c$$

Binary Operations - Set Differences

Computation Exercises


$$\begin{aligned}
 (5) \quad (A - B) \cup (A \cap B) &= (A \cap B^c) \cup (A \cap B) \\
 &= A \cap (B^c \cup B) \\
 &= A \cap U \\
 &= A
 \end{aligned}$$

$$\begin{aligned}
 (6) \quad (B - A) \cup (A \cap B) &= (B \cap A^c) \cup (A \cap B) \\
 &= \underbrace{(B \cap A^c) \cup (B \cap A)}_{\cap A^c} \\
 &= B \cap (A^c \cup A) \\
 &= B \cap U \\
 &= B
 \end{aligned}$$


$$\begin{aligned}
 (7) \quad A - (A - B) &= A - (A \cap B^c) \\
 &= A \cap (A \cap B^c)^c \\
 &= A \cap (A^c \cup B) \\
 &= (A \cap A^c) \cup (A \cap B) \\
 &= \emptyset \cup (A \cap B) \\
 &= A \cap B
 \end{aligned}$$

Binary Operations - Set Differences

set difference는 relative complement라고 부르기도 한다.



$$U - A = U \cap A^c$$




$$A - B = A \cap B^c$$

$$B - A = B \cap A^c$$

Binary Operations - Set Differences

Special Case



$$A \cap B = \emptyset$$

$$A - B = A$$

$$B - A = B$$

Binary Operations - Set Differences

The Algebraic Properties

Anti-commutativity $A - B \neq B - A$

Anti-associativity $A - (B - C) \neq (A - B) - C$

Distributive Law

$$\begin{aligned}(1) \quad C - (A \cap B) &= C \cap (A \cap B)^c \\ &= C \cap (A^c \cup B^c) \\ &= (C \cap A^c) \cup (C \cap B^c) \\ &= (C - A) \cup (C - B)\end{aligned}$$

$$\begin{aligned}(2) \quad C - (A \cup B) &= C \cap (A \cup B)^c \\ &= C \cap (A^c \cap B^c) \\ &= (C \cap A^c) \cap (C \cap B^c) \\ &= (C - A) \cap (C - B)\end{aligned}$$


$$\cup \Rightarrow \cap, \cap \Rightarrow \cup$$

Binary Operations - Symmetric Differences

집합 A, B 에 대해 $A - B$ 와 $B - A$ 의 union

$$A \triangle B = \{x \mid (A - B) \cup (B - A)\}$$

$$= \{x \mid [(x \in A) \vee (x \in B)] \wedge (x \notin A \cap B)\}$$



$$A \triangle B = (A - B) \cup (B - A)$$

$$= (A \cap B^c) \cup (B \cap A^c)$$

$$A \cap B^c = X$$

$$= X \cup (B \cap A^c)$$

$$= (X \cup B) \cap (X \cup A^c)$$

$$= [(A \cap B^c) \cup B] \cap [(A \cap B^c) \cup A^c]$$

$$= [(A \cup B) \cap (B^c \cup B)] \cap [(A \cup A^c) \cap (B^c \cup A^c)]$$

$$= [(A \cup B) \cap U] \cap [U \cap (B \cap A)^c]$$

$$= (A \cup B) \cap (B \cap A)^c$$

$$= (A \cup B) - (A \cap B)$$

오류정해내기

Binary Operations - Cartesian Product

집합 A, B 에서 원소 a, b 들을 각각 뽑아 뽑아 (a, b) 를 만들 때, 모든 (a, b) 들의 집합을 $A \times B$ 라 한다.

$$A \times B = \{(a, b) \mid (a \in A) \wedge (b \in B)\}$$


$$A = \{a_1, a_2, \dots, a_m\}$$

$$B = \{b_1, b_2, \dots, b_n\}$$

| | b_1 | b_2 | \dots | b_n |
|----------|--------------|--------------|----------|--------------|
| a_1 | (a_1, b_1) | (a_1, b_2) | \dots | (a_1, b_n) |
| a_2 | (a_2, b_1) | (a_2, b_2) | \dots | (a_2, b_n) |
| \vdots | \vdots | \vdots | \ddots | \vdots |
| a_m | (a_m, b_1) | (a_m, b_2) | \dots | (a_m, b_n) |

Cardinality $\rightarrow m$

Cardinality $\rightarrow n$



$$A \times B = \{(a_1, b_1), (a_1, b_2), \dots, (a_1, b_n) \\ (a_2, b_1), (a_2, b_2), \dots, (a_2, b_n) \\ \vdots \\ (a_m, b_1), (a_m, b_2), \dots, (a_m, b_n)\}$$

$$|A \times B| = m \times n = mn \text{ 개}$$

Binary Operations - Cartesian Product

ex.1)

두개의 object가 많으면 쌍을 만들지! 

$$\underline{A = \{0, 1, 2\}}, \quad \underline{B = \{a, b\}} \longrightarrow A \times B$$

| | a | b |
|---|-------|-------|
| 0 | (0,a) | (0,b) |
| 1 | (1,a) | (1,b) |
| 2 | (2,a) | (2,b) |

$$A \times B = \{(0, a), (0, b), (1, a), (1, b), (2, a), (2, b)\}$$

Binary Operations - Cartesian Product

ex.2)

$$A = \{0, 1, 2\} \longrightarrow A \times A = A^2$$

| | 0 | 1 | 2 |
|---|-------|-------|-------|
| 0 | (0,0) | (0,1) | (0,2) |
| 1 | (1,0) | (1,1) | (1,2) |
| 2 | (2,0) | (2,1) | (2,2) |

$$|A| = 3$$

↓

$$|A^2| = 3^2$$

$$A \times B = \{(0,0), (0,1), (0,2)\}$$

$$(1,0), (1,1), (1,2)$$

$$(2,0), (2,1), (2,2)\}$$




Binary Operations - Cartesian Product


ex.3)

$$\mathbb{R} \longrightarrow \mathbb{R}^2 = \{(x, y) \mid x, y \in \mathbb{R}\}$$


$$\mathbb{R} \longrightarrow \mathbb{R}^3 = \{(x, y, z) \mid x, y, z \in \mathbb{R}\}$$



$$\mathbb{R} \Rightarrow \xrightarrow{\text{9등 분할}}$$




... \Rightarrow



Disjoint Sets

집합 A, B 에 대해 $A \cap B = \emptyset$ 일 때 A, B 는 disjoint set이다.

disjoint는 **mutually exclusive**라고도 부른다.



Cardinality

$$|A \cap B| = 0$$


$$\star |A \cup B| = |A| + |B|$$

Partitions of Sets

Universal set(U)안에 n 개의 집합 A_1, A_2, \dots, A_n 이 있고, U 의 모든 원소들이 모두 단 하나의 A_i 에만 포함될 때, $\{A_1, A_2, \dots, A_n\}$ 를 U 의 partition이라 부른다.


$$A_i \cap A_j = \emptyset, i \neq j$$

$$\bigcup_{i=1}^n A_i = U$$




$$A \cap B = \emptyset$$

$$A \cup B = U$$



$$|A| + |B| = |U|$$



$$|A_1| + |A_2| + \dots + |A_n| = |U|$$

Partitions of Sets

Non-uniqueness of Partitions

Partition of Sets is **NOT UNIQUE**

$$A = \{x \mid 1 \leq x \in \mathbb{N} \leq 20\}$$

$$A_1 = \{x \mid 1 \leq x \in \mathbb{N} \leq 10\}$$

$$A_2 = \{x \mid 10 < x \in \mathbb{N} \leq 20\}$$

$$A_1 \cap A_2 = \emptyset$$

$$A_1 \cup A_2 = A$$

The $\{A_1, A_2\}$ is a partition of A

$$A_3 = \{x \mid (1 \leq x \in \mathbb{N} \leq 20) \wedge (x = 2n)\}$$

$$A_4 = \{x \mid (1 \leq x \in \mathbb{N} \leq 20) \wedge (x = 2n + 1)\}$$

$$A_3 \cap A_4 = \emptyset$$

$$A_3 \cup A_4 = A$$

The $\{A_3, A_4\}$ is a partition of A

$$A_5 = \{x \mid (1 \leq x \in \mathbb{N} \leq 20) \wedge (x \bmod 3 = 0)\}$$

$$A_6 = \{x \mid (1 \leq x \in \mathbb{N} \leq 20) \wedge (x \bmod 3 = 1)\}$$

$$A_7 = \{x \mid (1 \leq x \in \mathbb{N} \leq 20) \wedge (x \bmod 3 = 2)\}$$

$$A_5 \cap A_6 = \emptyset$$

$$A_5 \cap A_7 = \emptyset$$

$$A_6 \cap A_7 = \emptyset$$


$$A_5 \cup A_6 \cup A_7 = A$$

The $\{A_5, A_6, A_7\}$ is a partition of A

Partitions of Sets

Making a Complete Set with Partitions

$$B = \bigcup_{i=1}^n (A_i \cap B)$$



Partitions of Sets

ex.1)

A_i : 나이(x)가 $10 \cdot i \leq x \leq 10 \cdot (i + 1)$ 인 사람들의 집합

G : 안경을 쓴 사람들의 집합

| | G | G^c |
|----------|-----------------|-------|
| A_0 | $A_0 \cap G$ | |
| A_1 | $A_1 \cap G$ | |
| | | |
| A_{10} | $A_{10} \cap G$ | |

$$G = \bigcup_{i=1}^{10} (A_i \cap G)$$

$$G^c = \bigcup_{i=1}^{10} (A_i \cap G^c)$$

CLOSING

Basic Algebra

Chap.2 Sets