

# Willmore type inequalities in geodesic balls of hyperbolic space

Xiaoxiang Chai (Korea Institute for Advanced Study)

CoSAN conference, Korea University, Feb 20-23, 2023

# Outline

Mean curvature

Mean curvature type flow

Free boundary surface

First result: Convergence of the flow

Second result: Monotonicity of a Willmore type quantity

Proof: Monotonicity of a Willmore type quantity

# Mean curvature

- ▶ Mean curvature is the local variation of area under normal variations

# Mean curvature

- ▶ Mean curvature is the local variation of area under normal variations
- ▶ Let  $\Sigma$  be a hypersurface,  $\nu$  be a chosen unit normal to  $\Sigma$

# Mean curvature

- ▶ Mean curvature is the local variation of area under normal variations
- ▶ Let  $\Sigma$  be a hypersurface,  $\nu$  be a chosen unit normal to  $\Sigma$
- ▶ Let  $\Sigma_t$  be a differentiable family of hypersurfaces deforming  $\Sigma_0 = \Sigma$ , let  $X$  be the variational vector field (we choose compactly supported variations), then

$$\frac{d}{dt} \text{vol}(\Sigma_t)|_{t=0} = \int_{\Sigma_0} H \langle X, \nu \rangle$$

# Mean curvature

- ▶ Mean curvature is the local variation of area under normal variations
- ▶ Let  $\Sigma$  be a hypersurface,  $\nu$  be a chosen unit normal to  $\Sigma$
- ▶ Let  $\Sigma_t$  be a differentiable family of hypersurfaces deforming  $\Sigma_0 = \Sigma$ , let  $X$  be the variational vector field (we choose compactly supported variations), then

$$\frac{d}{dt} \text{vol}(\Sigma_t)|_{t=0} = \int_{\Sigma_0} H \langle X, \nu \rangle$$

- ▶ The above formula is also known as **the first variation of area**

# Pointwise definition

- ▶ Let  $\{e_i\}$  be an orthonormal frame tangent to  $\Sigma$

# Pointwise definition

- ▶ Let  $\{e_i\}$  be an orthonormal frame tangent to  $\Sigma$
- ▶ The second fundamental form is defined to be  $\bar{\nabla}\nu$ , in components,  $h_{ij} = \langle \bar{\nabla}_{e_i}\nu, e_j \rangle$



# Pointwise definition

- ▶ Let  $\{e_i\}$  be an orthonormal frame tangent to  $\Sigma$
- ▶ The second fundamental form is defined to be  $\bar{\nabla}\nu$ , in components,  $h_{ij} = \langle \bar{\nabla}_{e_i}\nu, e_j \rangle$
- ▶ The mean curvature is just the trace of  $h$ :  $H = \sum_i \langle \nabla_{e_i}\nu, e_i \rangle$

# Mean curvature flow

1. Brakke 78 weak version (varifold)

# Mean curvature flow

1. Brakke 78 weak version (varifold)
2. Huisken 84 The convex surface in  $\mathbb{R}^n$  contracts to a round point in finite time under mean curvature flow

$$\partial_t x = -H\nu.$$

# Mean curvature flow

1. Brakke 78 weak version (varifold)
2. Huisken 84 The convex surface in  $\mathbb{R}^n$  contracts to a round point in finite time under mean curvature flow

$$\partial_t x = -H\nu.$$

3. Stahl 96 mean curvature flow with free boundary

## Inverse mean curvature flow

1. Urbas 91, Gerhardt 90 The surface (star-shaped, strictly mean convex) expands to a round sphere under **inverse mean curvature flow**

$$\partial_t x = \frac{1}{H} \nu.$$

# Inverse mean curvature flow

1. Urbas 91, Gerhardt 90 The surface (star-shaped, strictly mean convex) expands to a round sphere under **inverse mean curvature flow**

$$\partial_t x = \frac{1}{H} \nu.$$

- Gerhardt also studied fully nonlinear flows in spheres, hyperbolic space

## Inverse mean curvature flow

1. Urbas 91, Gerhardt 90 The surface (star-shaped, strictly mean convex) expands to a round sphere under **inverse mean curvature flow**

$$\partial_t x = \frac{1}{H} \nu.$$

- ▶ Gerhardt also studied fully nonlinear flows in spheres, hyperbolic space
2. Huisken-Ilmanen 01 Inverse mean curvature flow to prove a Penrose inequality; most importantly the Hawking mass

$$m_H(\Sigma) = |\Sigma|^{1/2} \left( 1 - \frac{1}{16\pi} \int_{\Sigma} H^2 \right)$$

is monotone increasing under IMCF starting from a 2-sphere.

# Inverse mean curvature flow

1. **Urbas 91, Gerhardt 90** The surface (star-shaped, strictly mean convex) expands to a round sphere under **inverse mean curvature flow**

$$\partial_t x = \frac{1}{H} \nu.$$

- ▶ Gerhardt also studied fully nonlinear flows in spheres, hyperbolic space
2. **Huisken-Ilmanen 01** Inverse mean curvature flow to prove a Penrose inequality; most importantly the Hawking mass

$$m_H(\Sigma) = |\Sigma|^{1/2} \left( 1 - \frac{1}{16\pi} \int_{\Sigma} H^2 \right)$$

is monotone increasing under IMCF starting from a 2-sphere.

3. **Neves 07, Wang-Hung 14** starting from convex hypersurface the IMCF does not converge. Converges in stronger conditions: principal curvatures  $\kappa_i > 1$ . The Penrose inequality asymptotically hyperbolic is still open; locally hyperbolic case is settled by **Lee-Neves 13**.



# Free boundary surface

- ▶ Given manifold  $M$  with nonempty boundary  $\partial M$ ,  $\eta$  be the unit outward normal of  $\partial M$  in  $M$

# Free boundary surface

- ▶ Given manifold  $M$  with nonempty boundary  $\partial M$ ,  $\eta$  be the unit outward normal of  $\partial M$  in  $M$
- ▶ if  $\partial\Sigma \subset \partial M$  and  $\langle \eta, \nu \rangle = 0$  everywhere along  $\partial\Sigma$ , then  $\Sigma$  is called a **free boundary surface**.

# Free boundary surface

- ▶ Given manifold  $M$  with nonempty boundary  $\partial M$ ,  $\eta$  be the unit outward normal of  $\partial M$  in  $M$
- ▶ if  $\partial\Sigma \subset \partial M$  and  $\langle \eta, \nu \rangle = 0$  everywhere along  $\partial\Sigma$ , then  $\Sigma$  is called a **free boundary surface**.
- ▶ If the mean curvature of  $\Sigma$  is zero, we say that  $\Sigma$  is a **free boundary minimal surface**

## Example: Critical catenoid

- ▶ Put a large sphere centered at the center of the neck of a standard catenoid in  $\mathbb{R}^3$ ;


## Example: Critical catenoid

- ▶ Put a large sphere centered at the center of the neck of a standard catenoid in  $\mathbb{R}^3$ ;
- ▶ Shrinking the radius of the sphere until it is orthogonally intersecting the catenoid;

## Example: Critical catenoid

- ▶ Put a large sphere centered at the center of the neck of a standard catenoid in  $\mathbb{R}^3$ ;
- ▶ Shrinking the radius of the sphere until it is orthogonally intersecting the catenoid;
- ▶ The piece inside the sphere is called critical catenoid.

# Critical catenoid



# Geometric inequalities in the unit ball

1. Wang-Xia 19 Minkowski formula, Heintz-Karcher



# Geometric inequalities in the unit ball

1. Wang-Xia 19 Minkowski formula, Heintz-Karcher
2. Lambert-Scheuer 16, 17 Inverse mean curvature flow of convex, free boundary surface converges to a flat disk and a Willmore type inequality; a Willmore type quantity

$$\frac{1}{2}|M_t|^{\frac{2-n}{n}} \int_{M_t} H^2 + \omega_n^{\frac{2-n}{n}} |\partial M_t|$$

is monotonically decreasing.

# Geometric inequalities in the unit ball

1. Wang-Xia 19 Minkowski formula, Heintz-Karcher
2. Lambert-Scheuer 16, 17 Inverse mean curvature flow of convex, free boundary surface converges to a flat disk and a Willmore type inequality; a Willmore type quantity

$$\frac{1}{2}|M_t|^{\frac{2-n}{n}} \int_{M_t} H^2 + \omega_n^{\frac{2-n}{n}} |\partial M_t|$$

is monotonically decreasing.

3. Scheuer-Wang-Xia 22 Alexandrov-Fenchel type inequality which involves higher order mean curvature; Wang-Weng-Xia 22 capillary.

## Setup in hyperbolic space $\mathbb{H}^{n+1}$

- ▶  $B := B^{n+1}$  be the ball of radius  $\rho_0$

## Setup in hyperbolic space $\mathbb{H}^{n+1}$

- ▶  $B := B^{n+1}$  be the ball of radius  $\rho_0$
- ▶  $M$  be a free boundary convex surface inside  $B$

## Setup in hyperbolic space $\mathbb{H}^{n+1}$

- ▶  $B := B^{n+1}$  be the ball of radius  $\rho_0$
- ▶  $M$  be a free boundary convex surface inside  $B$
- ▶ We are going to generalize **Lambert-Scheuer 16, 17** result to  $M$

## Theorem: Convergence of the flow

**Theorem** Let  $M$  be a strictly convex, free boundary hypersurface in the ball  $B$ , then the inverse mean curvature flow  $M_t$  converges to a totally geodesic disk in finite time.

# Monotonicity of a Willmore type quantity: Theorem

- **Theorem (C. 22)** Suppose that  $M$  is strictly convex, then  $m_H(M)$  given by

$$m_H(M) = |M|^{\frac{2-n}{n}} \int_M (H^2 - n^2) + \Lambda |\partial M|$$

is monotone under IMCF with free boundary (assume for now strict convexity is preserved and that  $|M_t| \leq \lambda$ ).

# Monotonicity of a Willmore type quantity: Theorem

- **Theorem (C. 22)** Suppose that  $M$  is strictly convex, then  $m_H(M)$  given by

$$m_H(M) = |M|^{\frac{2-n}{n}} \int_M (H^2 - n^2) + \Lambda |\partial M|$$

is monotone under IMCF with free boundary (assume for now strict convexity is preserved and that  $|M_t| \leq \lambda$ ).

- $\lambda$  volume of  $n$ -dimensional hyperbolic geodesic ball of radius  $\rho_0$  (**note** dimension  $n$  here!)



# Monotonicity of a Willmore type quantity: Theorem

- **Theorem (C. 22)** Suppose that  $M$  is strictly convex, then  $m_H(M)$  given by

$$m_H(M) = |M|^{\frac{2-n}{n}} \int_M (H^2 - n^2) + \Lambda |\partial M|$$

is monotone under IMCF with free boundary (assume for now strict convexity is preserved and that  $|M_t| \leq \lambda$ ).

- $\lambda$  volume of  $n$ -dimensional hyperbolic geodesic ball of radius  $\rho_0$  (**note** dimension  $n$  here!)
- $\Lambda = 2 \coth \rho_0 \lambda^{(2-n)/n}$

## Lower bound of the Willmore type quantity

- **Theorem (C. 22)** Assuming that the flow preserves strict convexity, and converges to a totally geodesic disk, then

$$|M|^{\frac{2-n}{n}} \int_M (H^2 - n^2) + \Lambda |\partial M| \geq -n^2 \lambda^{\frac{2}{n}} + \Lambda \omega_{n-1} \sinh^{n-1} \rho_0.$$

## Some notations

- ▶  $\bar{T}$  largest time such that  $M_t$  is strictly convex for all  $0 < t < \bar{T}$

## Some notations

- ▶  $\bar{T}$  largest time such that  $M_t$  is strictly convex for all  $0 < t < \bar{T}$
- ▶ Maximal existence time:  $T^*$  largest time such that  $M_t$  is mean convex for all  $0 < t < T^*$

## Key lemma from convex geometry in spheres

( Makowski-Scheuer 15 ) If  $\partial M_{\mathcal{T}}$  is weakly convex in  $\mathbb{S}^n$ , then it lies in an open hemisphere or it is the equator.

# Idea

- ▶ ( easy ) If  $\partial M_T$  is some equator, then using mean convexity of  $M_T$ , free boundary condition, maximum principle,  $M_T$  has to be a totally geodesic disk. ( $T$  being  $\bar{T}$  or  $T^*$  is OK)

# Idea

- ▶ ( easy ) If  $\partial M_T$  is some equator, then using mean convexity of  $M_T$ , free boundary condition, maximum principle,  $M_T$  has to be a totally geodesic disk. ( $T$  being  $\bar{T}$  or  $T^*$  is OK)
- ▶ If  $\partial M_{\bar{T}}$  is contained in some hemisphere, then

# Idea

- ▶ ( easy ) If  $\partial M_T$  is some equator, then using mean convexity of  $M_T$ , free boundary condition, maximum principle,  $M_T$  has to be a totally geodesic disk. ( $T$  being  $\bar{T}$  or  $T^*$  is OK)
- ▶ If  $\partial M_{\bar{T}}$  is contained in some hemisphere, then
  1. Lower bound of the mean curvature  $H$  before  $\bar{T}$  (flow loses convexity)



# Idea

- ▶ ( easy ) If  $\partial M_T$  is some equator, then using mean convexity of  $M_T$ , free boundary condition, maximum principle,  $M_T$  has to be a totally geodesic disk. ( $T$  being  $\bar{T}$  or  $T^*$  is OK)
- ▶ If  $\partial M_{\bar{T}}$  is contained in some hemisphere, then
  1. Lower bound of the mean curvature  $H$  before  $\bar{T}$  (flow loses convexity)
  2.  $\bar{T} = T^*$

# Idea

- ▶ ( easy ) If  $\partial M_T$  is some equator, then using mean convexity of  $M_T$ , free boundary condition, maximum principle,  $M_T$  has to be a totally geodesic disk. ( $T$  being  $\bar{T}$  or  $T^*$  is OK)
- ▶ If  $\partial M_{\bar{T}}$  is contained in some hemisphere, then
  1. Lower bound of the mean curvature  $H$  before  $\bar{T}$  (flow loses convexity)
  2.  $\bar{T} = T^*$
  3. If the  $\partial M_t$  is not an equator, then the flow can be continued.

# Idea

- ▶ ( easy ) If  $\partial M_T$  is some equator, then using mean convexity of  $M_T$ , free boundary condition, maximum principle,  $M_T$  has to be a totally geodesic disk. ( $T$  being  $\bar{T}$  or  $T^*$  is OK)
- ▶ If  $\partial M_{\bar{T}}$  is contained in some hemisphere, then
  1. Lower bound of the mean curvature  $H$  before  $\bar{T}$  (flow loses convexity)
  2.  $\bar{T} = T^*$
  3. If the  $\partial M_t$  is not an equator, then the flow can be continued.
  4. Which means at maximal existence time, the flow  $M_t$  converges to a totally geodesic disk.

# Evolution equations (Huisken 86)

1.  $\partial_t g_{ij} = \frac{2}{H} h_{ij}$

# Evolution equations (Huisken 86)

1.  $\partial_t g_{ij} = \frac{2}{H} h_{ij}$
2.  $\partial_t \sqrt{g} = \sqrt{g}$

# Evolution equations (Huisken 86)

1.  $\partial_t g_{ij} = \frac{2}{H} h_{ij}$
2.  $\partial_t \sqrt{g} = \sqrt{g}$
3.  $\partial_t \nu = -\nabla \frac{1}{H}$

## Evolution equations (Huisken 86)

1.  $\partial_t g_{ij} = \frac{2}{H} h_{ij}$
2.  $\partial_t \sqrt{g} = \sqrt{g}$
3.  $\partial_t \nu = -\nabla \frac{1}{H}$
4.  $\partial_t h_{ij} = -\bar{R}(\partial_i, \nu, \nu, \partial_j) \frac{1}{H} + \frac{h_i^k h_{jk}}{H} - \nabla_i \nabla_j \frac{1}{H}$

## Evolution equations (Huisken 86)

1.  $\partial_t g_{ij} = \frac{2}{H} h_{ij}$
2.  $\partial_t \sqrt{g} = \sqrt{g}$
3.  $\partial_t \nu = -\nabla \frac{1}{H}$
4.  $\partial_t h_{ij} = -\bar{R}(\partial_i, \nu, \nu, \partial_j) \frac{1}{H} + \frac{h_i^k h_{jk}}{H} - \nabla_i \nabla_j \frac{1}{H}$
5.  $\partial_t H = -\Delta \frac{1}{H} - (\text{Ric}(\nu) + |A|^2) \frac{1}{H}$



# Evolution of coordinates

- ▶ Hyperbolic space  $\mathbb{H}^{n+1}$  is realized as the upper half of the hyperboloid in the Minkowski space  $\mathbb{M}^{n+1,1}$

# Evolution of coordinates

- ▶ Hyperbolic space  $\mathbb{H}^{n+1}$  is realized as the upper half of the hyperboloid in the Minkowski space  $\mathbb{M}^{n+1,1}$
- ▶ Denote  $z$  as the hyperboloidal coordinates

# Evolution of coordinates

- ▶ Hyperbolic space  $\mathbb{H}^{n+1}$  is realized as the upper half of the hyperboloid in the Minkowski space  $\mathbb{M}^{n+1,1}$
- ▶ Denote  $z$  as the hyperboloidal coordinates
- ▶ By IMCF:  $\partial_t z^\alpha = \frac{1}{H} \nu^\alpha$

# Evolution of coordinates

- ▶ Hyperbolic space  $\mathbb{H}^{n+1}$  is realized as the upper half of the hyperboloid in the Minkowski space  $\mathbb{M}^{n+1,1}$
- ▶ Denote  $z$  as the hyperboloidal coordinates
- ▶ By IMCF:  $\partial_t z^\alpha = \frac{1}{H} \nu^\alpha$
- ▶ By Hessian decomposition

$$\Delta z^\alpha = n z^\alpha - H \nu^\alpha$$

# Boundary derivative of mean curvature

► (Stahl 96)

$$\nabla_{\eta} \frac{1}{H} = \langle \nabla_{\eta} (\frac{1}{H} \nu), \nu \rangle \quad (1)$$

$$= \langle \nabla_{\frac{1}{H} \nu} \eta, \nu \rangle \quad (2)$$

$$= \frac{1}{H} A_{\partial B}(\nu, \nu) \quad (3)$$

# Boundary derivative of mean curvature

► (Stahl 96)

$$\nabla_{\eta} \frac{1}{H} = \langle \nabla_{\eta} (\frac{1}{H} \nu), \nu \rangle \quad (1)$$

$$= \langle \nabla_{\frac{1}{H} \nu} \eta, \nu \rangle \quad (2)$$

$$= \frac{1}{H} A_{\partial B}(\nu, \nu) \quad (3)$$

►  $\nabla_{\eta} H = -H \coth \rho_0$

# Boundary derivative of mean curvature

► (Stahl 96)

$$\nabla_{\eta} \frac{1}{H} = \langle \nabla_{\eta} (\frac{1}{H} \nu), \nu \rangle \quad (1)$$

$$= \langle \nabla_{\frac{1}{H} \nu} \eta, \nu \rangle \quad (2)$$

$$= \frac{1}{H} A_{\partial B}(\nu, \nu) \quad (3)$$

►  $\nabla_{\eta} H = -H \coth \rho_0$

► Boundary derivative of  $z^0$ :  $\nabla_{\eta} \log z^0 = \tanh \rho_0$

# Boundary derivative of mean curvature

► (Stahl 96)

$$\nabla_\eta \frac{1}{H} = \langle \nabla_\eta (\frac{1}{H} \nu), \nu \rangle \quad (1)$$

$$= \langle \nabla_{\frac{1}{H} \nu} \eta, \nu \rangle \quad (2)$$

$$= \frac{1}{H} A_{\partial B}(\nu, \nu) \quad (3)$$

►  $\nabla_\eta H = -H \coth \rho_0$

► Boundary derivative of  $z^0$ :  $\nabla_\eta \log z^0 = \tanh \rho_0$

► Boundary derivative of  $z^i$ :  $\nabla_\eta z^i = \coth \rho_0 z^i$



# Curvature estimates

- **Proposition** Up to  $\bar{T}$ ,

$$\sup_{x \in M_t} |A| \leq C$$

with  $C$  depending only on the initial data (including  $\rho_0$ ).

# Curvature estimates

- **Proposition** Up to  $\bar{T}$ ,

$$\sup_{x \in M_t} |A| \leq C$$

with  $C$  depending only on the initial data (including  $\rho_0$ ).

- Use evolution of the mean curvature  $\log H$ ; because of  $\text{Ric} = -n$ , need to add  $\log z_0$ , i.e. we consider

$$\log H + \log z_0$$

# Curvature estimates

- ▶ **Proposition** Up to  $\bar{T}$ ,

$$\sup_{x \in M_t} |A| \leq C$$

with  $C$  depending only on the initial data (including  $\rho_0$ ).

- ▶ Use evolution of the mean curvature  $\log H$ ; because of  $\text{Ric} = -n$ , need to add  $\log z_0$ , i.e. we consider

$$\log H + \log z_0$$

- ▶ Boundary derivatives and then by maximum principle.

# Height estimate

**Theorem** Let  $M$  be a strictly convex hypersurface, then  $z^1 \geq \delta > 0$  for all  $M$  where the constant  $\delta$  depends only on  $\sup_M |A|$  and the distance of  $\partial M$  to some equator of  $\mathbb{S}^n(\rho_0)$ .

**Comments** The proof is long; and uses hyperbolic trigonometry and convex geometry.

## Lower bound of the mean curvature

**Proposition** Let  $M_t$  be the solution to the mean curvature flow with free boundary, if  $\partial M_{\bar{T}}$  is positive distance from the equator, then

$$\sup_{M_t, t \in [0, \bar{T})} \frac{1}{H} \leq c$$

where  $c$  depends only on  $M_0$  and the distance of  $\partial M_{\bar{T}}$  to the equator.

## Proof of the lower bound

1.  $f(q) = -\log(\Lambda - q)$ ,  $q = \lambda z^1 + z^0$ ,  
 $F(z) = -\log(\Lambda - (\lambda z^1 + z^0))$  with  $0 < \Lambda < \frac{1}{\cosh \rho_0}$  and  
 $\lambda < -1 - \frac{\cosh \rho_0}{\delta}$  where  $\delta$  is the number in the height estimate.

## Proof of the lower bound

1.  $f(q) = -\log(\Lambda - q)$ ,  $q = \lambda z^1 + z^0$ ,  
 $F(z) = -\log(\Lambda - (\lambda z^1 + z^0))$  with  $0 < \Lambda < \frac{1}{\cosh \rho_0}$  and  
 $\lambda < -1 - \frac{\cosh \rho_0}{\delta}$  where  $\delta$  is the number in the height estimate.
2. Let  $\xi = -\log H + F(z)$

# Proof of the lower bound

1.  $f(q) = -\log(\Lambda - q)$ ,  $q = \lambda z^1 + z^0$ ,  
 $F(z) = -\log(\Lambda - (\lambda z^1 + z^0))$  with  $0 < \Lambda < \frac{1}{\cosh \rho_0}$  and  
 $\lambda < -1 - \frac{\cosh \rho_0}{\delta}$  where  $\delta$  is the number in the height estimate.
2. Let  $\xi = -\log H + F(z)$ 
  - easy to check  $\nabla_{\eta} \xi < 0$



# Proof of the lower bound

1.  $f(q) = -\log(\Lambda - q)$ ,  $q = \lambda z^1 + z^0$ ,  
 $F(z) = -\log(\Lambda - (\lambda z^1 + z^0))$  with  $0 < \Lambda < \frac{1}{\cosh \rho_0}$  and  
 $\lambda < -1 - \frac{\cosh \rho_0}{\delta}$  where  $\delta$  is the number in the height estimate.
2. Let  $\xi = -\log H + F(z)$ 
  - ▶ easy to check  $\nabla_\eta \xi < 0$
  - ▶ Use evolution to get an inequality

$$-\frac{n\Lambda}{(n-q)H^2} + 2f' \frac{\partial q}{\partial \zeta^\alpha} \nu^\alpha \frac{1}{H} + |A|^2/H^2 \geq 0$$

at maximum point of  $\xi$ .

# Proof of the lower bound

1.  $f(q) = -\log(\Lambda - q)$ ,  $q = \lambda z^1 + z^0$ ,  
 $F(z) = -\log(\Lambda - (\lambda z^1 + z^0))$  with  $0 < \Lambda < \frac{1}{\cosh \rho_0}$  and  
 $\lambda < -1 - \frac{\cosh \rho_0}{\delta}$  where  $\delta$  is the number in the height estimate.
2. Let  $\xi = -\log H + F(z)$ 
  - ▶ easy to check  $\nabla_\eta \xi < 0$
  - ▶ Use evolution to get an inequality

$$-\frac{n\Lambda}{(n-q)H^2} + 2f' \frac{\partial q}{\partial \zeta^\alpha} \nu^\alpha \frac{1}{H} + |A|^2/H^2 \geq 0$$

at maximum point of  $\xi$ .

- ▶ the equality implies an bound on  $1/H$

## Maximal existence time is also maximal convexity time

**Theorem** The strict convexity is preserved up to  $T^*$ .

**Observation** If  $\bar{T} < T^*$ , then the smallest principal curvature  $\kappa_1$  reaches zero when the flow  $M_t$  accross the time  $\bar{T}$

# Proof

- ▶ If  $\partial M_{\bar{T}}$  is an equator, **easy**.

# Proof

- ▶ If  $\partial M_{\bar{T}}$  is an equator, **easy**.
- ▶ If not, then  $H \geq c^{-1}$  for all  $t \in [0, \bar{T})$  and  $z^1 \geq \delta > 0$ .

# Proof

- ▶ If  $\partial M_{\bar{T}}$  is an equator, **easy**.
- ▶ If not, then  $H \geq c^{-1}$  for all  $t \in [0, \bar{T})$  and  $z^1 \geq \delta > 0$ .
- ▶ Let  $\tilde{H} = \sum_{i=1}^n \frac{1}{\kappa_i}$  and define (this is the same with **Lambert-Scheuer 16**)

$$\phi = \log \tilde{H} - (n+1) \log z^1 - \alpha t.$$

# Proof

- ▶ If  $\partial M_{\bar{T}}$  is an equator, **easy**.
- ▶ If not, then  $H \geq c^{-1}$  for all  $t \in [0, \bar{T})$  and  $z^1 \geq \delta > 0$ .
- ▶ Let  $\tilde{H} = \sum_{i=1}^n \frac{1}{\kappa_i}$  and define (this is the same with **Lambert-Scheuer 16**)

$$\phi = \log \tilde{H} - (n+1) \log z^1 - \alpha t.$$

- ▶ Boundary derivative of  $\phi$  less than zero.

# Proof

- ▶ If  $\partial M_{\bar{T}}$  is an equator, **easy**.
- ▶ If not, then  $H \geq c^{-1}$  for all  $t \in [0, \bar{T})$  and  $z^1 \geq \delta > 0$ .
- ▶ Let  $\tilde{H} = \sum_{i=1}^n \frac{1}{\kappa_i}$  and define (this is the same with **Lambert-Scheuer 16**)

$$\phi = \log \tilde{H} - (n+1) \log z^1 - \alpha t.$$

- ▶ Boundary derivative of  $\phi$  less than zero.
- ▶ At maximum point,

$$0 \leq c + \frac{(n+1)^2}{H^2(z^1)^2} |\nabla z^1|^2 - \alpha.$$

However, this cannot always be true; just pick a big  $\alpha$ .



# Proof

- ▶ If  $\partial M_{\bar{T}}$  is an equator, **easy**.
- ▶ If not, then  $H \geq c^{-1}$  for all  $t \in [0, \bar{T})$  and  $z^1 \geq \delta > 0$ .
- ▶ Let  $\tilde{H} = \sum_{i=1}^n \frac{1}{\kappa_i}$  and define (this is the same with **Lambert-Scheuer 16**)

$$\phi = \log \tilde{H} - (n+1) \log z^1 - \alpha t.$$

- ▶ Boundary derivative of  $\phi$  less than zero.
- ▶ At maximum point,

$$0 \leq c + \frac{(n+1)^2}{H^2(z^1)^2} |\nabla z^1|^2 - \alpha.$$

However, this cannot always be true; just pick a big  $\alpha$ .

- ▶ Then for large  $\alpha$ , the supremum of  $\phi$  would be decreasing which implies a bound on  $\tilde{H}$  contradicting the definition of  $\bar{T}$

# Monotonicity of a Willmore type quantity: Theorem

- **Theorem (C. 22)** Suppose that  $M$  is strictly convex, then  $m_H(M)$  given by

$$m_H(M) = |M|^{\frac{2-n}{n}} \int_M (H^2 - n^2) + \Lambda |\partial M|$$

is monotone under IMCF with free boundary (assume for now strict convexity is preserved and that  $|M_t| \leq \lambda$ ).

# Monotonicity of a Willmore type quantity: Theorem

- **Theorem (C. 22)** Suppose that  $M$  is strictly convex, then  $m_H(M)$  given by

$$m_H(M) = |M|^{\frac{2-n}{n}} \int_M (H^2 - n^2) + \Lambda |\partial M|$$

is monotone under IMCF with free boundary (assume for now strict convexity is preserved and that  $|M_t| \leq \lambda$ ).

- $\lambda$  volume of  $n$ -dimensional hyperbolic geodesic ball of radius  $\rho_0$  (**note** dimension  $n$  here!)

# Monotonicity of a Willmore type quantity: Theorem

- **Theorem (C. 22)** Suppose that  $M$  is strictly convex, then  $m_H(M)$  given by

$$m_H(M) = |M|^{\frac{2-n}{n}} \int_M (H^2 - n^2) + \Lambda |\partial M|$$

is monotone under IMCF with free boundary (assume for now strict convexity is preserved and that  $|M_t| \leq \lambda$ ).

- $\lambda$  volume of  $n$ -dimensional hyperbolic geodesic ball of radius  $\rho_0$  (**note** dimension  $n$  here!)
- $\Lambda = 2 \coth \rho_0 \lambda^{(2-n)/n}$

## Lower bound of the Willmore type quantity

- **Theorem (C. 22)** Assuming that the flow preserves strict convexity, and converges to a totally geodesic disk, then

$$|M|^{\frac{2-n}{n}} \int_M (H^2 - n^2) + \Lambda |\partial M| \geq -n^2 \lambda^{\frac{2}{n}} + \Lambda \omega_{n-1} \sinh^{n-1} \rho_0.$$

## Lower bound of the Willmore type quantity

- **Theorem (C. 22)** Assuming that the flow preserves strict convexity, and converges to a totally geodesic disk, then

$$|M|^{\frac{2-n}{n}} \int_M (H^2 - n^2) + \Lambda |\partial M| \geq -n^2 \lambda^{\frac{2}{n}} + \Lambda \omega_{n-1} \sinh^{n-1} \rho_0.$$

- A comment: The definition of  $m_H(M)$  depends the supporting hypersurface of the boundary. If  $\partial M$  lies on the horosphere, or an equidistant hypersurface, then  $m_H$  is different.

## Proof of monotonicity

- Use the evolution equation

$$\begin{aligned} & \partial_t \int_{M_t} (H^2 - n^2) \sqrt{g} \\ &= \int_{M_t} (H^2 - n^2) \sqrt{g} + 2 \int_{M_t} H \left( -\Delta \frac{1}{H} - (-n + |A|^2) \frac{1}{H} \right) \sqrt{g}. \end{aligned}$$

# Proof of monotonicity

- Use the evolution equation

$$\begin{aligned} & \partial_t \int_{M_t} (H^2 - n^2) \sqrt{g} \\ &= \int_{M_t} (H^2 - n^2) \sqrt{g} + 2 \int_{M_t} H \left( -\Delta \frac{1}{H} - (-n + |A|^2) \frac{1}{H} \right) \sqrt{g}. \end{aligned}$$

- Integration on the Laplacian term

$$\begin{aligned} & - \int_{M_t} H \Delta \frac{1}{H} \\ &= -2 \int_{\partial M_t} H \langle \nabla \frac{1}{H}, \eta \rangle + 2 \int_{M_t} \langle \nabla H, \nabla \frac{1}{H} \rangle \\ &\leq -2 \int_{\partial M_t} H \langle \nabla \frac{1}{H}, \eta \rangle \end{aligned}$$



# Proof of monotonicity

- Use the evolution equation

$$\begin{aligned} & \partial_t \int_{M_t} (H^2 - n^2) \sqrt{g} \\ &= \int_{M_t} (H^2 - n^2) \sqrt{g} + 2 \int_{M_t} H \left( -\Delta \frac{1}{H} - (-n + |A|^2) \frac{1}{H} \right) \sqrt{g}. \end{aligned}$$

- Integration on the Laplacian term

$$\begin{aligned} & - \int_{M_t} H \Delta \frac{1}{H} \\ &= -2 \int_{\partial M_t} H \langle \nabla \frac{1}{H}, \eta \rangle + 2 \int_{M_t} \langle \nabla H, \nabla \frac{1}{H} \rangle \\ &\leq -2 \int_{\partial M_t} H \langle \nabla \frac{1}{H}, \eta \rangle \end{aligned}$$

- $\langle \nabla \frac{1}{H}, \eta \rangle = \coth \rho_0 / H$

# Proof of monotonicity

- Use the evolution equation

$$\begin{aligned} & \partial_t \int_{M_t} (H^2 - n^2) \sqrt{g} \\ &= \int_{M_t} (H^2 - n^2) \sqrt{g} + 2 \int_{M_t} H \left( -\Delta \frac{1}{H} - (-n + |A|^2) \frac{1}{H} \right) \sqrt{g}. \end{aligned}$$

- Integration on the Laplacian term

$$\begin{aligned} & - \int_{M_t} H \Delta \frac{1}{H} \\ &= -2 \int_{\partial M_t} H \langle \nabla \frac{1}{H}, \eta \rangle + 2 \int_{M_t} \langle \nabla H, \nabla \frac{1}{H} \rangle \\ &\leq -2 \int_{\partial M_t} H \langle \nabla \frac{1}{H}, \eta \rangle \end{aligned}$$

- $\langle \nabla \frac{1}{H}, \eta \rangle = \coth \rho_0 / H$
- $|A|^2 \geq \frac{H^2}{n}$  (by diagonalize  $A$ )

## Proof of monotonicity: adding a correction term

► So

$$\partial_t \int_{M_t} (H^2 - n^2) \leq \frac{n-2}{n} \int_{M_t} (H^2 - n^2) - 2 \coth \rho_0 |\partial M_t|$$

## Proof of monotonicity: adding a correction term

► So

$$\partial_t \int_{M_t} (H^2 - n^2) \leq \frac{n-2}{n} \int_{M_t} (H^2 - n^2) - 2 \coth \rho_0 |\partial M_t|$$

► Let  $q(t) = |M_t|^{\frac{2-n}{n}} \int_{M_t} (H^2 - n^2) + \Lambda |\partial M_t|$

## Change rate of $q(t)$



$$q'(t) \leq \Lambda \partial_t |\partial M_t| - 2 \coth \rho_0 |M|^{(2-n)/n} |\partial M_t|$$

## Change rate of $q(t)$



$$q'(t) \leq \Lambda \partial_t |\partial M_t| - 2 \coth \rho_0 |M|^{(2-n)/n} |\partial M_t|$$



$$\partial_t |\partial M_t| = \int_{\partial M_t} \frac{H_{\partial M, \partial B}}{H} \leq |\partial M_t|$$

## Change rate of $q(t)$



$$q'(t) \leq \Lambda \partial_t |\partial M_t| - 2 \coth \rho_0 |M|^{(2-n)/n} |\partial M_t|$$

▶  $\partial_t |\partial M_t| = \int_{\partial M_t} \frac{H_{\partial M, \partial B}}{H} \leq |\partial M_t|$

▶  $H_{M, B} = H_{\partial M, \partial B} + A_M(\eta, \eta)$

## Change rate of $q(t)$



$$q'(t) \leq \Lambda \partial_t |\partial M_t| - 2 \coth \rho_0 |M|^{(2-n)/n} |\partial M_t|$$

▶  $\partial_t |\partial M_t| = \int_{\partial M_t} \frac{H_{\partial M, \partial B}}{H} \leq |\partial M_t|$

▶  $H_{M, B} = H_{\partial M, \partial B} + A_M(\eta, \eta)$

▶  $|M| \leq \lambda$



## Change rate of $q(t)$



$$q'(t) \leq \Lambda \partial_t |\partial M_t| - 2 \coth \rho_0 |M|^{(2-n)/n} |\partial M_t|$$

▶  $\partial_t |\partial M_t| = \int_{\partial M_t} \frac{H_{\partial M, \partial B}}{H} \leq |\partial M_t|$

▶  $H_{M, B} = H_{\partial M, \partial B} + A_M(\eta, \eta)$

▶  $|M| \leq \lambda$

▶ So  $q' \leq 0$

Thank you

Thank you!