

Computational Many-Body Physics

apl. Prof. Dr. R. Bulla

SS 2022

Exercise Sheet No. 5

Please upload your solutions to ILIAS until **Thursday, June 23, 16:00**.

Exercise 1: Spin correlations of the one-dimensional Heisenberg model

(11 points)

Here we focus on the isotropic Heisenberg model in dimension $d = 1$ with open boundary conditions

$$H = -J \sum_{i=1}^{N-1} \vec{S}_i \cdot \vec{S}_{i+1} ,$$

with $J = -1$ (the antiferromagnetic case). We are interested in the spin correlations between sites 1 and n , $\chi_{1n} = \langle \vec{S}_1 \cdot \vec{S}_n \rangle$, for both zero and finite temperature.

- a) Calculate $\chi_{1n} = \langle \psi_g | \vec{S}_1 \cdot \vec{S}_n | \psi_g \rangle$ for the ground state $|\psi_g\rangle$ of a Heisenberg chain with $N = 10$ sites and $n = 1, \dots, 10$. Note that for an even number of sites N , the ground state of the antiferromagnetic Heisenberg chain is non-degenerate. (5 points)
- b) Calculate the temperature dependence of the spin correlation

$$\chi_{1n}(T) = \frac{1}{Z} \sum_l \langle l | \vec{S}_1 \cdot \vec{S}_n | l \rangle e^{-\beta E_l} ,$$

with $Z = \sum_l e^{-\beta E_l}$ the partition function, $\beta = 1/(k_B T)$ (k_B can be set to 1), and $|l\rangle$ the eigenstates of H with eigenenergies E_l , for temperatures $T = 0.5, 2$, and 10. (4 points)

- c) Show numerically that, in the limit $T \rightarrow 0$, the finite-temperature spin-correlation $\chi_{1n}(T)$ of part b) corresponds to the zero-temperature spin-correlation of part a). (2 points)

Exercise 2: Reduced density matrix and entanglement entropy

(5 points)

Consider a two-site system (with sites A and B) with a two-dimensional basis for each site: $\{|i\rangle\} = \{|\uparrow\rangle_A, |\downarrow\rangle_A\}$ for site A and $\{|j\rangle\}$ for site B accordingly. A given state $|\psi\rangle$ can be expressed in this basis as

$$|\psi\rangle = \sum_{i=1}^2 \sum_{j=1}^2 \psi_{ij} |i\rangle |j\rangle . \quad (1)$$

Here we want to calculate the reduced density matrices ρ for the following three states:

$$\begin{aligned} |\psi\rangle_1 &= |\uparrow\rangle_A |\downarrow\rangle_B , \\ |\psi\rangle_2 &= \frac{1}{\sqrt{2}} (|\uparrow\rangle_A |\downarrow\rangle_B - |\downarrow\rangle_A |\uparrow\rangle_B) , \\ |\psi\rangle_3 &= \frac{1}{2} (|\uparrow\rangle_A + |\downarrow\rangle_A) (|\uparrow\rangle_B + |\downarrow\rangle_B) . \end{aligned}$$

- a) Write the states $|\psi\rangle_i$ ($i = 1, 2, 3$) in the form given by eq. (1), i.e. determine the matrix $\bar{\psi}$ with matrix elements $(\bar{\psi})_{ij} = \psi_{ij}$. (1 point)

The reduced density operator $\hat{\rho}$ is defined as

$$\hat{\rho}_A = \text{Tr}_B (|\psi\rangle\langle\psi|) = \sum_{j=1}^2 \langle j|\psi\rangle\langle\psi|j\rangle ,$$

with the matrix elements $\rho_{ii'} = \langle i|\hat{\rho}|i'\rangle = \sum_j \psi_{ij}\psi_{i'j}$.

- b) Calculate the reduced density matrices (i.e. the matrix elements $\rho_{ii'}$) for the states $|\psi\rangle_i$ ($i = 1, 2, 3$). (3 points)

We can now proceed with calculating the entanglement entropy S_e :

$$S_e = -\text{Tr}_A [\hat{\rho}_A \ln \hat{\rho}_A] = -\sum_{\alpha} w_{\alpha} \ln w_{\alpha} ,$$

with w_{α} the eigenvalues of the reduced density matrix. The entanglement entropy is a measure of the entanglement between subsystems A and B of a quantum system; this can now be tested on the three states $|\psi\rangle_i$, $i = 1, 2, 3$, given above.

- c) Calculate the entanglement entropy S_e for the states $|\psi\rangle_i$. (1 point)