

Completely regular growth of solutions to linear differential equations with exponential polynomial coefficients

Xing-Yu Li

Abstract

Consider the linear differential equation

$$f^{(n)} + A_{n-1}f^{(n-1)} + \cdots + A_0f = 0,$$

where the coefficients are exponential polynomials. We prove that every finite-order transcendental entire solution has completely regular growth and obtain asymptotic formulas for $\log |f|$ on almost every ray. This answers a question of Heittokangas et al. [11, p. 33]. The proof combines logarithmic derivative estimates with asymptotic integration along rays. We express the angular densities of zeros in terms of the indicator. We also extend the result beyond exponential polynomial coefficients under a sectorial Laurent–Puisseux condition.

Keywords: Linear differential equations, exponential polynomials, entire solutions of finite order, completely regular growth, asymptotic integration.

2020 MSC: Primary 34M10; Secondary 30D15, 34M30.

1 Introduction

Complex oscillation and the growth of solutions of linear differential equations with entire coefficients have been studied extensively. Exponential polynomials play an important role in oscillation and zero distribution; see the survey by Heittokangas, Ishizaki, Tohge and Wen [11].

An *exponential polynomial* is an entire function of the form

$$G = P_1e^{Q_1} + \cdots + P_l e^{Q_l}, \quad (1.1)$$

where P_j, Q_j are polynomials and $l \in \mathbb{N}$. An *exponential sum* is an exponential polynomial of the form

$$G(z) = F_1(z)e^{\lambda_1 z} + \cdots + F_m(z)e^{\lambda_m z}, \quad (1.2)$$

where $m \in \mathbb{N}$, the F_j are polynomials, and $\lambda_1, \dots, \lambda_m \in \mathbb{C}$ are distinct constants. The λ_j are called *frequencies* (or *leading coefficients*), and the F_j are called *multipliers*.

For an entire function f of finite positive order $\rho(f)$, the Phragmén–Lindelöf indicator is

$$h_f(\theta) = \limsup_{r \rightarrow \infty} r^{-\rho(f)} \log |f(re^{i\theta})|, \quad 0 \leq \theta < 2\pi.$$

Throughout the paper, completely regular growth is taken with respect to the constant proximate order $\rho(r) \equiv \rho(f)$, and finite nonzero type is required. We say that f has *completely regular growth* (*c.r.g.*) if there are Euclidean discs $D(z_k, r_k)$ satisfying

$$\sum_{|z_k| \leq r} r_k = o(r) \quad (1.3)$$

such that

$$\log |f(re^{i\theta})| = (h_f(\theta) + o(1))r^{\rho(f)}, \quad re^{i\theta} \notin \bigcup_k D(z_k, r_k), \quad (1.4)$$

as $r \rightarrow \infty$, uniformly in θ ; see [12, pp. 139–140] or [17, pp. 6–8].

Every transcendental exponential polynomial is of *c.r.g.*; see [10, Lemma 1.3]. Petrenko [16, pp. 104–112] and Steinmetz [18] independently proved that transcendental entire solutions of linear differential equations with polynomial coefficients are of *c.r.g.*

When at least one entire coefficient is transcendental, a theorem of Frei [7] shows that there are at most $n - 1$ linearly independent solutions of finite order. Gol'dberg and Ostrovskii stated a classical problem in the book [9, p. 300]:

(Gol'dberg-Ostrovskii) *Suppose that f is a finite order entire transcendental solution of*

$$f^{(n)} + a_{n-1}f^{(n-1)} + \cdots + a_1f' + a_0f = 0,$$

whose coefficients are entire functions of c.r.g. Is f also of c.r.g.?

Several positive results for second-order equations are discussed in [10, Section 4]. In particular, they recall cases with coefficients polynomial in e^z for which finite order transcendental solutions are exponential sums and hence of *c.r.g.* Wen, Gundersen and Heittokangas [20] studied the dual structure of exponential polynomial coefficients and solutions of n -th order equations. Li, Wang and Wen [14] considered exponential sum coefficients with rational frequencies and constant multipliers. They obtained representations of finite order entire solutions as exponential sums, and existence criteria when the nonzero frequencies have the same sign. In this case the *c.r.g.* conclusion follows from the representation.

Bergweiler [3, Theorems 1.1 and 1.2] constructed a second-order equation with coefficients of *c.r.g.* having a finite-order entire solution which is not of *c.r.g.* This raises the question of when *c.r.g.* of the coefficients is inherited by finite-order solutions. In particular, Heittokangas et al. [11, p. 33, Problem 5] asked whether this holds when all coefficients are exponential polynomials:

If the coefficients $a_0(z), \dots, a_{n-1}(z)$ of

$$f^{(n)} + a_{n-1}f^{(n-1)} + \cdots + a_1f' + a_0f = 0, \quad (1.5)$$

are exponential polynomials, and if it possesses a transcendental entire solution f of finite order of growth, then is it true that f is of c.r.g.?

Using Wiman–Valiron theory, Droletz [6] proved that the possible finite orders of transcendental entire solutions to (1.5) form a finite set of positive rational numbers, and that each such solution has order equal to its lower order.

Our proof is based on asymptotic estimates along rays. We combine Wasow's sectorial asymptotic theory for rational systems with an integral-equation argument controlling

exponentially small perturbations on almost every ray, in the spirit of [4]. This yields an affirmative answer to Heittokangas et al.'s Problem 5. The ray estimates also give an alternative derivation of the order conclusions proved by Droletz.

Theorem 1.1 *Suppose that the coefficients $a_0, \dots, a_{n-1}$ of (1.5) are exponential polynomials. Every transcendental entire solution of finite order has positive rational order, finite nonzero type, and completely regular growth.*

Outline of the proof of Theorem 1.1. Let f be a transcendental entire solution of finite order. Absorb the constant terms in the exponents into their polynomial multipliers, and combine terms with the same exponent. The equation becomes

$$\sum_{\nu=1}^N e^{Q_\nu(z)} \sum_{k=0}^n a_{k,\nu}(z) f^{(k)}(z) = 0,$$

where the Q_ν are distinct polynomials with $Q_\nu(0) = 0$ and the $a_{k,\nu}$ are polynomials. Outside finitely many directions, one exponential factor dominates all the others exponentially. Gundersen's estimate gives polynomial bounds for $f^{(k)}/f$ on almost every complete ray tail. Dividing by the largest exponential factor and by the nonzero polynomial multiplying the highest derivative in its group gives

$$f^{(m)} + P_{m-1}f^{(m-1)} + \dots + P_0f = \epsilon_\theta(r)f, \quad z = re^{i\theta},$$

where the P_k are rational and $|\epsilon_\theta(r)| \leq Cr^K e^{-cr}$ with $c > 0$. The case $m = 0$ would give $\epsilon_\theta = 1$, so $m \geq 1$. Constants and starting radii here may depend on the fixed angle θ .

The equation with zero right-hand side has rational coefficients. Wasow's theorem gives a fundamental matrix for its first-order system on a narrow sector containing the ray, outside finitely many further directions. It has the form $Y_0 = Te^Q$, where Q is diagonal with entries that are polynomials in a fractional power of z , and T and T^{-1} have polynomial bounds. This transformation reduces the system to a diagonal system with an integrable perturbation on the fixed ray. Lemma 2.4 compares its solutions with the diagonal system. Together with the derivative bounds, it gives

$$\log |f(re^{i\theta})| = \Re q_\theta(re^{i\theta}) + O(\log r).$$

The functions q_θ come from a finite list determined by the polynomial coefficient equations.

Finite order and the logarithmic derivative estimate exclude any selected expression of degree greater than the order of f . A Phragmén–Lindelöf argument then gives positive rational order and finite nonzero type. Levin's theorem yields *c.r.g.* in the whole plane from the limits on the remaining dense set of rays.

Section 2 collects the growth and asymptotic tools. Section 3 treats exponential sums and general exponential-polynomial coefficients, and derives consequences for indicators and zeros. Section 4 contains the proofs. Section 5 considers entire coefficients beyond the exponential polynomial class. Finitely many rays divide a neighbourhood of infinity into sectors S_ν . On each sector, the coefficients, including $a_n \equiv 1$, are assumed to have Laurent–Puisseux asymptotic expansions of the form

$$\frac{a_k(z)}{W_\nu(z)} = \sum_{j=j_\nu}^J c_{k,\nu,j} z^{-j/p_\nu} + O(|z|^{-(J+1)/p_\nu}), \quad 0 \leq k \leq n,$$

for every integer $J \geq j_\nu$, uniformly as $z \rightarrow \infty$ on every closed subsector of S_ν . Here W_ν is nonvanishing and holomorphic on S_ν , $p_\nu \geq 1$ and j_ν are integers, and a branch of z^{1/p_ν} is fixed. We require $(c_{k,\nu,j})_{0 \leq k \leq n, j \geq j_\nu} \neq 0$ for each ν . Under this condition, Proposition 5.1 proves that every transcendental entire solution of finite order has *c.r.g.* Corollaries 5.2 and 5.3 apply this criterion to integral coefficients whose normalized expansions involve integer powers and, respectively, possibly rational fractional powers.

2 Basic concepts and lemmas

2.1 Completely regular growth on rays

A set covered by discs satisfying (1.3) is called a C_0 -set. The relative measure of a measurable set $E \subset (0, \infty)$ is

$$m^*(E) = \lim_{r \rightarrow \infty} \frac{\text{mes}(E \cap (0, r))}{r},$$

when this limit exists. In particular, a set of finite linear measure has zero relative measure.

We shall use the constant proximate order $\rho(r) \equiv \rho$. For this purpose we first need to know that the function has order $\rho > 0$ and finite nonzero type, that is,

$$0 < \limsup_{r \rightarrow \infty} \frac{\log M(r, f)}{r^\rho} < \infty, \quad M(r, f) = \max_{|z|=r} |f(z)|.$$

With respect to this same proximate order, f is of *c.r.g.* on the ray $\arg z = \theta$ if

$$\lim_{r \rightarrow \infty, r \notin E_\theta} \frac{\log |f(re^{i\theta})|}{r^\rho}$$

exists and is finite, for some set E_θ of zero relative measure. For a set of rays one requires a common exceptional set and uniform convergence.

Lemma 2.1 (Levin) *Let f be an entire function of order $\rho > 0$ and finite nonzero type. If f is of *c.r.g.* on rays whose arguments form an everywhere dense set, with respect to the same proximate order $\rho(r) \equiv \rho$, then f is of *c.r.g.* in the whole plane.*

This is the corollary to the theorem on sets of rays in [12, Chapter III, Section 1, pp. 140–142]. The type assumption permits the use of the constant proximate order; see also [12, p. 32].

2.2 Asymptotic comparison on rays

Considering the differential equation

$$f^{(n)} + a_{n-1}f^{(n-1)} + \cdots + a_1f' + a_0f = 0 \tag{2.1}$$

with polynomial coefficients, Petrenko [16, pp. 104–112] has shown that its transcendental entire solutions are of *c.r.g.* Steinmetz [18, Theorem 1] proved a more general result for the equation

$$f^{(n)} + R_{n-1}f^{(n-1)} + \cdots + R_1f' + R_0f = 0, \tag{2.2}$$

where $R_0, \dots, R_{n-1}$ are rational functions.

Lemma 2.2 (Steinmetz) *Let f be a non-rational meromorphic solution of (2.2), and let $\arg z = \theta$ be an arbitrary direction. There are $p \in \mathbb{N}$, $h > 0$, and a polynomial Q such that*

$$\log |f(z)| = \Re Q(z^{1/p}) + O(\log |z|) \quad (2.3)$$

as $z \rightarrow \infty$ in $\theta \leq \arg z \leq \theta + h$, possibly outside

1. *discs $|z - z_\mu| < |z_\mu|^{1-\varepsilon}$, where $\varepsilon > 0$ and the counting function of the sequence $\{z_\mu\}$ is $O(\log r)$;*
2. *a logarithmic semi-strip $0 \leq \arg z - \theta < \log^+ |z|/|z|^{1/p}$.*

The latter occurs only for finitely many directions.

These exceptional directions are called Stokes rays; see [18] and [19, Section 15].

Vector norms are maximum norms and matrix norms are the induced norms. A branch of each fractional power is fixed on the sector under consideration.

In asymptotic estimates on a fixed ray $\arg z = \theta$, the constants in the O -terms and the starting radius may depend on θ . These estimates are not asserted to be uniform in θ unless this is stated explicitly.

The next lemma follows from Wasow's theorem [19, Theorem 19.1, pp. 111–112] and the construction in its proof.

Lemma 2.3 *Let $A(z)$ be a rational m -by- m matrix. Near any fixed direction, and for sufficiently large $|z|$, the system*

$$Y' = A(z)Y$$

has a fundamental matrix on a sufficiently narrow sector of the form

$$Y_0(z) = T(z)e^{Q(z)}. \quad (2.4)$$

Here

$$Q(z) = \text{diag}(G_1(z^{1/p}), \dots, G_m(z^{1/p})),$$

where $p \in \mathbb{N}$ and the G_j are polynomials with $G_j(0) = 0$. A branch of $z^{1/p}$ is fixed on the sector. The matrix T is holomorphic and invertible. On every closed subsector there are constants $C, K, R > 0$ such that

$$\|T(z)\| + \|T(z)^{-1}\| \leq C|z|^K, \quad |z| \geq R. \quad (2.5)$$

Moreover,

$$T^{-1}AT - T^{-1}T' = Q'. \quad (2.6)$$

The integer p and the polynomials $G_1, \dots, G_m$ can be fixed independently of the sector, with the diagonal entries reordered if needed. If a branch of $z^{1/p}$ has been fixed, the other choices are $e^{2\pi i \ell/p} z^{1/p}$, where $\ell = 0, \dots, p-1$. Thus the possible diagonal entries belong to the finite family

$$G_j(e^{2\pi i \ell/p} z^{1/p}), \quad 1 \leq j \leq m, \quad 0 \leq \ell < p.$$

Proof. Write $A(z) = z^d B(z)$, where $d \geq 1$ is an integer and B is holomorphic at infinity. By [19, Theorem 19.1, pp. 111–112] and its construction in Section 19, an actual fundamental matrix is

$$Y_0(z) = H(z)z^G e^{Q(z)},$$

where $\mathbf{G}$ is constant and $\mathbf{H}$ has an asymptotic Laurent series in $z^{-1/p}$, with finitely many positive powers allowed. The matrix of these formal Laurent series is invertible. The construction uses matrices with invertible leading terms and diagonal matrices of rational powers of z , followed by the regular singular reduction; see [19, pp. 58–60, 95–98, 101–111]. Their products are invertible over the field of formal Laurent series in $z^{-1/p}$. Hence, on a closed subsector,

$$\det \mathbf{H}(z) = cz^{\ell/p}(1 + o(1)), \quad c \neq 0,$$

for an integer ℓ . The entries of $\mathbf{H}$ are bounded by powers of $|z|$, and the adjugate formula gives the same bound for $\mathbf{H}^{-1}$. The matrices $z^{\mathbf{G}}$ and $z^{-\mathbf{G}}$ also have polynomial bounds on a bounded argument interval: their Jordan blocks contain only powers of z multiplied by polynomials in $\log z$. Thus $\mathbf{T} = \mathbf{H}z^{\mathbf{G}}$ satisfies (2.5). Constant terms in the diagonal polynomials can be absorbed into $\mathbf{T}$, so these polynomials may be taken to have zero constant term. In Wasow's construction, the integer p and the polynomials G_j are obtained from the rational system before the sectorial fundamental matrices are constructed. They can therefore be kept fixed when the sector changes; the different choices of $z^{1/p}$ give the finite family displayed above. Differentiating (2.4) gives

$$\mathbf{T}' + \mathbf{T}\mathbf{Q}' = \mathbf{A}\mathbf{T},$$

which proves (2.6). □

Lemma 2.4 *Let $q_1, \dots, q_m$ be complex-valued C^1 functions on a half-line. Suppose each difference $q_i - q_j$ is either identically zero or has real part eventually strictly monotone and tending to $+\infty$ or $-\infty$. If $\mathbf{R}$ is continuous and $\int_{r_0}^{\infty} \|\mathbf{R}(r)\| dr < \infty$, then the system*

$$\mathbf{Z}' = (\text{diag}(q'_1, \dots, q'_m) + \mathbf{R})\mathbf{Z}$$

has a fundamental matrix

$$\mathbf{Z}(r) = (\mathbf{I} + o(1)) \text{diag}(e^{q_1(r)}, \dots, e^{q_m(r)}). \quad (2.7)$$

Proof. Fix a column j , and write it as $e^{q_j} \mathbf{u}$. Put $\Delta_i = q_i - q_j$. Choose r_0 large enough that all the relevant real differences are monotone and

$$a := \int_{r_0}^{\infty} \|\mathbf{R}(s)\| ds < \frac{1}{2}.$$

On the Banach space of bounded continuous vectors on $[r_0, \infty)$, with the supremum norm, consider

$$u_i(r) = \begin{cases} \int_{r_0}^r e^{\Delta_i(r) - \Delta_i(s)} (\mathbf{R}(s) \mathbf{u}(s))_i ds, & \Re \Delta_i \rightarrow -\infty, \\ -\int_r^{\infty} e^{\Delta_i(r) - \Delta_i(s)} (\mathbf{R}(s) \mathbf{u}(s))_i ds, & \Re \Delta_i \rightarrow +\infty, \\ \delta_{ij} - \int_r^{\infty} (\mathbf{R}(s) \mathbf{u}(s))_i ds, & \Delta_i \equiv 0. \end{cases} \quad (2.8)$$

Every exponential kernel has modulus at most 1 on its integration interval. The integrals define bounded continuous functions, and the norm of the linear part of this map is at most a . It is therefore a contraction and has a unique fixed point.

The integrals to infinity tend to zero, being bounded by $\|\mathbf{u}\|_{\infty} \int_r^{\infty} \|\mathbf{R}(s)\| ds$. For a forward integral, fix $S > r_0$ and split it into $[r_0, S]$ and $[S, r]$. On the first interval the

kernel tends uniformly to zero as $r \rightarrow \infty$, because $\Re \Delta_i(r) \rightarrow -\infty$. The second integral is bounded by $\|\mathbf{u}\|_\infty \int_S^\infty \|\mathbf{R}(s)\| ds$. First let $r \rightarrow \infty$, and then let $S \rightarrow \infty$. It follows that $\mathbf{u}(r) \rightarrow \mathbf{e}_j$.

Differentiation of the forward integrals is immediate. For a backward integral, write it as $-e^{\Delta_i(r)} \int_r^\infty e^{-\Delta_i(s)} (\mathbf{R}(s)\mathbf{u}(s))_i ds$. The integral converges absolutely, so the fundamental theorem of calculus gives the same derivative. Thus each fixed point satisfies

$$u'_i = \Delta'_i u_i + (\mathbf{R}\mathbf{u})_i.$$

The columns $e^{q_j} \mathbf{u}$ solve the differential system. Their normalized matrix tends to $\mathbf{I}$, so it is invertible for large r . This proves (2.7). $\square$

Lemma 2.5 *Let*

$$f^{(m)} + P_{m-1}(z)f^{(m-1)} + \cdots + P_0(z)f = \epsilon_\theta(r)f, \quad z = re^{i\theta},$$

hold on a ray tail, where $m \geq 1$, the P_k are rational, and ϵ_θ is continuous with $|\epsilon_\theta(r)| \leq Cr^K e^{-cr}$ for some $c > 0$. Suppose f is holomorphic near the ray, is eventually nonzero there, and

$$|f^{(k)}(re^{i\theta})/f(re^{i\theta})| \leq Cr^K, \quad 1 \leq k < m.$$

There is a finite set of directions, determined by the rational equation, outside which

$$\log |f(re^{i\theta})| = \Re G_\theta((re^{i\theta})^{1/p}) + O(\log r). \quad (2.9)$$

The polynomial G_θ belongs to a finite list, with the branches fixed on finitely many argument intervals. A common denominator p may be chosen for this finite list.

Proof. Put $\mathbf{x} = (f, f', \dots, f^{(m-1)})^\top$, and let $\mathbf{A}$ be the companion matrix of the rational equation with zero right-hand side. Its last row is $(-P_0, \dots, -P_{m-1})$ and its superdiagonal entries are 1. Let $\mathbf{E} = \mathbf{e}_m \mathbf{e}_1^\top$. Solving the scalar equation for $f^{(m)}$ and using $dz/dr = e^{i\theta}$ gives the exact system

$$\frac{d\mathbf{x}}{dr} = e^{i\theta}(\mathbf{A}(re^{i\theta}) + \epsilon_\theta(r)\mathbf{E})\mathbf{x}. \quad (2.10)$$

Apply Lemma 2.3 to the unperturbed rational system $\mathbf{Y}' = \mathbf{A}\mathbf{Y}$ on a whole narrow sector containing this ray. Set $\mathbf{x}(re^{i\theta}) = \mathbf{T}(re^{i\theta})\mathbf{z}(r)$. The chain rule and (2.6) give

$$\frac{d\mathbf{z}}{dr} = \left[\frac{d}{dr} \mathbf{Q}(re^{i\theta}) + \mathbf{R}_\theta(r) \right] \mathbf{z}, \quad \mathbf{R}_\theta = e^{i\theta} \mathbf{T}^{-1} \epsilon_\theta \mathbf{E} \mathbf{T}. \quad (2.11)$$

By (2.5), $\mathbf{R}_\theta$ is integrable.

Write the diagonal entries of $\mathbf{Q}$ as $q_1(z), \dots, q_m(z)$. Exclude the directions where the leading term of a nonzero q_i or a nonzero $q_i - q_j$ has zero real part. There are only finitely many such directions for all choices of the functions q_i given by Lemma 2.3. On each remaining ray, $\Re(q_i - q_j)$ is either identically zero with $q_i = q_j$, or is eventually strictly monotone and tends to $+\infty$ or $-\infty$. Lemma 2.4 gives, for the particular solution $\mathbf{x}$,

$$\mathbf{x}(re^{i\theta}) = \mathbf{T}(re^{i\theta})(\mathbf{I} + o(1)) \text{diag}(e^{q_1(re^{i\theta})}, \dots, e^{q_m(re^{i\theta})}) \mathbf{c},$$

where $\mathbf{c} \neq 0$. The polynomial bounds for $\mathbf{T}$ and its inverse imply

$$\log \|\mathbf{x}(re^{i\theta})\| = \max_{c_i \neq 0} \Re q_i(re^{i\theta}) + O(\log r).$$

The hypotheses on the derivatives give $|f| \leq \|x\| \leq Cr^K|f|$, so the same formula holds for $\log|f|$. The real parts $\Re q_i(re^{i\theta})$ have a fixed ordering for large r . Thus the maximum is one fixed $q_i = G_\theta(z^{1/p})$ on this ray tail, which proves (2.9). $\square$

Lemma 2.6 *Let f be a transcendental entire function of finite order ρ . Suppose there is a dense set $\mathbb{M}$ of directions and a finite list of polynomials in fractional powers of z , on fixed argument intervals, such that for every $\theta \in \mathbb{M}$,*

$$\log |f(re^{i\theta})| = \Re q_\theta(re^{i\theta}) + O(\log r), \quad (2.12)$$

with q_θ from this list. Suppose also that for each $\delta > 0$,

$$|\log |f(re^{i\theta})|| \leq C_{\theta,\delta} r^{\rho+\delta} \quad (2.13)$$

on every such ray tail. If the highest term of a nonzero selected q_θ is az^d , suppose also that $\Re(ae^{id\theta}) \neq 0$. Then ρ is positive and rational, f has finite nonzero type, and f is of c.r.g.

Proof. For $q(z) = G(z^{1/p}) \neq 0$, define its degree to be $\deg G/p$; the zero function is assigned degree 0. If a selected q_θ had degree $d > \rho$, then (2.12) would contradict (2.13), by choosing $0 < \delta < d - \rho$. Thus every selected degree is at most ρ . Let σ be the largest degree actually selected on $\mathbb{M}$. This maximum exists because the list is finite, and $\sigma \leq \rho$.

If $0 < \tau \leq \rho$ and every ray in a dense set satisfies $|f(re^{i\theta})| \leq \exp(C_\theta r^\tau)$ eventually, then

$$\log M(r, f) = O(r^\tau). \quad (2.14)$$

To see this, choose finitely many of these rays around the circle, with each consecutive angular gap $\beta < \pi/(\rho + 1)$. Choose one inner radius beyond the finitely many starting radii. In each exterior sector between adjacent rays fix a branch of z^τ , and let θ_0 be its middle argument. Then

$$\Re(e^{-i\tau\theta_0} z^\tau) = r^\tau \cos(\tau(\arg z - \theta_0)) \geq c_0 r^\tau$$

on the closed sector, with $c_0 > 0$. For sufficiently large K , the function

$$f(z) \exp(-K e^{-i\tau\theta_0} z^\tau)$$

is bounded on both boundary rays and on the fixed inner circular arc. Its growth order is at most ρ , which is smaller than π/β . The sectorial Phragmén–Lindelöf theorem makes it bounded in the sector. The finitely many sectors give (2.14). If instead the bounds on each ray are polynomial, choose this same finite grid and one integer N larger than their exponents. Apply the argument to $f(z)/z^N$ on the exterior sectors. Then $M(r, f) = O(r^N)$, and Cauchy's estimates show that f is a polynomial.

If $\sigma = 0$, (2.12) gives polynomial bounds on all the rays in $\mathbb{M}$, a contradiction. Hence $\sigma > 0$. The same formula gives $|f(re^{i\theta})| \leq \exp(C_\theta r^\sigma)$. By (2.14), $\rho \leq \sigma$ and f has finite type at order σ . Consequently $\rho = \sigma$ is positive and rational.

This type is nonzero. Otherwise no selected q_θ of degree ρ could have a positive real leading coefficient, since one such ray already gives $\log M(r, f) \geq cr^\rho$ for large r . All selected q_θ of degree ρ would therefore have leading terms with negative real parts on their rays, and f would be bounded there. All other selected degrees belong to a finite list below ρ . If its largest degree is positive, (2.14) would give an order strictly below ρ ; if there is

no positive degree, the polynomial version would make f a polynomial. Both conclusions are impossible.

Finally, division of (2.12) by r^ρ gives a finite limit on every ray in $\mathbb{M}$. The constant ρ is now a proximate order of f . Lemma 2.1 proves *c.r.g.* $\square$

3 Coefficient equations and ray asymptotics

3.1 Exponential sum coefficients

We first consider the linear differential equation

$$f^{(n)}(z) + A_{n-1}(z)f^{(n-1)}(z) + \cdots + A_1(z)f'(z) + A_0(z)f(z) = 0 \quad (3.1)$$

with exponential sum coefficients $A_i(z)$ for $i = 0, 1, \dots, n-1$ of the form (1.2). The equation (3.1) can be transformed into a normal form

$$\sum_{j=0}^{N_0} e^{\lambda_j z} (a_{n,j}(z)f^{(n)}(z) + a_{n-1,j}(z)f^{(n-1)}(z) + \cdots + a_{0,j}(z)f(z)) = 0, \quad (3.2)$$

The notation used below is as follows.

1. *Frequency groups.* The $a_{k,j}$ are polynomials, and the distinct frequencies λ_j include 0. The term $f^{(n)}$ belongs to this zero-frequency group: $a_{n,j} = 1$ when $\lambda_j = 0$, and $a_{n,j} = 0$ otherwise. Groups in which all coefficients vanish are omitted.

2. *Coefficient equations.* For each frequency λ_j , we call

$$a_{n,j}(z)f^{(n)} + a_{n-1,j}(z)f^{(n-1)} + \cdots + a_{0,j}(z)f = 0$$

the coefficient differential equation at λ_j . The given solution need not satisfy these equations separately.

3. *The convex hull.* Put $W = \{\bar{\lambda}_j : 0 \leq j \leq N_0\}$. Write the vertices of its convex hull as

$$\tilde{W} = \{\bar{\omega}_j : 0 \leq j \leq N'_0\},$$

in anticlockwise order. There are $N'_0 + 1$ vertices, with indices read cyclically. Number the frequency groups so that $\lambda_j = \omega_j$ for $0 \leq j \leq N'_0$. We use conjugated frequencies because $\Re(\lambda z)$ is the Euclidean scalar product of $\bar{\lambda}$ and z .

4. *Critical rays.* Let s_j join $\bar{\omega}_j$ to $\bar{\omega}_{j+1}$. The outward normal to this side has argument η_j . The ray $\arg z = \eta_j$ is called a critical ray.
5. *Vertex sectors and boundary regions.* For sufficiently small $\varepsilon > 0$, put

$$\begin{aligned} \bar{S}_j &= \{z \neq 0 : \eta_{j-1} \leq \arg z \leq \eta_j\}, \\ S_j(\varepsilon) &= \{z \neq 0 : \eta_{j-1} + \varepsilon \leq \arg z \leq \eta_j - \varepsilon\}, \\ T_j(\varepsilon) &= \{z \neq 0 : |\arg z - \eta_j| < \varepsilon\}. \end{aligned}$$

Arguments are chosen continuously across the sector in question. The frequency ω_j is dominant in the interior of $\bar{S}_j$. The set $S_j(\varepsilon)$ stays away from its boundary rays, while $T_j(\varepsilon)$ is a small angular neighbourhood of a critical ray.




Figure 1: The conjugate-frequency hull, its critical rays, and a vertex sector.

Figure 1 shows this construction. If the hull is a segment, its two normal rays bound the two vertex half-planes. If it is a single point, there are no critical rays and the coefficient equation is the whole equation.

Lemma 3.1 *If $\bar{\lambda} \in W$, then for every $z \in S_j(\varepsilon)$,*

$$\Re((\omega_j - \lambda)z) \geq |z| |\omega_j - \lambda| \sin \varepsilon.$$

In particular, for a fixed interior ray of a vertex sector and $\lambda \neq \omega_j$, the right-hand side is at least $c_\theta |z|$ for some $c_\theta > 0$.

See also Dickson [5].

Our conclusions are as follows.

Theorem 3.2 *Every finite order transcendental entire solution f of (3.1) has positive rational order, finite nonzero type, and c.r.g. There is a set $\mathcal{E}_f \subset [0, 2\pi)$ of measure zero such that, for every $\theta \notin \mathcal{E}_f$ in a vertex sector at $\bar{\omega}_j$,*

$$\log |f(re^{i\theta})| = \Re G_{j\theta}((re^{i\theta})^{1/p_j}) + O(\log r), \quad r \rightarrow \infty,$$

on the complete ray tail. Here $p_j \in \mathbb{N}$ and $G_{j\theta}$ is a polynomial from a finite list determined by the coefficient differential equation at the frequency ω_j ; its branch is fixed on the argument interval. The polynomial, the error constant and the starting radius may depend on θ .

Remark. The error estimate above is on each fixed ray. We do not claim this $O(\log r)$ error uniformly in a sector. Completely regular growth still gives the uniform $o(r^{\rho(f)})$ estimate (1.4) outside a C_0 -set.

Example 3.3 The function $f(z) = e^{-4z/3}(1 - 7e^z)$ from [14, Example 3.6] solves

$$\left(f''' - \frac{4}{3}f' + \frac{16}{27}f\right) + e^z(3f'' - 2f' - f) = 0.$$

The two coefficient expressions, divided by f , are $-7e^z/(1 - 7e^z)$ and $7/(1 - 7e^z)$. They are exponentially small in closed subsectors of the left and right half-planes, respectively. Thus the order of the dominant coefficient equation changes between sectors: it is three in the left half-plane and two in the right half-plane.

3.2 Exponential polynomial coefficients

We now consider

$$f^{(n)} + G_{n-1}(z)f^{(n-1)} + \cdots + G_1(z)f' + G_0(z)f = 0, \quad (3.3)$$

where

$$G_m(z) = \sum_{t=1}^{l_m} P_{m,t}(z)e^{Q_{m,t}(z)}, \quad 0 \leq m < n, \quad (3.4)$$

and $P_{m,t}, Q_{m,t}$ are polynomials. Zero coefficients are allowed. We first group the terms with the same exponential factor. We then compare the coefficients in their exponents, starting with the highest degree.

1. *Grouping equal exponents.* Absorb the constant term of each exponent into its polynomial multiplier, combine equal exponents, and include the exponent 0 from $f^{(n)}$. After omitting zero groups, write the equation as

$$\sum_{\nu=1}^N e^{Q_\nu(z)} \left\{ \sum_{m=0}^n a_{m,\nu}(z)f^{(m)} \right\} = 0. \quad (3.5)$$

Here $Q_\nu(0) = 0$, the Q_ν are distinct, and the $a_{m,\nu}$ are polynomials. Each coefficient group is nonzero.

2. *Grouping by degree.* If all Q_ν have degree at most one, Theorem 3.2 applies. Otherwise put $s = \max_\nu \deg Q_\nu \geq 2$ and write

$$Q_\nu(z) = q_{\nu,s}z^s + q_{\nu,s-1}z^{s-1} + \cdots + q_{\nu,1}z.$$

Thus $q_{\nu,d}$ means only the coefficient of z^d in Q_ν ; missing coefficients are set equal to zero. First group the terms with the same $q_{\nu,s}$. Within each group, group the terms with the same $q_{\nu,s-1}$, and continue down to $q_{\nu,1}$. At each step, the coefficients of all higher powers are held fixed. Figure 2 illustrates this finite procedure. Only one branch is expanded in the figure; each leaf represents one of the polynomials Q_ν .

3. *Fundamental coefficient equations.* After the coefficient of z has been fixed, each group contains a single exponent Q_ν . We call

$$\sum_{m=0}^n a_{m,\nu}(z)f^{(m)} = 0 \quad (3.6)$$

its fundamental coefficient differential equation. These equations have polynomial coefficients.




Figure 2: Successive grouping of the exponents.

The angular construction uses the same notation at each step.

1. *Fix the higher-degree coefficients.* At the step concerning z^d , consider one of the groups already formed (the full set of exponents at the first step). Apply the convex-hull construction of Section 3.1 to the conjugates of the remaining values $q_{\nu,d}$ in this group. For this fixed step, use the same local notation $W, \bar{\omega}_j, \eta_j, S_j(\varepsilon), T_j(\varepsilon)$ as before, now in the variable $w = z^d$.
2. *Return to the z -plane.* A critical ray $\arg w = \eta_j$ has the inverse images

$$\arg z = \frac{\eta_j + 2\pi\ell}{d}, \quad \ell = 0, \dots, d-1.$$

In particular, $S_j(\varepsilon)$ has d inverse-image sectors:

$$\frac{\eta_{j-1} + \varepsilon + 2\pi\ell}{d} \leq \arg z \leq \frac{\eta_j - \varepsilon + 2\pi\ell}{d}, \quad \ell = 0, \dots, d-1.$$

Their angular widths are divided by d , and consecutive sectors are rotated by $2\pi/d$. A single remaining value introduces no further sector restriction. For a segment hull, use the two half-planes as in Section 3.1.

3. *Continue within the selected sectors.* At each subsequent step, intersect these inverse-image sectors with the sector already selected, and retain the nonempty intersections. On an interior ray, the procedure selects the same Q_ν as direct comparison of these polynomials. The proof below makes this comparison directly.

Figure 3 illustrates these steps for $d = 3$. Under $w = z^3$, the sector $30^\circ \leq \arg w \leq 120^\circ$ has three inverse-image sectors, with argument intervals $[10^\circ, 40^\circ]$, $[130^\circ, 160^\circ]$, and $[250^\circ, 280^\circ]$. If an earlier step has selected $20^\circ \leq \arg z \leq 60^\circ$ (dashed boundaries in the last panel), only the shaded sector $20^\circ \leq \arg z \leq 40^\circ$ remains. The arcs indicate angular ranges only.

Lemma 3.4 *For finitely many polynomials $Q_1, \dots, Q_N$ with nonconstant pairwise differences, there is a finite set of directions such that, outside this set, a unique Q_ν is dominant for large r on $z = re^{i\theta}$. For every other Q_μ there is $c_\theta > 0$ such that*

$$\Re(Q_\mu(re^{i\theta}) - Q_\nu(re^{i\theta})) \leq -c_\theta r \quad (3.7)$$

for all sufficiently large r . The dominant index is constant on every component of the remaining angular set, and the estimate is uniform on closed subintervals.




Figure 3: Inverse-image sectors and their intersection.

Proof. For each difference write its highest nonzero term as cz^d , where $d \geq 1$. Exclude the finitely many directions where $\Re(ce^{id\theta}) = 0$. On every remaining angular interval the real parts of all pairwise differences have fixed signs for large r . They therefore give a strict ordering and a unique largest exponent. On a closed subinterval the real leading coefficients are bounded away from zero. For a difference with negative real leading term, its lower terms are $O(r^{d-1})$, so its real part is at most $-cr^d \leq -cr$ for large r . Taking the minimum over the finite set of differences proves the assertion. $\square$

Now a similar conclusion to Theorem 3.2 follows.

Theorem 3.5 *Every finite order transcendental entire solution f of (3.3) has positive rational order, finite nonzero type, and c.r.g. There is a set $\mathcal{E}_f \subset [0, 2\pi)$ of measure zero such that, for every $\theta \notin \mathcal{E}_f$,*

$$\log |f(re^{i\theta})| = \Re G_\theta((re^{i\theta})^{1/N_\theta}) + O(\log r), \quad r \rightarrow \infty, \quad (3.8)$$

on the complete ray tail, where $N_\theta \in \mathbb{N}$ and G_θ is a polynomial. The functions $G_\theta(z^{1/N_\theta})$ belong to a finite list determined by the fundamental coefficient differential equations. The branch is fixed on its argument interval. The polynomial, the error constant and the starting radius may depend on θ .

Remark. The conclusion of c.r.g. is the uniform statement (1.4) outside a C_0 -set. The error $O(\log r)$ in (3.8) is on fixed rays. Corollary 3.6 describes the highest-degree terms that determine the indicator. It does not describe changes in the lower-degree terms of $G_\theta(z^{1/N_\theta})$.

Remark. The same growth conclusion holds for every finite order transcendental entire solution of

$$A_n f^{(n)} + A_{n-1} f^{(n-1)} + \cdots + A_0 f = 0, \quad (3.9)$$

where $A_0, \dots, A_n$ are exponential polynomials and $A_n \neq 0$. Indeed, group the terms as in (3.5) and, on each remaining ray, divide by the largest exponential factor and by the polynomial multiplying the highest derivative in its group. The latter polynomial has only finitely many zeros, and the other groups give the same exponentially small right-hand side as in (4.8). A group of order zero would give $1 = O(r^K e^{-cr})$, which is impossible. Lemmas 2.5 and 2.6 therefore apply without change. Droletz's results on the possible orders already include these equations [6].

3.3 Indicators and zeros

The rays in the following statement are directions where the formula for the indicator may change.

Corollary 3.6 *Fix an equation (3.9), where $A_0, \dots, A_n$ are exponential polynomials and $A_n \not\equiv 0$.*

1. *Only finitely many pairs $(\rho(f), h_f)$ occur among its transcendental entire solutions of finite order. There are finitely many rays $\arg z = \vartheta_j$, $1 \leq j \leq s$, depending only on the equation, such that each such solution of order ρ has*

$$h_f(\theta) = \Re(ce^{i\rho\theta})$$

between consecutive rays. The constant c depends on the solution and the interval; it may be zero.

2. *Outside any fixed angular neighbourhoods of these rays, the number of zeros in $|z| \leq r$, counted with multiplicities, is $o(r^\rho)$. In particular, this estimate holds in every closed angular sector separated from these rays.*

The angular densities are given more precisely as follows. Let $n_f(r; \alpha, \beta)$ count, with multiplicities, the zeros in $0 < |z| \leq r$, $\alpha < \arg z < \beta$. Represent the angles by $0 \leq \vartheta_1 < \dots < \vartheta_s < 2\pi$. If $0 \leq \alpha < \beta \leq 2\pi$ and neither boundary ray is among the listed rays, then

$$n_f(r; \alpha, \beta) = \frac{r^\rho}{2\pi\rho} \sum_{\alpha < \vartheta_j < \beta} (h'_f(\vartheta_j + 0) - h'_f(\vartheta_j - 0)) + o(r^\rho). \quad (3.10)$$

Every difference in the sum is nonnegative. For a particular solution, some of these differences may be zero. The estimate in the corollary concerns angular density; it does not assert that all zeros lie on the listed rays.

4 Proofs

4.1 Proof of Theorem 3.2

Proof. Let f be a transcendental entire solution of finite order ρ . By Gundersen [8, Corollary 1, p. 89], applied to $\{(k, 0) : 1 \leq k \leq n\}$, for each $\delta > 0$ there is an angular exceptional set of measure zero such that

$$\left| \frac{f^{(k)}(re^{i\theta})}{f(re^{i\theta})} \right| \leq r^{k(\rho-1+\delta)}, \quad 1 \leq k \leq n, \quad (4.1)$$

on every remaining ray, for all sufficiently large r . Apply this with $\delta = 1/\ell$, $\ell \in \mathbb{N}$, and take the countable union of the angular exceptional sets. We obtain a single null set $\mathcal{E}_G$ for which (4.1) holds for every $\delta > 0$. The starting radius may depend on θ and δ .

Each such ray is eventually zero-free. Indeed, if a zero of multiplicity s lay on its tail at $z_0 = r_0 e^{i\theta}$, then $f'(z)/f(z) = s/(z - z_0) + O(1)$ near z_0 . Approaching r_0 on this ray contradicts (4.1) for $k = 1$. On the zero-free tail,

$$\left| \frac{d}{dr} \log |f(re^{i\theta})| \right| \leq \left| \frac{f'(re^{i\theta})}{f(re^{i\theta})} \right| \leq r^{\rho-1+\delta}.$$

Integration from a fixed point gives the two-sided bound

$$|\log |f(re^{i\theta})|| \leq C_{\theta,\delta} r^{\rho+\delta}. \quad (4.2)$$

Fix a direction outside $\mathcal{E}_G$ and the critical rays of the frequency hull. Suppose its dominant frequency is ω_j . Put

$$m_j = \max\{k : a_{k,j} \neq 0\}, \quad b_j = a_{m_j,j}, \quad P_{k,j} = a_{k,j}/b_j \quad (0 \leq k < m_j).$$

The polynomial b_j has no zeros beyond a fixed radius. Dividing the normal form (3.2) by $e^{\omega_j z} b_j(z) f(z)$ gives, on this ray tail, the exact identity

$$\begin{aligned} \epsilon_{j,\theta}(r) &:= \sum_{k=0}^{m_j} \frac{a_{k,j}(z)}{b_j(z)} \frac{f^{(k)}(z)}{f(z)} \\ &= - \sum_{\lambda_\ell \neq \omega_j} e^{(\lambda_\ell - \omega_j)z} \sum_{k=0}^n \frac{a_{k,\ell}(z)}{b_j(z)} \frac{f^{(k)}(z)}{f(z)}, \quad z = re^{i\theta}. \end{aligned} \quad (4.3)$$

The rational multipliers are bounded by powers of r . By Lemma 3.1 and (4.1),

$$|\epsilon_{j,\theta}(r)| \leq C_\theta r^{K_\theta} e^{-c_\theta r}, \quad c_\theta > 0. \quad (4.4)$$

If there is only one frequency group, the right-hand side of (4.3) is zero, and the same bound is immediate. The case $m_j = 0$ is impossible, since its left-hand side would be 1. Thus $1 \leq m_j \leq n$.

We have proved that the given solution satisfies

$$f^{(m_j)} + P_{m_j-1,j} f^{(m_j-1)} + \dots + P_{0,j} f = \epsilon_{j,\theta}(r) f \quad (4.5)$$

on this ray. In general m_j is smaller than n . The function $\epsilon_{j,\theta}$ is continuous on this ray tail.

Apply Lemma 2.5 to (4.5). The rational coefficient equation is $f^{(m_j)} + \sum_{k < m_j} P_{k,j} f^{(k)} = 0$. There are finitely many such equations, and Lemma 2.5 excludes only finitely many additional directions. Together with $\mathcal{E}_G$ and the critical rays, they form a set $\mathcal{E}_f$ of measure zero. For every $\theta \notin \mathcal{E}_f$, we obtain

$$\log |f(re^{i\theta})| = \Re G_{j\theta}((re^{i\theta})^{1/p_j}) + O(\log r). \quad (4.6)$$

The functions $G_{j\theta}(z^{1/p_j})$ come from a finite list. Their nonzero leading terms have nonzero real parts on these rays, by the choice of the additional excluded directions. The set of remaining directions is dense. Equations (4.6) and (4.2) meet every hypothesis of Lemma 2.6. It follows that f has positive rational order, finite nonzero type, and *c.r.g.* This proves the theorem. $\square$

4.2 Proof of Theorem 3.5

Proof. Let f be a transcendental entire solution of finite order ρ . Use the same angular estimate (4.1), with a single null set $\mathcal{E}_G$, and hence the same zero-free ray tails and the two-sided estimate (4.2). Write the equation in the collected form (3.5). By Lemma 3.4, outside a finite set of directions one Q_ν has strictly largest real part along each ray tail.

This is the leaf selected by the successive coefficient equations in Section 3.2. Indeed, at each step only exponents with the same higher-degree coefficients are retained. If $Q_\nu - Q_\mu$ has highest term cz^d , its real part on the chosen ray has the sign of $\Re(ce^{id\theta})$ for large r . Thus the successive comparisons select the same exponent as the direct comparison in Lemma 3.4.

Fix such a direction outside $\mathcal{E}_G$, and put

$$m_\nu = \max\{k : a_{k,\nu} \neq 0\}, \quad b_\nu = a_{m_\nu,\nu}, \quad P_{k,\nu} = a_{k,\nu}/b_\nu \quad (0 \leq k < m_\nu).$$

The polynomial b_ν is nonzero and has only finitely many zeros. Increase the starting radius so that both b_ν and f are nonzero on the ray. Division of (3.5) by $e^{Q_\nu(z)}b_\nu(z)f(z)$ then gives

$$\begin{aligned} \epsilon_{\nu,\theta}(r) &:= \sum_{k=0}^{m_\nu} \frac{a_{k,\nu}(z)}{b_\nu(z)} \frac{f^{(k)}(z)}{f(z)} \\ &= - \sum_{\mu \neq \nu} e^{Q_\mu(z) - Q_\nu(z)} \sum_{k=0}^n \frac{a_{k,\mu}(z)}{b_\nu(z)} \frac{f^{(k)}(z)}{f(z)}, \quad z = re^{i\theta}. \end{aligned} \tag{4.7}$$

The function $\epsilon_{\nu,\theta}$ is continuous on this ray tail. Each rational multiplier $a_{k,\mu}/b_\nu$ is bounded by a power of r , and (4.1) gives the same kind of bound for $f^{(k)}/f$. The sum contains only finitely many terms. Lemma 3.4 therefore yields

$$|\epsilon_{\nu,\theta}(r)| \leq C_\theta r^{K_\theta} e^{-c_\theta r}, \quad c_\theta > 0. \tag{4.8}$$

If there is just one group, $\epsilon_{\nu,\theta} = 0$. As before, $m_\nu = 0$ would give $\epsilon_{\nu,\theta} \equiv 1$, which contradicts this estimate. Thus $m_\nu \geq 1$.

Thus

$$f^{(m_\nu)} + \sum_{k < m_\nu} P_{k,\nu} f^{(k)} = \epsilon_{\nu,\theta}(r) f.$$

Here $m_\nu \geq 1$, each $P_{k,\nu}$ is rational, and f is entire and eventually nonzero on the ray. Its derivative quotients satisfy (4.1), and the continuous error satisfies (4.8). Hence Lemma 2.5 applies. There are only finitely many fundamental coefficient equations, one for each exponent Q_ν . For each equation that lemma provides finitely many functions $G(z^{1/p})$ and excludes finitely many directions. Their union is still finite and depends only on the original equation.

Let $\mathcal{E}_f$ be the union of $\mathcal{E}_G$, the directions excluded by Lemma 3.4, and these additional directions. It has measure zero, and its complement is dense. For every $\theta \notin \mathcal{E}_f$, we have (3.8) on the whole ray tail. By the choice of the additional directions, every nonzero selected function has a highest term with nonzero real part on that ray. Together with the two-sided bound (4.2), valid for every $\delta > 0$, these are the hypotheses of Lemma 2.6. That lemma proves positive rational order, finite nonzero type, and *c.r.g.*, as required. $\square$

4.3 Proof of Corollary 3.6

Proof. Let f be a transcendental entire solution of the fixed equation of finite order $\rho = \rho(f)$. By Theorem 1.1 and Remark 3.2, f has *c.r.g.* Its indicator

$$h_f(\theta) = \limsup_{r \rightarrow \infty} r^{-\rho} \log |f(re^{i\theta})|$$

is continuous.

The fundamental coefficient equations depend only on the original equation. By Lemmas 2.3 and 2.5, there is a finite partition of $[0, 2\pi)$ into open argument intervals I_ν , $1 \leq \nu \leq M$, and their endpoints. On each interval there are finitely many functions

$$q_{\nu,j}(z) = G_{\nu,j}(z^{1/p_{\nu,j}}), \quad 1 \leq j \leq N_\nu,$$

with fixed branches. These functions are independent of f . On almost every ray with $\theta \in I_\nu$, one of them gives

$$\log |f(re^{i\theta})| = \Re q_{\nu,j}(re^{i\theta}) + O(\log r).$$

Put $d_{\nu,j} = \deg G_{\nu,j}/p_{\nu,j}$, with $d_{\nu,j} = 0$ when $G_{\nu,j} = 0$. Lemma 2.6 shows that ρ is the largest degree selected on the remaining rays. Hence

$$\rho \in \{d_{\nu,j} : d_{\nu,j} > 0, 1 \leq \nu \leq M, 1 \leq j \leq N_\nu\}.$$

This is a fixed finite set of positive rational numbers.

Fix one such ρ and an interval I_ν . For each $q_{\nu,j}$ of degree ρ , write its leading term on this interval as $c_{\nu,j}r^\rho e^{i\rho\theta}$ at $z = re^{i\theta}$. Set

$$C_{\nu,\rho} = \{0\} \cup \{c_{\nu,j} : d_{\nu,j} = \rho\}.$$

A selected function cannot have degree greater than ρ . Division of the ray estimate by r^ρ therefore gives

$$h_f(\theta) \in \{\Re(ce^{i\rho\theta}) : c \in C_{\nu,\rho}\}, \quad \theta \in I_\nu, \quad (4.9)$$

first on a dense set of directions; a selected function of smaller degree gives $c = 0$. Continuity of h_f and finiteness of $C_{\nu,\rho}$ extend (4.9) to every $\theta \in I_\nu$.

Two alternatives corresponding to distinct $c, d \in C_{\nu,\rho}$ are equal precisely at the solutions of

$$\Re((c - d)e^{i\rho\theta}) = 0.$$

There are only finitely many such points in I_ν . Divide I_ν at these points. On each resulting open interval the alternatives are pairwise distinct, so continuity forces h_f to equal one fixed alternative throughout that interval. There are finitely many intervals and finitely many choices on each; thus only finitely many indicators can occur for this order. Take the union of the interval endpoints and these equality points over the finitely many possible orders. These give the directions ϑ_j in the statement and prove the first assertion.

For angular intervals crossing 0, we use the 2π -periodic extension of h_f and the corresponding translates of the directions ϑ_j . The zero distribution theorem for such functions [12, Chapter III, Section 3, Theorem 3, pp. 152–155] gives

$$\lim_{r \rightarrow \infty} \frac{n_f(r; \alpha, \beta)}{r^\rho} = \frac{h'_f(\beta) - h'_f(\alpha) + \rho^2 \int_\alpha^\beta h_f(t) dt}{2\pi\rho}$$

when the indicator is differentiable at the endpoints. On each open interval of the finite partition, $h''_f + \rho^2 h_f = 0$. Integrate on the subintervals between α and β and add. The derivatives at the interior endpoints give

$$h'_f(\beta) - h'_f(\alpha) + \rho^2 \int_\alpha^\beta h_f(t) dt = \sum_{\alpha < \vartheta_j < \beta} (h'_f(\vartheta_j + 0) - h'_f(\vartheta_j - 0)).$$

This proves (3.10). Choose α and β on opposite sides of a single ϑ_j . Since the zero count is nonnegative, its limiting density shows that the corresponding jump is nonnegative. If the interval contains no ϑ_j , the sum is empty and $n_f(r; \alpha, \beta) = o(r^\rho)$. Cover the complement of the prescribed neighbourhoods by finitely many open angular intervals containing no listed ray. Applying this estimate to these intervals proves the second assertion. $\square$

5 An extension via Laurent–Puisseux asymptotics

The proof also applies to coefficients with Laurent–Puisseux asymptotic expansions on finitely many sectors. We state a sufficient condition for this extension. Put $a_n \equiv 1$ in (1.5).

Let finitely many rays divide a neighbourhood of infinity into open sectors S_ν . On each sector, fix a nonvanishing holomorphic function W_ν , an integer $p_\nu \geq 1$, and a branch of z^{1/p_ν} . Suppose that there are an integer j_ν and constants $c_{k,\nu,j}$ such that, for every integer $J \geq j_\nu$,

$$\frac{a_k(z)}{W_\nu(z)} = \sum_{j=j_\nu}^J c_{k,\nu,j} z^{-j/p_\nu} + O(|z|^{-(J+1)/p_\nu}), \quad 0 \leq k \leq n, \quad (5.1)$$

on every closed subsector of S_ν . The constants in the estimate and the starting radius may depend on J and the closed subsector. The function W_ν , the branch and all the coefficients $c_{k,\nu,j}$ are fixed independently of the solution. We require that, on each sector, at least one of these formal series is nonzero. These are Laurent–Puisseux asymptotic expansions: finitely many positive powers are allowed, and the series need not converge. A zero series means that the corresponding quotient is $O(|z|^{-L})$ for every $L > 0$. The condition includes $k = n$, so it also applies to $1/W_\nu$.

Exponential polynomial coefficients satisfy this condition: group the terms as in (3.5) and choose W_ν to be the largest exponential factor on each of the resulting sectors.

Proposition 5.1 *Suppose that the coefficients of (1.5) are entire and satisfy the above conditions. Then every transcendental entire solution of finite order has positive rational order, finite nonzero type and c.r.g. On almost every ray it also satisfies*

$$\log |f(re^{i\theta})| = \Re q_\theta(re^{i\theta}) + O(\log r), \quad (5.2)$$

where the q_θ are chosen from a finite list of polynomials in fractional powers of z , with branches fixed on finitely many argument intervals. The list depends only on the coefficient expansions.

Proof. Let f be such a solution. On each sector write

$$\widehat{a}_{k,\nu}(z) = \sum_{j=j_\nu}^{\infty} c_{k,\nu,j} z^{-j/p_\nu}, \quad m = \max\{k : \widehat{a}_{k,\nu} \neq 0\}.$$

Choose a ray outside Gundersen’s angular exceptional set. On its tail, f is nonzero and all $f^{(k)}/f$, $1 \leq k \leq n$, have polynomial bounds, as in the proof of Theorem 3.2. If $m = 0$, division of the equation by $W_\nu f$ would equate the nonzero leading term of a_0/W_ν to a quantity smaller than every negative power of r . This is impossible. Thus $m \geq 1$.

The nonzero leading term of a_m/W_ν shows that a_m has no zeros sufficiently far out on each closed subsector. Hence, on the chosen ray,

$$f^{(m)} + \sum_{k=0}^{m-1} \frac{a_k}{a_m} f^{(k)} = \epsilon_\theta f, \quad \epsilon_\theta = - \sum_{k=m+1}^n \frac{a_k}{a_m} \frac{f^{(k)}}{f}. \quad (5.3)$$

Each coefficient with $k > m$ has zero expansion. Therefore $\epsilon_\theta(r) = O(r^{-L})$ for every $L > 0$. The companion matrix A of the left-hand side has a Laurent–Puiseux asymptotic expansion, denoted by $\hat{A}$.

By the Turrittin–Levelt formal reduction theorem, this series admits a formal reduction at infinity after a finite ramification; see [1] and [2, Theorem 4]. For an ordinary point in the local variable $x = z^{-1/p_\nu}$, use its formal power series fundamental matrix. After replacing p_ν by a multiple if necessary, this gives a formally invertible Laurent–Puiseux matrix $\hat{H}$, a constant matrix G , and a diagonal matrix Q whose entries are polynomials in z^{1/p_ν} with zero constant terms. The formal change $Y = \hat{H}V$ reduces $Y' = \hat{A}Y$ to $V' = (Q' + G/z)V$. Hence

$$\hat{A}\hat{H} - \hat{H}' = \hat{H}(Q' + G/z), \quad GQ = QG. \quad (5.4)$$

The diagonal polynomials and the further choice of fractional powers are fixed by this formal system.

Let H_N be a sufficiently long finite truncation of $\hat{H}$ and put

$$D_N = AH_N - H'_N - H_N(Q' + G/z).$$

By (5.4) and the asymptotic expansion of A , for any prescribed $L > 0$ the truncation can be chosen so that $D_N = O(|z|^{-L})$ on each closed subsector. Indeed, the coefficients up to any fixed order vanish by (5.4) once sufficiently many terms are retained. For all sufficiently long truncations the leading powers of the entries of H_N and the leading nonzero term of its determinant are fixed. Thus H_N is invertible far out, and the adjugate formula gives polynomial bounds for H_N and H_N^{-1} with an exponent independent of these truncations. The matrices z^G and z^{-G} have polynomial bounds as in Lemma 2.3.

Set $T_N = H_N z^G$. Direct differentiation gives

$$T_N^{-1}AT_N - T_N^{-1}T'_N = Q' + S_N, \quad S_N = z^{-G}H_N^{-1}D_N z^G. \quad (5.5)$$

Choose L large enough to absorb the polynomial bounds in this last product. Then $S_N = O(|z|^{-2})$. Also, T_N and its inverse have polynomial bounds.

For $\mathbf{x} = (f, f', \dots, f^{(m-1)})^\top$, use (5.3) and set $\mathbf{x} = T_N \mathbf{z}$ on the ray. With $E = \mathbf{e}_m \mathbf{e}_1^\top$, this gives

$$\frac{d\mathbf{z}}{dr} = \left[\frac{d}{dr} Q(re^{i\theta}) + R_\theta(r) \right] \mathbf{z}, \quad R_\theta = e^{i\theta} (S_N + T_N^{-1} \epsilon_\theta E T_N).$$

Here every matrix is evaluated at $re^{i\theta}$, and $\int_R^\infty \|R_\theta(r)\| dr < \infty$. After excluding the finitely many directions determined by the diagonal polynomials, Lemma 2.4 applies. The norm comparison in the proof of Lemma 2.5 then gives (5.2). There are only finitely many sectors and finitely many choices of branches, so the list of polynomials is finite. The two-sided bound for $\log |f|$ follows from the same logarithmic derivative estimate as in the proof of Theorem 3.2. Lemma 2.6 now proves the remaining conclusions. $\square$

We give two coefficient classes satisfying the conditions of Proposition 5.1.

Corollary 5.2 *Suppose that*

$$a_k(z) = E_k(z) + \sum_{\ell=1}^s \int_{a_\ell}^{b_\ell} w_{k\ell}(t) e^{\alpha_\ell t z} dt, \quad 0 \leq k < n, \quad (5.6)$$

with $a_n \equiv 1$. Here the E_k are exponential polynomials, s is finite, $a_\ell < b_\ell$ are real numbers, $\alpha_\ell \in \mathbb{C} \setminus \{0\}$, and each weight $w_{k\ell}$ is a function of t , holomorphic in a complex neighbourhood of $[a_\ell, b_\ell]$. Then every transcendental entire solution of (1.5) of finite order satisfies the conclusions of Proposition 5.1.

Proof. Each integral is entire and of exponential type. For every positive integer N , integration by parts gives

$$\begin{aligned} \int_a^b w(t) e^{\alpha t z} dt &= \sum_{j=0}^{N-1} \frac{(-1)^j}{(\alpha z)^{j+1}} (w^{(j)}(b) e^{\alpha b z} - w^{(j)}(a) e^{\alpha a z}) \\ &\quad + O\left(\frac{|e^{\alpha a z}| + |e^{\alpha b z}|}{|z|^{N+1}}\right), \end{aligned} \quad (5.7)$$

on every closed subsector of $\Re(\alpha z) > 0$ or $\Re(\alpha z) < 0$. The constant may depend on N , α , w and the subsector. The remainder follows by integrating the bounded function $w^{(N)}$ against $e^{\alpha t z}$; the integral of its modulus is bounded by a constant times $(|e^{\alpha a z}| + |e^{\alpha b z}|)/|z|$.

Put $u = \alpha_\ell t$ and group the resulting segments by their supporting lines through the origin. On each line, subdivide overlapping segments and add the analytic weights, including their orientations. Regard the weight as zero outside the segments. If all derivatives agree across an endpoint, the adjacent weights agree near it and can be joined. After joining such pieces and deleting zero weights, each remaining nonzero endpoint contributes a nonzero series in at least one component. Different supporting lines have no common nonzero endpoint. The exponential factor 1 is retained because $a_n = 1$.

Group the endpoint exponential factors with those of the E_k , combining exponents that differ by constants. Discard factors whose series are zero in every component. A polynomial multiplier cannot cancel a nonzero series of negative powers. On each sector outside finitely many directions, choose W_ν to be the largest remaining exponential factor. Its coefficient series are not all zero, while the other factors are smaller than every negative power after division by W_ν . Thus (5.1) holds with $p_\nu = 1$. Proposition 5.1 applies. $\square$

A simple special case is given by the *Gaussian weight* $w(t) = e^{-t^2}$. Taking $s = 1$, $a_1 = 0$, $b_1 = 1$, $\alpha_1 = 1$ and $w_{k1}(t) = d_k e^{-t^2}$ in (5.6) gives

$$a_k(z) = E_k(z) + d_k H(z), \quad 0 \leq k < n, \quad H(z) = \int_0^1 e^{tz - t^2} dt,$$

where the d_k are constants. Formula (5.7) gives

$$e^{-z} H(z) \sim \frac{e^{-1}}{z} \quad (\Re z > 0), \quad H(z) \sim -\frac{1}{z} \quad (\Re z < 0),$$

on closed subsectors of the respective half-planes. The function H is not an exponential polynomial. Indeed, if it were, so would $e^{-z} H$ be. On a ray in the right half-plane outside finitely many directions, the latter would be asymptotic to $P(z) e^{Q(z)}$, where P is a nonzero polynomial and Q is a polynomial. If Q is nonconstant, its leading term has nonzero real

part on the chosen ray. Neither case is compatible with $e^{-z}H(z) \sim e^{-1}/z$. Thus (5.6) includes coefficients outside the class in Theorem 1.1.

The next class also allows fractional powers in the coefficient expansions. The exponent of each integral kernel is a polynomial in z and t of the form $Q_\ell(z)t^{m_\ell}$.

Corollary 5.3 *Suppose that*

$$a_k(z) = E_k(z) + \sum_{\ell=1}^s \int_0^1 w_{k\ell}(t) e^{Q_\ell(z)t^{m_\ell}} dt, \quad 0 \leq k < n, \quad (5.8)$$

with $a_n \equiv 1$. Here the E_k are exponential polynomials, s is finite, the Q_ℓ are nonconstant complex polynomials, the m_ℓ are positive integers, and the weights $w_{k\ell}$ are functions of t , holomorphic in complex neighbourhoods of $[0, 1]$. Then every transcendental entire solution of (1.5) of finite order satisfies the conclusions of Proposition 5.1.

Proof. We first obtain asymptotic expansions associated with the endpoints $t = 0$ and $t = 1$ for

$$I(\lambda) = \int_0^1 w(t) e^{\lambda t^m} dt.$$

Put

$$v(u) = \frac{1}{m} u^{1/m-1} w(u^{1/m}), \quad 0 < u \leq 1.$$

Then $I(\lambda) = \int_0^1 v(u) e^{\lambda u} du$. The function v is analytic near $u = 1$ and integrable at $u = 0$. If $\Re \lambda \geq c|\lambda|$ for a fixed $c > 0$, split this integral at a fixed point $0 < \delta < 1$. After multiplication by $e^{-\lambda}$, the first part is exponentially small. Repeated integration by parts in the second part gives

$$e^{-\lambda} I(\lambda) = \sum_{j=0}^{N-1} (-1)^j v^{(j)}(1) \lambda^{-j-1} + O(|\lambda|^{-N-1}). \quad (5.9)$$

If $\Re \lambda \leq -c|\lambda|$, expand w at $t = 0$. Integration of its Taylor polynomial gives

$$I(\lambda) = \sum_{j=0}^{N-1} \frac{w^{(j)}(0)}{j!} \frac{\Gamma((j+1)/m)}{m(-\lambda)^{(j+1)/m}} + O(|\lambda|^{-(N+1)/m}), \quad (5.10)$$

where the powers of $-\lambda$ have their principal values. Indeed, the Taylor remainder is bounded by a constant times t^N near zero, and its integral is $O(|\lambda|^{-(N+1)/m})$. The part away from zero and the tails introduced by extending the Taylor integrals to infinity are exponentially small. Both estimates hold uniformly for the indicated ranges of λ , for every positive integer N . They are the endpoint forms of Watson's lemma [15, Section 2.4(i)].

Substitute $\lambda = Q_\ell(z)$. Outside the finitely many directions where the leading term of some $Q_\ell(re^{i\theta})$ has zero real part, the sign is fixed and $|\Re Q_\ell(z)| \geq c|Q_\ell(z)|$ on closed subsectors far out. Formula (5.9) gives an exponential factor $e^{Q_\ell(z)}$ with an expansion in negative integer powers of z . Formula (5.10) gives a Laurent–Puiseux expansion in z ; a common denominator is a multiple of all the m_ℓ . All the integrals in (5.8) are entire.

Before comparing the exponential factors, combine polynomials Q_ℓ that differ by constants. If $Q_\ell = Q + c_\ell$, multiply the corresponding density $v_{k\ell}(u)$ by $e^{c_\ell u}$ and add the densities. The sum is analytic near $u = 1$ and on $0 < u < 1$. If all the coefficients in its

asymptotic expansion (5.9) vanish, all its derivatives at 1 vanish. It is then zero throughout $(0, 1)$, so the whole integral in that component is zero. Delete any group that is zero in every component. A nonzero remaining endpoint series has only negative integer powers of z , and cannot be cancelled by a polynomial multiplier in the E_k . Group the factors from the E_k in the same way, absorbing constant factors into their polynomial multipliers.

The remaining exponential factors, those in the E_k and the factor 1 form a finite list. On each sector outside finitely many comparison directions, choose the largest factor as W_ν . If $W_\nu \neq 1$, its coefficient series is nonzero in at least one component. If $W_\nu = 1$, the series for a_n/W_ν is 1. Equations (5.9) and (5.10) therefore verify all the conditions of Proposition 5.1. $\square$

For example, taking $Q(z) = z$ and $m = 2$ gives

$$J(z) = \int_0^1 e^{zt^2} dt,$$

with

$$e^{-z}J(z) \sim \frac{1}{2z} \quad (\Re z > 0), \quad J(z) \sim \frac{\sqrt{\pi}}{2(-z)^{1/2}} \quad (\Re z < 0),$$

on closed subsectors of the respective half-planes. Thus fractional powers already occur for an entire coefficient of exponential type. The choices $Q(z) = \alpha z$ and $Q(z) = \alpha z^m$ include the kernels $e^{\alpha z t^m}$ and $e^{\alpha (tz)^m}$, respectively.

Acknowledgements

The author thanks Zhi-Tao Wen for providing helpful references, and Walter Bergweiler for his comments and suggestions on an earlier version of the manuscript during the author's visit to Kiel University. The author acknowledges support from the China Scholarship Council (CSC No. 202508440485) during the revision of this paper.

AI assistance and formal verification

OpenAI's ChatGPT and Codex were used to assist with literature searches, to develop and check parts of the arguments, to revise the exposition and LaTeX source, and to develop and debug the Lean formalizations. The author is responsible for the mathematical content, references and final presentation.

All numbered theorem, proposition, and corollary statements in this manuscript have been formalized in Lean 4. The fixed source version and verification report are cited in [13]. All 320 modules passed the recorded verification, which audited 1,734 distinct declarations; the sources contain no `sorry`.

References

- [1] D. G. Babbitt and V. S. Varadarajan, *Formal reduction theory of meromorphic differential equations: a group theoretic view*, Pacific J. Math. **109** (1983), no. 1, 1–80.
- [2] M. Barkatou, F. Á. Carnicero-Martín and F. Sanz Sánchez, *Turrittin's theorem revisited. The real case*, arXiv:2305.08211 (2023).

- [3] W. Bergweiler, *A question of Gol'dberg and Ostrovskii concerning linear differential equations with coefficients of completely regular growth*, Proc. Amer. Math. Soc. **151** (2023), no. 5, 2097–2101.
- [4] G. Cotti, D. Guzzetti and D. Masoero, *Asymptotic solutions for linear ODEs with not-necessarily meromorphic coefficients: A Levinson type theorem on complex domains, and applications*, J. Differential Equations **428** (2025), 1–58.
- [5] D. G. Dickson, *Expansions in series of solutions of linear difference-differential and infinite order differential equations with constant coefficients*, Mem. Amer. Math. Soc., no. 23, American Mathematical Society, Providence, RI, 1957.
- [6] J. Drolet, *Über das Wachstum der Lösungen linear-homogener Differentialgleichungen mit Exponentialpolynomen als Koeffizienten*, Ph.D. thesis, Universität Siegen, 1984.
- [7] M. Frei, *Über die Lösungen linearer Differentialgleichungen mit ganzen Funktionen als Koeffizienten*, Comment. Math. Helv. **35** (1961), 201–222.
- [8] G. G. Gundersen, *Estimates for the logarithmic derivative of a meromorphic function, plus similar estimates*, J. London Math. Soc. (2) **37** (1988), no. 1, 88–104.
- [9] V. P. Havin and N. K. Nikolski (eds.), *Linear and complex analysis problem book 3, Part II*, Lecture Notes in Mathematics, vol. 1574, Springer-Verlag, Berlin, 1994.
- [10] J. Heittokangas, I. Laine, K. Tohge and Z.-T. Wen, *Completely regular growth solutions of second order complex linear differential equations*, Ann. Acad. Sci. Fenn. Math. **40** (2015), no. 2, 985–1003.
- [11] J. Heittokangas, K. Ishizaki, K. Tohge and Z.-T. Wen, *Value distribution of exponential polynomials and their role in the theories of complex differential equations and oscillation theory*, Bull. Lond. Math. Soc. **55** (2023), no. 1, 1–77.
- [12] B. Ja. Levin, *Distribution of zeros of entire functions*, revised ed., Translations of Mathematical Monographs, vol. 5, American Mathematical Society, Providence, RI, 1980.
- [13] X.-Y. Li, *Lean formalization of completely regular growth for differential equations*, Lean 4 source code and verification report, version 1.0.1 (2026), fixed Git commit fd0cb44a2f83f6fa926184e25dc6cfb218da317c, <https://github.com/xyli9596/crg-growth-and-zero-distribution/tree/fd0cb44a2f83f6fa926184e25dc6cfb218da317c>.
- [14] X.-Y. Li, J. Wang and Z.-T. Wen, *Representation of finite order solutions to linear differential equations with exponential sum coefficients*, J. Differential Equations **423** (2025), 118–132.
- [15] National Institute of Standards and Technology, *NIST Digital Library of Mathematical Functions*, <https://dlmf.nist.gov/>.
- [16] V. P. Petrenko, *Entire curves*, Vishcha Shkola, Kharkov, 1984 (in Russian).

- [17] L. I. Ronkin, *Functions of completely regular growth*, Mathematics and its Applications (Soviet Series), vol. 81, Kluwer Academic Publishers, Dordrecht, 1992.
- [18] N. Steinmetz, *Exceptional values of solutions of linear differential equations*, Math. Z. **201** (1989), no. 3, 317–326.
- [19] W. Wasow, *Asymptotic expansions for ordinary differential equations*, Pure and Applied Mathematics, vol. 14, Interscience Publishers, New York, 1965.
- [20] Z.-T. Wen, G. G. Gundersen and J. Heittokangas, *Dual exponential polynomials and linear differential equations*, J. Differential Equations **264** (2018), no. 1, 98–114.

Xingyu Li
Department of Mathematics
The University of Manchester
Oxford Road, Manchester, M13 9PL, United Kingdom
xingyu.li@manchester.ac.uk