

First-Order Logic: More Semantics

Yuting Wang

John Hopcroft Center for Computer Science
Shanghai Jiao Tong University

Nov 28, 2022

What is the equivalent definition of “tautological implication” in first-order logic?

Logical Implication

Definition

Let Γ be a set of wffs and φ a wff. Γ **logically implies** φ , written as

$$\Gamma \models \varphi$$

if for every structure \mathfrak{A} and every assignment $s : V \rightarrow |\mathfrak{A}|$,

if \mathfrak{A} satisfies Γ with s , then \mathfrak{A} satisfies φ with s .

Logical Implication

Definition

Let Γ be a set of wffs and φ a wff. Γ **logically implies** φ , written as

$$\Gamma \models \varphi$$

if for every structure \mathfrak{A} and every assignment $s : V \rightarrow |\mathfrak{A}|$,

if \mathfrak{A} satisfies Γ with s , then \mathfrak{A} satisfies φ with s .

Remark

$\Gamma \models \varphi$ is also read as:

- ▶ φ is a logical consequence of Γ , or
- ▶ Γ semantically implies φ , or
- ▶ φ is a semantic consequence of Γ .

Logical Implication for Sentences

Theorem

For a set of sentences Σ and a sentence σ , $\Sigma \models \sigma$ iff for every model \mathfrak{A} of Σ , \mathfrak{A} is a model of σ .

Scorates Again

Question

Assume the following premises:

- ▶ All men are mortal.
- ▶ Socrates is a man.

We can derive the conclusion:

- ▶ Socrates is mortal.

How do we express this reasoning using logical implication?

Scorates Again

Question

Assume the following premises:

- ▶ All men are mortal.
- ▶ Socrates is a man.

We can derive the conclusion:

- ▶ Socrates is mortal.

How do we express this reasoning using logical implication?

Answer

Let \mathbb{L} be the first-order language with 1-ary predicate symbols:

Scorates Again

Question

Assume the following premises:

- ▶ All men are mortal.
- ▶ Socrates is a man.

We can derive the conclusion:

- ▶ Socrates is mortal.

How do we express this reasoning using logical implication?

Answer

Let \mathbb{L} be the first-order language with 1-ary predicate symbols:

- ▶ \dot{P} for asserting a being is a man;

Scorates Again

Question

Assume the following premises:

- ▶ All men are mortal.
- ▶ Socrates is a man.

We can derive the conclusion:

- ▶ Socrates is mortal.

How do we express this reasoning using logical implication?

Answer

Let \mathbb{L} be the first-order language with 1-ary predicate symbols:

- ▶ \dot{P} for asserting a being is a man;
- ▶ \dot{Q} for asserting a being is mortal;

Scorates Again

Question

Assume the following premises:

- ▶ All men are mortal.
- ▶ Socrates is a man.

We can derive the conclusion:

- ▶ Socrates is mortal.

How do we express this reasoning using logical implication?

Answer

Let \mathbb{L} be the first-order language with 1-ary predicate symbols:

- ▶ \dot{P} for asserting a being is a man;
 - ▶ \dot{Q} for asserting a being is mortal;
- and a constant symbol \dot{c} denoting Socrates.

Scorates Again

Question

Assume the following premises:

- ▶ All men are mortal.
- ▶ Socrates is a man.

We can derive the conclusion:

- ▶ Socrates is mortal.

How do we express this reasoning using logical implication?

Answer

Let \mathbb{L} be the first-order language with 1-ary predicate symbols:

- ▶ \dot{P} for asserting a being is a man;
- ▶ \dot{Q} for asserting a being is mortal;

and a constant symbol \dot{c} denoting Socrates.

Let $\Sigma = \{\forall x(\dot{P}x \rightarrow \dot{Q}x), \dot{P}\dot{c}\}$. Then

$$\Sigma \models \dot{Q}\dot{c}.$$

Logical Equivalence

As before, we write $\alpha \models \beta$ for $\{\alpha\} \models \beta$.

Definition

α and β are **logically equivalent**, written as $\alpha \models \beta$, if $\alpha \models \beta$ and $\beta \models \alpha$.

Logical Equivalence

As before, we write $\alpha \models \beta$ for $\{\alpha\} \models \beta$.

Definition

α and β are **logically equivalent**, written as $\alpha \models \beta$, if $\alpha \models \beta$ and $\beta \models \alpha$.

Example

$$\forall x \forall y (\dot{P}x \rightarrow \neg \dot{Q}y) \models \forall x \forall y (\neg(\dot{P}x \wedge \dot{Q}y))$$

Valid Wffs

Some wffs are satisfied in every structure under every assignment s .

Definition

Let φ be a wff in the language \mathbb{L} . φ is **valid** if $\emptyset \models \varphi$, written as $\models \varphi$.

Valid Wffs

Some wffs are satisfied in every structure under every assignment s .

Definition

Let φ be a wff in the language \mathbb{L} . φ is **valid** if $\emptyset \models \varphi$, written as $\models \varphi$.

Lemma

φ is valid iff $\models_{\mathfrak{A}} \varphi[s]$ for every structure \mathfrak{A} for \mathbb{L} and every assignment function s for \mathfrak{A} .

Valid Wffs

Some wffs are satisfied in every structure under every assignment s .

Definition

Let φ be a wff in the language \mathbb{L} . φ is **valid** if $\emptyset \models \varphi$, written as $\models \varphi$.

Lemma

φ is valid iff $\models_{\mathfrak{A}} \varphi[s]$ for every structure \mathfrak{A} for \mathbb{L} and every assignment function s for \mathfrak{A} .

Corollary

A sentence σ is valid iff it is true in every structure.

Examples

Which of the following are valid?

Examples

Which of the following are valid?

► $x \dot{=} x$

Examples

Which of the following are valid?

▶ $x \dot{=} x$ YES

Examples

Which of the following are valid?

▶ $x \dot{=} x$ YES

▶ $\exists x \, x \dot{=} x$

Examples

Which of the following are valid?

▶ $x \dot{=} x$ YES

▶ $\exists x \, x \dot{=} x$ YES

Examples

Which of the following are valid?

- ▶ $x \doteq x$ YES
- ▶ $\exists x \, x \doteq x$ YES
- ▶ $\forall x \exists y \, x \neq y$

Examples

Which of the following are valid?

- ▶ $x \doteq x$ YES
- ▶ $\exists x \, x \doteq x$ YES
- ▶ $\forall x \exists y \, x \neq y$ NO

Examples

Which of the following are valid?

- ▶ $x \doteq x$ YES
- ▶ $\exists x \, x \doteq x$ YES
- ▶ $\forall x \exists y \, x \not\doteq y$ NO
- ▶ $\dot{P}_x \vee \neg \dot{P}_x$

Examples

Which of the following are valid?

- ▶ $x \dot{=} x$ YES
- ▶ $\exists x x \dot{=} x$ YES
- ▶ $\forall x \exists y x \dot{\neq} y$ NO
- ▶ $\dot{P}x \vee \neg \dot{P}x$ YES

Examples

Which of the following are valid?

- ▶ $x \doteq x$ YES
- ▶ $\exists x \, x \doteq x$ YES
- ▶ $\forall x \exists y \, x \not\doteq y$ NO
- ▶ $\dot{P}x \vee \neg \dot{P}x$ YES
- ▶ $\neg \exists x \, x \not\doteq x$

Examples

Which of the following are valid?

- ▶ $x \doteq x$ YES
- ▶ $\exists x x \doteq x$ YES
- ▶ $\forall x \exists y x \not\doteq y$ NO
- ▶ $\dot{P}_x \vee \neg \dot{P}_x$ YES
- ▶ $\neg \exists x x \not\doteq x$ YES

More Examples

Which of the following are valid?

More Examples

Which of the following are valid?

► $\forall x(\dot{P}x \rightarrow \exists y\dot{P}y)$

More Examples

Which of the following are valid?

► $\forall x(\dot{P}x \rightarrow \exists y\dot{P}y)$ YES

More Examples

Which of the following are valid?

▶ $\forall x(\dot{P}x \rightarrow \exists y\dot{P}y)$ YES

▶ $\dot{P}x \rightarrow \exists x\dot{P}x$

More Examples

Which of the following are valid?

▶ $\forall x(\dot{P}x \rightarrow \exists y\dot{P}y)$ YES

▶ $\dot{P}x \rightarrow \exists x\dot{P}x$ YES

More Examples

Which of the following are valid?

▶ $\forall x(\dot{P}x \rightarrow \exists y\dot{P}y)$ YES

▶ $\dot{P}x \rightarrow \exists x\dot{P}x$ YES

▶ $\dot{P}x \rightarrow \forall x\dot{P}x$

More Examples

Which of the following are valid?

▶ $\forall x(\dot{P}x \rightarrow \exists y\dot{P}y)$ YES

▶ $\dot{P}x \rightarrow \exists x\dot{P}x$ YES

▶ $\dot{P}x \rightarrow \forall x\dot{P}x$ NO

More Examples

Which of the following are valid?

▶ $\forall x(\dot{P}x \rightarrow \exists y\dot{P}y)$ YES

▶ $\dot{P}x \rightarrow \exists x\dot{P}x$ YES

▶ $\dot{P}x \rightarrow \forall x\dot{P}x$ NO

▶ $\exists x\dot{P}x \rightarrow \forall x\dot{P}x$

More Examples

Which of the following are valid?

▶ $\forall x(\dot{P}x \rightarrow \exists y\dot{P}y)$ YES

▶ $\dot{P}x \rightarrow \exists x\dot{P}x$ YES

▶ $\dot{P}x \rightarrow \forall x\dot{P}x$ NO

▶ $\exists x\dot{P}x \rightarrow \forall x\dot{P}x$ NO

More Examples

Which of the following are valid?

▶ $\forall x(\dot{P}x \rightarrow \exists y\dot{P}y)$ YES

▶ $\dot{P}x \rightarrow \exists x\dot{P}x$ YES

▶ $\dot{P}x \rightarrow \forall x\dot{P}x$ NO

▶ $\exists x\dot{P}x \rightarrow \forall x\dot{P}x$ NO

▶ $\exists x(\dot{P}x \rightarrow \forall x\dot{P}x)$

More Examples

Which of the following are valid?

▶ $\forall x(\dot{P}x \rightarrow \exists y\dot{P}y)$ YES

▶ $\dot{P}x \rightarrow \exists x\dot{P}x$ YES

▶ $\dot{P}x \rightarrow \forall x\dot{P}x$ NO

▶ $\exists x\dot{P}x \rightarrow \forall x\dot{P}x$ NO

▶ $\exists x(\dot{P}x \rightarrow \forall x\dot{P}x)$ YES

An Algorithm for Determining Validity?

Question

Is there an algorithm for determining validity?

An Algorithm for Determining Validity?

Question

Is there an algorithm for determining validity?

In other words, is there an algorithm that on input a wff φ will give an output of “yes” if φ is valid and output “no”, otherwise?

Satisfiability

Definition

- ▶ The wff φ is **satisfiable** if there is some structure \mathfrak{A} and some assignment $s : V \rightarrow |\mathfrak{A}|$ such that $\models_{\mathfrak{A}} \varphi[s]$.

Satisfiability

Definition

- ▶ The wff φ is **satisfiable** if there is some structure \mathfrak{A} and some assignment $s : V \rightarrow |\mathfrak{A}|$ such that $\models_{\mathfrak{A}} \varphi[s]$.
- ▶ The set Γ of wffs is **satisfiable** if there is some structure \mathfrak{A} and some assignment $s : V \rightarrow |\mathfrak{A}|$ such that $\models_{\mathfrak{A}} \varphi[s]$ for every φ in Γ .

Satisfiability

Definition

- ▶ The wff φ is **satisfiable** if there is some structure \mathfrak{A} and some assignment $s : V \rightarrow |\mathfrak{A}|$ such that $\models_{\mathfrak{A}} \varphi[s]$.
- ▶ The set Γ of wffs is **satisfiable** if there is some structure \mathfrak{A} and some assignment $s : V \rightarrow |\mathfrak{A}|$ such that $\models_{\mathfrak{A}} \varphi[s]$ for every φ in Γ .

Theorem

φ is not satisfiable iff $\neg\varphi$ is

Satisfiability

Definition

- ▶ The wff φ is **satisfiable** if there is some structure \mathfrak{A} and some assignment $s : V \rightarrow |\mathfrak{A}|$ such that $\models_{\mathfrak{A}} \varphi[s]$.
- ▶ The set Γ of wffs is **satisfiable** if there is some structure \mathfrak{A} and some assignment $s : V \rightarrow |\mathfrak{A}|$ such that $\models_{\mathfrak{A}} \varphi[s]$ for every φ in Γ .

Theorem

φ is not satisfiable iff $\neg\varphi$ is valid.

Is There a Compactness Theorem for First-Order Logic?

Question

Is the following statement true?

For every first-order language \mathbb{L} , and every set Γ of wffs of \mathbb{L} , if every finite subset of Γ is satisfiable then Γ is satisfiable.

Is There a Compactness Theorem for First-Order Logic?

Question

Is the following statement true?

For every first-order language \mathbb{L} , and every set Γ of wffs of \mathbb{L} , if every finite subset of Γ is satisfiable then Γ is satisfiable.

Answer

Yes! But we have to wait for a while to see the answer.

How can we characterize relations in *structures* by looking at wffs in first-order logic?

Relations Defined by Wffs

Definition

Let

- ▶ \mathfrak{A} be a structure, and
- ▶ φ be a wff and n be such that the variables occurring free in φ are included among v_1, \dots, v_n .

The n -ary relation **defined by φ in \mathfrak{A}** is

$$\{(a_1, \dots, a_n) \mid \models_{\mathfrak{A}} \varphi[a_1, \dots, a_n]\}$$

Examples

Example

Examples

Example

- ▶ Let $\mathfrak{R} = (\mathbb{R}, <, +, \times, 0, 1)$. The 1-ary relation $\{a \in \mathbb{R} \mid 0 \leq a\}$ is defined by

$$\exists v_2, v_1 \dot{=} v_2 \times v_2$$

in \mathfrak{R} ;

Examples

Example

- ▶ Let $\mathfrak{R} = (\mathbb{R}, <, +, \times, 0, 1)$. The 1-ary relation $\{a \in \mathbb{R} \mid 0 \leq a\}$ is defined by

$$\exists v_2, v_1 \dot{=} v_2 \times v_2$$

in \mathfrak{R} ;

- ▶ Let $\mathfrak{N} = (\mathbb{N}, <, +, \times, 0, 1)$. The 2-ary relation $\{(a, b) \mid a < b\}$ is defined by

$$\exists v_3 (v_1 \dot{+} (1 \dot{+} v_3) \dot{=} v_2)$$

in \mathfrak{N} .

Definable Relations

Definition

- ▶ The relation R is **definable in the structure \mathfrak{A}** if there is some wff φ that defines it in \mathfrak{A} .

Definable Relations

Definition

- ▶ The relation R is **definable in the structure \mathfrak{A}** if there is some wff φ that defines it in \mathfrak{A} .
- ▶ Let f be a n -ary function f whose domain is a subset of $\underbrace{|\mathfrak{A}| \times \dots \times |\mathfrak{A}|}_n$ and whose range is a subset of $|\mathfrak{A}|$. **f is definable in \mathfrak{A}** if the $(n+1)$ -ary relation

$$\{(a_1, \dots, a_n, b) \mid f(a_1, \dots, a_n) = b\}$$

is definable in \mathfrak{A} .

Examples

Example

Let $\mathfrak{N} = (\mathbb{N}, <, +, \times, 0, 1)$.

Examples

Example

Let $\mathfrak{N} = (\mathbb{N}, <, +, \times, 0, 1)$.

- ▶ $v_1 \dot{+} v_2 \doteq v_3$ defines $\{(a, b, c) \mid a + b = c\}$, which is the same as the function f , where $f(a, b) = a + b$.

Examples

Example

Let $\mathfrak{N} = (\mathbb{N}, <, +, \times, 0, 1)$.

- ▶ $v_1 \dot{+} v_2 \doteq v_3$ defines $\{(a, b, c) \mid a + b = c\}$, which is the same as the function f , where $f(a, b) = a + b$.
- ▶ $v_1 \dot{+} v_3 \doteq v_2$ defines $\{(a, b, c) \mid a + c = b\}$, which is the same as the function f , where

Examples

Example

Let $\mathfrak{N} = (\mathbb{N}, <, +, \times, 0, 1)$.

- ▶ $v_1 \dot{+} v_2 \doteq v_3$ defines $\{(a, b, c) \mid a + b = c\}$, which is the same as the function f , where $f(a, b) = a + b$.
- ▶ $v_1 \dot{+} v_3 \doteq v_2$ defines $\{(a, b, c) \mid a + c = b\}$, which is the same as the function f , where

$$f(a, b) = \begin{cases} b - a & \text{if } a \leq b \\ \text{Undefined} & \text{Otherwise} \end{cases}$$

What Relations are Definable in a Structure?

Proposition

Let \mathfrak{A} be a structure for \mathbb{L} .

What Relations are Definable in a Structure?

Proposition

Let \mathfrak{A} be a structure for \mathbb{L} .

- ▶ $|\mathfrak{A}|$ is definable

What Relations are Definable in a Structure?

Proposition

Let \mathfrak{A} be a structure for \mathbb{L} .

- ▶ $|\mathfrak{A}|$ is definable (by $v_1 \dot{=} v_1$, if $\dot{=}$ is in \mathbb{L});

What Relations are Definable in a Structure?

Proposition

Let \mathfrak{A} be a structure for \mathbb{L} .

- ▶ $|\mathfrak{A}|$ is definable (by $v_1 \dot{=} v_1$, if $\dot{=}$ is in \mathbb{L});
- ▶ \emptyset is definable

What Relations are Definable in a Structure?

Proposition

Let \mathfrak{A} be a structure for \mathbb{L} .

- ▶ $|\mathfrak{A}|$ is definable (by $v_1 \dot{=} v_1$, if $\dot{=}$ is in \mathbb{L});
- ▶ \emptyset is definable (by $v_1 \dot{\neq} v_1$, if $\dot{=}$ is in \mathbb{L});

What Relations are Definable in a Structure?

Proposition

Let \mathfrak{A} be a structure for \mathbb{L} .

- ▶ $|\mathfrak{A}|$ is definable (by $v_1 \dot{=} v_1$, if $\dot{=}$ is in \mathbb{L});
- ▶ \emptyset is definable (by $v_1 \dot{\neq} v_1$, if $\dot{=}$ is in \mathbb{L});
- ▶ $=$ is definable

What Relations are Definable in a Structure?

Proposition

Let \mathfrak{A} be a structure for \mathbb{L} .

- ▶ $|\mathfrak{A}|$ is definable (by $v_1 \dot{=} v_1$, if $\dot{=}$ is in \mathbb{L});
- ▶ \emptyset is definable (by $v_1 \dot{\neq} v_1$, if $\dot{=}$ is in \mathbb{L});
- ▶ $=$ is definable (by $v_1 \dot{=} v_2$, if $\dot{=}$ is in \mathbb{L});

What Relations are Definable in a Structure?

Proposition

Let \mathfrak{A} be a structure for \mathbb{L} .

- ▶ $|\mathfrak{A}|$ is definable (by $v_1 \doteq v_1$, if \doteq is in \mathbb{L});
- ▶ \emptyset is definable (by $v_1 \not\doteq v_1$, if \doteq is in \mathbb{L});
- ▶ $=$ is definable (by $v_1 \doteq v_2$, if \doteq is in \mathbb{L});
- ▶ for every n -ary predicate symbol \dot{P} , $\dot{P}^{\mathfrak{A}}$ is definable

What Relations are Definable in a Structure?

Proposition

Let \mathfrak{A} be a structure for \mathbb{L} .

- ▶ $|\mathfrak{A}|$ is definable (by $v_1 \dot{=} v_1$, if $\dot{=}$ is in \mathbb{L});
- ▶ \emptyset is definable (by $v_1 \dot{\neq} v_1$, if $\dot{=}$ is in \mathbb{L});
- ▶ $=$ is definable (by $v_1 \dot{=} v_2$, if $\dot{=}$ is in \mathbb{L});
- ▶ for every n -ary predicate symbol \dot{P} , $\dot{P}^{\mathfrak{A}}$ is definable (by $\dot{P}_{v_1 \dots v_n}$);

What Relations are Definable in a Structure?

Proposition

Let \mathfrak{A} be a structure for \mathbb{L} .

- ▶ $|\mathfrak{A}|$ is definable (by $v_1 \dot{=} v_1$, if $\dot{=}$ is in \mathbb{L});
- ▶ \emptyset is definable (by $v_1 \dot{\neq} v_1$, if $\dot{=}$ is in \mathbb{L});
- ▶ $=$ is definable (by $v_1 \dot{=} v_2$, if $\dot{=}$ is in \mathbb{L});
- ▶ for every n -ary predicate symbol \dot{P} , $\dot{P}^{\mathfrak{A}}$ is definable (by $\dot{P}_{v_1 \dots v_n}$);
- ▶ for every n -ary function symbol f , $f^{\mathfrak{A}}$ is definable

What Relations are Definable in a Structure?

Proposition

Let \mathfrak{A} be a structure for \mathbb{L} .

- ▶ $|\mathfrak{A}|$ is definable (by $v_1 \dot{=} v_1$, if $\dot{=}$ is in \mathbb{L});
- ▶ \emptyset is definable (by $v_1 \dot{\neq} v_1$, if $\dot{=}$ is in \mathbb{L});
- ▶ $=$ is definable (by $v_1 \dot{=} v_2$, if $\dot{=}$ is in \mathbb{L});
- ▶ for every n -ary predicate symbol \dot{P} , $\dot{P}^{\mathfrak{A}}$ is definable (by $\dot{P}v_1 \dots v_n$);
- ▶ for every n -ary function symbol f , $f^{\mathfrak{A}}$ is definable (by $fv_1 \dots v_n \dot{=} v_{n+1}$);

What Relations are Definable in a Structure?

Proposition

Let \mathfrak{A} be a structure for \mathbb{L} .

- ▶ $|\mathfrak{A}|$ is definable (by $v_1 \dot{=} v_1$, if $\dot{=}$ is in \mathbb{L});
- ▶ \emptyset is definable (by $v_1 \dot{\neq} v_1$, if $\dot{=}$ is in \mathbb{L});
- ▶ $=$ is definable (by $v_1 \dot{=} v_2$, if $\dot{=}$ is in \mathbb{L});
- ▶ for every n -ary predicate symbol \dot{P} , $\dot{P}^{\mathfrak{A}}$ is definable (by $\dot{P}v_1 \dots v_n$);
- ▶ for every n -ary function symbol f , $f^{\mathfrak{A}}$ is definable (by $fv_1 \dots v_n \dot{=} v_{n+1}$);
- ▶ for every constant symbol c , the singleton $\{c^{\mathfrak{A}}\}$ is definable

What Relations are Definable in a Structure?

Proposition

Let \mathfrak{A} be a structure for \mathbb{L} .

- ▶ $|\mathfrak{A}|$ is definable (by $v_1 \dot{=} v_1$, if $\dot{=}$ is in \mathbb{L});
- ▶ \emptyset is definable (by $v_1 \dot{\neq} v_1$, if $\dot{=}$ is in \mathbb{L});
- ▶ $=$ is definable (by $v_1 \dot{=} v_2$, if $\dot{=}$ is in \mathbb{L});
- ▶ for every n -ary predicate symbol \dot{P} , $\dot{P}^{\mathfrak{A}}$ is definable (by $\dot{P}v_1 \dots v_n$);
- ▶ for every n -ary function symbol f , $f^{\mathfrak{A}}$ is definable (by $fv_1 \dots v_n \dot{=} v_{n+1}$);
- ▶ for every constant symbol c , the singleton $\{c^{\mathfrak{A}}\}$ is definable (by $v_1 \dot{=} c$).

What Relations are Definable in a Structure?

Proposition

Let \mathfrak{A} be a structure for \mathbb{L} .

- ▶ $|\mathfrak{A}|$ is definable (by $v_1 \dot{=} v_1$, if $\dot{=}$ is in \mathbb{L});
- ▶ \emptyset is definable (by $v_1 \dot{\neq} v_1$, if $\dot{=}$ is in \mathbb{L});
- ▶ $=$ is definable (by $v_1 \dot{=} v_2$, if $\dot{=}$ is in \mathbb{L});
- ▶ for every n -ary predicate symbol \dot{P} , $\dot{P}^{\mathfrak{A}}$ is definable (by $\dot{P}v_1 \dots v_n$);
- ▶ for every n -ary function symbol f , $f^{\mathfrak{A}}$ is definable (by $fv_1 \dots v_n \dot{=} v_{n+1}$);
- ▶ for every constant symbol c , the singleton $\{c^{\mathfrak{A}}\}$ is definable (by $v_1 \dot{=} c$).

What happens if $\dot{=}$ is not in \mathbb{L} ?

Relations Definable in a Structure

Proposition

- ▶ If P and Q are n -ary relations that are definable in \mathfrak{A} , then so are: the complement of P , $P \cup Q$, $P \cap Q$, $P \setminus Q$.

Relations Definable in a Structure

Proposition

- ▶ If P and Q are n -ary relations that are definable in \mathfrak{A} , then so are: the complement of P , $P \cup Q$, $P \cap Q$, $P \setminus Q$.
- ▶ If the $n + 1$ -ary relation R is definable in \mathfrak{A} then so are the n -ary relations

$$\{(a_1, \dots, a_n) \mid \text{there exists } b \in |\mathfrak{A}|, (a_1, \dots, a_n, b) \in R\}$$

$$\{(a_1, \dots, a_n) \mid \text{there exists } b \in |\mathfrak{A}|, (b, a_1, \dots, a_n) \in R\}$$

Relations Definable in a Structure

Proposition

- ▶ If P and Q are n -ary relations that are definable in \mathfrak{A} , then so are: the complement of P , $P \cup Q$, $P \cap Q$, $P \setminus Q$.
- ▶ If the $n + 1$ -ary relation R is definable in \mathfrak{A} then so are the n -ary relations

$$\{(a_1, \dots, a_n) \mid \text{there exists } b \in |\mathfrak{A}|, (a_1, \dots, a_n, b) \in R\}$$

$$\{(a_1, \dots, a_n) \mid \text{there exists } b \in |\mathfrak{A}|, (b, a_1, \dots, a_n) \in R\}$$

In particular, if R is a binary relation that is definable in \mathfrak{A} then $\text{dom}(R)$ and $\text{rng}(R)$ is definable.

Definable Subsets of \mathfrak{N}

Which of the following subsets of \mathbb{N} are definable in $\mathfrak{N} = (\mathbb{N}, <)$?

Definable Subsets of \mathfrak{N}

Which of the following subsets of \mathbb{N} are definable in $\mathfrak{N} = (\mathbb{N}, <)$?

► \emptyset .

Definable Subsets of \mathfrak{N}

Which of the following subsets of \mathbb{N} are definable in $\mathfrak{N} = (\mathbb{N}, <)$?

► \emptyset . YES

Definable Subsets of \mathfrak{N}

Which of the following subsets of \mathbb{N} are definable in $\mathfrak{N} = (\mathbb{N}, <)$?

▶ \emptyset . YES

▶ \mathbb{N} .

Definable Subsets of \mathfrak{N}

Which of the following subsets of \mathbb{N} are definable in $\mathfrak{N} = (\mathbb{N}, <)$?

- ▶ \emptyset . YES
- ▶ \mathbb{N} . YES

Definable Subsets of \mathfrak{N}

Which of the following subsets of \mathbb{N} are definable in $\mathfrak{N} = (\mathbb{N}, <)$?

- ▶ \emptyset . YES
- ▶ \mathbb{N} . YES
- ▶ $\{0\}$.

Definable Subsets of \mathfrak{N}

Which of the following subsets of \mathbb{N} are definable in $\mathfrak{N} = (\mathbb{N}, <)$?

- ▶ \emptyset . YES
- ▶ \mathbb{N} . YES
- ▶ $\{0\}$. YES. Let $\varphi = \forall v_2 (v_1 < v_2 \vee v_1 = v_2)$

Definable Subsets of \mathfrak{N}

Which of the following subsets of \mathbb{N} are definable in $\mathfrak{N} = (\mathbb{N}, <)$?

- ▶ \emptyset . YES
- ▶ \mathbb{N} . YES
- ▶ $\{0\}$. YES. Let $\varphi = \forall v_2 (v_1 < v_2 \vee v_1 = v_2)$
- ▶ $\{1\}$.

Definable Subsets of \mathfrak{N}

Which of the following subsets of \mathbb{N} are definable in $\mathfrak{N} = (\mathbb{N}, <)$?

- ▶ \emptyset . YES
- ▶ \mathbb{N} . YES
- ▶ $\{0\}$. YES. Let $\varphi = \forall v_2 (v_1 < v_2 \vee v_1 = v_2)$
- ▶ $\{1\}$. YES.

Definable Subsets of \mathfrak{N}

Which of the following subsets of \mathbb{N} are definable in $\mathfrak{N} = (\mathbb{N}, <)$?

- ▶ \emptyset . YES
- ▶ \mathbb{N} . YES
- ▶ $\{0\}$. YES. Let $\varphi = \forall v_2 (v_1 < v_2 \vee v_1 = v_2)$
- ▶ $\{1\}$. YES.
 - ▶ Let $\varphi(x)$ be the result of replacing v_1 in φ with x ;

Definable Subsets of \mathfrak{N}

Which of the following subsets of \mathbb{N} are definable in $\mathfrak{N} = (\mathbb{N}, <)$?

- ▶ \emptyset . YES
- ▶ \mathbb{N} . YES
- ▶ $\{0\}$. YES. Let $\varphi = \forall v_2 (v_1 < v_2 \vee v_1 = v_2)$
- ▶ $\{1\}$. YES.
 - ▶ Let $\varphi(x)$ be the result of replacing v_1 in φ with x ;
 - ▶ We obtain the defining wff as follows:

Definable Subsets of \mathfrak{N}

Which of the following subsets of \mathbb{N} are definable in $\mathfrak{N} = (\mathbb{N}, <)$?

- ▶ \emptyset . YES
- ▶ \mathbb{N} . YES
- ▶ $\{0\}$. YES. Let $\varphi = \forall v_2 (v_1 < v_2 \vee v_1 = v_2)$
- ▶ $\{1\}$. YES.
 - ▶ Let $\varphi(x)$ be the result of replacing v_1 in φ with x ;
 - ▶ We obtain the defining wff as follows:

$$a = 1 \iff a \neq 0 \text{ and } \forall b \in \mathbb{N}, (b < a \implies b = 0)$$

Definable Subsets of \mathfrak{N}

Which of the following subsets of \mathbb{N} are definable in $\mathfrak{N} = (\mathbb{N}, <)$?

- ▶ \emptyset . **YES**
- ▶ \mathbb{N} . **YES**
- ▶ $\{0\}$. **YES**. Let $\varphi = \forall v_2 (v_1 \dot{<} v_2 \vee v_1 \dot{=} v_2)$
- ▶ $\{1\}$. **YES**.
 - ▶ Let $\varphi(x)$ be the result of replacing v_1 in φ with x ;
 - ▶ We obtain the defining wff as follows:

$$\begin{aligned} a = 1 &\iff a \neq 0 \text{ and } \forall b \in \mathbb{N}, (b < a \implies b = 0) \\ &\iff \models_{\mathfrak{N}} \neg \varphi[a] \text{ and } \models_{\mathfrak{N}} \forall v_3 (v_3 \dot{<} v_1 \rightarrow \varphi(v_3))[a] \end{aligned}$$

Definable Subsets of \mathfrak{N}

Which of the following subsets of \mathbb{N} are definable in $\mathfrak{N} = (\mathbb{N}, <)$?

- ▶ \emptyset . **YES**
- ▶ \mathbb{N} . **YES**
- ▶ $\{0\}$. **YES**. Let $\varphi = \forall v_2 (v_1 \dot{<} v_2 \vee v_1 \dot{=} v_2)$
- ▶ $\{1\}$. **YES**.
 - ▶ Let $\varphi(x)$ be the result of replacing v_1 in φ with x ;
 - ▶ We obtain the defining wff as follows:

$$\begin{aligned} a = 1 &\iff a \neq 0 \text{ and } \forall b \in \mathbb{N}, (b < a \implies b = 0) \\ &\iff \models_{\mathfrak{N}} \neg \varphi[a] \text{ and } \models_{\mathfrak{N}} \forall v_3 (v_3 \dot{<} v_1 \rightarrow \varphi(v_3))[a] \\ &\iff \models_{\mathfrak{N}} \neg \varphi \wedge \forall v_3 (v_3 \dot{<} v_1 \rightarrow \varphi(v_3))[a] \end{aligned}$$

Definable Subsets of \mathfrak{N}

Which of the following subsets of \mathbb{N} are definable in $\mathfrak{N} = (\mathbb{N}, <)$?

- ▶ \emptyset . YES
- ▶ \mathbb{N} . YES
- ▶ $\{0\}$. YES. Let $\varphi = \forall v_2 (v_1 \dot{<} v_2 \vee v_1 \dot{=} v_2)$
- ▶ $\{1\}$. YES.
 - ▶ Let $\varphi(x)$ be the result of replacing v_1 in φ with x ;
 - ▶ We obtain the defining wff as follows:

$$\begin{aligned} a = 1 &\iff a \neq 0 \text{ and } \forall b \in \mathbb{N}, (b < a \implies b = 0) \\ &\iff \models_{\mathfrak{N}} \neg \varphi[a] \text{ and } \models_{\mathfrak{N}} \forall v_3 (v_3 \dot{<} v_1 \rightarrow \varphi(v_3))[a] \\ &\iff \models_{\mathfrak{N}} \neg \varphi \wedge \forall v_3 (v_3 \dot{<} v_1 \rightarrow \varphi(v_3))[a] \end{aligned}$$

So the formula $\neg \varphi \wedge \forall v_3 (v_3 \dot{<} v_1 \rightarrow \varphi(v_3))$ defines $\{1\}$.

More Definable Subset of \mathfrak{A}

- ▶ $\{n\}$ for each $n \in \mathbb{N}$.

More Definable Subset of \mathfrak{A}

- ▶ $\{n\}$ for each $n \in \mathbb{N}$. YES

More Definable Subset of \mathfrak{A}

- ▶ $\{n\}$ for each $n \in \mathbb{N}$. YES
- ▶ Every finite subset of \mathbb{N} .

More Definable Subset of \mathfrak{A}

- ▶ $\{n\}$ for each $n \in \mathbb{N}$. YES
- ▶ Every finite subset of \mathbb{N} . YES

More Definable Subset of \mathfrak{A}

- ▶ $\{n\}$ for each $n \in \mathbb{N}$. YES
- ▶ Every finite subset of \mathbb{N} . YES
- ▶ Every cofinite subset of \mathbb{N} .

More Definable Subset of \mathfrak{A}

- ▶ $\{n\}$ for each $n \in \mathbb{N}$. YES
- ▶ Every finite subset of \mathbb{N} . YES
- ▶ Every cofinite subset of \mathbb{N} . YES

More Definable Subset of \mathfrak{A}

- ▶ $\{n\}$ for each $n \in \mathbb{N}$. YES
- ▶ Every finite subset of \mathbb{N} . YES
- ▶ Every cofinite subset of \mathbb{N} . YES
- ▶ Every subset of \mathbb{N} .

More Definable Subset of \mathfrak{A}

- ▶ $\{n\}$ for each $n \in \mathbb{N}$. YES
- ▶ Every finite subset of \mathbb{N} . YES
- ▶ Every cofinite subset of \mathbb{N} . YES
- ▶ Every subset of \mathbb{N} . NO.

How Many Relations are Definable?

Lemma

- (1) Given a structure \mathfrak{A} , the set of definable relations is *enumerable*;
- (2) Not every subset of \mathbb{N} is definable.

How Many Relations are Definable?

Lemma

- (1) Given a structure \mathfrak{A} , the set of definable relations is *enumerable*;
- (2) Not every subset of \mathbb{N} is definable.

Proof.

For (1), note that the set of wffs is enumerable, and every wff may define only one relation.



How Many Relations are Definable?

Lemma

- (1) Given a structure \mathfrak{A} , the set of definable relations is *enumerable*;
- (2) Not every subset of \mathbb{N} is definable.

Proof.

For (1), note that the set of wffs is enumerable, and every wff may define only one relation.

For (2), note that the set of all subsets of \mathbb{N} is uncountable. Therefore, some subset may not match a wff. □

Definable Subsets in General

Which of the following subsets of \mathbb{R} are definable in $\mathfrak{R} = (\mathbb{R}, <)$?

Definable Subsets in General

Which of the following subsets of \mathbb{R} are definable in $\mathfrak{R} = (\mathbb{R}, <)$?

► \emptyset .

Definable Subsets in General

Which of the following subsets of \mathbb{R} are definable in $\mathfrak{R} = (\mathbb{R}, <)$?

► \emptyset . YES

Definable Subsets in General

Which of the following subsets of \mathbb{R} are definable in $\mathfrak{R} = (\mathbb{R}, <)$?

- ▶ \emptyset . YES
- ▶ \mathbb{R} .

Definable Subsets in General

Which of the following subsets of \mathbb{R} are definable in $\mathfrak{R} = (\mathbb{R}, <)$?

- ▶ \emptyset . YES
- ▶ \mathbb{R} . YES

Definable Subsets in General

Which of the following subsets of \mathbb{R} are definable in $\mathfrak{R} = (\mathbb{R}, <)$?

- ▶ \emptyset . YES
- ▶ \mathbb{R} . YES
- ▶ Anything else?

Definable Subsets in General

Which of the following subsets of \mathbb{R} are definable in $\mathfrak{A} = (\mathbb{R}, <)$?

- ▶ \emptyset . YES
- ▶ \mathbb{R} . YES
- ▶ Anything else?

Question

More generally, given a first-order language \mathbb{L} and a structure \mathfrak{A} for \mathbb{L} , how do we figure out which relations in \mathfrak{A} are definable?

Given any wff φ , how do we relate its satisfactions
in different structures?

Homomorphisms

Definition

Let \mathfrak{A} and \mathfrak{B} be structures for \mathbb{L} . A **homomorphism from \mathfrak{A} to \mathfrak{B}** is a function $h : |\mathfrak{A}| \rightarrow |\mathfrak{B}|$ such that:

Homomorphisms

Definition

Let \mathfrak{A} and \mathfrak{B} be structures for \mathbb{L} . A **homomorphism from \mathfrak{A} to \mathfrak{B}** is a function $h : |\mathfrak{A}| \rightarrow |\mathfrak{B}|$ such that:

- ▶ for every n -ary predicate symbol R , other than \doteq , and $a_1, \dots, a_n \in |\mathfrak{A}|$,

$$(a_1, \dots, a_n) \in R^{\mathfrak{A}} \iff (h(a_1), \dots, h(a_n)) \in R^{\mathfrak{B}};$$

Homomorphisms

Definition

Let \mathfrak{A} and \mathfrak{B} be structures for \mathbb{L} . A **homomorphism from \mathfrak{A} to \mathfrak{B}** is a function $h : |\mathfrak{A}| \rightarrow |\mathfrak{B}|$ such that:

- ▶ for every n -ary predicate symbol R , other than \doteq , and $a_1, \dots, a_n \in |\mathfrak{A}|$,

$$(a_1, \dots, a_n) \in R^{\mathfrak{A}} \iff (h(a_1), \dots, h(a_n)) \in R^{\mathfrak{B}};$$

- ▶ for every n -ary function symbol f and $a_1, \dots, a_n \in |\mathfrak{A}|$,

$$h(f^{\mathfrak{A}}(a_1, \dots, a_n)) = f^{\mathfrak{B}}(h(a_1), \dots, h(a_n));$$

Homomorphisms

Definition

Let \mathfrak{A} and \mathfrak{B} be structures for \mathbb{L} . A **homomorphism from \mathfrak{A} to \mathfrak{B}** is a function $h : |\mathfrak{A}| \rightarrow |\mathfrak{B}|$ such that:

- ▶ for every n -ary predicate symbol R , other than \doteq , and $a_1, \dots, a_n \in |\mathfrak{A}|$,

$$(a_1, \dots, a_n) \in R^{\mathfrak{A}} \iff (h(a_1), \dots, h(a_n)) \in R^{\mathfrak{B}};$$

- ▶ for every n -ary function symbol f and $a_1, \dots, a_n \in |\mathfrak{A}|$,

$$h(f^{\mathfrak{A}}(a_1, \dots, a_n)) = f^{\mathfrak{B}}(h(a_1), \dots, h(a_n));$$

- ▶ for every constant symbol c ,

$$h(c^{\mathfrak{A}}) = c^{\mathfrak{B}}.$$

Homomorphisms

Definition

- ▶ h is a homomorphism of \mathfrak{A} **onto** \mathfrak{B} if h is a homomorphism from \mathfrak{A} to \mathfrak{B} and h maps \mathfrak{A} onto \mathfrak{B} ;

Homomorphisms

Definition

- ▶ h is a homomorphism of \mathfrak{A} **onto** \mathfrak{B} if h is a homomorphism from \mathfrak{A} to \mathfrak{B} and h maps \mathfrak{A} onto \mathfrak{B} ;
- ▶ A homomorphism h from \mathfrak{A} into \mathfrak{B} is an **isomorphism** if h is one-to-one;

Homomorphisms

Definition

- ▶ h is a homomorphism of \mathfrak{A} **onto** \mathfrak{B} if h is a homomorphism from \mathfrak{A} to \mathfrak{B} and h maps \mathfrak{A} onto \mathfrak{B} ;
- ▶ A homomorphism h from \mathfrak{A} into \mathfrak{B} is an **isomorphism** if h is one-to-one;
- ▶ The structures \mathfrak{A} and \mathfrak{B} are **isomorphic**, written $\mathfrak{A} \cong \mathfrak{B}$, if there is some isomorphism of \mathfrak{A} onto \mathfrak{B} ;

Homomorphisms

Definition

- ▶ h is a homomorphism of \mathfrak{A} **onto** \mathfrak{B} if h is a homomorphism from \mathfrak{A} to \mathfrak{B} and h maps \mathfrak{A} onto \mathfrak{B} ;
- ▶ A homomorphism h from \mathfrak{A} into \mathfrak{B} is an **isomorphism** if h is one-to-one;
- ▶ The structures \mathfrak{A} and \mathfrak{B} are **isomorphic**, written $\mathfrak{A} \cong \mathfrak{B}$, if there is some isomorphism of \mathfrak{A} onto \mathfrak{B} ;
- ▶ An **automorphism** of \mathfrak{A} is an isomorphism of \mathfrak{A} onto \mathfrak{A} .

Examples

Example

Let $\mathfrak{A} = (\mathbb{N}, <^{\mathbb{N}}, +^{\mathbb{N}})$ and $\mathfrak{B} = (\mathbb{E}, <^{\mathbb{E}}, +^{\mathbb{E}})$.

Here \mathbb{E} is the set of even non-negative integers, $<^{\mathbb{E}}$ is the “less than” relation on \mathbb{E} , etc.

Then h is an isomorphism of \mathfrak{A} onto \mathfrak{B} , where for all $n \in \mathbb{N}$,

$$h(n) =$$

Examples

Example

Let $\mathfrak{A} = (\mathbb{N}, <^{\mathbb{N}}, +^{\mathbb{N}})$ and $\mathfrak{B} = (\mathbb{E}, <^{\mathbb{E}}, +^{\mathbb{E}})$.

Here \mathbb{E} is the set of even non-negative integers, $<^{\mathbb{E}}$ is the “less than” relation on \mathbb{E} , etc.

Then h is an isomorphism of \mathfrak{A} onto \mathfrak{B} , where for all $n \in \mathbb{N}$,

$$h(n) = 2n.$$

Examples

Example

Let $\mathfrak{A} = (\mathbb{N}, <^{\mathbb{N}}, +^{\mathbb{N}})$ and $\mathfrak{B} = (\mathbb{O}, <^{\mathbb{O}}, +^{\mathbb{O}})$.

Here \mathbb{O} is the set of odd non-negative integers, $<^{\mathbb{O}}$ is the “less than” relation on \mathbb{O} , etc.

- Then an isomorphism of \mathfrak{A} onto \mathfrak{B} is:

Examples

Example

Let $\mathfrak{A} = (\mathbb{N}, <^{\mathbb{N}}, +^{\mathbb{N}})$ and $\mathfrak{B} = (\mathbb{O}, <^{\mathbb{O}}, +^{\mathbb{O}})$.

Here \mathbb{O} is the set of odd non-negative integers, $<^{\mathbb{O}}$ is the “less than” relation on \mathbb{O} , etc.

- Then an isomorphism of \mathfrak{A} onto \mathfrak{B} is: There is NONE!

Examples

Example

Let $\mathfrak{A} = (\mathbb{N}, <^{\mathbb{N}}, +^{\mathbb{N}})$ and $\mathfrak{B} = (\mathbb{O}, <^{\mathbb{O}}, +^{\mathbb{O}})$.

Here \mathbb{O} is the set of odd non-negative integers, $<^{\mathbb{O}}$ is the “less than” relation on \mathbb{O} , etc.

- ▶ Then an isomorphism of \mathfrak{A} onto \mathfrak{B} is: There is NONE!
- ▶ In fact, \mathfrak{B} is not even a structure, because

Examples

Example

Let $\mathfrak{A} = (\mathbb{N}, <^{\mathbb{N}}, +^{\mathbb{N}})$ and $\mathfrak{B} = (\mathbb{O}, <^{\mathbb{O}}, +^{\mathbb{O}})$.

Here \mathbb{O} is the set of odd non-negative integers, $<^{\mathbb{O}}$ is the “less than” relation on \mathbb{O} , etc.

- ▶ Then an isomorphism of \mathfrak{A} onto \mathfrak{B} is: There is NONE!
- ▶ In fact, \mathfrak{B} is not even a structure, because \mathbb{O} is not closed under addition.

Automorphisms of $\mathfrak{R} = (\mathbb{R}, <)$

Let $\mathfrak{R} = (R, <)$. Which of the following functions h are automorphisms of \mathfrak{R} ?

Automorphisms of $\mathfrak{R} = (\mathbb{R}, <)$

Let $\mathfrak{R} = (R, <)$. Which of the following functions h are automorphisms of \mathfrak{R} ?

- The identity function.

Automorphisms of $\mathfrak{R} = (\mathbb{R}, <)$

Let $\mathfrak{R} = (R, <)$. Which of the following functions h are automorphisms of \mathfrak{R} ?

- The identity function. YES

Automorphisms of $\mathfrak{R} = (\mathbb{R}, <)$

Let $\mathfrak{R} = (R, <)$. Which of the following functions h are automorphisms of \mathfrak{R} ?

- ▶ The identity function. YES
- ▶ $h(a) = a + 3$

Automorphisms of $\mathfrak{R} = (\mathbb{R}, <)$

Let $\mathfrak{R} = (R, <)$. Which of the following functions h are automorphisms of \mathfrak{R} ?

- ▶ The identity function. YES
- ▶ $h(a) = a + 3$ YES

Automorphisms of $\mathfrak{R} = (\mathbb{R}, <)$

Let $\mathfrak{R} = (R, <)$. Which of the following functions h are automorphisms of \mathfrak{R} ?

- ▶ The identity function. YES
- ▶ $h(a) = a + 3$ YES
- ▶ $h(a) = a - 3$

Automorphisms of $\mathfrak{R} = (\mathbb{R}, <)$

Let $\mathfrak{R} = (R, <)$. Which of the following functions h are automorphisms of \mathfrak{R} ?

- ▶ The identity function. YES
- ▶ $h(a) = a + 3$ YES
- ▶ $h(a) = a - 3$ YES

Automorphisms of $\mathfrak{R} = (\mathbb{R}, <)$

Let $\mathfrak{R} = (R, <)$. Which of the following functions h are automorphisms of \mathfrak{R} ?

- ▶ The identity function. YES
- ▶ $h(a) = a + 3$ YES
- ▶ $h(a) = a - 3$ YES
- ▶ $h(a) = 2a$

Automorphisms of $\mathfrak{R} = (\mathbb{R}, <)$

Let $\mathfrak{R} = (R, <)$. Which of the following functions h are automorphisms of \mathfrak{R} ?

- ▶ The identity function. YES
- ▶ $h(a) = a + 3$ YES
- ▶ $h(a) = a - 3$ YES
- ▶ $h(a) = 2a$ YES

Automorphisms of $\mathfrak{R} = (\mathbb{R}, <)$

Let $\mathfrak{R} = (R, <)$. Which of the following functions h are automorphisms of \mathfrak{R} ?

▶ The identity function. YES

▶ $h(a) = a + 3$ YES

▶ $h(a) = a - 3$ YES

▶ $h(a) = 2a$ YES

▶ $h(a) = -a$

Automorphisms of $\mathfrak{R} = (\mathbb{R}, <)$

Let $\mathfrak{R} = (R, <)$. Which of the following functions h are automorphisms of \mathfrak{R} ?

▶ The identity function. YES

▶ $h(a) = a + 3$ YES

▶ $h(a) = a - 3$ YES

▶ $h(a) = 2a$ YES

▶ $h(a) = -a$ NO

Automorphisms of $\mathfrak{R} = (\mathbb{R}, <)$

Let $\mathfrak{R} = (R, <)$. Which of the following functions h are automorphisms of \mathfrak{R} ?

- ▶ The identity function. YES
- ▶ $h(a) = a + 3$ YES
- ▶ $h(a) = a - 3$ YES
- ▶ $h(a) = 2a$ YES
- ▶ $h(a) = -a$ NO
- ▶ $h(a) = k \times a + l$

Automorphisms of $\mathfrak{R} = (\mathbb{R}, <)$

Let $\mathfrak{R} = (R, <)$. Which of the following functions h are automorphisms of \mathfrak{R} ?

- ▶ The identity function. YES
- ▶ $h(a) = a + 3$ YES
- ▶ $h(a) = a - 3$ YES
- ▶ $h(a) = 2a$ YES
- ▶ $h(a) = -a$ NO
- ▶ $h(a) = k \times a + l$ YES provided $k > 0$

Automorphisms of $\mathfrak{R} = (\mathbb{R}, <)$

Let $\mathfrak{R} = (R, <)$. Which of the following functions h are automorphisms of \mathfrak{R} ?

- ▶ The identity function. YES
- ▶ $h(a) = a + 3$ YES
- ▶ $h(a) = a - 3$ YES
- ▶ $h(a) = 2a$ YES
- ▶ $h(a) = -a$ NO
- ▶ $h(a) = k \times a + l$ YES provided $k > 0$
- ▶ $h(a) = a^3$

Automorphisms of $\mathfrak{R} = (\mathbb{R}, <)$

Let $\mathfrak{R} = (R, <)$. Which of the following functions h are automorphisms of \mathfrak{R} ?

- ▶ The identity function. YES
- ▶ $h(a) = a + 3$ YES
- ▶ $h(a) = a - 3$ YES
- ▶ $h(a) = 2a$ YES
- ▶ $h(a) = -a$ NO
- ▶ $h(a) = k \times a + l$ YES provided $k > 0$
- ▶ $h(a) = a^3$ YES

Automorphisms of $\mathfrak{R} = (\mathbb{R}, <)$

Let $\mathfrak{R} = (R, <)$. Which of the following functions h are automorphisms of \mathfrak{R} ?

- ▶ The identity function. YES
- ▶ $h(a) = a + 3$ YES
- ▶ $h(a) = a - 3$ YES
- ▶ $h(a) = 2a$ YES
- ▶ $h(a) = -a$ NO
- ▶ $h(a) = k \times a + l$ YES provided $k > 0$
- ▶ $h(a) = a^3$ YES
- ▶ $h(a) = a^2$

Automorphisms of $\mathfrak{R} = (\mathbb{R}, <)$

Let $\mathfrak{R} = (R, <)$. Which of the following functions h are automorphisms of \mathfrak{R} ?

- ▶ The identity function. YES
- ▶ $h(a) = a + 3$ YES
- ▶ $h(a) = a - 3$ YES
- ▶ $h(a) = 2a$ YES
- ▶ $h(a) = -a$ NO
- ▶ $h(a) = k \times a + l$ YES provided $k > 0$
- ▶ $h(a) = a^3$ YES
- ▶ $h(a) = a^2$ NO

Automorphisms of $\mathfrak{N} = (\mathbb{N}, <)$

Let $\mathfrak{N} = (\mathbb{N}, <)$.

Automorphisms of $\mathfrak{N} = (\mathbb{N}, <)$

Let $\mathfrak{N} = (\mathbb{N}, <)$.

- ▶ The identity function is an automorphism of \mathfrak{N} ;

Automorphisms of $\mathfrak{N} = (\mathbb{N}, <)$

Let $\mathfrak{N} = (\mathbb{N}, <)$.

- ▶ The identity function is an automorphism of \mathfrak{N} ; what about others?

Automorphisms of $\mathfrak{N} = (\mathbb{N}, <)$

Let $\mathfrak{N} = (\mathbb{N}, <)$.

- ▶ The identity function is an automorphism of \mathfrak{N} ; what about others?
- ▶ Suppose h is an automorphism of \mathfrak{N} , $h(0) =$

Automorphisms of $\mathfrak{N} = (\mathbb{N}, <)$

Let $\mathfrak{N} = (\mathbb{N}, <)$.

- ▶ The identity function is an automorphism of \mathfrak{N} ; what about others?
- ▶ Suppose h is an automorphism of \mathfrak{N} , $h(0) = 0$

Automorphisms of $\mathfrak{N} = (\mathbb{N}, <)$

Let $\mathfrak{N} = (\mathbb{N}, <)$.

- ▶ The identity function is an automorphism of \mathfrak{N} ; what about others?
- ▶ Suppose h is an automorphism of \mathfrak{N} , $h(0) = 0$
- ▶ Suppose h is an automorphism of \mathfrak{N} , $h(1) =$

Automorphisms of $\mathfrak{N} = (\mathbb{N}, <)$

Let $\mathfrak{N} = (\mathbb{N}, <)$.

- ▶ The identity function is an automorphism of \mathfrak{N} ; what about others?
- ▶ Suppose h is an automorphism of \mathfrak{N} , $h(0) = 0$
- ▶ Suppose h is an automorphism of \mathfrak{N} , $h(1) = 1$

Automorphisms of $\mathfrak{N} = (\mathbb{N}, <)$

Let $\mathfrak{N} = (\mathbb{N}, <)$.

- ▶ The identity function is an automorphism of \mathfrak{N} ; what about others?
- ▶ Suppose h is an automorphism of \mathfrak{N} , $h(0) = 0$
- ▶ Suppose h is an automorphism of \mathfrak{N} , $h(1) = 1$
- ▶ In general, if h is an automorphism of \mathfrak{N} , $h(n) =$

Automorphisms of $\mathfrak{N} = (\mathbb{N}, <)$

Let $\mathfrak{N} = (\mathbb{N}, <)$.

- ▶ The identity function is an automorphism of \mathfrak{N} ; what about others?
- ▶ Suppose h is an automorphism of \mathfrak{N} , $h(0) = 0$
- ▶ Suppose h is an automorphism of \mathfrak{N} , $h(1) = 1$
- ▶ In general, if h is an automorphism of \mathfrak{N} , $h(n) = n$

Automorphisms of $\mathfrak{N} = (\mathbb{N}, <)$

Let $\mathfrak{N} = (\mathbb{N}, <)$.

- ▶ The identity function is an automorphism of \mathfrak{N} ; what about others?
- ▶ Suppose h is an automorphism of \mathfrak{N} , $h(0) = 0$
- ▶ Suppose h is an automorphism of \mathfrak{N} , $h(1) = 1$
- ▶ In general, if h is an automorphism of \mathfrak{N} , $h(n) = n$
- ▶ Therefore, the identity function is the **only** automorphism of \mathfrak{N} .

Substructures

A special kind of isomorphisms:

Definition

Let $\mathfrak{A} = (A, \dots)$ and $\mathfrak{B} = (B, \dots)$ be structures for \mathbb{L} . \mathfrak{A} is a substructure of \mathfrak{B} (written $\mathfrak{A} \subseteq \mathfrak{B}$) if:

Substructures

A special kind of isomorphisms:

Definition

Let $\mathfrak{A} = (A, \dots)$ and $\mathfrak{B} = (B, \dots)$ be structures for \mathbb{L} . \mathfrak{A} is a substructure of \mathfrak{B} (written $\mathfrak{A} \subseteq \mathfrak{B}$) if:

- ▶ $A \subseteq B$;

Substructures

A special kind of isomorphisms:

Definition

Let $\mathfrak{A} = (A, \dots)$ and $\mathfrak{B} = (B, \dots)$ be structures for \mathbb{L} . \mathfrak{A} is a substructure of \mathfrak{B} (written $\mathfrak{A} \subseteq \mathfrak{B}$) if:

- ▶ $A \subseteq B$;
- ▶ for every k -ary predicate symbol P :

$$P^{\mathfrak{A}} = P^{\mathfrak{B}} \cap A^k$$

(Note this is not the same as saying $P^{\mathfrak{A}} \subseteq P^{\mathfrak{B}}$);

Substructures

A special kind of isomorphisms:

Definition

Let $\mathfrak{A} = (A, \dots)$ and $\mathfrak{B} = (B, \dots)$ be structures for \mathbb{L} . \mathfrak{A} is a substructure of \mathfrak{B} (written $\mathfrak{A} \subseteq \mathfrak{B}$) if:

- ▶ $A \subseteq B$;
- ▶ for every k -ary predicate symbol P :

$$P^{\mathfrak{A}} = P^{\mathfrak{B}} \cap A^k$$

(Note this is not the same as saying $P^{\mathfrak{A}} \subseteq P^{\mathfrak{B}}$);

- ▶ for every k -ary function symbol f and every k -tuple (a_1, \dots, a_k) of elements of A :

$$f^{\mathfrak{A}}(a_1, \dots, a_k) = f^{\mathfrak{B}}(a_1, \dots, a_k);$$

Substructures

A special kind of isomorphisms:

Definition

Let $\mathfrak{A} = (A, \dots)$ and $\mathfrak{B} = (B, \dots)$ be structures for \mathbb{L} . \mathfrak{A} is a substructure of \mathfrak{B} (written $\mathfrak{A} \subseteq \mathfrak{B}$) if:

- ▶ $A \subseteq B$;
- ▶ for every k -ary predicate symbol P :

$$P^{\mathfrak{A}} = P^{\mathfrak{B}} \cap A^k$$

(Note this is not the same as saying $P^{\mathfrak{A}} \subseteq P^{\mathfrak{B}}$);

- ▶ for every k -ary function symbol f and every k -tuple (a_1, \dots, a_k) of elements of A :

$$f^{\mathfrak{A}}(a_1, \dots, a_k) = f^{\mathfrak{B}}(a_1, \dots, a_k);$$

- ▶ for every constant symbol c , $c^{\mathfrak{A}} = c^{\mathfrak{B}}$.

Examples

Example

Let

► $\mathfrak{N} = (\mathbb{N}, <^{\mathbb{N}}, +^{\mathbb{N}}, \times^{\mathbb{N}})$

► $\mathfrak{E} = (\mathbb{E}, <^{\mathbb{E}}, +^{\mathbb{E}}, \times^{\mathbb{E}}).$

Then \mathfrak{E} is a substructure of \mathfrak{N} .

Examples

Example

Let

▶ $\mathfrak{N} = (\mathbb{N}, <^{\mathbb{N}}, +^{\mathbb{N}}, \times^{\mathbb{N}})$

▶ $\mathfrak{E} = (\mathbb{E}, <^{\mathbb{E}}, +^{\mathbb{E}}, \times^{\mathbb{E}}).$

Then \mathfrak{E} is a substructure of \mathfrak{N} .

Question

Let

▶ $\mathfrak{A} = (\{0, 1, 2, 3\}, P^{\mathfrak{A}})$, where $P^{\mathfrak{A}} = \{0, 1, 2\}$;

▶ $\mathfrak{B} = (\{0, 1\}, P^{\mathfrak{B}})$, where $P^{\mathfrak{B}} = \{0\}$.

Is \mathfrak{B} is a substructure of \mathfrak{A} ?

Examples

Example

Let

► $\mathfrak{N} = (\mathbb{N}, <^{\mathbb{N}}, +^{\mathbb{N}}, \times^{\mathbb{N}})$

► $\mathfrak{E} = (\mathbb{E}, <^{\mathbb{E}}, +^{\mathbb{E}}, \times^{\mathbb{E}})$.

Then \mathfrak{E} is a substructure of \mathfrak{N} .

Question

Let

► $\mathfrak{A} = (\{0, 1, 2, 3\}, P^{\mathfrak{A}})$, where $P^{\mathfrak{A}} = \{0, 1, 2\}$;

► $\mathfrak{B} = (\{0, 1\}, P^{\mathfrak{B}})$, where $P^{\mathfrak{B}} = \{0\}$.

Is \mathfrak{B} is a substructure of \mathfrak{A} ?

Answer

No. Because $P^{\mathfrak{A}} \cap \{0, 1\} = \{0, 1\} \neq \{0\} = P^{\mathfrak{B}}$.

Notation: Function Composition

Definition

If f and g are functions, then $f \circ g$ is the composition of f and g .
That is,

$$f \circ g(a) = f(g(a)).$$

Notation: Function Composition

Definition

If f and g are functions, then $f \circ g$ is the composition of f and g . That is,

$$f \circ g(a) = f(g(a)).$$

Example

Suppose $s : V \rightarrow |\mathfrak{A}|$ is an assignment function for \mathfrak{A} , and h is a homomorphism from \mathfrak{A} to \mathfrak{B} . Then $h \circ s$ is an assignment function for \mathfrak{B} .

The Value of Terms Under a Homomorphism

Lemma

Let \mathfrak{A} and \mathfrak{B} be structures for the language \mathbb{L} .

Let h be a homomorphism from \mathfrak{A} to \mathfrak{B} , and $s : V \rightarrow |\mathfrak{A}|$ be an assignment for \mathfrak{A} . Then for every term t of \mathbb{L} ,

$$h(\bar{s}(t)) = \overline{h \circ s}(t).$$

The Value of Terms Under a Homomorphism

Lemma

Let \mathfrak{A} and \mathfrak{B} be structures for the language \mathbb{L} .

Let h be a homomorphism from \mathfrak{A} to \mathfrak{B} , and $s : V \rightarrow |\mathfrak{A}|$ be an assignment for \mathfrak{A} . Then for every term t of \mathbb{L} ,

$$h(\bar{s}(t)) = \overline{h \circ s}(t).$$

Proof.

By induction on t .



The Homomorphism Theorem

Theorem (The Homomorphism Theorem)

Let h be a homomorphism from \mathfrak{A} to \mathfrak{B} and s be an assignment function for \mathfrak{A} . The statement

$$\models_{\mathfrak{A}} \varphi[s] \iff \models_{\mathfrak{B}} \varphi[h \circ s]$$

The Homomorphism Theorem

Theorem (The Homomorphism Theorem)

Let h be a homomorphism from \mathfrak{A} to \mathfrak{B} and s be an assignment function for \mathfrak{A} . The statement

$$\models_{\mathfrak{A}} \varphi[s] \iff \models_{\mathfrak{B}} \varphi[h \circ s]$$

(a) is true for every quantifier-free wff φ not containing $\dot{=}$;

The Homomorphism Theorem

Theorem (The Homomorphism Theorem)

Let h be a homomorphism from \mathfrak{A} to \mathfrak{B} and s be an assignment function for \mathfrak{A} . The statement

$$\models_{\mathfrak{A}} \varphi[s] \iff \models_{\mathfrak{B}} \varphi[h \circ s]$$

- (a) is true for every quantifier-free wff φ not containing $\dot{=}$;
- (b) is true for every quantifier-free wff φ if h is one-to-one;

The Homomorphism Theorem

Theorem (The Homomorphism Theorem)

Let h be a homomorphism from \mathfrak{A} to \mathfrak{B} and s be an assignment function for \mathfrak{A} . The statement

$$\models_{\mathfrak{A}} \varphi[s] \iff \models_{\mathfrak{B}} \varphi[h \circ s]$$

- (a) is true for every quantifier-free wff φ not containing $\dot{=}$;
- (b) is true for every quantifier-free wff φ if h is one-to-one;
- (c) is true for every wff φ not containing $\dot{=}$ if h is onto;

The Homomorphism Theorem

Theorem (The Homomorphism Theorem)

Let h be a homomorphism from \mathfrak{A} to \mathfrak{B} and s be an assignment function for \mathfrak{A} . The statement

$$\models_{\mathfrak{A}} \varphi[s] \iff \models_{\mathfrak{B}} \varphi[h \circ s]$$

- (a) is true for every quantifier-free wff φ not containing \doteq ;
- (b) is true for every quantifier-free wff φ if h is one-to-one;
- (c) is true for every wff φ not containing \doteq if h is onto;
- (d) is true for every wff φ if h is an isomorphism of \mathfrak{A} onto \mathfrak{B} (i.e., $\mathfrak{A} \cong \mathfrak{B}$).

The Homomorphism Theorem

Theorem (The Homomorphism Theorem)

Let h be a homomorphism from \mathfrak{A} to \mathfrak{B} and s be an assignment function for \mathfrak{A} . The statement

$$\models_{\mathfrak{A}} \varphi[s] \iff \models_{\mathfrak{B}} \varphi[h \circ s]$$

- (a) is true for every quantifier-free wff φ not containing \doteq ;
- (b) is true for every quantifier-free wff φ if h is one-to-one;
- (c) is true for every wff φ not containing \doteq if h is onto;
- (d) is true for every wff φ if h is an isomorphism of \mathfrak{A} onto \mathfrak{B} (i.e., $\mathfrak{A} \cong \mathfrak{B}$).

Proof.

By induction on φ .



Corollaries of the Homomorphism Theorem

Corollary

If $\mathfrak{A} \cong \mathfrak{B}$, then $\mathfrak{A} \equiv \mathfrak{B}$.

Corollaries of the Homomorphism Theorem

Corollary

If $\mathfrak{A} \cong \mathfrak{B}$, then $\mathfrak{A} \equiv \mathfrak{B}$.

Question

Do you think the converse is true?

Corollaries of the Homomorphism Theorem

Corollary

If $\mathfrak{A} \cong \mathfrak{B}$, then $\mathfrak{A} \equiv \mathfrak{B}$.

Question

Do you think the converse is true?

Answer

No.

Corollaries of the Homomorphism Theorem

Corollary

If $\mathfrak{A} \cong \mathfrak{B}$, then $\mathfrak{A} \equiv \mathfrak{B}$.

Question

Do you think the converse is true?

Answer

No. Take $\mathfrak{A} = (\mathbb{R}, <)$ and $\mathfrak{B} = (\mathbb{Q}, <)$ as a counter example.

Corollaries of the Homomorphism Theorem

Corollary (Automorphism Theorem)

Let h be an automorphism of \mathfrak{A} . Let R be an n -ary relation on $|\mathfrak{A}|$ that is definable in \mathfrak{A} . For every n -tuple (a_1, \dots, a_n) of elements of \mathfrak{A} :

$$(a_1, \dots, a_n) \in R \iff (h(a_1), \dots, h(a_n)) \in R.$$

Corollaries of the Homomorphism Theorem

Corollary (Automorphism Theorem)

Let h be an automorphism of \mathfrak{A} . Let R be an n -ary relation on $|\mathfrak{A}|$ that is definable in \mathfrak{A} . For every n -tuple (a_1, \dots, a_n) of elements of \mathfrak{A} :

$$(a_1, \dots, a_n) \in R \iff (h(a_1), \dots, h(a_n)) \in R.$$

We often use this lemma to show certain relations are not definable:

Corollaries of the Homomorphism Theorem

Corollary (Automorphism Theorem)

Let h be an automorphism of \mathfrak{A} . Let R be an n -ary relation on $|\mathfrak{A}|$ that is definable in \mathfrak{A} . For every n -tuple (a_1, \dots, a_n) of elements of \mathfrak{A} :

$$(a_1, \dots, a_n) \in R \iff (h(a_1), \dots, h(a_n)) \in R.$$

We often use this lemma to show certain relations are not definable:

Example

Let $\mathfrak{R} = (\mathbb{R}, <)$. Its subset \mathbb{N} is not definable in \mathfrak{R} .

Corollaries of the Homomorphism Theorem

Corollary (Automorphism Theorem)

Let h be an automorphism of \mathfrak{A} . Let R be an n -ary relation on $|\mathfrak{A}|$ that is definable in \mathfrak{A} . For every n -tuple (a_1, \dots, a_n) of elements of \mathfrak{A} :

$$(a_1, \dots, a_n) \in R \iff (h(a_1), \dots, h(a_n)) \in R.$$

We often use this lemma to show certain relations are not definable:

Example

Let $\mathfrak{R} = (\mathbb{R}, <)$. Its subset \mathbb{N} is not definable in \mathfrak{R} because $h(a) = a^3$ is an automorphism of \mathfrak{R} .