

Causal Inference

Lab 5: Propensity Score, Matching, and Weighting

Xiangyu Song

Washington University in St. Louis

February 13, 2026

Propensity Score

Definition (Rosenbaum and Rubin (1983))

Propensity score is defined as the conditional probability of receiving a treatment given pre-treatment covariates X :

$$e(X) = \Pr(D = 1 \mid X) = \mathbb{E}[D \mid X],$$

where $X = (X_1, \dots, X_p)$ is the collection of p covariates.

- Propensity score is a probability
 - ▶ Analogous to a summary statistic of the assignment mechanism.

Balancing Property of Propensity Score

- Property 1. The propensity score $e(X)$ balances the distribution of all X between the treatment groups:

$$D \perp\!\!\!\perp X \mid e(X)$$

- Conditional on the PS, treatment is randomized.
- Definition: A balancing score $b(X)$ is a function of the covariates such that

$$Z \perp\!\!\!\perp X \mid b(X)$$

- Propensity score is a balancing score.

Balancing Scores

- Property 2. Let $b(X)$ be a function of X , $b(X)$ is a balancing score if and only if $b(X)$ is finer than $e(X)$ in the sense that $e(X) = f(b(X))$ for some function f .

Proof sketch:

- ⇐ ▶ Suppose $b(X)$ is finer than $e(X)$, prove $\Pr(D = 1 \mid X, b(X)) = \Pr(D = 1 \mid b(X))$.
 - ▶ LHS: $\Pr(D = 1 \mid X, b(X)) = \Pr(D = 1 \mid X) = e(X)$.
 - ▶ RHS: $\Pr(D = 1 \mid b(X)) = \mathbb{E}[\mathbb{E}[D \mid X, b(X)] \mid b(X)] = \mathbb{E}[\mathbb{E}[D \mid X] \mid b(X)] = \mathbb{E}[e(X) \mid b(X)] = e(X)$.
- ⇒ ▶ Suppose $b(X)$ is a balancing score but not finer than $e(X)$. Then $\exists X_1, X_2$ such that $e(X_1) \neq e(X_2)$ but $b(X_1) = b(X_2)$
 - ▶ $\Pr(D = 1 \mid X_1, b(X_1)) = \Pr(D = 1 \mid X_1) \neq \Pr(D = 1 \mid X_2) = \Pr(D = 1 \mid X_2, b(X_2))$: $D \not\perp\!\!\!\perp X \mid b(X)$
 - ▶ $b(X)$ cannot be a balancing score, so proof by contradiction, $b(X)$ must be finer than $e(X)$ to be a balancing score.

Balancing Scores and Propensity Score

- Rosenbaum and Rubin (1983) showed that $e(X)$ is the coarsest balancing score, X is the finest balancing score.
- Knowledge of the balancing score could create a conditionally randomized trial, which supports valid causal inference (creating an ideal randomized experiment)
- Mostly focus on the coarsest balancing score $e(X)$, but the general theory holds for other specification of $b(X)$.

Remarks on the Balancing Property

1. If a subclass of units or a matched treatment-control pair is homogeneous in $e(X)$, then the treatment and control units have the same distribution of X .
2. If a subclass of units or a matched treatment-control pair is homogeneous in both $e(X)$ and certain X , the other components of X within those refined class is also balanced – practical implication: estimating causal estimand in subpopulation, e.g., male of female group
3. The balancing property is a statement on the distribution of X , NOT on assignment mechanism or potential outcomes.

Propensity Score: Unconfoundedness

- Property 3. If D is unconfounded given X , then D is unconfounded given $b(X)$, i.e.,

$$(Y(1), Y(0)) \perp\!\!\!\perp D \mid X \implies (Y(1), Y(0)) \perp\!\!\!\perp D \mid b(X)$$

- Proof. Sufficient to show

$$\Pr(D = 1 \mid Y(1), Y(0), b(X)) = \Pr(D = 1 \mid b(X))$$

- Recall from the previous proof, RHS $\Pr(D = 1 \mid b(X)) = e(X)$.
- LHS by LIE,

$$\begin{aligned} \mathbb{E} [\Pr(D = 1 \mid Y(1), Y(0), X) \mid Y(1), Y(0), b(X)] \\ = \mathbb{E} [e(X) \mid Y(1), Y(0), b(X)] = e(X), \end{aligned}$$

because $b(X)$ is finer than $e(X)$

Propensity Score: Unconfoundedness

- Property 4. If D is unconfounded given X , then D is unconfounded given $e(X)$, i.e.,

$$(Y(1), Y(0)) \perp\!\!\!\perp D \mid X \implies (Y(1), Y(0)) \perp\!\!\!\perp D \mid e(X)$$

1. Given a vector of covariates that ensure unconfoundedness, adjustment for differences in propensity scores removes all biases associated with differences in the covariates
2. $e(X)$ can be viewed as a summary score of the observed covariates
3. Causal inference can be drawn through stratification, matching, weighting, etc. using the scalar $e(X)$ instead of high dimensional covariates (dimension reduction).

Propensity Score: Remarks

- Controlling for propensity score (or more generally, balancing score)
 - ▶ Create a conditionally randomized trial $(Y(1), Y(0)) \perp\!\!\!\perp D \mid e(X)$
 - ▶ Balances covariates between groups $D \perp\!\!\!\perp X \mid e(X)$
- Propensity score balances the observed covariates but does not generally balance unobserved covariates
- Adjusting for differences in $e(X)$ can remove bias due to X
- In most observational studies, the propensity score $e(X)$ is unknown and thus needs to be estimated

Propensity Score Analysis of Causal Effects

1. Estimate $e(X)$ by logistic regression or machine learning methods
2. Given $\hat{e}(X)$, estimate the causal effects through
 - ▶ Subclassification
 - ▶ Matching
 - ▶ Regression
 - ▶ Weighting
 - ▶ Mixed procedure of the above

Propensity Score Matching

- Special case of matching: the distance metric is the (estimated) propensity score
- 1-to- k nearest neighbor matching is common when the control group is large compared to treatment group
- Pros: robust, popular, vast literature, and intuitive to understand
- Cons: inefficient (check Abadie and Imbens (2006)) for closed form variance expression

Propensity Score Weighting

- Most common approach for causal inference with observational data
- Intuitive: reweighting to mimic randomized treatment assignment
- Flexible: extends to binary, multi-valued, and continuous treatments
- Transparent: balance diagnostics and effective sample size
- Limitations: instability with extreme weights

Inverse Probability Weighting (IPW)

- Easy to show

$$\mathbb{E} \left[\frac{DY}{e(X)} - \frac{(1-D)Y}{1-e(X)} \right] = \tau^{\text{ATE}}.$$

- Observe

$$\mathbb{E} \left[\frac{DY}{e(X)} \right] = \mathbb{E} \left[\frac{1}{e(X)} \mathbb{E} [DY(1) \mid X] \right] = \mathbb{E} \left[\frac{\mathbb{E} [D \mid X]}{e(X)} \mathbb{E} [Y(1) \mid X] \right]$$

Inverse Probability Weighting (IPW), cont.

- Define the inverse probability weights (IPW)

$$\begin{cases} w_1(X_i) = \frac{1}{e(X_i)}, & D_i = 1 \\ w_0(X_i) = \frac{1}{1-e(X_i)}, & D_i = 0 \end{cases}$$

- An unbiased moment-type estimator of ATE: difference in the mean of the weighted outcomes between groups

$$\begin{aligned} \hat{\tau}_{\text{IPW},1} &= \frac{1}{N} \left(\sum_{i=1}^N \frac{Y_i D_i}{e(X_i)} - \sum_{i=1}^N \frac{Y_i (1 - D_i)}{1 - e(X_i)} \right) \\ &= \frac{1}{N} \sum_{i=1}^N (Y_i D_i w_1(X_i) - Y_i (1 - D_i) w_0(X_i)) \end{aligned}$$

Normalize the Weights

- When using any weighting methods, a good practice is to normalize the weights - sum of the total weights within one group should be 1
- Divide each unit's weight by the sum of all weights in that group $\frac{w_i}{\sum_{i:D=d} w_i}$ for $d = 0, 1$, i.e., the Hajek estimator:

$$\hat{\tau}_{\text{IPW}, 2} = \frac{\sum_{i=1}^N Y_i D_i w_1(X_i)}{\sum_{i=1}^N D_i w_1(X_i)} - \frac{\sum_{i=1}^N Y_i (1 - D_i) w_0(X_i)}{\sum_{i=1}^N (1 - D_i) w_0(X_i)}.$$

- Reduce variance: $\text{Var}(\hat{\tau}_{\text{IPW}, 2}) \leq \text{Var}(\hat{\tau}_{\text{IPW}, 1})$, and lead to more stable estimate (Hirano et al. 2003)